Which Factors Are Predicting the Mortality in Patients With Covid 19 in the Intensive Care Unit?

Murat Parpucu

Saglik Bilimleri Universitesi Gulhane Saglik Bilimleri Enstitusu

Sema Turan (✉ semakultufan@yahoo.com)

Ankara Şehir Hastanesi: Ankara Şehir Hastanesi <https://orcid.org/0000-0003-2443-0390>

Hayriye Cankar Dal

Ankara City Hospital: Ankara Şehir Hastanesi

Damla Tosun

Ankara City Hospital: Ankara Şehir Hastanesi

Kudret Yasemin Yalnız

Ankara City Hospital: Ankara Şehir Hastanesi

Dilek Kazancı

Ankara City Hospital: Ankara Şehir Hastanesi

Research

Keywords: Covid 19, Intensive Care Unit, Mortality in Patients, APACHE II value

Posted Date: March 1st, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-258591/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: COVID 19 infection is a global health problem; clinical and laboratory parameters have been developed to predict this disease-related mortality/morbidity. Some of these parameters are clinical parameters, while some are laboratory parameters. This study aims to determine whether APACHE II, GCS, age, presence of comorbidity, and absolute lymphocyte count effectively predict mortality in patients admitted to intensive care due to COVID 19.

Method: We have included 108 PCR-positive COVID 19 patients admitted to the intensive care between 1 October and 31 November 2020 in our research. Demographic characteristics of all patients, APACHE II values within the first 24 hours of admission to intensive care, the Glasgow Coma Scale, the presence of comorbidity, lymphocyte count during intensive care admission, duration of intensive care stay, and the mortality rates were recorded.

Results: The average age of 108 individuals evaluated in the study was 67 ± 13.61 years, and 56.5% of the patient group consisted of the geriatric age range. Seventy (64.8%) of the patients were female, eighty-nine (82.4%) patients had at least one comorbidity. In the multivariate analysis, it was determined that lymphocyte value, APACHE II score, and the presence of any comorbidity are independent prognostic factors for mortality when accepted to ICU.

Conclusion: In our study, we have determined that age, APACHE II value, presence of comorbidity, and baseline lymphocyte counts are independent predictors of mortality.

Trial Registry: SEMA TURAN-2021-01-13T10_31_59

Registry address: <https://bilimselarastirma.saglik.gov.tr>

Introduction

The SARS CoV 2 epidemic that emerged in Wuhan, China, is a member of the coronavirus family and has affected all world countries and caused a pandemic throughout 2020 (1). According to this infection spread table generated by the SARS CoV 2 virus, 10–20% of the patients require intensive care (2). While the disease-related hospital mortality rate is 4.3–11%, the mortality rate among the patients admitted to intensive care varies between centers and is around 30–60% (3). In COVID 19 patients admitted to intensive care with respiratory failure, clinical deterioration may rapidly deteriorate. Patients are lost due to severe ARDS and subsequent multiple organ dysfunction (4). COVID 19 infection is a global health problem, and an effective scoring system established with clinical and laboratory parameters to predict this disease-related mortality/morbidity has not been defined. Scoring systems developed to predict patients' prognosis during admission to intensive care and evaluate the treatment's effectiveness is widely used in non-COVID patients. APACHE II, one of the most commonly used scoring systems, is known to be a successful scoring system in terms of mortality prediction (5). There is insufficient information that these scoring systems effectively predict mortality in COVID 19 patients, and research on

this subject continues. Again, the decrease in the number of lymphocytes in COVID 19-associated viral infection is thought to be the result of direct binding of the virus with ACE2 receptors on lymphocytes and apoptosis caused by a cytokine storm. In a recent meta-analysis published by Lui et al., 35–75% of the patients developed lymphopenia, and it states that mortality increased significantly in these patients (6). Fan et al. reported that a lymphocyte count of $< 0.6 \times 10^9 / L$ was associated with a poor prognosis in an analysis of 67 COVID 19 patients during their admission to an intensive care unit (7, 8).

In this study, we aim to determine which factors effectively predict mortality in patients admitted to intensive care due to COVID 19.

Methods

Study design and candidates:

This study was conducted in intensive care units of Ankara City Hospital. 3. These intensive care units, where patients were monitored and treated as level intensive care, were intensive care units separated as COVID 19 intensive care units due to the pandemic. Patients admitted to intensive care between 1 October – 31 November 2020 were included in our study. After our hospital's ethics committee's approval, all patient data were obtained from electronic medical records and patient follow-up forms.

The demographic characteristics of all patients, APACHE II values within the first 24 hours of admission to intensive care, Glasgow Coma Scale, presence of comorbidity, lymphocyte count during intensive care admission, length of stay in intensive care, and presence of mortality were recorded.

In this study, while 65 years and over were accepted as advanced age, the average APACHE value for APACHE II was obtained by univariate analysis. As a result of this analysis, it was planned to compare those with APACHE II values of 19 and above and those below. Again, patients with a value of 10 and below for the Glasgow coma scale were accepted as patients in the precoma state. The cut-off value for the baseline lymphocyte count was accepted as $0.8 \times 10^9 / L$, based on the national COVID – 19 science committee guidelines.

Statistical Analysis

Statistical analyses were performed using Statistical Package for Social Sciences (IBM SPSS Inc, Chicago, IL, USA) version 20.0. Descriptive statistics were expressed as mean $\pm$ standard deviation or median (min-max) for continuous variables and number/percentage for categorical variables. Chi-square test was used for the categorical parameters, and Anova Table Test was used for continuous parameters. Variables with a p-value < 0.05 in the univariate analysis were included in the multivariate analysis after the correlation between the factors was determined for exitus. Multivariate Backward Stepwise Cox Proportional Hazard Regression Analysis was used to determine the effects of variables effective on mortality. P-value < 0.05 was considered statistically significant for the results.

Results

The average age of the 108 individuals evaluated in the study was 67 ± 13.61 years, ranging from 27 to 89. The geriatric group comprised 56.5% of the patient group. Seventy (64.8%) of the patients were female, eighty-nine (82.4%) patients had at least one comorbidity. There were hypertension in 46 (42.6%) patients, diabetes mellitus in 39 (36.1%) patients, cardiovascular disease in 31 (28.7%) patients, chronic obstructive pulmonary disease in 18 (16.7%) patients, chronic renal failure in 9 (8.3%) patients, history of cerebrovascular disease 8 (7.4%) patients, 8 (7.4%) patients with a history of malignancy, and 4 (3.7%) asthma bronchiale (Table 1).

Table 1
Demographic, clinical, characteristics of patients

Characteristics		Mean $\pm$ SD	Median (range)
Age (year)		67 $\pm$ 13.61	69 (27–89)
APACHE II		20.17 $\pm$ 11.55	19 (1–50)
Glasgow Coma Scale Score		11.78 $\pm$ 4.14	13.5 (3–15)
ICU length of stay (day)		9.56 $\pm$ 7.43	7 (1–39)
Lymphocyte count at ICU admission ($\times 10^9$ /l)		774.5 $\pm$ 589.28	605 (80-3910)
		n	Percentage
Geriatric patient	< 65 years	47	43.5
	$\geq$ 65 years	61	56.5
Glasgow Coma Scale Score	$\leq$ 10	31	28.7
	> 10	77	71.3
Gender	Female	38	35.2
	Male	70	64.8
Any Co-morbidity	Absent	19	17.6
	Present	89	82.4
Hypertension	Absent	62	57.4
	Present	46	42.6
Diabetes Mellitus	Absent	69	63.9
	Present	39	36.1
Chronic Obstructive Pulmonary Disease	Absent	90	83.3
	Present	18	16.7
Asthma Bronchiale	Absent	104	96.3
	Present	4	3.7
Cardiovascular Disease	Absent	77	71.3
	Present	31	28.7
History of Malignity	Absent	100	92.6
	Present	8	7.4

Characteristics		Mean ± SD	Median (range)
Chronic Renal Failure	Absent	99	91.7
	Present	9	8.3
Cerebrovascular Disease	Absent	100	92.6
	Present	8	7.4
Other Co-morbidity ¹	Absent	79	73.1
	Present	29	26.9
Mortality	No	46	42.6
	Yes	62	57.4

The study group's median APACHE II value was 19 (range; 1–50), and the median Glasgow Coma Scale Score value was 13.5 (range; 3–15). Glasgow Coma Scale Score value was 10 in 31 (28.7%) patients. When admitted to ICU, the median lymphocyte value was $605 \times 10^9 /L$ and varied between $80 \times 10^9 /L$ and $3910 \times 10^9 /L$. It was observed that the patients stayed in the ICU for an average of 9.56 ± 7.43 days, 46 (42.6%) patients were discharged from the ICU, and 62 (57.4%) patients died (Table 1).

In the univariate analysis, it was seen that age, APACHE II score, Glasgow Coma Scale Score, lymphocyte value when accepted to ICU, presence of any comorbidity determined mortality (Table 2).

Table 2

Factors predicting the mortality in patient with COVID 19 in intensive care unit, univariate analysis

Features		Mortality		<i>P</i> Value
		No	Yes	
		n (%)	n (%)	
Age ¹	< 65 years	30 (63.8)	17 (36.2)	< 0.001
	≥ 65 years	16 (26.2)	45 (73.8)	
APACHE II ²	≤ 19	43 (79.6)	11 (20.4)	< 0.001
	> 19	3 (5.6)	51 (94.4)	
Glasgow Coma Scale Score ³	≤ 10	3 (9.7)	28 (90.3)	< 0.001
	> 10	43 (55.8)	34 (44.2)	
Gender	Female	14 (36.8)	24 (63.2)	0.373
	Male	32 (45.7)	38 (54.3)	
Lymphocyte count at ICU admission (10 ⁹ /l) ⁴	≤ 800	24 (32.4)	50 (67.6)	0.002
	> 800	22 (64.7)	12 (35.3)	
Any Co-morbidity	Absent	13 (68.4)	6 (31.6)	0.012
	Present	33 (37.1)	56 (62.9)	
Hypertension	Absent	31 (50)	31 (50)	0.071
	Present	15 (32.6)	31 (67.4)	
Diabetes Mellitus	Absent	35 (50.7)	34 (49.3)	0.023
	Present	11 (28.2)	28 (71.8)	
Chronic Obstructive Pulmonary Disease	Absent	42 (46.7)	48 (53.3)	0.056
	Present	4 (22.2)	14 (77.8)	
Asthma Bronchiale	Absent	44 (42.3)	60 (57.7)	0.760
	Present	2 (50)	2 (50)	
Cardiovascular Disease	Absent	35 (45.5)	42 (54.5)	0.343
	Present	11 (35.5)	20 (64.5)	
History of Malignity	Absent	43 (43)	57 (57)	0.762
	Present	3 (37.5)	5 (62.5)	
Chronic Renal Failure	Absent	43 (43.4)	56 (56.6)	0.557

	Present	3 (33.3)	6 (66.7)	
Cerebrovascular Disease	Absent	43 (43)	57 (57)	0.762
	Present	3 (37.5)	5 (62.5)	
Other Co-morbidity ⁵	Absent	36 (45.6)	43 (54.4)	0.302
	Present	10 (34.5)	19 (65.5)	
¹ : Geriatric age				
² : Median value				
³ : The cut-off value for Glasgow Coma Scale Score was selected 10 which is defined comatose patient				
⁴ :				
⁵ : Other co-morbidity;				
ICU: Intensive Care Unit				

Factors associated with mortality were evaluated by correlation in univariate analysis, and it was found that there was only a correlation between the presence of any comorbidity and diabetes mellitus. Therefore, excluding diabetes mellitus, age (≤ 65 years vs. <65 years), lymphocyte value ($\leq 800 \times 10^9 /L$ vs. $> 800 \times 10^9 /L$), APACHE II score (> 19 vs. ≤ 19) when admitted to ICU), the presence of any comorbidity (present vs. absent), and the Glasgow Coma Scale Score (≤ 10 vs. > 10) for multivariate analysis to determine death. In the multivariate analysis, it was determined that lymphocyte value, APACHE II score, and the presence of any comorbidity are independent prognostic factors for mortality when accepted to ICU. Death increased 76 times in those with APACHE II score > 19 (95% Confidence Interval: 10.851-533.783; $p < 0.001$). This ratio was 7 (95% Confidence Interval: 1.520-33.827; $p = 0.013$) for those with a lymphocyte count of $\leq 800 \times 10^9 /L$ and 8 for those with any comorbidity (95% Confidence Interval: 1.015–64.151; $p = 0.048$) (Table 3).

Table 3
Factors predicting exitus, multivariate analysis

Factors	Odds Ratio	95% Confidence Interval	P Value
Age (≥ 65 years vs. <65 years) ¹	2.4	0.625–9.212	0.202
Lymphocyte count ($\leq 800 \times 10^9/l$ vs. $>800 \times 10^9/l$) ²	7.171	1.520-33.827	0.013
APACHE II (> 19 vs. ≤ 19) ³	76.105	10.851-533.783	< 0.001
Co-Morbidity (<i>present</i> vs. <i>absent</i>)	8.068	1.015–64.151	0.048
Glasgow Coma Scale Score (≤ 10 vs. >10) ⁴	1.487	0.154–14.374	0.732
¹ : Geriatric age was selected for analysis			
² :			
³ : Median value			
⁴ : The cut-off value for Glasgow Coma Scale Score was selected 10 which is defined comatose patient			

Discussion

Our primary aim in this study was to evaluate whether age, APACHE II value, presence of comorbidity, GCS, and baseline lymphocyte count were successful in predicting mortality in COVID 19 patients. Our study determined that the baseline lymphocyte value, APACHE II score, and the presence of any comorbidity are independent prognostic factors for mortality.

For many years, many scoring systems have been used in intensive care units to predict mortality and morbidity. Among these systems, APACHE II is accepted as the most successful scoring system in predicting mortality in all intensive care types and different patient groups (9, 10). In some articles, it has been reported to be successful in COVID 19 patients (11, 12). A study conducted by Zou et al., they stated that the APACHE II value of 17 or above in COVID 19 patients was an independent predictor for hospital mortality (13). Our study determined that the APACHE II value of > 19 and above is an independent risk factor, and death is 76 times more common in these cases. APACHE II scoring, which is calculated by taking the worst values during admission to intensive care, acts as an early warning system for physicians' high scores following these cases. These scoring systems are widely used in centers where intensive patient admissions are made during the pandemic, and they are instrumental in planning the treatment process of cases with high APACHE II values.

Although there is not enough information about using the Glasgow coma scale, which is frequently preferred in neurological examination in intensive care, in COVID – 19 cases, the GCS data can be based on studies using APACHE II since it is a parameter of APACHE II scoring. Our study showed that GCS being 10 or less has a significant relationship with mortality. In 28.7% of the cases, the GCS was 10 or less, and the mortality was 90.3%.

In the COVID 19 outbreak, it was observed that mortality rates were different in different age groups. It is observed that pulmonary physiology, pathology, and functions change in the presence of lung infection with aging. Therefore, in elderly individuals, response to the disease and tolerability deteriorate, and the mortality rate increases (14). Studies on advanced age COVID 19 patients have shown an increased risk of death (15–18). In our study, 56.5% of the cases were 65 years old and above, and the mortality rate in these cases was significantly higher compared to patients aged 65 years or younger.

The presence of concomitant diseases in COVID 19 cases complicates the clinical picture. In their study by Chen et al., in which they evaluated the epidemiological and clinical characteristics of the cases they followed up with COVID 19 viral infection, they determined the presence of chronic disease in 51% of the cases and stated that cardiovascular, cerebrovascular disease, and diabetes mellitus were the most common accompanying diseases. They indicate in their research that mortality is higher in cases with comorbidity (19). In our research, we observed that 82.4% of our cases had at least one concomitant disease. The most common accompanying disease was hypertension (42.6%), followed by diabetes mellitus (36.1%). In our study, unlike Chen et al., the accompanying cardiovascular disease rate was in the 3rd rank with 28.7%. While any comorbidity's presence increased the risk of death eight times, mortality was found to be statistically significantly higher in patients with diabetes mellitus compared to those without diabetes mellitus.

The absolute value of lymphocytes decreases in COVID 19 associated viral infection. The reason for this decrease is related to the effect of 2019-nCoV on SARS-CoV lymphocytes, especially T lymphocytes. Virus particles spread to the respiratory mucosa and initiate a cytokine storm in the body. This situation stimulates the immune system and causes changes in peripheral white blood cells and immune cells such as lymphocytes. Some patients progress rapidly and pass away by developing ARDS, septic shock, and multiple organ failure. For this reason, early detection and timely treatment of critical cases are vital. The decrease in the absolute lymphocyte count during admission to intensive care is a laboratory parameter that supports clinicians' diagnosis during the diagnosis of COVID 19. In comparison, Huang (20) and colleagues stated that their absolute lymphocyte count was $< 1.0 \times 10^9 / L$ in 63% of their patients, Fan et al. showed that absolute lymphocyte count $< 0.6 \times 10^9 / L$ had a significant correlation with mortality (7). In our study, when we evaluated the mortality relation of absolute lymphocyte count $< 0.8 \times 10^9 / L$, we determined that lymphocyte count $< 0.8 \times 10^9 / L$ was an independent predictor for mortality. Absolute lymphocyte count $< 0.8 \times 10^9 / L$ increased the risk of death seven times.

Frater et al. state that there is some geographic variation in the percentage of COVID-19 patients presenting with lymphopenia in their article evaluating COVID 19 and clinical, hematological laboratory

findings (21). For example, an article from Singapore reporting several COVID-19 patients describes a much lower percentage of lymphopenia patients, as in a retrospective analysis of COVID-19 patients from Zhejiang Province, located ~ 450 miles from Wuhan (7, 22). In contrast, in studies reported from Italy, lymphopenia is common in most patients admitted to the emergency room (23). The reasons for these and similar discrepancies are unclear, although they are probably multifactorial. Due to viral genomic mutations, it is possible that the immunological response to the virus will change as the pandemic spreads to other countries. Another possibility is that testing patients is not uniform, and the degree of lymphopenia can vary depending on the time of admittance. In our study, we have observed that 74 of 108 patients had an absolute lymphocyte value $< 0.8 \times 10^9 / L$.

There were some limitations in our study. The first is that it is a retrospective study, and the second is that there is no long-term (28 days or 6 months) data when determining intensive care and hospital mortality. Besides, we think that further studies should be conducted that comparative studies of baseline absolute lymphocyte count with data from different countries may help determine the cut-off value for lymphopenia.

Conclusion

In our study, we determined that age, APACHE II value, presence of comorbidity, and initial lymphocyte count are independent predictors of mortality. We concluded that studies with more patients and other clinical / laboratory data related to COVID-19 would be beneficial.

Declarations

Ethics approval and consent to participate:

Trial Registry:

SEMA TURAN-2021-01-13T10_31_59

Registry address:

<https://bilimselarastirma.saglik.gov.tr>

Consent for publication:

No consent form since this study is retrospective research study

Availability of data and materials :

All data is available our hospital medical data bank

Competing interest:

The authors declare that they have no competing interests

Funding:

No funding

Acknowledgements:

No applicable

References

1. Gralinski LE, Menachery VD: Return of the coronavirus: 2019-nCoV. *Viruses* 2020; 12:135
2. Immovilli P, Morelli N, Antonucci E, Radaelli G, Barbera M, Guidetti D. COVID-19 mortality and ICU admission: the Italian experience. *Crit Care*. 2020;24(1):228.
3. Mejdoubi M, Kyndt X, Djennaoui M. ICU admissions and in-hospital deaths linked to COVID-19 in the Paris region are correlated with previously observed ambient temperature. *PLoS One*. 2020;15(11):e0242268.
4. Zhang X, Li S, Niu S. ACE2 and COVID-19 and the resulting ARDS. *Postgrad Med J*. 2020;96(1137):403-407.
5. Lee H, Lim CW, Hong HP, Ju JW, Jeon YT, Hwang JW, Park HP. Efficacy of the APACHE II score at ICU discharge in predicting post-ICU mortality and ICU readmission in critically ill surgical patients. *Anaesth Intensive Care*. 2015;43(2):175-86.
6. Lippi G, Plebani M. Laboratory abnormalities in patients with COVID-2019 infection. *Clin Chem Lab Med*. 2020.
7. Fan BE, Chong VCL, Chan SSW, et al. Hematologic parameters in patients with COVID-19 infection. *Am J Hematol*. 2020.
8. Pourbagheri-Sigaroodi A, Bashash D, Fateh F, Abolghasemi H. Laboratory findings in COVID-19 diagnosis and prognosis. *Clin Chim Acta*. 2020 Nov;510:475-482.
9. Moreno RP, Nassar AP Jr. Is APACHE II a useful tool for clinical research? *Rev Bras Ter Intensiva*. 2017;29(3):264-267.
10. Zimmerman JE, Kramer AA, McNair DS, Malila FM. Acute Physiology and Chronic Health Evaluation (APACHE) IV: hospital mortality assessment fortoday's critically ill patients. *Crit Care Med*. 2006;34(5):1297-310.
11. Wendel Garcia PD, Fumeaux T, Guerci P, Heuberger DM, Montomoli J, Roche-Campo F, Schuepbach RA, Hilty MP; RISC-19-ICU Investigators. Prognostic factors associated with mortality risk and disease progression in 639 critically ill patients with COVID-19 in Europe: Initial report of the international RISC-19-ICU prospective observational cohort. *EClinicalMedicine*. 2020;25:100449.
12. Grasselli G, Zangrillo A, Zanella A, et al. Baseline Characteristics and Outcomes of 1591 Patients Infected With SARS-CoV-2 Admitted to ICUs of the Lombardy Region, Italy. *JAMA*.

2020;323(16):1574–1581.

13. Zou X, Li S, Fang M, Hu M, Bian Y, Ling J, Yu S, Jing L, Li D, Huang J. Acute Physiology and Chronic Health Evaluation II Score as a Predictor of Hospital Mortality in Patients of Coronavirus Disease 2019. *Crit Care Med*. 2020;48(8):e657-e665.
14. Miller EJ, Linge HM. Age-related changes in immunological and physiological responses following pulmonary challenge. *Int J Mol Sci* 2017; 18: E1294
15. Yang X, Yu Y, Xu J, et al. Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *Lancet Respir Med* 2020; 8: 475–481.
16. Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020; 395: 1054–1062.
17. Wu C, Chen X, Cai Y, et al. Risk factors associated with acute respiratory distress syndrome and death in patients with coronavirus disease 2019 pneumonia in Wuhan, China. *JAMA Intern Med* 2020; in press [<https://doi.org/10.1001/jamainternmed.2020.0994>].
18. Du RH, Liang LR, Yang CQ, et al. Predictors of mortality for patients with COVID-19 pneumonia caused by SARS-CoV-2: a prospective cohort study. *Eur Respir J* 2020; 55: 2000524.
19. Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, Qiu Y, Wang J, Liu Y, Wei Y, Xia J, Yu T, Zhang X, Zhang L. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet*. 2020 Feb 15;395(10223):507-513. doi: 10.1016/S0140-6736(20)30211-7. Epub 2020 Jan 30. PMID: 32007143; PMCID: PMC7135076.
20. Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet*. 2020;395(10223):497-506.
21. Frater JL, Zini G, d'Onofrio G, Rogers HJ. COVID-19 and the clinical hematology laboratory. *Int J Lab Hematol*. 2020 Jun;42 Suppl 1:11-18. doi: 10.1111/ijlh.13229. PMID: 32311826; PMCID: PMC7264622.
22. Xu XW, Wu XX, Jiang XG, et al. Clinical findings in a group of patients infected with the 2019 novel coronavirus (SARS-Cov-2) outside of Wuhan, China: retrospective case series. *BMJ*. 2020;368:m606.
23. Buoro S, Di Marco F, Rizzi M, et al. Papa Giovanni XXIII Bergamo Hospital at the time of the COVID-19 outbreak: letter from the warfront *Int J Lab Hematol*. 2020;42(Suppl. 1):8–10