

# Control of Panama Disease of Banana by Intercropping with Chinese Chive (*Allium tuberosum* Rottler): Cultivar Differences

Zhen-Fang Li

Fujian Agriculture and Forestry University

Tong Wang

Fujian Agriculture and Forestry University

Chenlin He

Fujian Agriculture and Forestry University

Kelin Cheng

Fujian Agriculture and Forestry University

Rensen Zeng

Fujian Agriculture and Forestry University

Yuanyuan Song (✉ [yyuansong@163.com](mailto:yyuansong@163.com))

---

## Research article

**Keywords:** Banana, Fusarium wilt disease, Chinese chive, Intercropping, 2-Methyl-2-pentenal, Cultivar

**Posted Date:** May 21st, 2020

**DOI:** <https://doi.org/10.21203/rs.3.rs-25892/v1>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

**Version of Record:** A version of this preprint was published on September 17th, 2020. See the published version at <https://doi.org/10.1186/s12870-020-02640-9>.

# Abstract

## Highlights

- | Intercropping of banana with Chinese chive can effectively reduce Panama disease
- | Selection of appropriate cultivars of Chinese chive for intercropping is critical
- | 2-Methyl-2-pentenal (MP) is the key compound for the pathogen inhibition
- | MP content and antifungal activity are key indices for cultivar selection

**Background** Panama disease (Fusarium wilt disease) caused by *Fusarium oxysporum* f. sp. *cubense* race 4 (FOC) severely threatens banana (*Musa* spp.) production worldwide. Intercropping of banana with *Allium* plants has shown a potential to reduce Panama disease. In this study six cultivars of Chinese chive (*Allium tuberosum* Rottler) were selected to compare their differences in antifungal activity and active compounds. Three cultivars Duokang Fujiu 11, Fujiuhuang 2, and Duokang Sijiqing with higher levels of antifungal compounds were further used for intercropping with banana in the pots and field to compare their effects on growth and disease incidence of banana.

**Methods and Results** The six cultivars showed significant differences in antifungal activity against FOC mycelia growth in both leaf volatiles and aqueous leachates. The aqueous leachates displayed stronger antifungal activity than the volatiles. Contents of three main antifungal compounds dipropyl trisulfide (DPT), dimethyl trisulfide (DMT), and 2-methyl-2-pentenal (MP) in volatiles and aqueous leachates varied considerably among cultivars. Pot and field experiments showed that intercropping with three selected Chinese chive cultivars significantly improved banana vegetative growth, increased photosynthetic characteristics and yield but decreased disease incidence of Panama disease.

**Conclusions** Our results indicate that intercropping with Chinese chive shows potential to reduce banana Panama disease and selection of appropriate cultivars is vital for effective disease control.

## Background

Banana (*Musa* spp.) is one of the world's most important crops and it serves as a staple food in many developing tropical and subtropical countries (Li et al., 2019). It ranks next to rice, wheat, and maize in terms of its importance as a staple food crop (Singh et al., 2016). Over 100 million metric tons of the crop are produced annually (Paul et al., 2017). However, this crop production is severely hampered by Panama disease (also known as Fusarium wilt disease) caused by *Fusarium oxysporum* f. sp. *cubense* race 4 (FOC) (Segura et al., 2018), leading to extensive failure of commercial plantations in all banana-producing areas, including Asia, Central and South America, Africa, and Australia (Ploetz et al., 2015; Xue et al., 2015; Dong et al., 2019). The soil-borne fungal pathogen colonizes and occludes the host xylem to cause a vascular wilt (Butler, 2013; Ploetz et al., 2015; Dong et al., 2019). So far not yet a practically effective approach of controlling this notorious disease in the field is available. Soil sterilization using

fungicides has been used only for intensive agriculture in greenhouses on a limited scale (Shen et al., 2015; Shen et al., 2018). Environmentally friendly cultural practices and fungicides are urgently needed (Sun et al., 2015; Fu et al., 2017; Xiong et al., 2017). Fungicides are not able effectively to control this soil-borne pathogen in the field, for a variety of ecological reasons (Shen et al., 2013). Nowadays, cultural practices have been tried with little success, for there has not been a suitably productive cultivar to selection or breeding for reducing the incidence of diseases caused by FOC (Wei et al., 2017). Recently, crop rotation and microorganisms have shown promise for reducing the disease severity (Trivedi et al., 2017; Pattison et al., 2018).

Chinese farmers have practiced intercropping and rotation for disease control over two thousand years (Han et al., 2016; Zhu & Morel et al., 2019). Intercropping, which is defined as the growing of two or more crop species in proximity simultaneously during a growing season, is an agricultural practice that not only increases crop yield and stress resilience, but also reduces the incidence of microbial diseases. Chemicals produced by one plant in the intercropping system, can either directly inhibit microbial pathogens in the soils, or indirectly trigger resistance through the induction of the plant defense system in neighboring plants (Li et al., 2018; Yu et al., 2019). The *Allium* plants are characterized with production of many antimicrobial substances (Yu 1999; Benkeblia 2004; Sharifi-Rad et al., 2016; Liaqat et al., 2019). They are among the oldest medicinal herbs, used as antibacterial, antifungal, antioxidant and cytotoxic drugs. In addition to many organosulphur compounds, mainly disulphides and trisulphides, *Allium* plants also produce non-volatile compounds such as spirostanol, furostanol and cholesterol saponin with antimicrobial, antifungal, antiplatelet aggregating, and cytotoxic activities (Hu et al., 2009; Zuo et al., 2015; Katarzyna et al., 2016; Gao et al., 2018; Jannat et al., 2019). Many *Allium* species such as garlic, onion and Chinese chive (*Allium tuberosum*) are widely cultivated for their spice and medicinal properties. Chinese chive is an important vegetable for making dumpling and kimchi in many Asian countries such as China and Korea. Chinese chive is now gaining increasingly attention due to its diverse important pharmacological activities (Huang et al., 2012; Jannat et al., 2019). Although many studies have been conducted for food and pharmacological purposes (Lanzotti et al., 2014; Sharifi-Rad et al., 2016;), *Allium* plants as biological control agents for crop disease in the intercropping system and underlying mechanisms are scarcely addressed (Zhang et al., 2013; Dania et al., 2019). The sequential cropping systems are becoming increasingly popular in all management practices for their environmental friendliness.

Recent studies indicated that banana intercropping with Chinese chive can effectively control Fusarium wilt of banana (Huang et al., 2012; Zhang et al., 2013; Wibowo et al., 2015). Our previous study showed that Chinese chive exhibited strong inhibitory effects on FOC through the release of organic volatiles, among which 2-methyl-2-pentenal, dimethyl trisulfide, dimethyl disulfide, dipropyl disulfide, and dipropyl trisulfide are the main active antifungal compounds (Zhang et al., 2013). However, crop varieties may markedly differ in their ability to produce antimicrobial chemicals (Soni et al., 2019).

In this paper, we selected six Chinese chive cultivars with obviously morphological difference and distinct characteristics to compare their antifungal activity against FOC and active compounds in the leaf

volatiles and aqueous leachates of aerial parts. Three cultivars with higher contents of three antifungal compounds including dipropyl trisulfide (DPT), dimethyl trisulfide (DMT), 2-methyl-2-pentenal (MP) in the volatiles or aqueous leachates were further used in pot experiments to evaluate their potential in reducing disease incidence of *Fusarium* wilt disease of banana.

## Methods

### Chemicals

Three antifungal compounds including dipropyl trisulfide (DPT), dimethyl trisulfide (DMT), and 2-methyl-2-pentenal (MP) were obtained from Sigma-Aldrich (St. Louis, MO, USA). All solvents used were analytical or HPLC grade.

### Plant Materials

The banana plant in this study was “Williams” (AAA genome, Cavendish subgroup) banana cultivar that is highly susceptible to *Fusarium oxysporum* f. sp. *cubense* race 4 (Warman, 2018). The six Chinese chive cultivars were collected from Fujian province (China). The cultivar Duokang Fujiu 11 (DKF) shows a strong resistance to stress and a strong special spicy odor. The Fujiubao (FJB) shows the strongest spicy odor among the six varieties but lowest stress resistance. The Fujiuhuang 2 (FJH) shows a moderate spicy odor and stress resistance, its leaves are yellow (the rest were green). The Duokang Sijiqing (DKS) shows highly resistance but almost no spicy odor. The Futaijiu 1 (FTJ) shows a strong spicy odor but weak resistance and garlic sprout. The Zigenchun Zaohong (ZGC) shows a medium level of stress resistance and spicy odor but red roots (roots of the rest are white).

### Foc Fungal Materials

The original fungal inocula of *Fusarium oxysporum* f. sp. *cubense* race 4 (FOC) were kindly provided by Professor Zide Jiang of College of Agriculture at South China Agricultural University in Guangzhou (Zhang et al., 2013). The strain was maintained in Potato Dextrose Agar (PDA), and the inocula were cultured in PDA at 25 °C in darkness. PDA medium was prepared and sterilized by autoclaving at 121 °C for 20 min (Dong et al., 2019).

The FOC conidial suspension was prepared by incubating FOC in PDA broth on a shaker at 200 rpm and 28 °C for 5 days, followed by filtration with two layers of gauze to remove the mycelia. The conidial concentration was adjusted to  $1 \times 10^6$  conidia per milliliter (Sun, 2019).

## Inhibitory Effect Of Leaf Volatiles

To compare the inhibitory effect of the volatiles from the six Chinese chive cultivars, fresh leaves were cut into 1 cm segments or ground into powder in liquid nitrogen, respectively. Then the leaves were added in a four-room petri dish (90 mm in diameter) and inoculated with two 4 mm discs of FOC inocula. The concentration of volatiles was 0.6 g FW in the petri dish. Similar petri dish without Chinese chive leaves were used as controls. Six replicates were prepared for each treatment. Each six petri dish containing the same sample were placed in the middle of a 4,000 ml glass jar (the height and the diameters of top and bottom were 12.5, 22, and 19 cm, respectively).

The experiment was conducted in a climate-controlled room at 25 °C in darkness. Colony diameters of the inoculated FOC were measured with a Vernier caliper when the colony of the control had reached the peripheral wall.

## **Inhibitory Effect Of Aqueous Leachates Of The Aerial Parts**

To examine the inhibitory effect of aqueous leachates of the six Chinese chive cultivars on FOC colony growth, the fresh aerial parts (50 g) were also ground into powder in liquid nitrogen and soaked in 100 mL distilled water at 25 °C for 20 min for extractions by an ultrasonic agitation, respectively (Zuo et al 2016). Then aqueous solution was centrifuged at 10000 rpm for 5 min. After centrifugation, the supernatant was filtered first through normal filter paper (0.45 µm, Xingya Purification Factory, Shanghai, China) and then through a microfilter of pore size 0.22 µm (Whatman puradisc™25AS polyethersulfone membrane, catalog NO.6794 – 2514, England). The final filtrate was our original leachates with a concentration of 0.5 g FW·mL<sup>-1</sup>.

The filtered leachates (5 mL) of each cultivar and fungal culture medium (5 mL) were mixed and added in petri dishes (90 mm in diameter) and inoculated with a 4 mm discs of FOC. The concentration of final culture medium was 0.25 g FW·mL<sup>-1</sup>. Similar petri dish without Chinese chive leachate but with the same volume of distilled water were used as controls. Six replicates were prepared for each treatment.

The experiment was conducted in a climate-controlled room at 25 °C in darkness. Colony diameters of the inoculated FOC were measured with a Vernier caliper when the colony of the control had reached the peripheral wall.

## **Antifungal Activities Of Dpt, Dmt And Mp**

Based on our previous result, three compounds including DPT, DMT, and MP were used to test the antifungal activities against FOC colony growth. Four beakers (250 mL, 75 mm in diameter) filled with 10 mL fungal culture medium of PDA were placed in a 4,000 mL glass jar (the height and the diameters of top and bottom were 12.5, 22, and 19 cm, respectively), and inoculated with 4 mm disc of FOC. A piece of degreasing cotton filled with 100 µL the antifungal compounds (DPT, DMT, and MP) was placed in the middle of the jar. The final concentration of the three compounds was 25 µL·L<sup>-1</sup>. Similar beakers in each

glass jar without antifungal compound were used as controls. The glass jars were covered carefully with two layers of cellophane to stop the volatiles from escaping the (Nandakumar et al., 2018). Five replicates were prepared for each treatment. Thus, each treatment had 20 beakers placed in 5 glass jars.

The experiment was conducted in an incubator at 25 °C. Colony diameters of the inoculated FOC were measured with a ruler from the bottom by placing the glass jar on a high platform starting from 48 h post inoculation until 5 day when the colony of the control (no antifungal compound) breakers had grown close to the peripheral wall. **Quantification of DPT, DMT and MP in the volatiles or aqueous leachates**

To quantify DPT, DMT and MP in the volatiles released from Chinese chive cultivars, the volatiles were trapped according to the methods described by Zhang et al (Zhang et al., 2013). The leaves of Chinese chives (150 g) were cut with 1 cm segments and added into a 500 mL flask. The volatiles were trapped through a glass tube (3 mm inner diam, 14 cm long) packed with 150 mg of Tenax TA (2.6–diphenylene oxide polymer resin) for 5 h. Trapped volatiles were eluted from each tube with 1 mL *n*-hexane and then were used for analysis by a gas chromatography system (GC) as described by Nandakumar, R. et al. (2018).

To quantify DPT, DMT and MP in the aqueous leachates of Chinese chive cultivars, the leaves (50 g) were ground into powder in liquid nitrogen and extracted with 100 mL distilled water at 25 °C for 20 min through an ultrasonic agitation, respectively. Then aqueous solution was centrifuged at 10000 rpm for 5 min. The supernatant (60 mL) was consecutively partitioned with dichloromethane (100 mL). The dichloromethane layer was separated with a separatory funnel, followed by evaporation under vacuum. The substance of dichloromethane phase was finally dissolved with dichloromethane to 5 mL. The final solution was filtrated by a micro filter with pore size 0.22 µm for and ready for GC analyses.

GC analyses were performed on an Agilent GC 7890B system (Agilent Technologies, Palo Alto, CA, USA) with a flame ionization detector (FID) and a capillary column Agilent 19091J-413 HP-5 (5% phenylmethyl siloxane, 30.0 m × 320 µm × 0.25 µm). The injection temperature was 150 °C, and the oven temperature was raised from 35 °C (2 min hold) to 250 °C at a rate of 10 °C/min. Two microliters of sample were injected, and nitrogen was used as the carrier gas at a flow rate of 1.0 ml/min.

### **Banana intercropping with three Chinese chive cultivars in the pots**

Based on the results of antifungal activities and quantity of the three active compounds, three cultivars (FJH, DKS, and DKF) of Chinese chive were selected to intercrop with banana in pots (the height and the diameters were 34 and 40 cm) filled with soils from the field of monoculture banana plantation for six years in Moxi Village, Tianbao Town of Nanjing County, Fujian Province (China, 24°29'14" north latitude, 117°29'3" longitude east). Thirty days after intercropping with Chinese chive under greenhouse conditions (25 ± 4 °C; 16/8 h light/dark photoperiod), each pot was watered with 100 ml *Fusarium conidial* suspension ( $1 \times 10^6$  spores·mL<sup>-1</sup>) prepared as described above. The monoculture banana in the same soil served as a control (CK). The treatments and controls consisted of 10 plants per replicate.

Sixty days after fungal inoculation, plant height, stem circumference, photosynthesis parameters and disease indices were recorded. Net photosynthetic rate and stomatal conductance of the bananas were measured with Li-6400 (LI-COR, Lincoln, NE, USA). At the end of the experiment, each plant was harvested to assess the disease indices (Cheng et al., 2020).

### **Banana intercropping with three Chinese chive cultivars in the field**

Seeds of three Chinese chive cultivars (FJH, DKS, and DKF) were sown in an experimental field in Moxi Village, Tianbao Town of Nanjing County, Fujian Province (China), in March of 2016. Ten months after planting, the Chinese chive plants were then transplanted (at a spacing of  $15 \times 10 \text{ cm}^2$ ) in a Fusarium wilt disease frequently-occurred field in the village in January of 2017. Before the chives were planted, the field was divided into 20 plots each with a size of  $2.5 \times 25 \text{ m}^2$  (Yu 1999; Huang et al., 2016). Then, ten plants (at a spacing of  $2.5 \times 2.25 \text{ m}^2$ ) of banana cultivar “Williams” were intercropped with the Chinese chive cultivars (FJH, DKS, and DKF) in the field. The field experiment design was randomized complete block design with four different treatments with 5 plots as replications. The planting site was prepared and managed using established cultivation practices, without the application of either herbicide or fungicides. Plants were fertilized twice a year by using NPK (N:  $\text{P}_2\text{O}_5$ : KCL = 10:3:2) with dose of  $3 \text{ kg}\cdot\text{plot}^{-1}$ . The incidence of Fusarium wilt disease of banana was examined every 30 days. When the banana ripened, fruit weight of each plant was weighed.

## **Statistical Analysis**

Antifungal activities of three antifungal compounds, inhibitory effect of volatiles or aqueous leachates of the six Chinese chive cultivars on FOC were presented as inhibition rate (%) compared to control using the following formula: inhibition rate (%) = [(colony diameters of control - colony diameters of treatment) / colony diameters of control]  $\times 100$ .

All data were subjected to analysis of variance using the Statistical Analysis System Program (SPSS 9.0). Each value was expressed as the mean  $\pm$  the standard error (SE). Statistical significance was analyzed using Student's *t*-test and one-way analysis of variance (ANOVA). The precision of the method was confirmed by least-significant difference (LSD, %). The values were considered significant when the *P* value was  $< 0.05$ .

## **Results**

### **Inhibitory effect of volatiles from the six cultivars**

Fresh leaves of the six Chinese chive cultivars were treated in two ways: cut into 1 cm segments to acquire the naturally released volatiles, or ground into powder in liquid nitrogen to acquire maximally released volatiles. Bioassays showed that both the sliced leaves treatment and the ground-leaves treatment strongly inhibited the mycelial growth of FOC (Fig. 1).

In the sliced leaves treatment, the volatiles from all the six Chinese chive cultivars significantly inhibited mycelial growth of FOC (Fig. 1A). FJH cultivar with yellow leaves displayed the maximum inhibition (61%), which was much higher than the other five cultivars. The DKS and ZGC showed 25 and 22% inhibition, respectively. The other three cultivars showed marginal inhibitory effect ( $\leq 12\%$ ).

In the ground-leaves treatment, the inhibitory effect of volatiles of all cultivars was much higher than that in sliced leaves treatments (Fig. 1B). Three cultivars (FJH, ZGC and DKS) displayed 60–62% inhibition against FOC mycelia growth, which were significantly higher than other three cultivars.

## Inhibitory Effects Of Aqueous Leachates Of The Six Cultivars

The FOC colony growth in the presence of aqueous leachates of six Chinese chive cultivars is shown in Fig. 2. The aqueous leachates exhibited significant higher inhibitory effects (inhibition rate ranging from 55–72%) on FOC colony growth than the volatiles. The largest antifungal effect (72%) was obtained from the Chinese chive cultivar DKS, followed by FJH and DKF with inhibition of 68% and 67%, respectively (Fig. 2).

## Antifungal Activity Of Dpt, Dmt And Mp

2-Methyl-2-pentenal (MP), dimethyl trisulfide (DMT) and dipropyl trisulfide (DPT) identified from Chinese chive significantly inhibited mycelial growth of FOC at a concentration of  $25 \mu\text{L}\cdot\text{L}^{-1}$  (Fig. 3). MP almost completely inhibited the mycelial growth (99% inhibition), while DPT and DMT showed approximate 30% inhibition.

### Contents of MP, DMT and DPT in the six cultivars

Since MP, DMT and DPT are important antifungal compounds in Chinese chive, and different chive species exhibited inconsistent inhibitory effect, we speculated that difference in antifungal activities among the six cultivars resulted from their difference in concentrations of DPT, DMT and MP. We, therefore, detected the contents of MP, DMT and DPT (Fig. 4) in the volatiles and aqueous leachates of the six Chinese chive cultivars.

The contents of MP, DMT and DPT significantly differed among the six Chinese chive cultivars. The highest concentrations of MP, the most active anti-FOC compound, was found in the cultivar FJH in both volatiles and leachates. The second most abundant MP was found in the volatiles of the cultivar ZGC and in the leachates of the cultivar DKS. DKS contained relative higher DPT, DMT and MP contents among all six chive species, while FJB and FTJ contained relative lower contents of these three antifungal substances (Fig. 4). For example, the content of DMT in the volatiles of FJH ( $4.454 \text{ g}\cdot\text{g}^{-1}$ ), similar to that in DKS, was nearly two times as the content in FJB ( $2.282 \mu\text{g}\cdot\text{g}^{-1}$ ). This might explain that FJH and DKS

had better performances in antifungal assays. Although DKF released less DMT and DPT than ZGC in aqueous leachates (Fig. 4), its anti-FOC activity was high, which might result from the higher contents of MP in aqueous leachates of DKF ( $0.794 \text{ g}\cdot\text{g}^{-1}$ ) that was significantly higher than ZGC ( $0.253 \text{ g}\cdot\text{g}^{-1}$ ) (Fig. 4).

## Intercropping Improved Banana Growth And Photosynthesis In The Pots

Based on antifungal activity against FOC, three Chinese chive cultivars DKF, FJH and DKS with higher contents of DPT, DMT and MP, were selected for intercropping with banana in pot experiment to examine their differential effects on Panama disease.

Ninety days after intercropping with Chinese chive, plant heights of banana plants were significantly improved by all three cultivars relative to monoculture control. Stem diameters were significantly improved by intercropping with FJH and DKS (Fig. 5).

As shown in Fig. 6, the photosynthetic characteristics of banana functional leaves including net photosynthetic rate ( $P_n$ ), stomatal conductance ( $G_s$ ), and transpiration rate ( $T_r$ ) in intercropping system were significantly higher than those in the monoculture system. Obviously, intercropping with all three cultivars significantly improved net photosynthetic rate and stomatal conductance of banana plants, but reduced intercellular  $\text{CO}_2$  concentration relative to monoculture banana plants (Fig. 6). Intercropping with FJH and DKS also significantly enhanced transpiration rate. Intercropping with DKF did not have obvious effect on transpiration rate. Furthermore, the cultivar FJH showed the strongest effects on the four photosynthetic parameters in the intercropping system.

### Intercropping reduced the Fusarium wilt incidence of banana in the pots

The occurrence of Panama disease of banana was significantly reduced after intercropping with all three cultivars of Chinese chive relative to monoculture banana plants (Fig. 7A). Intercropping with FJH, DKS and DKF reduced the disease indices by 65.7, 62.8 and 29.8%, respectively. The lowest disease indices (Fig. 7A) and very few yellow leaf symptoms were found in the banana plants intercropping with FJH.

### Intercropping reduced the Fusarium wilt disease incidence but increased banana yield in the field

Field experiments were conducted in 2017 and 2018 to evaluate the effects of intercropping with the three selected Chinese chive cultivars on banana resistance to Fusarium wilt, as well as fruit weight per plant and yield (Fig. 8). Intercropping with all three Chinese chives cultivars significantly reduced the disease incidence but increased banana yield in comparison with the monoculture control (Fig. 8A & 8C). The lowest disease incidence and highest yield were found in banana plants intercropping with the cultivar Fujiuhuang 2 (FJH), followed by the cultivar DKS in both 2017 and 2018. Intercropping with FJH and DKS increased the yield by 5.04- and 2.78-fold, respectively (Fig. 8C). Intercropping with DKF

displayed the weakest effect on both banana disease incidence and yield. However, the intercropping with FJH and DKS showed no obvious effects on fresh weight of each plant (Fig. 8B). Only DKF displayed a bit negative effect on fresh weight of each plant.

## Discussion

Panama disease severely affects banana production worldwide. Appropriate agricultural practices show potentials to successfully minimize the detrimental effect of the disease (Shen et al., 2015; Wang et al., 2015; Shen et al., 2018; Shen et al., 2019). Rotation and intercropping with Chinese chive have been demonstrated to be effective approaches to control the disease at both greenhouse and field levels (Huang et al., 2012; Zhang et al., 2013; Wibowo et al., 2015). This study showed that the cultivar selection is a key factor for successful control of the disease. Although all cultivars displayed obvious inhibition to the disease in both greenhouse and field conditions (Figs. 7 & 8), they showed significant difference in their ability to reduce the disease incidence. Two cultivars FJH and DKS exhibited much stronger inhibition against Panama disease. In particular, the cultivar FJH showed the highest inhibition to the disease and best improvement for the banana yield in both pot and field experiments (Figs. 7 & 8). Interestingly, in laboratory both volatiles and aqueous leachates of FJH showed the strongest inhibitory effect on the pathogen FOC among the six cultivars tested (Figs. 1 & 2). Meanwhile, the cultivar FJH produced the highest amount of 2-methyl-2-pentenal in both volatiles and aqueous leachates (Fig. 4A & 4B), which showed the strongest antifungal activity against FOC (Fig. 3; Zhang et al., 2013). All results from laboratory, greenhouse and field experiments suggested that the cultivar FJH is the best cultivar of Chinese chive for intercropping with banana to control Panama disease. The second better cultivar is DKS, which produced the second abundant 2-methyl-2-pentenal in the aqueous leachate (Fig. 4B), and showed the second strongest antifungal activity against FOC in the leachate (Fig. 2). Schulz et al. (1998) revealed the influence of the genetic background of the parent cultivars on the sulfur volatile composition in the *Allium* hybrid. Therefore, selection of appropriate cultivars of Chinese chive with strong anti-FOC activity for intercropping is key for successfully control of Panama disease. More cultivars should be screened to find suitable ones for intercropping with banana.

Volatile compounds released from host plants plays an important role in plant defense against microbial pathogens (Hammerbacher et al., 2019). *Allium* species are well-known to produce many volatile organosulfur compounds, which contribute the unique smell and taste of these plant species (Nandakumar et al., 2018; Marrelli et al., 2019). These compounds exhibit broad-spectrum biological activities towards microorganisms, plants and animals (Ozkan et al., 2013; Marrelli et al., 2019). Although this study showed that aqueous leachates displayed much higher anti-FOC activity, the active compounds in the aqueous leachates were volatiles and low polarity (Zhang et al., 2013; Lian et al., 2019), which is consistent with the result that the aqueous leachates had significantly more active compounds than volatiles (Fig. 4). The results indicated that anti-FOC compounds such as 2-methyl-2-pentenal and dimethyl trisulfide are easily released by Chinese chive via leaching and root exudation. Lian et al. (2019) demonstrated that the aqueous leachates of Chinese chive roots were more inhibitory to FOC than those of stems and leaves. Zhang et al. (2013) revealed that volatiles exuded by roots of

Chinese chive inhibited spore germination of FOC. It is likely that in the field most active compounds of Chinese chive are released into the rhizospheres and then these volatiles spread into the soils to exhibit anti-FOC activity.

Volatile sulfur compounds have been demonstrated to be inhibitory to *Fusarium* and other pathogenic fungi (Kocić-Tanackov et al., 2017). Non-sulfur-containing compound 2-methyl-2-pentenal, a degradation product of thiopropanal S-oxide (Ferary et al., 1996), displayed the strongest anti-FOC activity (Fig. 4). The chemical was also found in the onion (*Allium cepa* L.) essential oil and displayed antifungal activity to *Fusarium* (Kocić-Tanackov et al., 2017).

Anti-FOC activity and contents of 2-methyl-2-pentenal in the volatiles and aqueous leachates may serve as two desired parameters for preliminary selection of appropriate cultivars of Chinese chive for intercropping with banana.

## Conclusion

Panama disease (*Fusarium* wilt disease) caused by *Fusarium oxysporum* f. sp. *cubense* race 4 (FOC) severely threatens banana (*Musa* spp.) production worldwide. Intercropping of banana with Chinese chive can effectively reduce Panama disease.

Our study demonstrates that banana intercropping with Chinese chive can effectively reduce Panama disease and selection of appropriate cultivars of Chinese chive for intercropping is key for the success. The six cultivars showed significant differences in antifungal activity against FOC mycelia growth in both leaf volatiles and aqueous leachates. Pot and field experiments showed that intercropping with three selected Chinese chive cultivars significantly improved banana vegetative growth, increased photosynthetic characteristics and yield but decreased disease incidence of Panama disease. Among six cultivars used FJH and DKS are the most suitable cultivars for intercropping.

## Abbreviations

CK

Control; **DKF**:Duokang Fujiu 11; **DKS**:Duokang Sijiqing; **DMT**:Dimethyl trisulfide; **DPT**:Dipropyl trisulfide; **FJB**:Fujiubao; **FJH**:Fujiuhuang 2; **FOC**:*Fusarium oxysporum* f. sp. *cubense* race 4; **FTJ**:Futaijiu 1; **Gs**:Stomatal conductance; **MP**:2-Methyl-2-pentenal; **PDA**:Potato Dextrose Agar; **Pn**:Net photosynthetic rate; **SE**:Standard error; **Tr**:Transpiration rate; **ZGC**:Zigenchun Zaohong.

## Declarations

- **Ethics approval and consent to participate**

Not applicable.

- **Consent for publication**

Not applicable.

- **Availability of data and materials**

All data generated or analysed during this study are included in this published article

- **Competing interests**

The authors declare that they have no competing interests

- **Funding**

This research was supported by the Natural Science Foundation of Fujian Province of China (2019J01657, 2019J01375), the National Natural Science Foundation of China (31870361, 31670414) and Excellent Youth Science Foundation of Fujian Province of China (2017J06010).

- **Author's contribut**

ZF, RS, and YY conceived and designed the experiments and wrote the manuscript; ZF, TW, CL and KL carried out the experiments and analyzed data; All authors revised the manuscript.

- **Acknowledgments**

Not applicable.

## References

1. Benkeblia N. Antimicrobial activity of essential oil extracts of various onions (*Allium cepa*) and garlic (*Allium sativum*). LWT-Food Science Technology. 2004;37:263–8. <https://doi.org/10.1016/j.lwt.2003.09.001>.
2. Butler D. Fungus threatens top banana. Nature. 2013;504:195–6. <https://doi.org/10.1038/nmiddleeast.2013.241>.
3. Cheng CZ, Li D, Qi Q, Sun XL, Anue MR, David BM, Zhang YY, Hao XG, Zhang ZH, Lai ZX. The root endophytic fungus *Serendipita indica* improves resistance of banana to *Fusarium oxysporum* f. sp. *cubense* tropical race 4. Eur J Plant Pathol. 2020;156:87–100. <https://doi.org/10.1007/s10658-019-01863-3>.
4. Dania VO, Omidiora JA. Combination of biological control agents and garlic (*Allium sativum*) extract in reducing damping-off disease of tomato. Bangladesh Journal of Agricultural Research. 2019;44:553–67. <https://doi.org/10.3329/bjar.v44i3.43485>.
5. Dong HH, Fan HY, Lei ZX, Wu C, Zhou DB, Li HP. Histological and gene expression analyses in banana reveals the pathogenic differences between races 1 and 4 of banana Fusarium wilt

- pathogen. *Phytopathology*. 2019;109:1029–42. <https://doi.org/10.1094/phyto-10-18-0384-r>.
6. Ferary S, Auger J. What is the true odour of cut *Allium*? Complementarity of various hyphenated methods: Gas chromatography-mass spectrometry and high-performance liquid chromatography-mass spectrometry with particle beam and atmospheric pressure ionization interfaces in sulphenic acids rearrangement components discrimination. *J Chromatogr A*. 1996;750:63–74. [https://doi.org/10.1016/0021-9673\(96\)00405-0](https://doi.org/10.1016/0021-9673(96)00405-0).
  7. Gao Q, Li XB, Sun J, Xia ED, Tang F, Cao HQ, Xun H. Isolation and identification of new chemical constituents from Chinese chive (*Allium tuberosum*) and toxicological evaluation of raw and cooked Chinese chive. *Food Chem Toxicol*. 2018;112:400–11. <https://doi.org/10.1016/j.fct.2017.02.011>.
  8. Fu L, Penton CR, Ruan Y, Shen Z, Xue C, Li R, Shen Q. Inducing the rhizosphere microbiome by biofertilizer application to suppress banana *Fusarium* wilt disease. *Soil Biol Biochem*. 2017;104:39–48. <https://doi.org/10.1016/j.soilbio.2016.10.008>.
  9. Hammerbacher A, Coutinho TA, Gershenzon J. Roles of plant volatiles in defense against microbial pathogens and microbial exploitation of volatiles. *Plant Cell Environment*. 2019;42:2827–43. <https://doi.org/10.1111/pce.13602>.
  10. Han GY, Lang J, Sun Y, Wang YY, Zhu YY, Lu BR. Intercropping of rice varieties increases the efficiency of blast control through reduced disease occurrence and variability. *Journal of Integrative Agriculture*. 2016;15:795–802. [https://doi.org/10.1016/s2095-3119\(15\)61055-3](https://doi.org/10.1016/s2095-3119(15)61055-3).
  11. Hu G, Mao R, Ma Z. A new steroidal saponin from the seeds of *Allium tuberosum*. *Food Chem*. 2009;113:1066–8. <https://doi.org/10.1016/j.foodchem.2008.08.061>.
  12. Huang YH, Wang RC, Li CH, Zuo CW, Wei YR, Zhang L, Yi GJ. Control of *Fusarium* wilt in banana with Chinese leek. *Eur J Plant Pathol*. 2012;134:87–95. <https://doi.org/10.1007/s10658-012-0024-3>.
  13. Jannat K, Rahman T, Rahmatullah M. Traditional uses, phytochemicals and pharmacological properties of *Allium tuberosum* Rottler ex spreng. *Journal of Medicinal Plants Studies*. 2019;7:214–20.
  14. Katarzyna AS, Joanna T. Yielding and biological value of Garlic chives (*Allium tuberosum* Rottl. Ex Spreng.) depending on the type of Mulch. *Journal Elementol*. 2016;21:7–9. <https://doi.org/10.5601/jelem.2015.20.2.910>.
  15. Kocić-Tanackov S, Dimić G, Mojović L, Gvozdanović-Varga J, Djukić-Vuković A, Tomović V, ... Pejin J. Antifungal activity of the onion (*Allium cepa* L.) essential oil against *Aspergillus*, *Fusarium* and *Penicillium* species isolated from food. *J Food Process Preserv*. 2017;41:e13050. <https://doi.org/10.1111/jfpp.13050>.
  16. Lanzotti V, Scala F, Bonanomi G. Compounds from *Allium* species with cytotoxic and antimicrobial activity. *Phytochemistry reviews*. 2014;13:769–91. <https://doi.org/10.1007/s11101-014-9366-0>.
  17. Li XG, Boer WD, Zhang YN, Ding CF, Zhang TL, Wang XX. Suppression of soil-borne *Fusarium* pathogens of peanut by intercropping with the medicinal herb *Atractylodes lancea*. *Soil Biol Biochem*. 2018;116:120–30. <https://doi.org/10.1016/j.soilbio.2017.09.029>.

18. Li WB, Wang XL, Li CQ, Sun JB, Li SX, Peng M (2019) Dual species transcript profiling during the interaction between banana (*Musa acuminata*) and the fungal pathogen *Fusarium oxysporum* f. sp. *cubense*. BMC Genom 20:519–34. <https://doi.org/10.1186/s12864-019-5902-z>.
19. Lian FZ, Xue RR, Lin XH, Yang YR, Zhang H, Zeng RR, Song YY. Inhibitory effects of aqueous leachates and volatiles from *Allium tuberosum* and *Ageratum conyzoides* on *Fusarium oxysporum* f. sp. *Cubense* Journal of South China Agricultural University. 2019;40:40–6. (In Chinese).
20. Liaqat A, Zahoor T, Randhawa MA, Shahid M. Characterization and antimicrobial potential of bioactive components of sonicated extract from garlic (*Allium sativum*) against foodborne pathogens. J Food Process Preserv. 2019;43:e13936. <https://doi.org/10.1111/jfpp.13936>.
21. Marrelli M, Amodeo V, Statti G, Conforti F. Biological properties and bioactive components of *Allium cepa* L.: focus on potential benefits in the treatment of obesity and related comorbidities. Molecules. 2019;24:119–36. <https://doi.org/10.3390/molecules24010119>.
22. Mnayer D, Fabiano-Tixier AS, Petitcolas E, Hamieh T, Nehme N, Ferrant C, Fernandez X, Chemat F. Chemical composition, antibacterial and antioxidant activities of six essential oils from the *Alliaceae* Family. Molecules. 2014;19:20034–53. <https://doi.org/10.3390/molecules191220034>.
23. Nandakumar R, Eyrs G, Burritt DJ, Kebede B, Leus M, Oey I. Impact of pulsed electric fields on the volatile compounds produced in whole onions (*Allium cepa* and *Allium fistulosum*). Foods. 2018;7:183–9. <https://doi.org/10.3390/foods7110183>.
24. Ozkan O, Gul S, Kart A, Cicek BA, Kilic K. *In vitro* antimutagenicity of *Allium tuncelianum* ethanol extract against induction of chromosome aberration by mutagenic agent mitomycin C. Kafkas Universitesi Veteriner Fakultesi Dergisi. 2013;19:259–62. <https://doi.org/10.9775/kvfd.2012.7637>.
25. Pattison AB, Molina AB, Chao CP, Viljoen A, Lindsay SJ. Integrating management practices to support banana production in the presence of Fusarium wilt. Acta Hort. 2018;1196:129–35. <https://doi.org/10.17660/actahortic.2018.1196.15>.
26. Paul JY, Khanna H, Kleidon J, Hoang P, Geijskes J, Daniells J, Zaplin E, Rosenberg E, James A, Mlalazi B, Deo P, Arinaitwe G, Namanya P, Becker D, Tindamanyire J, Tushemereirwe W, Harding R, Dale J. Golden bananas in the field: elevated fruit pro-vitamin A from the expression of a single banana transgene. Plant Biotechnol J. 2017;5:520–32. <https://doi.org/10.1111/pbi.12650>.
27. Ploetz RC. Management of Fusarium wilt of banana: A review with special reference to tropical race 4. Crop Prot. 2015;73:7–15. <https://doi.org/10.1016/j.cropro.2015.01.007>.
28. Schulz H, Krüger H, Liebmann J, Peterka H. Distribution of volatile sulfur compounds in an interspecific hybrid between onion (*Allium cepa* L.) and leek (*Allium porrum* L.). J Agric Food Chem. 1998;46:5220–4. <https://doi.org/10.1021/jf9806208>.
29. Segura RA, Stoorvogel JJ, Samuels JZ, Sandoval JA. Managing the interactions between soil abiotic factors to alleviate the effect of Fusarium wilt in bananas. Acta Hort. 2018;1196:163–8. <https://doi.org/10.17660/actahortic.2018.1196.19>.
30. Sharifi-Rad J, Mnayer D, Tabanelli G, Stojanović-radić DD, Sharifi-Rad M, Yousaf Z, Vallone L, Setzer WN, Iriti M. Plants of the genus *Allium* as antibacterial agents: From tradition to pharmacy. Cell Mol

- Biol (Noisy-le-grand). 2016;62:57–68. <https://doi.org/10.1016/j.foodchem.2014.10.042>.
31. Shen ZZ, Zhong ST, Wang YG, Wang BB, Mei XL, Li R, Ruan YZ, Shen QR. Induced soil microbial suppression of banana fusarium wilt disease using compost and biofertilizers to improve yield and quality. *European Journal of Soil Biology*. 2013;57:1–8. <https://doi.org/10.1016/j.ejsobi.2013.03.006>.
  32. Shen ZZ, Ruan YZ, Chao X, Zhang J, Li R, Shen QR. Rhizosphere microbial community manipulated by 2 years of consecutive biofertilizer application associated with banana *Fusarium wilt* disease suppression. *Biol Fertil Soils*. 2015;51:553–62. <https://doi.org/10.1007/s00374-015-1002-7>.
  33. Shen ZZ, Xue C, Taylor PWJ, Ou YN, Wang BB, Zhao Y, Ruan YZ, Li R, Shen QR. Soil pre-fumigation could effectively improve the disease suppressiveness of biofertilizer to banana *Fusarium wilt* disease by reshaping the soil microbiome. *Biol Fertil Soils*. 2018;54:793–806. <https://doi.org/10.1007/s00374-018-1303-8>.
  34. Shen Z, Xue C, Penton CR, Thomashow LS, Zhang N, Wang B, Shen Q. Suppression of banana Panama disease induced by soil microbiome reconstruction through an integrated agricultural strategy. *Soil Biol Biochem*. 2019;128:164–74. <https://doi.org/10.1016/j.soilbio.2018.10.016>.
  35. Singh B, Singh JP, Kaur A, Singh N. Bioactive compounds in banana and their associated health benefits - A review. *Food Chem*. 2016;206:1–11. <https://doi.org/10.1016/j.foodchem.2016.03.033>.
  36. Soni B, Tseng TMP, Yue ZM. Identification and quantification of allelochemicals from selected sweet potato (*Ipomoea batatas* (L.) Lam.) cultivars. *American Journal of Plant Sciences*. 2019;10:2354–65. <https://doi.org/10.4236/ajps.2019.1012163>.
  37. Sun L, Song S, Fu L, Deng XH, Wang DS, Liang XL, Li R, Shen QR. Exploring a soil fumigation strategy based on ammonium bicarbonate to control *Fusarium wilts* of cucurbits. *Crop Prot*. 2015;70:53–60. <https://doi.org/10.1016/j.cropro.2015.01.004>.
  38. Sun JM, Zhang JZ, Fang H, Peng L, Wei SL, Li HS, Zheng SJ, Lu J. Comparative transcriptome analysis reveals resistance-related genes and pathways in *Musa acuminata* banana 'Guijiao 9' in response to *Fusarium wilt*. *Plant Physiol Biochem*. 2019;141:83–94. <https://doi.org/10.1016/j.plaphy.2019.05.022>.
  39. Trivedi P, Delgado-Baquerizo M, Trivedi C, Hamontsb K, Anderson IC, Singhbd BK. Keystone microbial taxa regulate the invasion of a fungal pathogen in agro-ecosystems. *Soil Biol Biochem*. 2017;111:10–4. <https://doi.org/10.1016/j.soilbio.2017.03.013>.
  40. Yu HJ, Chen SC, Zhang XX, Zhou XG, Wu FZ. Rhizosphere bacterial community in watermelon-wheat intercropping was more stable than in watermelon monoculture system under *Fusarium oxysporum* f. sp. *niveum* invasion. *Plant Soil*. 2019;445:369–81. <https://doi.org/10.1007/s11104-019-04321-5>.
  41. Wang B, Li R, Ruan Y, Ou Y, Zhao Y, Shen Q. Pineapple–banana rotation reduced the amount of *Fusarium oxysporum* more than maize–banana rotation mainly through modulating fungal communities. *Soil Biol Biochem*. 2015;86:77–86. <https://doi.org/10.1016/j.soilbio.2015.02.021>.
  42. Wei YX, Liu W, Hu W, Liu GY, Wu CJ, Liu W, Zeng HQ, He CZ, Shi HT. Genome-wide analysis of autophagy-related genes in banana highlights *MaATG8s* in cell death and autophagy in immune

response to *Fusarium* wilt. *Plant Cell Report*. 2017;36:1237–50. <https://doi.org/10.1007/s00299-017-2149-5>.

43. Wibowo A, Alboneh AR, Somala MUA, Subandiyah S, Pattison T, Molina A. Increasing soil suppressivity to *Fusarium* wilt of banana through banana intercropping with *Allium* spp. *Journal Perlindungan Tanaman Indonesia*. 2015;19:33–9. <https://doi.org/10.22146/jpti.16344>.
44. Xiong W, Li R, Ren Y, Liu C, Zhao Q, Wu H, Jousset A, Shen Q. Distinct roles for soil fungal and bacterial communities associated with the suppression of vanilla *Fusarium wilt* disease. *Soil Biol Biochem*. 2017;107:198–207. <https://doi.org/10.1016/j.soilbio.2017.01.010>.
45. Xue C, Penton CR, Shen ZZ, Zhang RF, Huang QW, Li R, Ruan YZ, Shen QR (2015) Manipulating the banana rhizosphere microbiome for biological control of Panama disease. *Sci Rep* 5: e11124. <https://doi.org/10.1038/srep11124>.
46. Yu JQ. Allelopathic suppression of *pseudomonas solanacearum* infection of tomato (*Lycopersicon esculentum*) in a tomato-Chinese Chive (*Allium tuberosum*) intercropping system. *J Chem Ecol*. 1999;25:2409–17.
47. Zhang H, Mallik A, Zeng R. Control of Panama disease of banana by rotating and intercropping with Chinese Chive (*Allium tuberosum* Rottler): Role of Plant Volatiles. *J Chem Ecol*. 2013;39:243–52. <https://doi.org/10.1007/s10886-013-0243-x>.
48. Zuo GW, Li CY, Li B, Wei YR, Hu CH, Yang QS, Yang J, Sheng O, Kuang RB, Deng GM, Biswas MK, Yi GJ. The toxic mechanism and bioactive components of Chinese leek root exudates acting against *Fusarium oxysporum* f. sp. *ubense*, tropical race 4. *Eur J Plant Pathol*. 2015;143:447–60. <https://doi.org/10.1007/s10658-015-0697-5>.

## Figures






Figure 1

Inhibitory effects of volatiles released from sliced leaves (A) and the ground-leaves (B) of the six Chinese chive cultivars on colony growth of *Fusarium oxysporum* f. sp. cubense race 4 (FOC). The six cultivars are Duokang Fujiu 11 (DKF), Fujiubao (FJB), Fujiuhuang 2 (FJH), Duokang Sijiqing (DKS), Futaijiu 1 (FTJ), and Zigenchun Zaohong (ZZH). The concentration of volatiles was 0.6 g FW in the four-room petri dish (90 mm in diameter). Data are means  $\pm$  SE (n= 12). Significant differences among treatments were indicated by different letters above the bars (P=0.01 by Duncan's multiple range test).




**Figure 2**

Inhibitory effects of aqueous leachates of aerial parts of six Chinese chive on colony growth of *Fusarium oxysporum* f. sp. cubense race 4 (FOC). The six cultivars were Futaijiu 1 (FTJ), Fujiubao (FJB), Fujiuhuang 2 (FJH), Duokang Sijiqing (DKS), Duokang Fujiu 11 (DKF) and Zigenchun Zaohong (ZGC). The concentration of aqueous leachates was 0.25 g FW·mL<sup>-1</sup>. Data are means  $\pm$  SE (n=6). Significant differences among treatments were indicated by different letters above the bars (P=0.01 by Duncan's multiple range test).



**Figure 3**

Antifungal activity of three compounds identified from Chinese chive against colony growth of *Fusarium oxysporum* f. sp. *cubense* race 4 (FOC). The final concentration of the three compounds, 2-methyl-2-pentenal (MP), dimethyl trisulfide (DMT) and dipropyl trisulfide (DPT) was 25  $\mu\text{L}\cdot\text{L}^{-1}$ . Data are means  $\pm$  SE (n=20). Significant differences among treatments were indicated by different letters above the bars (P=0.01 by Duncan's multiple range test).



**Figure 4**

Contents of 2-methyl-2-pentenal (MP), dimethyl trisulfide (DMT), and dipropyl trisulfide (DPT) in the volatiles (A, C, E) and aqueous leachates (B, D, F) of the six Chinese chive cultivars. The six cultivars were Futaijiu 1 (FTJ), Fujiubao (FJB), Fujiuhuang 2 (FJH), Duokang Sijiqing (DKS), Duokang Fujiu 11 (DKF) and Zigenchun Zaohong (ZGC). Data are means  $\pm$  SE (n= 3). Significant differences among treatments were indicated by different letters above the bars (P=0.01 by Duncan's multiple range test).




Figure 5

Vegetative growth of banana plants intercropping with three Chinese chive cultivars in pot experiment. (A) Plant height, (B) stem diameter. The three cultivars used for intercropping with banana were Fujiuhuang 2 (FJH), Duokang Sijiqing (DKS) and Duokang Fujiu 11 (DKF). Data are means  $\pm$  SE (n= 10). Significant differences among treatments were indicated by different letters above the bars (P=0.01 by Duncan's multiple range test).





Figure 6

Net photosynthetic rate (Pn, A), stomatal conductance (Gs, B), intercellular CO<sub>2</sub> concentration (Ci, C), and transpiration rate (Tr, D) of bananas plants intercropping with three Chinese chives cultivars in pot experiment. The three cultivars used for intercropping with banana were Fuji Huang 2 (FJH), Duokang Sijiqing (DKS) and Duokang Fuji 11 (DKF). The photosynthetic parameters were measured 60 days after FOC inoculation. Data are means  $\pm$  SE (n= 10). Significant differences among treatments were indicated by different letters above the bars (P=0.01 by Duncan's multiple range test).



(B)




Figure 7

Fusarium wilt disease incidence (A) and growth performance (B) of banana plants 60 days after inoculation with of *Fusarium oxysporum* f. sp. cubense race 4 (FOC) in monoculture and intercropping with three Chinese chive cultivars in the pots. The three cultivars used for intercropping with banana were Fuji Huang 2 (FJH), Duokang Sijiqing (DKS) and Duokang Fujiu 11 (DKF). Data are means  $\pm$  SE (n= 10). Significant differences among treatments were indicated by different letters above the bars (P=0.01 by Duncan's multiple range test).




Figure 8

Fusarium wilt disease incidence (A), fruit weight (B) and yield (C) of banana plants 30 days after FOC inoculation in monoculture and intercropping with three Chinese chive cultivars in the field. The three cultivars used for intercropping with banana were Fujiuhuang 2 (FJH), Duokang Sijiqing (DKS) and Duokang Fujiu 11 (DKF). Data are means  $\pm$  SE (n= 5). Significant differences among treatments were indicated by different letters above the bars (P=0.01 by Duncan's multiple range test).