

Nicotine Regulates Autophagy of Human Periodontal Ligament Cells Through $\alpha 7$ nAChR That Promotes Secretion of Inflammatory Factors IL-1 β and IL-8

yang du

Department of Pediatric Dentistry, School of Stomatology, Fourth Military Medical University

Kuan Yang

Department of Orthodontics, College of Stomatology, Xi'an Jiaotong University

Zhifei Zhou

Department of Stomatology, The General Hospital of Tibetan Military Region

Lizheng Wu

Department of Stomatology, Characteristic Medical Center of People's Armed Police Force

Lulu Wang

Department of Pediatric Dentistry, School of Stomatology, Fourth Military Medical University

Yujiang Chen

Department of Pediatric Dentistry, School of Stomatology, Fourth Military Medical University

Xin Ge

Department of Pediatric Dentistry, School of Stomatology, Fourth Military Medical University

Xiaoqing Wang (✉ wxjingpd@hotmail.com)

Department of Pediatric Dentistry, School of Stomatology, Fourth Military Medical University

Research Article

Keywords: Nicotine, autophagy, $\alpha 7$ nAChR, periodontitis

Posted Date: March 11th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-258935/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Nicotine is an important risk factor and the main toxic component associated with periodontitis. However, the mechanism of nicotine induced periodontitis is not clear. To investigate the mechanism through which nicotine regulates autophagy of human periodontal ligament cells (hPDLCs) through the $\alpha 7$ nicotinic acetylcholine receptor ($\alpha 7$ nAChR) and how autophagy further regulates the release of IL-1 β and IL-8 secretion in hPDLCs.

Methods: hPDLCs were obtained from root of extracted teeth and pre-incubated in alpha-bungarotoxin (α -BTX) or 3-Methyladenine (3-MA), followed by culturing in nicotine. We used a variety of experimental detection techniques including western blotting, immunofluorescence, enzyme-linked immunosorbent assay (ELISA), transmission electron microscopy (TEM) and RT-qPCR to assess the expression of the LC3 protein, autolysosome, and release of IL-1 β and IL-8 from hPDLCs.

Results: Western blots, immunofluorescence and TEM results found that the nicotine significantly increased the autophagy expression in hPDLCs that was time and concentration dependent and reversed by α -BTX treatment ($p < 0.05$). RT-qPCR and ELISA results revealed a noticeable rise in the release of inflammatory factors IL-1 β and IL-8 from hPDLCs in response to nicotine. RT-qPCR and ELISA results showed that nicotine can significantly up-regulate the release of inflammatory factors IL-1 β and IL-8 in hPDLCs, and this effect can be inhibited by 3-MA ($p < 0.05$).

Conclusions: Nicotine regulated autophagy of hPDLCs through $\alpha 7$ nAChR and in turn the regulation of the release of inflammatory factors IL-1 β and IL-8 by hPDLCs.

Background

Periodontitis is a chronic progressive infectious disease with a high prevalence of 45%–50% overall, with the most severe form affecting 11.2% of the world's population, being the sixth most common human disease [1, 2]. Tobacco smoking has been recognized as one of the main risk factors for the occurrence and development of periodontitis [3]. Nicotine is one of the most toxic substances in tobacco [4]. It can cause pathological changes in human periodontal tissues, promote alveolar bone resorption, and ultimately lead to tooth loss [5, 6].

$\alpha 7$ nicotinic acetylcholine receptor ($\alpha 7$ nAChR) is a predominant subunit of nicotinic acetylcholine receptors (nAChRs), as well as a potent target of the nicotine binding receptor [6]. Previously, we had demonstrated the functional expression of $\alpha 7$ nAChR in periodontal ligament (PDL) tissues and human periodontal ligament cells (hPDLCs). Nicotine can enhance the expression of $\alpha 7$ nAChR in PDL tissues and hPDLCs, activate inflammation-related signaling pathways, and further regulate the secretion of inflammatory cytokines such as IL-1 β and IL-8 [6-8].

Autophagy is an essential cellular mechanism which plays “housekeeping” role in normal physiological processes including removing of long lived, aggregated and misfolded proteins, clearing damaged

organelles, growth regulation and aging [9].

Autophagy is a dynamic multi-step process that involves the formation of autophagosomes, fusion of the autophagosome with the lysosome to form the autolysosome, and finally the degradation of the contents in the autolysosome [10, 11]. The measurement of fluorescently labeled LC3 puncta and autolysosomes in cells can be used as a method to quantify autophagy [10].

Autophagy could influence the pathogenesis of various inflammatory disorders [12, 13]. Recent studies have shown evidences of autophagy in periodontal tissue, which may be involved in periodontitis [9]. For instance, the expression of autophagy-related factors was elevated in periodontitis [14]. In addition, autophagy also exerted multiple effects on hPDLCs in different conditions [15]. In current years, the interaction between nicotine and autophagy has been discussed [16]. Additionally, nicotine regulated the autophagy process via nAChRs in other cell types [17]. So, we assume that autophagy may also participate in smoking-related periodontitis via the regulation of inflammation-related signaling that leads to inflammatory disorders and periodontal tissue damage.

In this study, we investigated the mechanism which nicotine regulated autophagy of hPDLCs through the $\alpha 7$ nAChR and how autophagy further regulated the secretion of inflammatory factor IL-1 β and IL-8 in hPDLCs. Our study provides experimental evidence for the pathological development of smoking-related periodontitis and sheds a new light on smoking related periodontitis.

Methods

Isolation and culturing of hPDLCs

The current study included caries free and periodontally healthy premolars ($n = 14$) that were extracted due to the orthodontic reasons from young patients (12 to 16 years). We received approval from the institutional Ethical Review Board at the School of Stomatology, the Fourth Military Medical University, China. The children's parents/guardians were informed about the purpose of the study and inclusion of the extracted teeth in the research and provided informed consent in written format. The extracted teeth were cleaned and stored in the DMEM (Hyclone) containing 15% fetal calf serum (FCS) (Hyclone) and antibiotics (Invitrogen). In order to avoid contamination from the gingival and pulpal tissues, PDL tissues were excised from the middle third of the root using a sharp scalpel. PDL tissues were seeded in a six-well plate (Hyclone) and cultured at 37°C in a humidified atmosphere with 95% air and 5% CO₂. The medium was changed every 3 days. After achieving the confluence, the cell layers were sub-cultured and used for further experiments.

Treatment with nicotine, α -BTX and 3-MA

Nicotine (Sigma) was administrated to hPDLCs either at variable concentrations (10^{-4} , 10^{-5} , 10^{-6} , 10^{-7} mol/L) for 12 h, or a fixed concentration (10^{-5} mol/L) for a variable period of time (3, 6, 9, 12 and 24 h). In order to prove that nicotine-induced autophagy of hPDLCs was mediated by $\alpha 7$ nAChR, we pretreated

hPDLs with α -BTX (10^{-8} mol/L, $\alpha 7$ nAChR specific receptor antagonist) (Tocris Bioscience), for 30 min before nicotine (10^{-5} mol/L, 12 h) stimulation, and examined autophagy expression. To further examine whether autophagy played a role in the secretion of inflammatory factors in hPDLs, we pretreated hPDLs with 3-methyladenine (3-MA) (10^{-3} mol/L, a PI3K inhibitor that effectively blocks autophagy) (Sigma), for 30 min before nicotine (10^{-5} mol/L, 12 h) stimulation, the expression of autophagy and the secretion of IL-1 β and IL-8 were detected.

Protein isolation and Western blot

Protein extraction was performed as described previously [18]. To prepare the total cell lysate, the cells (density, 1×10^6 cells/dish) were seeded and then treated with nicotine, α -BTX, or 3-MA various agents, cell proteins were isolated using the Nuclear Extract Kit (Sangon Biotech). The BCA kit (Thermo) was used to calculate protein concentration in the solution. Proteins (40 μ g /lane) were isolated with SDS-PAGE and then transferred to a PVDF membrane (Millipore). Following blocking with 5% skimmed milk dissolved in Tris-buffered saline containing 0.1% Tween-20 at 37°C for 1 h, the membranes were probed with primary antibodies against: LC3B, Beclin-1, GAPDH (Cell Signaling Technology) at 4°C overnight. After three washes, membranes were incubated with secondary antibody for 1 h at 37°C. An Infrared Imaging System (Odyssey) was used to assess blots. Representative results from one of three independent experiments were shown.

Immunofluorescence

For immunofluorescence, the cells were fixed in 4% paraformaldehyde for 30 min, washed using the PBS (Hyclone), permeabilized at room temperature using the 0.5 % Triton X-100 and blocked with BSA (Sigma) for 2 h. Cells were then incubated with LC3B primary antibody (Cell Signaling Technology) at the 1:200 dilution. Cells were washed using PBS and incubated with the FITC-labeled secondary antibody (1:1000). Cell nuclei stained by the Hoechst (Sigma) were analyzed using a fluorescence microscope (Olympus FV1000). In this study, the method of immunofluorescence point counting was described previously [19]. Representative results from one of three independent experiments are shown.

Transmission electron microscope assays (TEM)

The TEM samples as prepared as were described previously [20]. Briefly, cells were fixed using the 2.5% glutaraldehyde in 0.1 mol/L cacodylate buffer for 30 min. Representative ultra-thin sections were analyzed using standard TEM methods. Representative results from one of three independent experiments are shown.

Real-time quantitative polymerase chain reaction (RT-qPCR)

In order to detect IL-1 β and IL-8 expression, total RNA was isolated and RT-qPCR assay were performed as described previously [21]. Briefly, we used TRIZOL reagent (Takara) to extract the total RNA according to the manufacturer's instructions and the PrimeScript RT reagent Kit (Takara) to synthesize the cDNA. All

experiments were conducted in triplicate using Mastercycler realplex (Eppendorf AG) and GAPDH as an internal standard. The following primer sequences were used: human IL-1 β forward: 5'-ATGATGGCTTATTACAGTGGCAA-3' and reverse, 5'-GTCGGAGATTCGTAGCT

GGA-3'; human IL-8 forward: 5'-TTGCCAAGGAGTGCTAAAGAA-3' and reverse, 5'-GCCCTCTTCAAAAATTCTCC-3'; GAPDH forward 5'-ACCCACTCCTCCAC

CTTTG-3' and reverse, 5'-ATCTTGTGCTCTTGCTGGG-3'. Gene expression was calculated using the $2^{-\Delta\Delta C_q}$ method [22]. Experiments were performed in triplicate.

Enzyme linked immunosorbent assay (ELISA)

Cell culture supernatants were analyzed using ELISA for detecting IL-1 β and IL-8 release as described previously [23]. The highly sensitive ELISA kits from R&D systems (Minneapolis) were used to analyze IL-1 β and IL-8 concentrations and normalized to the number of cells. The supernatants were thawed once and all assays were run at the same time. Each set of experiments was performed in triplicate.

Statistical analysis

All data from triplicate experiments was analyzed using the SPSS software (Version 21, IBM) and presented as mean \pm standard deviation (SD). Student's t-test was used to compare the control and treatment groups, and multiple comparisons were performed using one-way ANOVA. $P < 0.05$ was considered to indicate a statistically significant difference.

Results

Nicotine increased LC3 fluorescence puncta and autolysosomes in hPDLCs in vivo

According to immunofluorescence results (Figure 1A-B), nicotine (10^{-5} mol/L) treatment of hPDLCs for 3 h significantly increased LC3 fluorescence puncta ($p < 0.001$). The increase peaked at 12 h of nicotine treatment ($p < 0.0001$) and decreased after that ($p < 0.05$). Using a nicotine application length to hPDLCs for 12 h, we observed dose-dependent increases in LC3 fluorescence puncta around the nucleus (Figure 1C-D). In our previous study, it was found that the nicotine concentration was greater than 10^{-5} mol/L, and the activity of hPDLCs was significantly reduced [24]. As nicotine (10^{-5} mol/L) treatment for 12 h significantly enhanced autophagy in hPDLCs, these conditions were chosen for subsequent experiments.

In subsequent experiments, we verified the formation of autolysosomes in hPDLCs using TEM under the action of nicotine at this concentration (10^{-5} mol/L) and time (12 h). TEM results (Figure 1E-F) showed that there were more autolysosomes in nicotine-treated hPDLCs than in the control group ($p < 0.001$). The results of this part indicate that nicotine can promote the formation of autophagy in hPDLCs.

Effects of nicotine on hPDLCs autophagy were mediated through $\alpha 7$ nAChR

In order to explore the role of $\alpha 7$ nAChR in nicotine-induced autophagy of hPDLCs, we pretreated hPDLCs with α -BTX (10^{-8} mol/L, an $\alpha 7$ nAChR specific antagonist), for 30 min before nicotine (10^{-5} mol/L, 12 h) stimulation, and examined autophagy related protein and autolysosomes expression. The autophagy expression of hPDLCs was detected by immunofluorescence, Western blot and TEM. According to immunofluorescence results (Figure 2A-B), nicotine treatment of hPDLCs significantly increased LC3 fluorescence puncta ($p \leq 0.0001$). Compared with the nicotine group, the LC3 fluorescence puncta in nicotine combined with α -BTX treatment group were significantly reduced ($p \leq 0.001$). There was no significant difference in the number of LC3 fluorescent puncta between the α -BTX alone group and the control group. According to Western blot (Figure 2C-E) and TEM (Figure 2F-G) results, we detected the LC3 and Beclin-1 protein expression and autolysosomes in hPDLCs and the results were consistent with immunofluorescence. Therefore, α -BTX treatment blocked the regulation of autophagy of hPDLCs by nicotine. This finding indicates that the effects of nicotine on the autophagy of hPDLCs are mediated by $\alpha 7$ nAChR.

Effects of 3-MA on the expression of nicotine-induced autophagy of hPDLCs

To further examine the association between nicotine and autophagy in hPDLCs, we pretreated hPDLCs with 3-MA (10^{-3} mol/L, a PI3K inhibitor that effectively blocks autophagy), for 30 min before nicotine stimulation, and examined autophagy related protein and autolysosomes expression. Western blot results (Figure 3A-C) showed that nicotine significantly increased the expression of LC3 and Beclin-1 in hPDLCs. Compared with the nicotine group, LC3 and Beclin-1 protein expression in nicotine combined with 3-MA treatment group was significant decreased. We also analyzed the expression of autolysosomes by TEM (Figure 3D-E), the results also showed a similar trend, indicating that 3-MA can inhibit nicotine-induced autophagy of hPDLCs. It further illustrates the role of PI3K pathway in nicotine-induced autophagy.

Nicotine may partly depend on the $\alpha 7$ nAChR-PI3K pathway to induce hPDLC to secrete IL-1 β and IL-8

In our previous study, we had repeatedly demonstrated that nicotine induced the production of IL-1 β and IL-8 in hPDLCs through $\alpha 7$ nAChR. To further investigate whether autophagy is involved in the secretion of inflammatory cytokines in hPDLCs. RT-qPCR and ELISA were used to detect the expression of inflammatory factors in hPDLCs after nicotine and/or 3-MA treatment. RT-qPCR results indicated that nicotine upregulated IL-1 β and IL-8 mRNA expression in hPDLCs (Figure 4A-B). Compared with the nicotine group, the mRNA expression of IL-1 β and IL-8 in nicotine combined with 3-MA treatment group was significantly decreased ($p \leq 0.001$, $p \leq 0.05$). IL-1 β and IL-8 mRNA expression was not significantly different between the 3-MA alone group and the control group ($p > 0.05$). ELISA results showed that nicotine can significantly promote the secretion of IL-1 β and IL-8 from hPDLCs (Figure 4C-D), Compared with the nicotine group, the secretion of IL-1 β and IL-8 in nicotine combined with 3-MA treatment group was significantly decreased ($p \leq 0.005$, $p \leq 0.0001$). There was no significant different in the secretion of IL-1 β and IL-8 between the 3-MA treatment group and the control group ($p > 0.05$). Collectively, these results indicate that nicotine-induced production of IL-1 β and IL-8 from hPDLCs is partially dependent on the $\alpha 7$ nAChR-PI3K pathway.

Discussion

This study explored the mechanism by which nicotine regulated autophagy of hPDLCs and thereby up-regulated the release of inflammatory factors from hPDLCs. We found that nicotine significantly enhanced the autophagy of hPDLCs and secretion of IL-1 β and IL-8 through α 7 nAChR thus aggravating the inflammatory response of the periodontal tissues and periodontitis.

Nicotine is an important risk factor and the main toxic component associated with periodontitis [25, 26]. It can change the morphology and structure of periodontal tissues cells and cause pathological changes in periodontal tissues [26, 27]. In recent years, studies have found that autophagy is closely related to periodontitis [15, 28]. To investigate the role of nicotine in autophagy of hPDLCs, hPDLCs were treated with different concentrations of nicotine for different time. We found nicotine-induced hPDLCs autophagy in a dose-and time-dependent manner (Fig. 1). Consistent with this, nicotine enhanced the autophagy expression in human cancer colon cells and ocular smooth muscle cells, and the autophagy expression showed a nicotine concentration and time-dependent [29, 30]. The results of our in vivo study revealed that nicotine can up-regulate the autophagy expression of hPDLCs.

α 7 nAChR is a predominant subunit of nAChRs, as well as a potent target of the nicotine binding receptor [24]. In previous studies, we had confirmed the functional expression of α 7 nAChR in PDL tissues and hPDLCs, and nicotine can up-regulate the expression of α 7 nAChR [7, 8, 31]. Some existing studies had confirmed the complex connection between nAChRs, nicotine and autophagy in other cell types [17, 29]. Huang et al [32] reported that nicotine up-regulated the expression of α 7 nAChR and autophagy in SH-SY-5Y cells, and α 7 nAChR specific antagonist (α -BTX) can inhibit nicotine-induced autophagy. Consistent with this, in this study we observed that α -BTX inhibited nicotine-induced autophagy (Fig. 2), suggesting that α 7 nAChR is related to nicotine-induced autophagy of hPDLCs. To our knowledge, this is the first report to prove that nicotine regulates autophagy in hPDLCs via α 7 nAChR.

Autophagy plays an important role in the occurrence and development of inflammation and immune response [33]. In recent years, more and more studies in vitro and in vivo have confirmed the role of autophagy in periodontitis [9], but there is no sufficient evidence to confirm whether the role of autophagy in periodontitis is protective or pathological [34, 35]. An et al [36] reported that autophagy disorder in periodontitis was associated with protection. Increased autophagy was required to protect periodontal ligament stem cells from apoptosis in inflammatory microenvironment. Alternatively, Tsuda et al [37] revealed that the overexpression of autophagy induced cell death in gingival epithelial cell line, suggesting its pathological involvement in periodontitis. In this study, we found that the autophagy of hPDLCs was significantly enhanced after the use of nicotine, and the release of IL-1 β and IL-8 was enhanced by autophagy (Fig. 4). This finding suggests that autophagy is related to the pathogenesis of smoking-related periodontitis.

Previously, we had repeatedly demonstrated nicotine induced the production of IL-1 β and IL-8 via α 7 nAChR in hPDLCs [6, 8]. Other related studies found that nicotine up-regulated the expression of IL-8 and IL-1 β in human gingival epithelial cells through the nAChR pathway [38]. IL-1 β is not only related to

smoking-related periodontitis, but increasing IL-1 β can trigger a series of inflammatory reactions and promote bone resorption [39]. IL-8 is associated with periodontal status, the level of IL-8 in gingival crevicular fluid is valuable in detecting the inflammation of periodontal tissue [40]. It is speculated that IL-1 β and IL-8 could be potential therapeutic targets for smoking-related periodontitis.

Autophagy plays a role in determining the fate of IL-1 β and IL-8 [41, 42]. Starved macrophages during inflammasome activation have been reported to secrete IL-1 β in an autophagy-dependent manner [43]. Furthermore, intracellular IL-1 β colocalized with LC3 puncta indicating the intersection between the autophagy process and secretion of IL-1 β [44]. Autophagy was required for toll-like receptor-mediated IL-8 production in intestinal epithelial cells [45]. ATG can promote the release of IL-8 in human airway epithelial cells, contributing to neutrophilic airway inflammation in the pathogenesis of adult asthma [46]. In our research, we found that nicotine up-regulated the expression of LC3, Beclin-1 protein and the secretion of inflammatory factors IL-1 β and IL-8. This result can be inhibited by 3-MA, which is a PI3K inhibitor that can effectively block autophagy (Figs. 3 and 4). PI3K is a complex signaling system. We preliminarily confirmed that the α 7 nAChR/PI3K pathway can activate autophagy by nicotine and promote the secretion of inflammatory cytokines IL-1 β and IL-8. The specific molecular mechanism remains to be further studied.

In summary, nicotine regulated the autophagy level of hPDL cells through α 7 nAChR, and significantly enhanced the secretion of IL-1 β and IL-8, thereby aggravating the inflammatory response of hPDL cells. Although we reported interesting findings, these results were based on the response of certain cells in vitro, rather than mimicking in vivo or clinical conditions. Animal models are needed for further studies to confirm the results of this study. In addition, further research should be conducted to explore the detailed molecular mechanisms involving nicotine- α 7 nAChR-autophagy pathway and smoking-related periodontitis.

Conclusions

Nicotine regulated autophagy of hPDL cells through α 7 nAChR and in turn the regulation of the release of inflammatory factors IL-1 β and IL-8 by hPDL cells. This study provides experimental evidence for the pathological development of smoking-related periodontitis and sheds new light on developing smoking related periodontitis.

Abbreviations

hPDL cells

human periodontal ligament cells; α 7 nAChR:alpha7 nicotinic acetylcholine receptor; α -BTX:alpha-bungarotoxin; 3-MA:3-Methyladenine; ELISA:enzyme-linked immunosorbent assay; TEM:transmission electron microscopy; PDL:periodontal ligament; FCS:fetal calf serum.

Declarations

Acknowledgments

The authors gratefully acknowledge the department of Microbiology, School of Preclinical Medicine, Fourth Military Medical University.

Funding

This research was funded by the National Natural Science Foundation of China, grant numbers 81670988.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable requests.

Authors' contributions

YD, ZFZ, LZW and XJW contributed to the conception and design of the study. YD contributed to the experiments, writing and revising the manuscript. KY and LLW prepared figures 1-2. YJC and XG prepared figures 3-4. Material preparation and data collection were performed by YD. The first draft of the manuscript was written by YD and all authors added comments to previous versions of the manuscript. All authors read and approved the final manuscript.

Funding

This study was funded by the National Natural Science Foundation of China (Code No. 81670988).

Ethics approval and consent to participate

All methods were carried out in accordance with relevant guidelines and regulations. Tissue sampling protocol was approved by the institutional Ethical Review Board of the School of Stomatology, the Fourth Military Medical University, China (approval no.IRB-REV-20160360). The children's parents/guardians were informed about the purpose of the study and inclusion of the extracted teeth in the research and provided informed consent in written format.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1. Sanz M, Marco Del Castillo A, Jepsen S, Gonzalez-Juanatey JR, D'Aiuto F, Bouchard P, Chapple I, Dietrich T, Gotsman I, Graziani F *et al*: **Periodontitis and cardiovascular diseases: Consensus report.** *J Clin Periodontol* 2020, **47**(3):268-288.
2. Kassebaum NJ, Smith AGC, Bernabé E, Fleming TD, Reynolds AE, Vos T, Murray CJL, Marcenes W: **Global, Regional, and National Prevalence, Incidence, and Disability-Adjusted Life Years for Oral Conditions for 195 Countries, 1990-2015: A Systematic Analysis for the Global Burden of Diseases, Injuries, and Risk Factors.** *Journal of dental research* 2017, **96**(4):380-387.
3. Leite FRM, Nascimento GG, Baake S, Pedersen LD, Scheutz F, López R: **Impact of Smoking Cessation on Periodontitis: A Systematic Review and Meta-analysis of Prospective Longitudinal Observational and Interventional Studies.** *Nicotine & tobacco research : official journal of the Society for Research on Nicotine and Tobacco* 2019, **21**(12):1600-1608.
4. Carson SJ, Burns J: **Impact of smoking on tooth loss in adults.** *Evidence-based dentistry* 2016, **17**(3):73-74.
5. Takeuchi-Igarashi H, Kubota S, Tachibana T, Murakashi E, Takigawa M, Okabe M, Numabe Y: **Matrix remodeling response of human periodontal tissue cells toward fibrosis upon nicotine exposure.** *Odontology* 2016, **104**(1):35-43.
6. Wu LZ, Duan DM, Liu YF, Ge X, Zhou ZF, Wang XJ: **Nicotine favors osteoclastogenesis in human periodontal ligament cells co-cultured with CD4(+) T cells by upregulating IL-1 β .** *International journal of molecular medicine* 2013, **31**(4):938-942.
7. Zhou Z, Liu F, Wang L, Zhu B, Chen Y, Yu Y, Wang X: **Inflammation has synergistic effect with nicotine in periodontitis by up-regulating the expression of $\alpha 7$ nAChR via phosphorylated GSK-3 β .** *Journal of cellular and molecular medicine* 2020, **24**(4):2663-2676.
8. Wu L, Zhou Y, Zhou Z, Liu Y, Bai Y, Xing X, Wang X: **Nicotine induces the production of IL-1 β and IL-8 via the $\alpha 7$ nAChR/NF- κ B pathway in human periodontal ligament cells: an in vitro study.** *Cellular physiology and biochemistry : international journal of experimental cellular physiology, biochemistry, and pharmacology* 2014, **34**(2):423-431.
9. Greabu M, Giampieri F, Imre MM, Mohora M, Totan A, Pituru SM, Ionescu E: **Autophagy, One of the Main Steps in Periodontitis Pathogenesis and Evolution.** *Molecules (Basel, Switzerland)* 2020, **25**(18).
10. Pugsley HR: **Quantifying autophagy: Measuring LC3 puncta and autolysosome formation in cells using multispectral imaging flow cytometry.** *Methods (San Diego, Calif)* 2017, **112**:147-156.
11. Ni HM, Bockus A, Wozniak AL, Jones K, Weinman S, Yin XM, Ding WX: **Dissecting the dynamic turnover of GFP-LC3 in the autolysosome.** *Autophagy* 2011, **7**(2):188-204.
12. Mizushima N, Levine B: **Autophagy in Human Diseases.** *The New England journal of medicine* 2020, **383**(16):1564-1576.
13. Jin M, Zhang Y: **Autophagy and Inflammatory Diseases.** *Advances in experimental medicine and biology* 2020, **1207**:391-400.

14. Bullon P, Cordero MD, Quiles JL, Ramirez-Tortosa Mdel C, Gonzalez-Alonso A, Alfonsi S, García-Marín R, de Miguel M, Battino M: **Autophagy in periodontitis patients and gingival fibroblasts: unraveling the link between chronic diseases and inflammation.** *BMC Med* 2012, **10**:122.
15. Mei YM, Li L, Wang XQ, Zhang M, Zhu LF, Fu YW, Xu Y: **AGEs induces apoptosis and autophagy via reactive oxygen species in human periodontal ligament cells.** *Journal of cellular biochemistry* 2019.
16. Xing R, Cheng X, Qi Y, Tian X, Yan C, Liu D, Han Y: **Low-dose nicotine promotes autophagy of cardiomyocytes by upregulating HO-1 expression.** *Biochemical and biophysical research communications* 2020, **522**(4):1015-1021.
17. Wang Z, Liu B, Zhu J, Wang D, Wang Y: **Nicotine-mediated autophagy of vascular smooth muscle cell accelerates atherosclerosis via nAChRs/ROS/NF-κB signaling pathway.** *Atherosclerosis* 2019, **284**:1-10.
18. Hahnvajanawong C, Sahakulboonyarak T, Boonmars T, Reutrakul V, Kerdsin A, Boueroy P: **Inhibitory effect of isomorellin on cholangiocarcinoma cells via suppression of NF-κB translocation, the phosphorylated p38 MAPK pathway and MMP-2 and uPA expression.** *Experimental and therapeutic medicine* 2021, **21**(2):151.
19. Wei W, An Y, An Y, Fei D, Wang Q: **Activation of autophagy in periodontal ligament mesenchymal stem cells promotes angiogenesis in periodontitis.** *Journal of periodontology* 2018, **89**(6):718-727.
20. Zhang Y, Zhang Y, Jin XF, Zhou XH, Dong XH, Yu WT, Gao WJ: **The Role of Astragaloside IV against Cerebral Ischemia/Reperfusion Injury: Suppression of Apoptosis via Promotion of P62-LC3-Autophagy.** *Molecules (Basel, Switzerland)* 2019, **24**(9).
21. Zhang S, Li Y, Tu Y: **Lidocaine attenuates CFA-induced inflammatory pain in rats by regulating the MAPK/ERK/NF-κB signaling pathway.** *Experimental and therapeutic medicine* 2021, **21**(3):211.
22. Livak KJ, Schmittgen TD: **Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method.** *Methods (San Diego, Calif)* 2001, **25**(4):402-408.
23. Ge X, Liu YF, Wong Y, Wu LZ, Tan L, Liu F, Wang XJ: **Impact of nicotine on the interplay between human periodontal ligament cells and CD4+ T cells.** *Human & experimental toxicology* 2016, **35**(9):983-990.
24. Wu L, Yang K, Gui Y, Wang X: **Nicotine-upregulated miR-30a arrests cell cycle in G1 phase by directly targeting CCNE2 in human periodontal ligament cells.** *Biochemistry and cell biology = Biochimie et biologie cellulaire* 2020, **98**(3):354-361.
25. Isik Andrikopoulos G, Farsalinos K, Poulas K: **Electronic Nicotine Delivery Systems (ENDS) and Their Relevance in Oral Health.** *Toxics* 2019, **7**(4).
26. Garcia VG, Gualberto ECJ, Ervolino E, Nagata MJH, de Almeida JM, Theodoro LH: **aPDT for periodontitis treatment in ovariectomized rats under systemic nicotine.** *Photodiagnosis and photodynamic therapy* 2018, **22**:70-78.
27. Holliday RS, Campbell J, Preshaw PM: **Effect of nicotine on human gingival, periodontal ligament and oral epithelial cells. A systematic review of the literature.** *Journal of dentistry* 2019, **86**:81-88.

28. Zhang X, Jin Y, Wang Q, Jian F, Li M, Long H, Lai W: **Autophagy-mediated regulation patterns contribute to the alterations of the immune microenvironment in periodontitis.** *Aging* 2020, **13**(1):555-577.
29. Yan HY, Wen X, Chen LZ, Feng YT, Liu HX, Qu W, Zhao WH, Xu DQ, Ping J: **Augmented autophagy suppresses thymocytes development via Bcl10/p-p65 pathway in prenatal nicotine exposed fetal mice.** *Ecotoxicology and environmental safety* 2021, **207**:111272.
30. Pelissier-Rota MA, Pelosi L, Meresse P, Jacquier-Sarlin MR: **Nicotine-induced cellular stresses and autophagy in human cancer colon cells: A supportive effect on cell homeostasis via up-regulation of Cox-2 and PGE(2) production.** *The international journal of biochemistry & cell biology* 2015, **65**:239-256.
31. Wang XJ, Liu YF, Wang QY, Tsuruoka M, Ohta K, Wu SX, Yakushiji M, Inoue T: **Functional expression of alpha 7 nicotinic acetylcholine receptors in human periodontal ligament fibroblasts and rat periodontal tissues.** *Cell and tissue research* 2010, **340**(2):347-355.
32. Hung SY, Huang WP, Liou HC, Fu WM: **Autophagy protects neuron from Abeta-induced cytotoxicity.** *Autophagy* 2009, **5**(4):502-510.
33. Deretic V, Levine B: **Autophagy balances inflammation in innate immunity.** *Autophagy* 2018, **14**(2):243-251.
34. Yang Y, Huang Y, Li W: **Autophagy and its significance in periodontal disease.** *Journal of periodontal research* 2021, **56**(1):18-26.
35. Jiang M, Li Z, Zhu G: **The role of autophagy in the pathogenesis of periodontal disease.** *Oral diseases* 2020, **26**(2):259-269.
36. An Y, Liu W, Xue P, Zhang Y, Wang Q, Jin Y: **Increased autophagy is required to protect periodontal ligament stem cells from apoptosis in inflammatory microenvironment.** *J Clin Periodontol* 2016, **43**(7):618-625.
37. Tsuda H, Ochiai K, Suzuki N, Otsuka K: **Butyrate, a bacterial metabolite, induces apoptosis and autophagic cell death in gingival epithelial cells.** *Journal of periodontal research* 2010, **45**(5):626-634.
38. Kashiwagi Y, Yanagita M, Kojima Y, Shimabukuro Y, Murakami S: **Nicotine up-regulates IL-8 expression in human gingival epithelial cells following stimulation with IL-1 β or P. gingivalis lipopolysaccharide via nicotinic acetylcholine receptor signalling.** *Archives of oral biology* 2012, **57**(5):483-490.
39. Cheng R, Wu Z, Li M, Shao M, Hu T: **Interleukin-1 β is a potential therapeutic target for periodontitis: a narrative review.** *International journal of oral science* 2020, **12**(1):2.
40. Lagdive SS, Marawar PP, Byakod G, Lagdive SB: **Evaluation and comparison of interleukin-8 (IL-8) level in gingival crevicular fluid in health and severity of periodontal disease: a clinico-biochemical study.** *Indian journal of dental research : official publication of Indian Society for Dental Research* 2013, **24**(2):188-192.

41. Claude-Taupin A, Bissa B, Jia J, Gu Y, Deretic V: **Role of autophagy in IL-1 β export and release from cells.** *Seminars in cell & developmental biology* 2018, **83**:36-41.
42. Korhonen E, Piippo N, Hytti M, Hyttinen JMT, Kaarniranta K, Kauppinen A: **SQSTM1/p62 regulates the production of IL-8 and MCP-1 in IL-1 β -stimulated human retinal pigment epithelial cells.** *Cytokine* 2019, **116**:70-77.
43. Torp MK, Yang K, Ranheim T, Husø Lauritzen K, Alfsnes K, Vinge LE, Aukrust P, Stensløykken KO, Yndestad A, Sandanger Ø: **Mammalian Target of Rapamycin (mTOR) and the Proteasome Attenuates IL-1 β Expression in Primary Mouse Cardiac Fibroblasts.** *Frontiers in immunology* 2019, **10**:1285.
44. Cadwell K: **Crosstalk between autophagy and inflammatory signalling pathways: balancing defence and homeostasis.** *Nature reviews Immunology* 2016, **16**(11):661-675.
45. Li YY, Ishihara S, Aziz MM, Oka A, Kusunoki R, Tada Y, Yuki T, Amano Y, Ansary MU, Kinoshita Y: **Autophagy is required for toll-like receptor-mediated interleukin-8 production in intestinal epithelial cells.** *International journal of molecular medicine* 2011, **27**(3):337-344.
46. Pham DL, Kim SH, Losol P, Yang EM, Shin YS, Ye YM, Park HS: **Association of autophagy related gene polymorphisms with neutrophilic airway inflammation in adult asthma.** *The Korean journal of internal medicine* 2016, **31**(2):375-385.

Figures




Figure 1

Nicotine increased LC3 fluorescence puncta and autolysosomes in hPDLCs. A: LC3 fluorescence puncta in hPDLCs after nicotine (10⁻⁵ mol/L) treatment at time (0, 3, 6, 12, 24 h) as determined by immunofluorescence using confocal laser scanning microscopy. Scale bars represent 100 μ m. B: Bar graph showing the number of LC3 fluorescence puncta per cell for at least 10 cells per group. C: LC3 fluorescence puncta in hPDLCs after nicotine (10⁻⁴, 10⁻⁵, 10⁻⁶, 10⁻⁷ mol/L) treatment at 12 h as

determined by immunofluorescence using confocal laser scanning microscopy. Scale bars represent 100 μm . D: Bar graph showing the number of LC3 fluorescence puncta per cell for at least 10 cells per group. E: The formation of autolysosomes in hPDLCs induced by nicotine (10-5 mol/L, 12 h) was observed by TEM. Magnification, 40000 \times . Scale bar, 0.5 μm . The red arrows represent autolysosomes. F: Quantitative analysis of autolysosomes in one cell, at least 3 cells in each group for statistics. (* $p \leq 0.05$, ** $p \leq 0.005$, *** $p \leq 0.001$, **** $p \leq 0.0001$).




Figure 2

Effects of nicotine and/or α -BTX on the expression of autophagy protein and autolysosomes in hPDLCs. A: LC3 fluorescence puncta in hPDLCs after nicotine and/or α -BTX as determined by immunofluorescence using confocal laser scanning microscopy. Scale bars represent 100 μm ; B. Bar graph showing the number of LC3 fluorescence puncta per cell for at least 10 cells per group; C: LC3 and Beclin-1 protein expression in hPDLCs assessed by Western blot analysis; D: LC3 protein quantitative analysis. Data were expressed as mean \pm SD from at least three independent experiments; E: Beclin-1 protein quantitative analysis. Data were expressed as mean \pm SD from at least three independent experiments; F: TEM was used to evaluate autophagy induced by nicotine (10⁻⁵ mol/L, 12 h) and/or α -BTX (10⁻⁸ mol/L, 12.5 h). Magnification, 40000 \times . Scale bar, 0.5 μm . The red arrows represent autolysosomes; G: Quantitative analysis of autolysosomes in one cell, at least 3 cells in each group for statistics (* $p < 0.05$, ** $p < 0.005$, *** $p < 0.001$, **** $p < 0.0001$).




Figure 3

Effects of nicotine and/or 3-MA in the expression of autophagy-related markers in hPDLs. A: LC3II and Beclin-1 expression in hPDLs assessed by Western blot analysis; B: LC3II protein quantitative analysis. Data were expressed as mean \pm SD from at least three independent experiments; C: Beclin-1 protein quantitative analysis. Data were expressed as mean \pm SD from at least three independent experiments; D: TEM was used to evaluate autophagy induced by nicotine (10⁻⁵ mol/L, 12 h) and/or α -BTX (10⁻⁸ mol/L,

12.5 h). Magnification, 40000 \times . Scale bar, 0.5 μ m. The red arrows represent autolysosome; E: Quantitative analysis of autolysosomes in one cell, at least 3 cells in each group for statistics. (* $p \leq 0.05$, ** $p \leq 0.005$, *** $p \leq 0.001$).




Figure 4

Effects of nicotine and/or 3-MA on the production of IL-1 β and IL-8. A: The relative expression of IL-1 β mRNA from hPDLCS after nicotine and/or 3-MA treatment; B: The relative expression of IL-8 mRNA from

hPDLCs after nicotine and/or 3-MA treatment; C: Release of IL-1 β from hPDLCs after nicotine and/or 3-MA application; D: Release of IL-8 from hPDLCs after nicotine and/or 3-MA application. Data from three independent experiments were presented as mean \pm SD (* $p \leq 0.05$, ** $p \leq 0.005$, *** $p \leq 0.001$, **** $p \leq 0.0001$).