

# Prevalence of Middle Mesial Canal and Radix Entomolaris of Mandibular First Permanent Molars in a Western Chinese Population: An In Vivo Cone-Beam Computed Tomographic Study

**Xin Qiao**

Sichuan University West China College of Stomatology

**Hualing Zhu**

Sichuan University West China College of Stomatology

**Yujia Yan**

Sichuan University West China College of Stomatology

**Jinglin Li**

Sichuan University West China College of Stomatology

**Jiayin Ren**

Sichuan University West China College of Stomatology

**Yuan Gao**

Sichuan University West China College of Stomatology

**Ling Zou** (✉ [zouling@scu.edu.cn](mailto:zouling@scu.edu.cn))

Sichuan University West China College of Stomatology <https://orcid.org/0000-0002-6022-2911>

---

## Research article

**Keywords:** the first permanent mandibular molar, root canal system, cone-beam computed tomographic (CBCT), middle mesial canals (MMC), radix entomolaris (RE)

**Posted Date:** May 6th, 2020

**DOI:** <https://doi.org/10.21203/rs.3.rs-25911/v1>

**License:** © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

---

**Version of Record:** A version of this preprint was published on August 17th, 2020. See the published version at <https://doi.org/10.1186/s12903-020-01218-z>.

# Abstract

**Background:** The aim of this study was to investigate the prevalence of the middle mesial canal (MMC) and radix entomolaris (RE) in mandibular first permanent molars in a western Chinese population using cone-beam computed tomography (CBCT).

**Methods:** A total of 1174 CBCT images of the mandibular first molars were collected from West China Hospital of Stomatology, Sichuan University. The following information was recorded and evaluated: the detection rate and location of the MMC and RE, the curvature of the RE, the canal configuration and bilateral symmetry.

**Results:** The detection rates of the MMC and RE were 3.41% and 25.04%, respectively, as calculated by individuals, and 1.79% and 22.15%, respectively, as calculated by total teeth. The average curvature in the buccolingual (BL) orientation ( $40.63 \pm 14.39^\circ$ ) was significantly larger than that in the mesiodistal (MD) orientation ( $17.64 \pm 7.82^\circ$ ) ( $p < 0.05$ ). Of 587 patients, 71.72% (421/587) had bilateral symmetry according to the root canal morphology. The incidence of three-rooted mandibular first molars was higher in males than in females, while the incidence of two-rooted mandibular first molars was higher in females than in males.

**Conclusions:** Our results showed that the RE could be detected in almost 1/4 of the western Chinese population; thus, RE detection requires special attention and careful assessment in endodontic treatment.

## Background

It is well known that successful endodontic treatment depends on many factors, among which the anatomy of the teeth and root canal system provides the anatomical basis for endodontic treatment [1, 2]. Missing a single root canal in endodontic treatment may lead to secondary or persistent apical periodontitis [3]. Mandibular first molars, the first permanent teeth that appear in our oral cavities, which are defined as “the key of the occlusion” and have a large number of pits and fissures on the occlusal surface, demonstrate a poor self-cleaning effect. Therefore, the mandibular first molars with a high risk of caries are the most likely to undergo endodontic treatment [4-6]. Traditionally, the most common form of a mandibular first molar has been briefly depicted as a two-rooted tooth with two canals in the mesial root and one or two canals in the distal root [1]. However, there are several aberrant canal morphology in the mandibular molars, and the most common variations include an additional distolingual root and canal [7] and a third canal in the mesial root termed as the middle mesial canal (MMC) [8].

Vertucci and William [9] as well as Barker et al. [10] first reported the presence of the MMC in mandibular molars in 1974. Since then, there have been multiple studies that have reported this kind of variation in the mesial root between the mesiobuccal canal (MBC) and mesiolingual canal (MLC), named the “intermediate”, “mesiocentral”, “middle mesial”, “third mesial” or “accessory mesial” root canal [1, 11-15]. Previous studies have reported a frequency of this canal ranging from 0.26%-46.15%, and this extra root has received special attention when endodontic treatment is needed [16, 17].

The presence of mandibular first molars with an third distolingual root was noticed earlier than the presence of the MMC, which was first reported in 1844 by Carabelli et al [18]. It has been found that the additional distolingual root termed the “radix entomolaris” (RE) is basically smaller and more curved than the mesiobuccal and distobuccal roots [19, 20]. Relevant literature [7, 21-25] demonstrates that the RE detection rate has an

obvious genetic and ethnic predilection: A maximum of 3% is discovered in African populations, whereas the frequency is less than 5% in Eurasian, Indian, European or Caucasian populations. In populations with Mongolian descent such as the Chinese, Malaysian, Eskimo, and American Indian populations, the frequency is higher than 5% (even up to 40%). In the populations mentioned above, the RE is considered to be a kind of normal racial structure rather than a variation. We all know that western China is a multiple ethnic region, therefore, it is of great clinical value to study the RE detection rate in this region representative of China.

The purpose of the study was to analyze the root number and morphology, the canal number and configuration, and the location of the MMC and RE in the first mandibular molars in a western Chinese population in order to provide additional imaging anatomical data analysis for clinicians and improve the success rate of endodontic treatment. The symmetry of homonym teeth and whether sex was related to root canal variation were also recorded.

## Methods

### Patients

The CBCT data were enrolled at the department of radiology of West China Hospital of Stomatology, Sichuan University from October 2017 to March 2018, from patients who have medical treatment needs (pulpitis, periapical periodontitis, periodontitis, impacted wisdom teeth, jaw fractures, etc.) The present study included 587 patients (237 males and 350 females) and 1174 bilateral mandibular first permanent molars, and the basic information of the patient's name, gender and age was recorded. The study was approved by the Medical Ethics Committee of West China Stomatological Hospital of Sichuan University with the approval number: WCHSIRB-ST-2017-85.

The study screened teeth according to the following criteria:

- (1) No trauma or defects;
- (2) No periapical lesions;
- (3) No root canal treatment or post- or crown restoration;
- (4) No root canals with open apices, resorption or calcification;
- (5) Good-quality CBCT images.

### Radiographic techniques

CBCT images were obtained using a CBCT device (3-dimension Accuitomo, J.MORITA MFG.CORP.Kyoto Japan) with those exposure parameters: 85 kVp, 4.5 mA, 17.5 seconds scan time, with voxel size of 0.125 mm and field of view of 60\*60mm for all images. All images were produced by an experienced technician according to the manufacturer's instructions using lowest dose radiation.

# Evaluation of the images

Using the i-Dixel software (One Volume Viewer 1.5.0) to reconstruct and measure the image. An oral radiologist and an endodontist concurrently analyzed all the images to reach an agreement on the findings of these images. The root canal system of the first permanent mandibular molars was observed from the medullary cavity to the root apical layers on the coronal, sagittal and cross-section views, and the following data of the left and right sides were measured and recorded: the root number and morphology, the canal number and configuration, the prevalence of the middle mesial root canal (MMC) and the radix entomolaris (RE) root canal, and the curvature of the RE in the BL direction and in the MD direction. The methods of Schneider [26] was used to measure the angle of the RE curvature.

Using SPSS 21.0 software (SPSS, Inc, Chicago, IL, USA) for statistical analysis. Using descriptive statistics to describe the number of different roots and root canals, as well as the detection rate of root canal morphology. The number of roots and canals, root canal morphology and detection rate of the MMC and RE in different groups according to sex and their bilateral symmetries were analyzed using the Chi-square test. Further, the curvature of the RE root in the buccolingual and mesiodistal directions were compared with the paired-samples t-test. Statistically significant differences were defined at  $p < 0.05$ .

## Results

### The detection rates and locations of the MMC and RE

Cases of mandibular first molars with RE or MMC in the axial section are showed in Figure 1. The detection rates of the MMC and RE were 3.41% and 25.04%, respectively, as calculated by individuals, and 1.79% and 22.15%, respectively, as calculated by the total teeth. The mean distances between the MMC and MBC orifice and between the MMC and MLC orifice were 1.84 mm and 1.95 mm, respectively, while the mean distances between the RE canal orifice and distobuccal canal orifice and between the RE canal orifice and mesiolingual canal orifice were 4.22 mm and 4.01 mm, respectively.

### The curvature of the RE

The curvature of the RE canal was more severe in the BL orientation ( $40.63 \pm 14.39^\circ$ ) than in the MD orientation ( $17.64 \pm 7.82^\circ$ ) ( $p < 0.05$ ). Most RE canals exhibited severe curvature in the BL orientation and moderate curvature or a straight appearance in the MD orientation.

### The bilateral symmetry

The numbers of roots and root canals are listed in Table 1, and the canal configurations according to Vertucci's classification are listed in Table 2.

Table 1  
The number of roots and root canals in the mandibular first molars.

Root number	Two root canals	Three root canals	Four root canals	Five root canals	Total
Two roots	53	619	219	3	894(76.15%)
Three roots	0	44	231	5	280(23.85%)
Total	53(4.51%)	663(56.47%)	450(38.33%)	8(0.68%)	1174(100%)

Table 2  
The canal configuration based on Vertucci's classification in the mandibular first molars.

Root	Type I (1)	Type II (2-1)	Type III (1-2-1)	Type IV (2-2)	Type V (1-2)	Type VI (2-1-2)	Type III (3)
Mesial root	63(5.37%)	235(20.02%)	48(4.09%)	773(65.84%)	26(2.21%)	8(0.68%)	21(1.79%)
Distal root	918(78.19%)	41(3.49%)	12(1.02%)	180(15.33%)	20(1.70%)	3(0.26%)	0(0)

Of the 587 patients, 91.14% (535/587) had bilateral symmetry in terms of the number of roots, 84.50% (496/587) had bilateral symmetry in terms of the number of root canals, and 71.72% (421/587) had bilateral symmetry in terms of the root canal morphology. The right mandibular first molar had a higher incidence of having two roots (80.4%), while the left molar had a higher incidence of having three roots (28.1%). The detection rate of the mandibular first molar with five canals was higher on the left side (3.2%) than on the right side (1.2%).

In addition, the frequency of bilateral occurrence of the MMC was only 0.05%, with no significant differences between the right and left sides. However, the frequency of bilateral occurrence of the RE was 76.87%, and there was a statistically significant difference between the right and left sides ( $p < 0.05$ ), as it was higher on the left side.

### The correlations of these parameters with sex

The incidence of three-rooted mandibular first molars was higher in males (27.2%) than in females (21.6%), while the incidence of two-rooted mandibular first molars was higher in females (78.4%) than in males (72.8%). The incidence of five-canaled molars was higher in men (2.7%) than in women (1.9%), while the incidence of three-canaled molars was higher in women (57.1%) than in men (51.7%). There was no significant difference in the incidence of the remaining number of root canals. In addition, the prevalence of the RE and MMC was not significantly different between males and females ( $p > 0.05$ ).

## Discussion

The most common form of a mandibular first molar is the presence of two roots with three canals: two canals located in the mesial root and one canal in the distal root. There is little difference between countries and ethnic

groups [5, 27-31]. In our study, the most common root canal configuration of the mandibular first molar was type IV (65.84%) in the mesial roots and type I (78.19%) in the distal roots. The result is in accordance with the findings of most earlier studies [5, 16, 27, 29-35].

As mentioned before, the detection rates of the MMC in mandibular first molars varied among studies, ranging from 0.26%-46.15% [16, 17]. This varied detection rates can be due to the varying ethnic groups and ages as well as the study design or methods of detection. In China, this anatomical variation has been found to range from 0.8% to 22% [36-39]. In the present study, the detection rate of the MMC was only 3.41%. The mean distances between the MMC and MBC orifice and the MMC and MLC orifice were 1.84 mm and 1.95 mm, respectively. These distances were obviously longer than those detected by Versiani et al. [13], who found that the average distances were 1.35 mm and 1.34 mm, respectively. In addition, Akbarzadeh et al. [40] showed that in molars with MMC, the mean distance between the MBC and MLC orifice was 3.1 mm, while the distance was 3.7 mm in those without MMC; But in a most recent study,[41] that the mean distance between the mesial canals was 3.643 mm in teeth with MMC, and 3.818 mm in teeth without MMC, and Weinberg et al. suggested that there is no significant difference between above-mentioned distance in molars with or without MMC. But these distances are also apparently shorter than those detected in our study. Therefore, clinicians can carefully explore the MMC within a wide range between the MBC and MLC to avoid missing root canals in western Chinese population as well as in Chinese populations. In addition, the analysis of data by Nosrat et al. [8] showed that the incidence of MMC among different age groups has significant difference. They found that the incidence of MMC was 32.1% , 23.8%, 3.8% in patients  $\leq 20$  years old, 21-40 years old, and  $>40$  years old, respectively. This finding suggests that clinicians should be more careful and spend more time searching for an MMC when handling younger patients.

In the clinic, MMCs are difficult to find and treat. Versiani et al. [13] reported that the mean minor diameter of the MMC (0.16 mm) orifice was significantly smaller than that of the MBC (0.46 mm) and MLC (0.50 mm) orifices; it was always too small to detect and was also prone to root strip perforation during instrumentation. In addition, according to Akbarzadeh et al. [40], there could be an isthmus in 87% of mandibular first molars with an recognizable MMC. Due to the existence of an isthmus in most cases with an MMC, it is difficult to completely clean up the microbial biofilm attached to the isthmus, which can lead to failure in endodontic treatment[42]. Interestingly, Tahmasbi et al. [42] also found that an MMC originated from a separate apical foramen in only 2.4% of the total cases. They proposed that the omission of an MMC in mandibular molars may not be severe and would not definitely contribute to the failure of endodontic treatment in contrast to the omission of a second MB canal in maxillary molars [43].

In terms of the RE, a previous study by Zhang et al. [44] in western Chinese population, in which CBCT images of 232 mandibular first molars were detected, showed a detection rate of 30% for distal extra roots, and all of the extra roots had a type I configuration. Another study in western Chinese population [32] found that of the 558 CBCT images of mandibular first permanent molars, 24.7% exhibited a distolingual root. The results of the present study showed a detection rate of 22.15% (260/1174) for the total teeth and a detection rate of 25.04% (147/587) for individuals, which is consistent with the results described above. These results suggest that the prevalence of a RE in western Chinese populations is relatively high, and one out of every four people may have a distolingual root. Therefore, clinicians should carefully and consciously explore the presence of distolingual roots during endodontic treatment.

According to Schneider's method, Chen et al. [25] found that most REs had a more severe curvature in the BL orientation ( $36.35^{\circ} \pm 9.38^{\circ}$ ) than in the MD orientation ( $9.24^{\circ} \pm 6.10^{\circ}$ ). The present study evaluated 487 mandibular first molars with an RE using Schneider's method and showed that the angles of the RE root curvature in the BL and MD orientations were  $40.69 \pm 14.37^{\circ}$  and  $17.58 \pm 7.84^{\circ}$ , respectively; it was also found that most RE canals exhibited severe curvature in the BL orientation and exhibited moderate curvature or a straight appearance in the MD orientation, which is in agreement with previous studies. It is usually known that there is an increased risk of instrument fracture with an increasing angle of curvature [45, 46]. Attention should be taken to avoid instrument fracture or perforation for the treatment of this additional unexpected canal since they are usually short and severely curved. The present study also found that the mean inter-orifice distances from the DLC to the DBC and from the DLC to the MLC were 4.22 and 4.01 mm, respectively. These distances are higher than those reported by Zhang et al. [29], in which the distances found between the two distal canal orifices in most (65.2%) mandibular first molars with an RE were 2.5-3.5 mm. This suggests that clinicians could try to explore the presence of an orifice of the RE canal within a larger range in western Chinese populations, and a highly trapezoidal cavity may be helpful for locating the RE canal orifice [47].

It is also necessary to learn the similarity and symmetry of the number and morphology of roots and root canals between the left and right sides if bilateral mandibular first molars are to be treated. Of the 587 patients we studied, 91.14% had bilateral symmetry in terms of the number of roots, 84.50% had bilateral symmetry in terms of the number of root canals, and 71.72% had bilateral symmetry in terms of the root canal morphology (both sides had the same canal configuration according to Vertucci's classification in each root). In addition, our study showed that the right molars had a higher incidence of having two roots (80.4%), while the left molars had a higher incidence of having three roots (28.1%), which is consistent with the results of Wang et al. [32]. In addition, the frequency of bilateral occurrence of the MMC was only 0.05%, with no significant differences between the right and left sides. The frequency of bilateral occurrence of the RE was 76.87%, and there was a statistically significant difference between the right and left sides. The incidence of a RE according to the side of occurrence still has some dispute. Some studies have found a right-side predominance [24, 48, 49], whereas other investigators have reported REs seen more frequently on the left side [32, 50-52]. The results of the current study support the latter evidence. These contradictory findings might be attributed to the ethnic backgrounds, sample sizes and methods used.

## Conclusions

In western Chinese populations, there are 3 or 4 root canals in most mandibular first permanent molars. Vertucci type I canal configurations was most prevalent in the distal while type IV was the most prevalent in the mesial roots. Although the detection rate of MMC was low, the detection rate of the RE was 25%. Therefore, clinicians should be careful enough to avoid overlooking the possible existence of a RE or an MMC.

## List Of Abbreviations

MMC☐middle mesial canal☐RE☐radix entomolaris☐CBCT☐cone-beam computed tomography☐BL☐buccolingual☐MD☐mesiodistal☐MBC☐mesiobuccal canal☐MLC☐mesiolingual canal☐

## Declarations

## Acknowledgements

Not applicable.

## Authors' contributions

Xin Qiao, Yujia Yan and Ling Zou contributed to conception or design, collected the data; JingLin Li and Jiayin Ren contributed to acquisition, analysis or interpretation of data. Hualing Zhu contributed to draft the manuscript; Ling Zou and Yuan Gao critically revised manuscript. All authors read and approved the final manuscript.

## Funding

This study has no funding.

## Availability of data and materials

All the datasets used and analyzed during the current study are available from the corresponding author on reasonable request.

## Ethics approval and consent to participate

The study was approved by the Medical Ethics Committee of West China Stomatological Hospital of Sichuan University with the approval number: WCHSIRB-ST-2017-85. Consent to participate was not applicable.

## Consent for publication

Not applicable.

## Competing interests

The authors declare that they have no competing interests.

## References


1. de Pablo OV, Estevez R, Peix Sanchez M, Heilborn C, Cohenca N. Root anatomy and canal configuration of the permanent mandibular first molar: a systematic review. *J Endod.* 2010;36(12):1919-31.
2. Toru N. Better success rate for root canal therapy when treatment includes obturation short of the apex. *Evid Based Dent.* 2005;6(2):45.
3. Reuben J, Velmurugan N, Kandaswamy D. The Evaluation of Root Canal Morphology of the Mandibular First Molar in an Indian Population Using Spiral Computed Tomography Scan: An In Vitro Study. *J Endod.* 2008;34(2):212-5.
4. Skidmore AE, Bjorndal AM. Root canal morphology of the human mandibular first molar. *Oral Surg Oral Med Oral Pathol.* 1971;32(5):778-84.
5. Silva EJ, Nejaim Y, Silva AV, Haiter-Neto F, Cohenca N. Evaluation of root canal configuration of mandibular molars in a Brazilian population by using cone-beam computed tomography: an in vivo study. *J Endod.*

- 2013;39(7):849-52.
6. Baugh D, Wallace J. Middle mesial canal of the mandibular first molar: a case report and literature review. *J Endod.* 2004;30(3):185-6.
  7. Huang RY, Cheng WC, Chen CJ, Lin CD, Lai TM, Shen EC, Chiang CY, Chiu HC, Fu E. Three-dimensional analysis of the root morphology of mandibular first molars with distolingual roots. *Int Endod J.* 2010;43(6):478-84.
  8. Nosrat A, Deschenes RJ, Tordik PA, Hicks ML, Fouad AF. Middle mesial canals in mandibular molars: incidence and related factors. *J Endod.* 2015;41(1):28-32.
  9. Vertucci FJ, Williams RG. Root canal anatomy of the mandibular first molar. *J N J Dent Assoc.* 1974;45(3):27.
  10. Barker BCW, Parsons KC, Mills PR, Williams GL. Anatomy of root canals. III. permanent mandibular molars. *Aust Dent J.* 1974;19(6):408-13.
  11. Weine FS. Case report: three canals in the mesial root of a mandibular first molar. *J Endod.* 1982;8(11):517-20.
  12. Bond JL, Hartwell GR, Donnelly JC, Portell FR. Clinical management of middle mesial root canals in mandibular molars. *J Endod.* 1988;14(6):312-4.
  13. Versiani MA, Ordinola-Zapata R, Keles A, Alcin H, Bramante CM, Pécora JD, Sousa-Neto MD. Middle mesial canals in mandibular first molars: A micro-CT study in different populations. *Arch Oral Biol.* 2016;61:130-7.
  14. Navarro LF, Luzi A, García AA, García AH. Third canal in the mesial root of permanent mandibular first molars: review of the literature and presentation of 3 clinical reports and 2 in vitro studies. *Med Oral Patol Oral Cir Bucal.* 2007;12(8):E605-9.
  15. Karapinar-Kazandag M, Basrani BR, Friedman S. The Operating Microscope Enhances Detection and Negotiation of Accessory Mesial Canals in Mandibular Molars. *J Endod.* 2010;36(8):1289-94.
  16. Kim SY, Kim BS, Woo J, Kim Y. Morphology of mandibular first molars analyzed by cone-beam computed tomography in a Korean population: variations in the number of roots and canals. *J Endod.* 2013;39(12):1516-21.
  17. Azim AA, Deutsch AS, Solomon CS. Prevalence of Middle Mesial Canals in Mandibular Molars after Guided Troughing under High Magnification: An In Vivo Investigation. *J Endod.* 2015;41(2):164-8.
  18. Carabelli G. *Systematisches Handbuch der Zahnheilkunde*, 2nd edn. Vienna, Austria: Braumüller and Seidel; 1844.
  19. Chandra SS, Chandra S, Shankar P, Indira R. Prevalence of radix entomolaris in mandibular permanent first molars: a study in a South Indian population. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2011;112(3):e77-82.
  20. Abella F, Mercade M, Duran-Sindreu F, Roig M. Managing severe curvature of radix entomolaris: three-dimensional analysis with cone beam computed tomography. *Int Endod J.* 2011;44(9):876-85.
  21. Calberson FL, De Moor RJ, Deroose CA. The Radix Entomolaris and Paramolaris: Clinical Approach in Endodontics. *J Endod.* 2007;33(1):58-63.
  22. De Moor RJ, Deroose CA, Calberson FL. The radix entomolaris in mandibular first molars: an endodontic challenge. *Int Endod J.* 2004;37(11):789-99.

23. Souza-Flamini LE, Leoni GB, Chaves JF, Versiani MA, Cruz-Filho AM, Pécora JD, Sousa-Neto MD. The radix entomolaris and paramolaris: a micro-computed tomographic study of 3-rooted mandibular first molars. *J Endod.* 2014;40(10):1616-21.
24. Tu MG, Tsai CC, Jou MJ, Chen WL, Chang YF, Chen SY, Cheng HW. Prevalence of Three-rooted Mandibular First Molars among Taiwanese Individuals. *J Endod.* 2007;33(10):1163-6.
25. Chen YC, Lee YY, Pai SF, Yang SF. The morphologic characteristics of the distolingual roots of mandibular first molars in a Taiwanese population. *J Endod.* 2009;35(5):643-5.
26. Schneider SW. A comparison of canal preparations in straight and curved root canals. *Oral Surg Oral Med Oral Pathol.* 1971;32(2):271-5.
27. Plotino G, Tocci L, Grande NM, Testarelli L, Messineo D, Ciotti M, Glassman G, D'ambrosio F, Gambarini G. Symmetry of root and root canal morphology of maxillary and mandibular molars in a white population: a cone-beam computed tomography study in vivo. *J Endod.* 2013;39(12):1545-8.
28. Caputo BV, Noro Filho GA, de Andrade Salgado DMR, Moura-Netto C, Giovani EM, Costa C. Evaluation of the Root Canal Morphology of Molars by Using Cone-beam Computed Tomography in a Brazilian Population: Part I. *J Endod.* 2016;42(11):1604-7.
29. Zhang X, Xiong S, Ma Y, Han T, Chen X, Wan F, Lu Y, Yan S, Wang Y. A Cone-Beam Computed Tomographic Study on Mandibular First Molars in a Chinese Subpopulation. *PLoS One.* 2015;10(8):e0134919.
30. Demirbuga S, Sekerci AE, Dincer AN, Cayabatmaz M, Zorba YO. Use of cone-beam computed tomography to evaluate root and canal morphology of mandibular first and second molars in Turkish individuals. *Med Oral Patol Oral Cir Bucal.* 2013:e737-44.
31. Mukhaimer RH. Evaluation of Root Canal Configuration of Mandibular First Molars in a Palestinian Population by Using Cone-Beam Computed Tomography: An Ex Vivo Study. *Int Sch Res Notices.* 2014;2014:583621.
32. Wang Y, Zheng QH, Zhou XD, Tang L, Wang Q, Zheng GN, Huang DM. Evaluation of the root and canal morphology of mandibular first permanent molars in a western Chinese population by cone-beam computed tomography. *J Endod.* 2010;36(11):1786-9.
33. Nur BG, Ok E, Altunsoy M, Aglarci OS, Colak M, Gungor E. Evaluation of the root and canal morphology of mandibular permanent molars in a south-eastern Turkish population using cone-beam computed tomography. *Eur J Dent.* 2014;8(2):154-9.
34. Celikten B, Tufenkci P, Aksoy U, Kalender A, Kermeoglu F, Dabaj P, Orhan K. Cone beam CT evaluation of mandibular molar root canal morphology in a Turkish Cypriot population. *Clin Oral Investig.* 2016;20(8):2221-6.
35. Mehraban N, Moudi E, Bijani A, Zahra Sadat M. Root and Canal Morphology of Mandibular Molars in a Selected Iranian Population Using Cone-Beam Computed Tomography. *Iran Endod J.* 2017;12(2):143.
36. Yong MA. Clinical feature and treatment of the middle mesial canals of mandibular molars. *Beijing J Stom.* 2016;24(06):341-4.
37. Chen G, Yao H, Tong C. Investigation of the root canal configuration of mandibular first molars in a Taiwan Chinese population. *Int Endod J.* 2009;42(11):1044-9.
38. Lili Y, Yan Z, Shijun Z, Shuai Z, Na W, Jie X, Shue H, Zhiyuan X. Clinical application of cone beam computed tomography combined with micro-ultrasound technique in treating three mesial canals in mandibular first

- molars. *West Chin J Stomatol.* 2017;35(4):384-8.
39. Gu Y, Lu Q, Wang H, Ding Y, Wang P, Ni L. Root Canal Morphology of Permanent Three-rooted Mandibular First Molars—Part I: Pulp Floor and Root Canal System. *J Endod.* 2010;36(6):990-4.
  40. Akbarzadeh N, Aminoshariae A, Khalighinejad N, Palomo JM, Syed A, Kulild JC, Sadeghi G, Mickel A. The Association between the Anatomic Landmarks of the Pulp Chamber Floor and the Prevalence of Middle Mesial Canals in Mandibular First Molars: An In Vivo Analysis. *J Endod.* 2017;43(11):1797-801.
  41. Weinberg EM, Pereda AE, Khurana S, Lotlikar PP, Falcon C, Hirschberg C. Incidence of Middle Mesial Canals Based on Distance between Mesial Canal Orifices in Mandibular Molars: A Clinical and Cone-beam Computed Tomographic Analysis. *J Endod.* 2020;46(1):40-3.
  42. Tahmasbi M, Jalali P, Nair MK, Barghan S, Nair UP. Prevalence of Middle Mesial Canals and Isthmi in the Mesial Root of Mandibular Molars: An In Vivo Cone-beam Computed Tomographic Study. *J Endod.* 2017;43(7):1080-3.
  43. Kulid JC, Peters DD. Incidence and configuration of canal systems in the mesiobuccal root of Maxillary first and second molars. *J Endod.* 1990;16(7):311-7.
  44. Zhang R, Wang H, Tian YY, Yu X, Hu T, Dummer PM. Use of cone-beam computed tomography to evaluate root and canal morphology of mandibular molars in Chinese individuals. *Int Endod J.* 2011;44(11):990-9.
  45. Grande NM, Plotino G, Pecci R, Bedini R, Malagnino VA, Somma F. Cyclic fatigue resistance and three-dimensional analysis of instruments from two nickel–titanium rotary systems. *Int Endod J.* 2006;39(10):755-63.
  46. Zelada G, Varela P, Martín B, Bahillo JG, Magán F, Ahn S. The Effect of Rotational Speed and the Curvature of Root Canals on the Breakage of Rotary Endodontic Instruments. *J Endod.* 2002;28(7):540-2.
  47. Jang JK, Peters O, Lee W, Son SA, Park JK, Kim HC. Incidence of three roots and/or four root canals in the permanent mandibular first molars in a Korean sub-population. *Clin Oral Investig.* 2013;17(1):105-11.
  48. Tu MG, Huang HL, Hsue SS, Hsu JT, Chen SY, Jou MJ, Tsai CC. Detection of Permanent Three-rooted Mandibular First Molars by Cone-Beam Computed Tomography Imaging in Taiwanese Individuals. *J Endod.* 2009;35(4):503-7.
  49. Song JS, Kim SO, Choi BJ, Choi HJ, Son HK, Lee JH. Incidence and relationship of an additional root in the mandibular first permanent molar and primary molars. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2009;107(1):e56-60.
  50. Gulabivala K, Opananon A, Ng YL, Alavi A. Root and canal morphology of Thai mandibular molars. *Int Endod J.* 2002;35(1):56-62.
  51. Loh HS. Incidence and features of three-rooted permanent mandibular molars. *Aust Dent J.* 1990;35(5):434-7.
  52. Gulabivala K, Aung TH, Alavi A, Ng YL. Root and canal morphology of Burmese mandibular molars. *Int Endod J.* 2001;34(5):359-70.

## Figures



**Figure 1**

Cases of mandibular first molars with RE or MMC in the axial section; the white circles indicate the examined tooth; the white arrows indicate REs; the black arrows indicate MMCs. CBCT images show (A-C) mandibular first molars with RE, (D-E) mandibular first molars with MMC, (F) mandibular first molars with both RE and MMC.