Comparison between CT and RT-PCR in a cohort of symptomatic patients with suspected COVID-19 pneumonia during the outbreak peak in Italy

Giulia Besutti (✉ Giulia.Besutti@ausl.re.it)

University of Modena and Reggio Emilia. Modena. Italy

Paolo Giorgi Rossi

AUSL-IRCCS di Reggio Emilia. Reggio Emilia. Italy

Valentina Iotti

AUSL-IRCCS di Reggio Emilia. Reggio Emilia. Italy

Lucia Spaggiari

AUSL-IRCCS di Reggio Emilia. Reggio Emilia. Italy

Riccardo Bonacini

AUSL-IRCCS di Reggio Emilia. Reggio Emilia. Italy

Andrea Nitrosi

AUSL-IRCCS di Reggio Emilia. Reggio Emilia. Italy

Marta Ottone

AUSL-IRCCS di Reggio Emilia. Reggio Emilia. Italy

Efrem Bonelli

AUSL-IRCCS di Reggio Emilia. Reggio Emilia. Italy

Tommaso Fasano

AUSL-IRCCS di Reggio Emilia. Reggio Emilia. Italy

Simone Canovi

AUSL-IRCCS di Reggio Emilia. Reggio Emilia. Italy

Rossana Colla

AUSL-IRCCS di Reggio Emilia. Reggio Emilia. Italy

Marco Massari

AUSL-IRCCS di Reggio Emilia. Reggio Emilia. Italy

Ivana Lattuada

AUSL-IRCCS di Reggio Emilia. Reggio Emilia. Italy

Laura Trabucco

AUSL-IRCCS di Reggio Emilia. Reggio Emilia. Italy

Pierpaolo Pattacini

AUSL-IRCCS di Reggio Emilia. Reggio Emilia. Italy

Research Article

Keywords: COVID-19; severe acute respiratory syndrome coronavirus 2; Pneumonia; Polymerase Chain Reaction; Tomography, X-Ray Computed

Posted Date: May 4th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-26275/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.
[Read Full License](#)

Version of Record: A version of this preprint was published at European Radiology on May 4th, 2020. See the published version at <https://doi.org/10.1007/s00330-020-07050-x>.

Abstract

Objective: To assess sensitivity and specificity of CT vs RT-PCR for the diagnosis of COVID-19 pneumonia in a prospective Italian cohort of symptomatic patients during the outbreak peak.

Methods: In this cross-sectional study we included all consecutive patients who presented to the ER between March 13th and 23rd for suspected COVID-19 and underwent both CT and RT-PCR within 3 days. Using a structured report, radiologists prospectively classified CTs in highly suggestive, suggestive, and non-suggestive of COVID-19 pneumonia. Ground-glass, consolidation, and visual extension of parenchymal changes were collected. Three different RT-PCR-based reference standard definitions were used. Oxygen saturation level, CRP, LDH, and blood cell counts were collected and compared between CT/RT-PCR classes.

Results: The study included 696 patients (41.4% women; age 59 ± 15.8 years): 423/454 (93%) patients with highly suggestive CT, 97/127 (76%) with suggestive CT, and 31/115 (27%) with non-suggestive CT had positive RT-PCR. CT sensitivity ranged from 73% to 77% and from 90% to 94% for high and low positivity threshold, respectively. Specificity ranged from 79% to 84% for high positivity threshold and was about 58% for low positivity threshold. PPV remained $\geq 90\%$ in all cases. Ground-glass was more frequent in patients with positive RT-PCR in all CT classes. Blood tests were significantly associated with RT-PCR and CT classes. Leukocytes, lymphocytes, neutrophils, and platelets decreased, CRP and LDH increased from non-suggestive to suggestive CT classes.

Conclusions: During the outbreak peak, CT presented high PPV and may be considered a good reference to recognize COVID-19 patients while waiting for RT-PCR confirmation.

Introduction

The clinical spectrum of SARS-CoV-2 infection ranges from asymptomatic infection to a disease (COVID-19) that can lead to potentially fatal respiratory failure [1,2].

The outbreak of SARS-CoV-2 has rapidly spread worldwide since the end of 2019, with Italy being one of the first and most affected countries facing the epidemic out of China [3,4].

The diagnosis of COVID-19 currently relies on reverse transcription-polymerase chain reaction (RT-PCR) conducted on oropharyngeal and/or nasopharyngeal swabs. However, while false positives are conceivably rare, false negatives can occur, even in patients with pneumonia, who may have negative nasal/oropharyngeal samples but positive lower airway samples. The true clinical sensitivity of RT-PCR is thus unknown [5,6]. Moreover, the huge demand for tests is compromising their availability in some areas [5] and creating frequent delays in diagnosis confirmation, with consequences on timely treatment, isolation, and contact tracing.

Computed tomography (CT) often shows typical findings in COVID-19 pneumonia, especially bilateral patchy ground-glass opacities and consolidation, with a predominantly peripheral distribution. Crazy-paving pattern, peripheral vessel enlargement, and findings of organizing pneumonia such as reverse halo sign have also been described [7-11]. However, CT may show no abnormalities, especially early after symptom onset [12], and CT findings are not specific, significantly overlapping with other infections [13].

The role of CT in the COVID-19 epidemic is debated. Its use as a screening tool has been proposed [14,15] but is highly discouraged by major radiology societies in Western countries [10,16]. In China the diagnostic criteria for COVID-19 include chest CT as major evidence [17], while European and American societies underline the need for RT-PCR for diagnostic confirmation [10,16], even though they suggest repetition of RT-PCR in cases of suggestive CT findings in symptomatic patients [10]. Finally, in patients with respiratory symptoms such as dyspnea and desaturation, CT may help stratify disease severity and patient prognosis [10]. The Fleischner Society has recently released a multinational consensus statement suggesting adapting the use of chest imaging to different clinical scenarios according to the severity of clinical features, pre-test probability of COVID-19, and resource constraints [18].

The aim of this study was to assess the sensitivity and specificity of CT vs RT-PCR for the diagnosis of COVID-19 pneumonia in a prospective Italian cohort of symptomatic patients presenting to the emergency room (ER) for suspected COVID-19 during the outbreak peak. We also compared relevant blood tests in classes of patients with different combinations of CT and RT-PCR results.

Methods

Setting

In the Reggio Emilia province (Northern Italy, 532,000 inhabitants, six hospitals), the first case of SARS-CoV-2 infection was diagnosed on February 27, 2020. Up to March 24, there were 1200 RT-PCR-confirmed cases and the epidemic was still spreading. The study was approved by the Area Vasta Emilia Nord Ethics Committee on 04/07/2020 (protocol number 2020/0045199). Patients' written consent to publish their images was obtained, and patients' informed consent to participate in the study was obtained whenever possible, given the retrospective nature of the study.

Study design

This was a cross-sectional study assessing sensitivity and specificity of CT for COVID-19 pneumonia at two different thresholds of suspicion, using RT-PCR as the reference standard.

Study population

All consecutive patients who presented to the Reggio Emilia province ERs between March 13th and March 23rd for suspected COVID-19 and underwent both CT and RT-PCR were eligible. Subjects with a time gap between CT and RT-PCR > 3 days were excluded.

During the COVID-19 outbreak, the diagnostic protocol for these patients included nasopharyngeal and oropharyngeal swabs for RT-PCR, blood tests, chest X-rays, and CT scan in cases of suggestive X-ray findings or negative X-rays but highly suggestive clinical features. A structured CT report was introduced on March 13th.

Reference standard

Two issues hamper the measure of CT accuracy for COVID-19. Firstly, the clinical sensitivity of RT-PCR, our reference standard, although not yet quantified, is not 100% [5]. Secondly, the target condition of our index test, i.e., CT, is viral pneumonia, while RT-PCR target condition is SARS-CoV-2 infection.

While the second issue cannot be easily solved, to overcome the first problem we used different definitions of reference standard: 1) the first RT-PCR within 3 days after CT; 2) the first RT-PCR and, if negative, repeated RT-PCR tests in the following 15 days; if not repeated, the patient was considered non-COVID19; 3) as in definition 2, but RT-PCR-negative patients who were not retested were classified as COVID19 or non-COVID19 in the same proportion as patients who were actually retested in that group of CT results.

A commercial One-Step Reverse Transcription RT-PCR (GeneFinder™ COVID -19 PLUS Real Real Amp Kit) was used and RT-PCR assay was performed on an Applied Biosystems 7500 Sequence Detection System.

CT acquisition technique

CT scans were performed using one of three scanners (128-slice Somatom Definition Edge, Siemens Healthcare; 64-slice Ingenuity, Philips Healthcare; 16-slice GE Brightspeed, GE Medical System) without contrast media injection, with the patient in supine position, during end-inspiration. Scanning parameters were: tube voltage 120 KV, automatic tube current modulation, collimation width 0.625 or 1.25 mm, acquisition slice thickness 2.5 mm, and interval 1.25 mm. Images were reconstructed with a high-resolution algorithm at slice thickness 1.0/1.25 mm. Patients wore face masks, and thorough decontamination of the room was performed after each patient.

CT analysis and structured reporting

During routine reporting, each radiologist completed both the usual radiology report as well as a structured report about the probability of COVID-19 pneumonia based on CT findings (highly suggestive, suggestive, non-suggestive) (Fig. 1), the presence/absence of ground-glass opacities and consolidations, and the extension of pulmonary lesions using a visual scoring system (< 20%, 20-40%, 40-60%, and > 60% of parenchymal involvement) (Fig. 2). Swab results were unknown when reporting, so radiologists were blinded to RT-PCR. However, they were frequently informed of blood test results and of patients' clinical features.

Blood tests

When available, C-reactive protein (CRP) level, LDH, total leukocyte, lymphocyte, neutrophil, and platelet counts measured on ER admission were collected. The tests were carried out in the Hospital Clinical Laboratories with routine automated methods. Oxygen saturation level (SpO₂) was also collected for patients who had it measured before being provided with oxygen support.

These tests were included since previously associated with COVID-19 diagnosis, severity and prognosis: increased CRP reflects host inflammatory response, along with increased total leukocyte and neutrophil counts; elevated LDH concentrations may be a sign of end-organ damage; decreased platelet count may be associated with an underlying coagulopathy; lymphopenia may represent a concomitant immune dysfunction and has been associated with increased disease severity and worse prognosis; SpO₂ provides information on lung damage and functionality [19,20].

Statistical analyses

CT scan sensitivity, specificity, and positive predictive value (PPV) for COVID-19 pneumonia were computed for two different thresholds: highly suggestive only, and highly suggestive plus suggestive findings.

Accuracy measures according to the three reference standard definitions reported above were calculated with relative 95% confidence intervals (CI) computed on the exact binomial distribution.

Distribution of clinical characteristics across groups of CT and RT-PCR results is reported. Associations between clinical characteristics and CT and RT-PCR classes were measured through Pearson's chi² and Fischer's exact distribution. For blood tests, we report mean ($\pm$ SD) and median (IQR) stratified by groups of CT and RT-PCR results. Comparisons between groups were conducted with one-way ANOVA and linear regression models adjusted for sex and age.

P values are reported as continuous measures and no prefixed significance threshold was used.

We used Stata 13.0 SE (Stata Corporation, Texas, TX) software package.

Results

Population

After excluding 6 patients with a > 3-day interval between CT and RT-PCR, we included 696 patients who underwent CT and RT-PCR in the Reggio Emilia provincial ERs between March 13th and 23rd for suspected COVID-19 (Fig. 3). Of these 696 patients, 288 (41.4%) were women. Overall mean age was 59 years (SD 15.8), 58.3 years (SD 16.5) for women, and 59.5 years (SD 15.2) for men.

Overall, 454 (65%) had CT findings which were judged by the reporting radiologist as highly suggestive of COVID-19, 127 (18%) as suggestive, and 115 (17%) as non-suggestive.

Among patients with highly suggestive CT findings, 423 (93%) had positive and 31 (7%) negative RT-PCR at the first swab performed on ER admission. Of the 127 patients with suggestive CT findings, 97 (76%) had positive and 30 (24%) negative RT-PCR at the first swab, while 31/115 (27%) patients with non-suggestive CT findings had positive RT-PCR at the first swab.

The proportion of women decreased with the increase in CT suspicion, while age was slightly higher in the group with suggestive CT (Table 1).

In 11/31 patients with CT findings highly suggestive of COVID-19 pneumonia and negative RT-PCR, the swabs were repeated within 15 days; 5 of these 11 patients (45%) had a positive result (Fig 4). Among patients with CT findings suggestive of COVID-19 pneumonia and negative RT-PCR, 8/30 had a second swab, 1 of whom (12.5%) resulting positive. Finally, among 84 patients with non-suggestive CT and negative RT-PCR, 19 were retested, 6 of whom (32%) resulting positive (Fig 4).

CT findings

Ground-glass opacities were present in almost all patients with highly suggestive (453/454) and suggestive (123/127) CT findings, and in 40% (44/115) of patients with non-suggestive findings. When considering subgroups of RT-PCR results, in all the three CT classes ground-glass opacities were more frequent in patients with positive RT-PCR.

Consolidation was present in 279/454 (61.5%) patients with highly suggestive, 76/115 (59.8%) patients with suggestive, and 41/115 (35.7%) patients with non-suggestive CT findings. The distribution of consolidation among subgroups of positive and negative RT-PCR patients in the three CT classes varied, being more frequent in negative RT-PCR patients in the intermediate CT class, i.e., suggestive.

A limited (< 20%) extension of pulmonary lesions was more frequently present in patients with suggestive and non-suggestive CT findings. Parenchymal extension was not estimated in 55/115 patients with non-suggestive findings; 41/55 had a completely normal CT scan, while 14/55 presented pleural effusion without parenchymal involvement. Interestingly, among the 31 patients with non-suggestive CT and positive RT-PCR, 13 patients had no CT abnormalities and 1 had only pleural effusion (Table 2).

CT diagnostic accuracy

The sensitivity, specificity, and PPV of CT at two different thresholds (highly suggestive findings only and highly suggestive plus suggestive findings) with respect to the three different reference standards described in the Methods section are reported in Table 3.

Estimation of RT-PCR sensitivity in patients with pneumonia

Our study can also provide a plausible estimate of RT-PCR sensitivity for SARS-CoV-2 in patients with pneumonia. In the hypothesis that all patients with highly suspicious CT have COVID-19 pneumonia, there would be 31 RT-PCR false negatives in this group, for a sensitivity of 93% (423/451). If we only

consider RT-PCR false negatives the proportion of patients who retested positive, false negatives would be approximatively 14 out of 451, for a sensitivity of 97%.

Blood tests

Means (SD) and medians (IQR) across CT classes are reported in Table 4.

Among RT-PCR-negative patients, the values of total leukocyte, lymphocyte, neutrophil, and platelet counts, and CRP level were higher than among RT-PCR-positive patients. Total leukocyte, lymphocyte, and platelets counts decreased, whereas CRP and LDH increased from non-suggestive to suggestive and/or highly suggestive CT. Patients experienced similar SpO2 values across CT and RT-PCR groups when the CT was suggestive or highly suggestive, while in patients with non-suggestive, CT SpO2 was higher, particularly for those who tested positive at RT-PCR.

Discussion

In a large sample of consecutive patients presenting to the ER for suspected pneumonia during the peak of the SARS-CoV-2 outbreak in Northern Italy, we estimated CT sensitivity for COVID-19 pneumonia to be between 73% and 77% when adopting a high positivity threshold, which corresponded to a specificity of between 79% and 84%. When adopting a lower positivity threshold, CT sensitivity was between 90% and 94%, but specificity decreased to 58%. Nevertheless, given the very high prevalence of COVID-19 during the epidemic peak, the proportion of patients with SARS-CoV-2 infection among the CT-positive patients (PPV) was always equal to or higher than 90%, whatever the positivity threshold adopted.

Despite the fact that none of the included blood tests could definitely guide diagnosis, they were significantly associated with RT-PCR and CT classes. Leukocyte, lymphocyte, neutrophil, and platelet counts decreased, and LDH increased both in RT-PCR-positive patients and in highly suggestive and/or suggestive CT scans. CRP decreased in RT-PCR-positive compared to RT-PCR-negative patients and increased in highly suggestive and suggestive CT scans. SpO2 was slightly lower in patients with suggestive and highly suggestive CT results.

A recent study has assessed CT diagnostic performance in Italy by using only one threshold (positive versus negative CT scan) and two RT-PCR within 24 hours as the reference standard in 158 patients. Results were similar to those obtained in our study with lower positivity threshold: sensitivity and specificity of 97% (95% CI, 88-99%) and 56% (95% CI, 45-66%), respectively [21].

In the largest study conducted in China on over 1000 patients, CT scan presented similar sensitivity (97%) but lower specificity and PPV (25% and 65%, respectively) [22]. Given the lower RT-PCR positive rate (59.2%) and high rate of RT-PCR-negative CT-positive patients who were eventually classified as probable COVID-19 according to their global clinical course (81%), it is plausible to think that the RT-PCR false negative rate was higher in the study by Ai et al., leading to underestimating CT specificity and PPV.

Accordingly, in our study, when applying different reference standards so as to reduce RT-PCR false negative rates, specificity rose to 84% and PPV to 96% at the higher CT suspicion level.

Furthermore, in China, CT scan has been proposed and is used as a screening tool in asymptomatic or mildly symptomatic patients [14,23], who are more likely to have low viral loads and false negative RT-PCR results [24]. Thus, the combination of an increase in clinical sensitivity of the reference standard, the application of CT only to symptomatic cases presenting in ER, and a greater knowledge of COVID-19 CT findings in the radiology community thanks to the pioneering studies from China may explain the higher CT specificity estimate in our study.

It is not surprising that in the outbreak peak phase, with very high disease incidence, the proportion of RT-PCR positive among non-suspicious CT was quite high, as the patient has not yet developed pneumonia or will never develop pneumonia even if infected. In fact, almost one half of these patients in our study had no CT abnormalities.

Limitations and strengths

The main limitation of this study, in common with all the others with a similar aim, is that CT and RT-PCR target two different conditions, e.g., COVID-19 pneumonia and SARS-CoV-2 infection; using one as the reference standard of the other thus introduces methodological challenges. Any RT-PCR false negative will result in an overestimation of CT false positives. Nevertheless, our estimate of RT-PCR sensitivity in patients with pneumonia ranges between 93% and 97%. Several biases can affect this estimate because retesting of RT-PCR-negative CT-positive cases was not systematic in our study.

Radiologists were not blinded to clinical presentation and blood tests, which is common in real practice, but it should be considered if the results were applied to settings where this information is not yet available. Also, as a consequence of how the reporting was structured and of the need to be rapid and concise during the epidemic phase of the disease, we collected only some of the CT findings of COVID-19. This approach is surely less accurate than the retrospective review performed by experienced thoracic radiologists adopted by most studies, but it is more representative of the real-life diagnostic process.

Implications for practice

The results of this study are not intended to produce evidence to be generalized to all clinical scenarios. Nevertheless, we show high sensitivity and high positive predictive value of CT for COVID-19 pneumonia in the epidemic setting.

The specific phase during which the study was conducted reflects one of the scenarios proposed by the Fleischner Society [18], characterized by a high number of symptomatic patients presenting to the ERs, high pre-test probability, and unavailability of rapid virological testing.

Further studies are needed to assess specificity and PPV in lower prevalence settings. Also, a structured report for suspected COVID-19 patients may help in monitoring the proportion of positive results and the

PPV of different positivity thresholds in different phases of the epidemic. Moreover, including CT from the pre-epidemic period [13] may help in assessing specificity to be projected in periods when COVID-19 could cyclically shift from endemic to epidemic phases.

Conclusions

During the outbreak peak, CT presented a high PPV and may thus be considered a good reference to help clinicians to recognize and triage COVID-19 patients while waiting for RT-PCR diagnostic confirmation. Our results also confirm that in case of negative RT-PCR and highly suggestive CT findings, RT-PCR should be repeated; the patient should remain isolated, given the high probability of RT-PCR false negatives in this group.

Abbreviations

SARS-CoV-2: severe acute respiratory syndrome coronavirus-2;

COVID-19: coronavirus disease 2019;

CT: computed tomography;

RT-PCR: reverse transcription-polymerase chain reaction;

CRP: C-Reactive Protein;

LDH: Lactate dehydrogenase;

PPV: positive predictive value;

SD: standard deviation;

IQR: interquartile range.

Declarations

Acknowledgment: We thank Jacqueline Costa for the English language editing.

Competing interests: The authors declare no competing interests.

Funding: The authors received no external funding for this research.

References

1. Guan WJ, Ni ZY, Hu Y, et al; China Medical Treatment Expert Group for Covid-19. Clinical Characteristics of Coronavirus Disease 2019 in China. N Engl J Med. 2020 Feb 28. doi:

10.1056/NEJMoa2002032.

2. Wu Z, McGoogan JM. Characteristics of and Important Lessons From the Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72 314 Cases From the Chinese Center for Disease Control and Prevention. *JAMA*. 2020 Feb 24. doi: 10.1001/jama.2020.2648.
3. Grasselli G, Zangrillo A, Zanella A, et al; COVID-19 Lombardy ICU Network. Baseline Characteristics and Outcomes of 1591 Patients Infected With SARS-CoV-2 Admitted to ICUs of the Lombardy Region, Italy. *JAMA*. 2020 Apr 6. doi: 10.1001/jama.2020.5394.
4. Saglietto A, D'Ascenzo F, Zoccai GB, De Ferrari GM. COVID-19 in Europe: the Italian lesson. *Lancet*. 2020;s0140-6736: 30690–5. doi:10.1016/S0140-6736(20)30690-5.
5. Patel R, Babady E, Theel ES, et al. Report from the American Society for Microbiology COVID-19 International Summit, 23 March 2020: Value of Diagnostic Testing for SARS-CoV-2/COVID-19. *mBio*. 2020 Mar 26;11(2). pii: e00722-20. doi:10.1128/mBio.00722-20. PubMed PMID: 32217609.
6. Winichakoon P, Chaiwarith R, Liwsrisakun C, et al. Negative nasopharyngeal and oropharyngeal swab does not rule out COVID-19. *J Clin Microbiol*. 26 February 2020. doi:10.1128/JCM.00297-20.
7. Wu J, Wu X, Zeng W et al. Chest CT Findings in Patients with Corona Virus Disease 2019 and its Relationship with Clinical Features. *Invest Radiol* 2020. doi:10.1097/RLI.0000000000000670.
8. Zhao W, Zhong Z, Xie X, et al. Relation Between Chest CT Findings and Clinical Conditions of Coronavirus Disease (COVID-19) Pneumonia: A Multicenter Study. *AJR Am J Roentgenol*. 2020 Mar 3:1-6. doi: 10.2214/AJR.20.22976.
9. Pan Y, Guan H, Zhou et al. Initial CT findings and temporal changes in patients with the novel coronavirus pneumonia (2019-nCoV): a study of 63 patients in Wuhan, China. *Eur Radiol*. 2020 Feb 13. doi: 10.1007/s00330-020-06731-x.
10. European Society of Radiology. COVID-19 patients and the Radiology Department – advice from ESR and ESTI. April, 2020. Available via <https://www.myesr.org/article/3209> (Accessed April 17, 2020).
11. Chen X, Tang Y, Mo Y, et al. A diagnostic model for coronavirus disease 2019 (COVID-19) based on radiological semantic and clinical features: a multi-center study. *Eur Radiol*. 2020 Apr 16. doi: 10.1007/s00330-020-06829-2.
12. Bernheim A, Mei X, Huang M et al. Chest CT Findings in Coronavirus Disease-19 (COVID-19): Relationship to Duration of Infection. *Radiology*. 2020 Feb 20:200463. doi: 10.1148/radiol.2020200463.
13. Bai HX, Hsieh B, Xiong Z, et al. Performance of radiologists in differentiating COVID-19 from viral pneumonia on chest CT. *Radiology*. 2020 Mar 10:200823. doi: 10.1148/radiol.2020200823.
14. Li K, Fang Y, Li W, et al. CT image visual quantitative evaluation and clinical classification of coronavirus disease (COVID-19). *Eur Radiol*. 2020 Mar 25. doi: 10.1007/s00330-020-06817-6.
15. Lee EYP, Ng MY, Khong PL. COVID-19 pneumonia: what has CT taught us? *Lancet Infect Dis*. 2020 Apr;20(4):384-385.

16. American College of Radiology. ACR recommendations for the use of chest radiography and computed tomography (CT) for suspected COVID-19 infection. March 11, 2020. Available via <https://www.acr.org/Advocacy-and-Economics/ACR-Position-Statements/Recommendations-for-Chest-Radiography-and-CT-for-Suspected-COVID19-Infection> (Accessed April 15, 2020).

17. Zu ZY, Jiang MD, Xu PP, et al. Coronavirus Disease 2019 (COVID-19): A Perspective from China. Radiology. 2020 Feb 21:200490. doi:10.1148/radiol.2020200490.

18. Rubin GD, Ryerson CJ, Haramati L, et al. The Role of Chest Imaging in Patient Management during the COVID-19 Pandemic: A Multinational Consensus Statement from the Fleischner Society. Radiology. 2020 Apr 7:201365. doi: 10.1148/radiol.2020201365.

19. Favaloro EJ, Lippi G. Recommendations for Minimal Laboratory Testing Panels in Patients with COVID-19: Potential for Prognostic Monitoring. Semin Thromb Hemost. 2020 Apr 12. doi: 10.1055/s-0040-1709498.

20. Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. Lancet. 2020 Mar 28;395(10229):1054-1062. doi: 10.1016/S0140-6736(20)30566-3.

21. Caruso D, Zerunian M, Polici M, et al. Chest CT Features of COVID-19 in Rome, Italy. Radiology. 2020 Apr 3:201237. doi: 10.1148/radiol.2020201237.

22. Ai T, Yang Z, Hou H, et al. Correlation of Chest CT and RT-PCR Testing in Coronavirus Disease 2019 (COVID-19) in China: A Report of 1014 Cases. Radiology. 2020 Feb 26:200642. doi: 10.1148/radiol.2020200642.

23. Fang Y, Zhang H, Xie J, et al. Sensitivity of Chest CT for COVID-19: Comparison to RT-PCR. Radiology. 2020 Feb 19:200432. doi: 10.1148/radiol.2020200432.

24. Lippi G, Simundic AM, Plebani M. Potential preanalytical and analytical vulnerabilities in the laboratory diagnosis of coronavirus disease 2019 (COVID-19). Clin Chem Lab Med. 2020 Mar 16. doi:10.1515/cclm-2020-0285.

Tables

Table 1: Age and sex in CT classes

		Highly suggestive CT		Suggestive CT		Non-suggestive CT		<i>P</i> *
		Positive RT-PCR (n=423)	Negative RT-PCR (n=31)	Positive RT-PCR (n=97)	Negative RT-PCR (n=30)	Positive RT-PCR (n=31)	Negative RT-PCR (n=84)	
		n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	
Age < 60 years	393 (56.5)	253 (59.8)	18 (58.1)	37 (38.1)	15 (50.0)	22 (71.0)	48 (57.1)	0.003
Age ≥ 60 years	303 (43.5)	170 (40.2)	13 (41.9)	60 (61.9)	15 (50.0)	9 (29.0)	36 (42.9)	
Male	408 (58.6)	261 (61.7)	20 (64.5)	55 (56.7)	17 (56.7)	16 (51.6)	39 (46.4)	0.152
Female	288 (41.4)	162 (38.3)	11 (35.5)	42 (43.3)	13 (43.3)	15 (48.4)	45 (53.6)	

Proportions by age and sex across CT and RT-PCR classes [n (% col)]. *Pearson's chi-squared *P* value for the hypothesis of independence in the two-way table.

Table 2: CT findings in CT classes

	Highly suggestive CT		Suggestive CT		Non-suggestive CT		<i>P</i> *
	Positive RT-PCR (n=423)	Negative RT-PCR (n=31)	Positive RT-PCR (n=97)	Negative RT-PCR (n=30)	Positive RT-PCR (n=31)	Negative RT-PCR (n=84)	
	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	
<i>Ground-glass</i>	423 (100)	30 (96.8)	95 (97.9)	28 (93.3)	14 (45.2)	30 (35.7)	<0.001
<i>Consolidation</i>	259 (61.2)	20 (64.5)	55 (56.7)	21 (70.0)	12 (38.7)	29 (34.5)	<0.001
<i>Extension:</i>							<0.001
< 20	138 (32.6)	11 (35.5)	66 (68.0)	15 (50.0)	14 (45.2)	39 (46.4)	
20-40	200 (47.3)	11 (35.5)	24 (24.7)	8 (26.7)	2 (6.5)	2 (2.4)	
40-60	59 (14.0)	5 (16.1)	1 (1.0)	3 (10.0)	0 (0)	0 (0)	
> 60	26 (6.2)	4 (12.9)	6 (6.2)	4 (13.3)	1 (3.2)	2 (2.4)	
Missing	0 (0)	0 (0)	0 (0)	0 (0)	14 (45.2)	41 (28.2)	

Proportions of CT findings across CT and RT-PCR classes [n (% col)]. *Fisher's exact test and *P* value for the hypothesis of independence in the two-way table.

Table 3: CT diagnostic accuracy

					Highly suggestive CT findings		
	TP	FP	TN	FN	Sensitivity (95% CI)	Specificity (95% CI)	PPV (95% CI)
Reference standard 1	423	31	114	128	76.77% (73.0%-80.2%)	78.62% (71.0%-85.0%)	93.17% (90.4%- 95.3%)
Reference standard 2	428	26	107	135	76.02% (72.3%-79.5%)	80.45% (72.7%-86.8%)	94.27% (91.7%-96.2%)
Reference standard 3	438	16	84	158	73.49% (69.8%-77.0%)	84.00% (75.3%-90.6%)	96.48% (94.3%-98.0%)
					Highly suggestive + suggestive CT findings		
	TP	FP	TN	FN	Sensitivity (95% CI)	Specificity (95% CI)	PPV (95% CI)
Reference standard 1	520	61	84	31	94.37% (92.1%-96.1%)	57.93% (49.5%-66.1%)	89.50% (86.7%-91.9%)
Reference standard 2	526	55	78	37	93.43% (91.1%-95.3%)	58.65% (49.8%-67.1%)	90.53% (87.9%-92.8%)
Reference standard 3	539	42	58	57	90.44% (87.8%-92.7%)	58.0% (47.7%-67.8%)	92.77% (90.4%-94.7%)

Sensitivity, specificity, and positive predictive value (PPV) with respective 95% confidence intervals (CI) for two levels of CT-based probability of COVID-19 pneumonia (highly suggestive only and highly suggestive plus suggestive findings), and for three reference standards: 1) the first RT-PCR; 2) plus repeated RT-PCR tests (within 15 days) in patients with negative first RT-PCR; if RT-PCR was not repeated the patient was considered negative; 3) as in the second scenario, but negative patients who were not retested with RT-PCR were classified as positive or negative in the same proportion of patients who were actually retested in each group of CT-based probability of COVID-19.

Table 4: blood tests in CT classes

	Highly suggestive CT		Suggestive CT		Non-suggestive CT		P ANOVA	P adj*
	Positive	Negative	Positive	Negative	Positive	Negative		
	RT-PCR (n=423)	RT-PCR (n=31)	RT-PCR (n=97)	RT-PCR (n=30)	RT-PCR (n=31)	RT-PCR (n=84)		
Total leukocytes (*10⁹/L)							< 0.001	< 0.001
Mean (SD)	5.4 (2.0)	8.4 (6.1)	5.3 (2.1)	8.5 (2.8)	6.1 (2.6)	9.3 (3.6)		
Median (IQR)	5.0 (4.0-6.5)	7.1 (5.0-8.7)	5.0 (3.8-6.4)	8.2 (6.2-10.9)	6.0 (3.8-7.3)	8.4 (6.9-11.0)		
Lymphocytes (*10⁹/L)							< 0.001	< 0.001
Mean (SD)	1.2 (0.5)	1.7 (1.4)	1.2 (0.6)	1.6 (1.6)	1.5 (0.6)	1.8 (0.8)		
Median (IQR)	1.1(0.8-1.4)	1.3 (1.0-1.7)	1.1 (0.8-1.6)	1.3 (1.0-1.6)	1.5 (1.0-1.9)	1.7 (1.2-2.5)		
Neutrophils (*10⁹/L)							<0.001	<0.001
Mean (SD)	3.9 (2.1)	6.0 (6.2)	3.6 (2.0)	6.2 (3.1)	3.9 (1.9)	6.7 (3.3)		
Median (IQR)	3.5 (2.6-4.7)	4.0 (2.7-5.7)	3.1 (2.3-4.3)	5.2 (3.8-9.5)	3.7 (2.9-4.9)	5.7 (4.3-8.4)		
Platelets (*10⁹/L)							< 0.001	< 0.001
Mean (SD)	193.1 (72.0)	241.6 (119.2)	192.2 (62.0)	220.3 (57.6)	223.8 (76.3)	257.4 (92.3)		
Median (IQR)	182.0 (148.0-223.0)	207.0 (181.0-269.5)	180 (148.0-221.0)	220.5 (193.0-253.0)	197.0 (156.0-273.0)	242.0 (200.0-296.0)		
CRP (mg/dL)							0.004	0.003
Mean (SD)	5.1 (5.2)	5.9 (5.6)	3.5 (4.6)	5.8 (6.9)	3.1 (6.0)	3.5 (4.8)		
Median (IQR)	3.2 (1.5-6.9)	5.2 (1.3-8.8)	1.9 (0.7-3.5)	3.7 (0.3-8.6)	0.6 (0.1-3.0)	1.3 (0.2-4.1)		
LDH (U/L)							0.003	< 0.001
Mean (SD)	525.4 (270.3)	469.5 (204.6)	456.8 (137.0)	418.0 (186.8)	401.5 (118.4)	423.3 (334.8)		

Median (IQR)	489.0 (394-606)	431.0 (366-553)	433.0 (372.6-520.0)	381.0 (320.0-484.0)	371.9 (351.0-459.5)	357.0 (299.0-441.0)		
SpO2 (%)							< 0.001	< 0.001
Mean (SD)	95.2 (3.2)	94.9 (3.5)	95.0 (3.7)	95.2 (3.3)	97.1 (2.1)	96.5 (2.5)		
Median (IQR)	95.7 (94.3-96.9)	95.6 (93.7-97.5)	95.9 (93.8-97.2)	95.3 (92.4-98.1)	97.8 (96.0-98.7)	96.9 (95.3-98.2)		

Blood tests expressed in mean (SD) and median (IQR) according to CT and RT-PCR classes. *P* adj*: *P* value for coef. of category in a linear regression adjusted for sex and age (< 60, ≥ 60). CRP: C-Reactive Protein. LDH: Lactate dehydrogenase. SpO2: oxygen saturation level.

Figures

Figure 1

Exemplification of classification of CT findings: a) CT findings highly suggestive of COVID-19 pneumonia, with bilateral interstitial involvement, patchy ground-glass opacities (arrow) and peripheral consolidations (*), confirmed by positive RT-PCR; b) CT findings suggestive of COVID-19 pneumonia, with unilateral peripheral consolidation and subtle ground-glass opacities, confirmed by positive RT-PCR; c) CT

findings non-suggestive of COVID-19 pneumonia, with mostly unilateral bronchial wall thickening, endobronchial secretions, tree-in-bud nodules, and consolidation, confirmed by negative RT-PCR.

Figure 2

Visual scoring system used to classify the extension of parenchymal involvement: < 20% (a), 20-40% (b), 40-60% (c), and > 60% (d).

Figure 3

Flowchart representing the study population and the subgroups with different combinations of CT and RT-PCR results.

Figure 4

Examples of discordant cases between CT and RT-PCR. a) focal polygonal consolidation without ground-glass opacity, considered as non-suggestive at CT scan but resulting in positive RT-PCR test. b) bilateral (mostly right) patchy ground-glass opacities, with small areas of consolidation, which was classified as highly suggestive, resulting in a first negative RT-PCR, followed by a positive RT-PCR performed 7 days later.