

Clinical Characteristics and Cardiovascular Implications of 116 Hospitalized Patients with COVID-19 in Wuhan, China: A single-Centered, Retrospective, Observational Study

Shiqiang Xiong

the General Hospital of Central Theater Command

Lin Liu

the General Hospital of Central Theater Command

Feng Lin

the General Hospital of Central Theater Command

Jinhu Shi

the General Hospital of Central Theater Command

Lei Han

the General Hospital of Central Theater Command

Huijian Liu

the general hospital of central theater command

Lewei He

the general hospital of central theater command

Qijun Jiang

the general hospital of central theater command

Zeyang Wang

the general hospital of central theater command

Wenbo Fu

the general hospital of central theater command

Zhigang Li

the general hospital of central theater command

Qing Lu

the general hospital of central theater command

Zhinan Chen

the general hospital of central theater command

Shifang Ding (✉ dfsmdwh@163.com)

<https://orcid.org/0000-0002-9129-9890>

Research article

Keywords: SARS-CoV-2, COVID-19, cardiovascular disease, hypertension, coronary heart disease

Posted Date: May 6th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-26358/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on October 22nd, 2020. See the published version at <https://doi.org/10.1186/s12879-020-05452-2>.

Abstract

Background Corona Virus Disease 2019 (COVID-19) caused by 2019 novel coronavirus (SARS-CoV-2), has become a global pandemic. Aged population with cardiovascular diseases are more likely to be infected with SARS-CoV-2 and result in more severe outcomes and elevated case-fatality rate. Meanwhile, cardiovascular diseases have a high prevalence in the middle-aged and elderly population. However, despite of several researches in COVID-19, cardiovascular implications related to it still remains largely unclear.

Methods 116 patients with laboratory-confirmed COVID-19 were enrolled, who admitted to the General Hospital of Central Theater Command (Wuhan, China) from January 20 to March 8, 2020. The demographic data, underlying comorbidities, clinical symptoms and signs, laboratory findings, chest computed tomography, treatment measures, and outcome data were collected from electronic medical records. Data were compared between non-severe and severe cases.

Results Of 116 hospitalized patients with COVID-19, the median age was 58.5 years (IQR, 47.0-69.0), and 36 (31.0%) were female. Hypertension (45 [38.8%]), diabetes (19 [16.4%]), and coronary heart disease (17 [14.7%]) were the most common coexisting conditions. Common symptoms included fever [99 (85.3%)], dry cough (61 [52.6%]), fatigue (60 [51.7%]), dyspnea (52 [44.8%]), anorexia (50 [43.1%]), and chest discomfort (50 [43.1%]). Lymphopenia (lymphocyte count, $1.0 \times 10^9/L$ [IQR, 0.7-1.3]) was observed in 66 patients (56.9%), and elevated lactate dehydrogenase (245.5 U/L [IQR, 194.3-319.8]) in 69 patients (59.5%). Compared with non-severe cases, severe cases were older (64.0 years [IQR, 53.0-76.0] vs 56.0 years [IQR, 37.0-64.0]), more likely to have comorbidities (35 [63.6%] vs 24 [39.3%]), and more likely to develop acute cardiac injury (19 [34.5%] vs 4 [6.6%]), acute heart failure (18 [32.7%] vs 3 [4.9%]), and ARDS (20 [36.4%] vs 0 [0%]). During hospitalization, the prevalence of new onset hypertension was significantly higher in severe patients (55.2% vs 19.0%) than in non-severe ones.

Conclusions We found that the infection of SARS-CoV-2 was more likely to occur in aged population with cardiovascular comorbidities. Cardiovascular complications, including new onset hypertension and heart injury were common in severe patients. More detailed researches in cardiovascular involvement in COVID-19 are urgently needed to further understand the disease.

Background

In early December 2019, a cluster of acute respiratory illness, now known as Corona Virus Disease 2019 (COVID-19) caused by 2019 novel coronavirus (SARS-CoV-2), occurred in Wuhan, Hubei Province, China [1–4]. COVID-19 has rapidly spread all over the world and become a global pandemic. Although several recent studies have described the clinical spectrum of COVID-19, including general epidemiological characteristics, clinical manifestations, and clinical outcomes of patients [4–7], cardiovascular implications of COVID-19 remain largely unclear.

Cardiovascular diseases have a high prevalence in middle-aged and elderly population.[8] aged population with comorbidities, such as cardiovascular diseases are more susceptible to COVID-19 and result in severe outcomes and elevated case-fatality rate [5, 9]. Acute cardiac injury is one of the common complications in COVID-19 patients [7]. These findings suggest that cardiovascular system is tightly implicated in COVID-19. By collecting data from 116 laboratory-confirmed cases who were admitted to the General Hospital of Central Theater Command, we sought to provide an up-to-date description of the clinical characteristics and cardiovascular status of patients with COVID-19. This study would not only identify the clinical manifestations with greater precision, but also provide assistance for non-cardiovascular specialists to manage COVID-19 patients with cardiovascular disorders.

Methods

Study Design and Participants

For this single-centered, retrospective, observational study, we recruited patients with laboratory-confirmed COVID-19 admitted to the General Hospital of Central Theater Command (Wuhan, China) from January 20 to March 8, 2020. A confirmed case with SARS-CoV-2 infection was defined as a positive result for real-time reverse-transcriptase polymerase-chain-reaction (RT-PCR) assay for pharyngeal swab specimens. The General Hospital of Central Theater Command is one of the major hospitals responsible for COVID-19 treatment designated by the government. At the time of admission, the severity of COVID-19 was defined as mild, moderate, severe and critical cases based on the 6th edition guideline issued by the National Health Commission of China. Mild patients were not admitted in this designated hospital. Moderate cases were having mild symptoms of respiratory infections with pneumonia. Severe cases were defined as dyspnea, respiratory frequency ≥ 30 /minute, blood oxygen saturation $\leq 93\%$, PaO₂/FiO₂ ratio ≤ 300 mmHg, and/or pulmonary inflammation progressing $\geq 50\%$ within 24 to 48 hours. Critical cases were those who exhibited respiratory failure, shock, and/or multiple organ dysfunction. In this study, moderate cases were categorized as the non-severe group; severe and critical cases were categorized as the severe group. Oral informed consent was obtained from patients. All the protocol in this study was approved by the Ethics Commission of the General Hospital of Central Theater Command ([2020]025 - 1). The clinical outcomes (ie, discharge, mortality, and length of stay) were monitored up to April 13, 2020, the final date of follow-up.

Data Collection

We obtained the demographic data, medical history, underlying comorbidities, clinical symptoms and signs, laboratory findings, chest computed tomography (CT), treatment measures, and outcome data from electronic medical records for all hospitalized patients with laboratory confirmed COVID-19. ARDS and shock were defined in accordance with the guidance of WHO for COVID-19.[10] The diagnosis of acute kidney injury was based on the highest serum creatinine level and urine output [11]. Cardiac injury was defined if the serum concentration of hypersensitive cardiac troponin T (cTnT) was above the upper

limit of the reference range (≈ 0.02 ng/mL). Acute heart failure was defined based on the typical symptoms that may be accompanied by signs caused by a structural and/or functional cardiac abnormality [12]. Hypertension is defined in adults as the results of systolic blood pressure ≥ 140 mmHg and/or diastolic blood pressure ≥ 90 mmHg three times on different days.

Statistical Analysis

Continuous variables were presented as means and standard error (SEM) or medians and interquartile ranges (IQR) values as appropriate. Categorical variables were expressed as the counts and percentages in each category. Independent group t-test was used for continuous variable, when the data conformed to normal distribution; otherwise, the Mann-Whitney test was applied. Data (not normal distributed) from repeated measures were compared using the generalized linear mixed model. The chi-square test and Fisher's exact test were applied for categorical variables as appropriate. Two-sided p-values less than 0.05 were considered statistically significant. All analyses were conducted with GraphPad Prism software (version 8.0).

Results

Baseline Characteristics

The study population included 116 hospitalized patients with laboratory-confirmed COVID-19 (Table 1). The median age was 58.5 years (IQR, 47.0–69.0), and 36 (31.0%) were females. Fifty-nine (50.9%) patients had at least one underlying disorder. Hypertension (45 [38.8%]), diabetes (19 [16.4%]), coronary heart disease (17 [14.7%]), and cerebrovascular diseases (8 [6.9%]) were the most common coexisting conditions (Table 1). Fever (85.3%), dry cough (52.6%), fatigue (51.7%), anorexia (43.1%), dyspnea (44.8%), and chest discomfort (43.1%) were the most common symptoms, whereas dizziness (6.0%), nasal obstruction (5.2%), abdominal pain (2.6%), hemoptysis (0.9%) were less common (Table 1). The median durations from first symptoms to dyspnea and hospital admission were 4.5 days (IQR, 0–9.0), and 8.0 days (IQR, 4.0–11.0) respectively (Table 1).

Table 1
Baseline Characteristics of Patients Infected With COVID-19

Total (N=116)	Disease severity		PValue	
Non-severe (n=61)	Severe (n=55)			
Age, median (IQR), y	58.5 (47.0-69.0)	56.0 (37.0-64.0)	64.0 (53.0-76.0)	0.001
0-14 y	0	0	0	-
15-49 y	33/116 (28.4)	24/61 (39.3)	9/55 (16.4)	-
50-64 y	41/116 (35.3)	23/61 (37.7)	18/55 (32.7)	-
≥65 y	42/116 (36.2)	14/61 (23.0)	28/55 (50.9)	-
Sex –No., %				
Female	36/116 (31.0)	19/61(31.1)	17/55 (30.9)	1.000
Male	80/116 (69.0)	42/61 (68.9)	38/55 (69.1)	
Comorbidities –No., %				
Any	59/116 (50.9)	24/61 (39.3)	35/55 (63.6)	0.009
COPD	1/116 (0.9)	0/61 (0.0)	1/55 (1.8)	0.290
Diabetes	19/116 (16.4)	8/61 (13.1)	11/55 (20.0)	0.317
Hypertension	45/116 (38.8)	19/61 (31.1)	26/55 (47.3)	0.075
Coronary heart disease	17/116 (14.7)	4/61 (6.6)	13/55 (23.6)	0.009
Cerebrovascular disease	8/116 (6.9)	1/61 (1.6)	7/55 (12.7)	0.019
Malignancy	4/116 (3.4)	0/61 (0.0)	4/55 (7.3)	0.032
Chronic kidney disease	0/116 (0.0)	0/61 (0.0)	0/55 (0.0)	-
Chronic liver disease	2/116 (1.7)	1/61 (1.6)	1/55 (1.8)	0.941
HIV infection	0/116	0/61 (0.0)	0/55 (0.0)	-

(0.0)

Signs and symptoms—No., %				
Fever	99/116 (85.3)	49/61 (80.3)	50/55 (91.0)	0.108
Fatigue	60/116 (51.7)	26/61 (42.6)	34/55 (61.9)	0.039
Dry cough	61/116 (52.6)	31/61 (50.8)	30/55 (54.5)	0.688
Anorexia	50/116 (43.1)	27/61 (44.3)	23/55 (41.8)	0.791
Myalgia	32/116 (27.6)	19/61 (31.1)	13/55 (23.6)	0.366
Dyspnea	52/116 (44.8)	21/61 (34.4)	31/55 (56.4)	0.018
Expectoration	31/116 (26.7)	17/61 (27.9)	14/55 (25.5)	0.769
Hemoptysis	1/116 (0.9)	0/61 (0.0)	1/55 (1.8)	0.290
Pharyngalgia	15/116 (12.9)	7/61 (11.5)	8/55 (14.5)	0.623
Nasal obstruction	6/116 (5.2)	4/61 (6.6)	2/55 (3.6)	0.478
Diarrhea	17/116 (14.7)	14/61 (23.0)	3/55 (5.5)	0.008
Nausea	10/116 (8.6)	4/61 (6.6)	6/55 (11.0)	0.404
Dizziness	7/116 (6.0)	2/61 (3.3)	5/55 (9.1)	0.189
Headache	6/116 (5.2)	5/61 (8.2)	1/55 (1.8)	0.121
Vomiting	5/116 (4.3)	3/61 (4.9)	2/55 (3.6)	0.734
Chill	24/116 (20.7)	13/61 (21.3)	11/55 (20.0)	0.901
Shiver	5/116 (4.3)	3/61 (4.9)	2/55 (3.6)	0.734
Abdominal pain	3/116 (2.6)	3/61 (4.9)	0/55 (0.0)	0.096
Chest discomfort	50/116	22/61 (36.1)	28/55	0.107

	(43.1)		(51.0)	
Palpitation	13/116 (11.2)	7/61 (11.5)	6/55 (11.0)	0.923
Sleep disorders	27/116 (23.3)	15/61 (24.6)	12/55 (21.8)	0.724
Onset of symptom to, median (IQR), d				
Hospital admission	8.0 [4.0- 11.0]	8.0 [4.0-11.0]	8.0 [4.0- 11.0]	0.910
Dyspnea	4.5 [0.0- 9.0]	3.0 [0.0-12.0]	6.0 [1.5- 8.5]	0.649
Heart rate on admission, median (IQR), bpm	86.0 (80.0- 98.0)	85.5 (80.0-97.5)	86.0 (80.0- 98.0)	0.958
Respiratory rate on admission, median (IQR)	19.0 (18.0- 23.0)	19.0 (18.0-20.0)	20.0 (18.0- 25.0)	0.030
Mean arterial pressure on admission, median (IQR), mmHg	96.7 (86.7- 103.3)	96.7 (86.7- 103.5)	96.7 (86.7- 103.3)	1.000
Temperature on admission, median (IQR), °C	36.8 (36.5- 37.5)	36.7 (36.5-37.3)	37.0 (36.4- 37.8)	0.138

On admission, 61 and 55 patients were categorized into non-severe and severe subgroups, respectively. The age differed significantly between the two groups (median age, non-severe vs severe, 56.0 years [IQR, 37.0–64.0] vs 64.0 years [IQR, 53.0–76.0]; $P < 0.001$). Severe cases were more prone to having underlying comorbidities, including coronary heart disease (13 [23.6%] vs 4 [6.6%], cerebrovascular diseases (7 [12.7%] vs 1 [1.6%]), and malignancy (4 [7.3%] vs 0 [0.0%]). Compared with the non-severe group, dyspnea and fatigue were more frequently reported in severe patients. Vital signs were recorded on the day of admission to hospital for all patients. Respiratory rate was higher in severe cases as compared with non-severe cases (20.0 [IQR, 18.0–25.0] vs 19.0 [IQR, 18.0–20.0; $P = 0.030$]). While, heart rate, mean arterial pressure, and body temperature showed no significant difference between the two groups (all $P \geq 0.05$)

Laboratory and radiologic findings at presentation

The most common pattern on chest CT was bilateral patchy shadowing (69.9%). These imaging alterations were more prominent in severe patients (47 [85.5%] vs 36 [59.0%]; $P = 0.002$) (Table 2).

Table 2

Chest Computed Tomographic Images Findings of Patients Infected With COVID-19 on Admission to Hospital

Abnormalities on chest CT – No./total No. (%)	Total (N=116)		Disease severity	PValue
	Non-severe (n=61)		Severe (n=55)	
Local patchy shadowing	19/116 (16.4)	14/61 (23.0)	5/55 (9.1)	0.044
Bilateral patchy shadowing	81/116 (69.9)	36/61 (59.0)	47/55 (85.5)	0.002
Interstitial abnormalities	4/116 (3.4)	2/61 (3.3)	2/55 (3.6)	0.916

There were numerous differences in laboratory findings between severe and non-severe cases (Table 3). Laboratory abnormalities were more obviously seen in severe cases, including lower counts of lymphocyte, T cells, CD4⁺ and CD8⁺ T cells, and elevated levels of neutrophil count, procalcitonin, c-reactive protein, interleukin 6, D-dimer, creatinine, blood urea nitrogen, lactate dehydrogenase, myoglobin, cTnT, and NT-proBNP (all P < 0.05). Hypokalemia (20.7%) was prevailing in both severe and non-severe patients, though no statistical difference was found between them.

Table 3

Laboratory Findings of Patients Infected With COVID-19 on Admission to Hospital

Laboratory findings	Disease severity				
	Normal Range	Total (N=116)	Non-severe (n=61)	Severe (n=55)	PValue
White blood cell count, × 10 ⁹ /L	3.5-9.5	4.9 (3.9-6.1)	5.0 (3.9-5.7)	4.9 (3.8-7.2)	0.3442
≥9.5 × 10 ⁹ /L, No./total No. (%)		9/116 (7.8)	1/61 (1.6)	8/55 (14.5)	0.0129
≥3.5 × 10 ⁹ /L, No./total No. (%)		17/116 (14.7)	9/61 (14.8)	8/55 (14.5)	1.0000
Neutrophil count, × 10 ⁹ /L	1.8-6.3	3.1 (2.2-4.5)	2.9 (2.1-3.8)	3.4 (2.5-5.8)	0.0134
Lymphocyte count, × 10 ⁹ /L	1.1-3.2	1.0 (0.7-1.3)	1.2 (0.9-1.6)	0.9 (0.6-1.1)	< 0.0001
≥1.1 × 10 ⁹ /L, No./total No. (%)		66/116 (56.9)	24/61(39.3)	42/55 (76.4)	< 0.0001
Total T cell count × 10 ⁶ /L	955-2860	590.5 (361.0-941.8)	915.0 (481.0-1246.0)	448 (238.5-644.0)	< 0.0001
≥955 × 10 ⁶ /L, No./total No. (%)		65/84 (77.4)	27/43 (62.8)	38/41 (92.7)	0.0014
CD4+ T cell count × 10 ⁶ /L	550-1440	335.0 (214.8-493.3)	458.0 (275.0-607.0)	262.0 (156.5-335.0)	0.0002
≥550 × 10 ⁶ /L, No./total No. (%)		66/84 (78.6)	29/43 (67.4)	37/41 (90.2)	0.0158
CD8+ T cell count × 10 ⁶ /L	320-1250	217.0 (111.3-359.8)	285.0 (144.0-435.0)	157.0 (78.5-297.0)	0.0052
≥320 × 10 ⁶ /L, No./total No. (%)		46/84 (54.8)	13/43 (30.2)	33/41 (80.5)	< 0.0001
Monocyte count, × 10 ⁹ /L	0.1-0.6	0.4 (0.3-0.6)	0.4 (0.4-0.6)	0.4 (0.2-0.6)	0.4886
Platelet count, × 10 ⁹ /L	125-350	177.0 (137.0-230.0)	179.0 (145.0-236.0)	161.5 (121.8-218.3)	0.1166
Haemoglobin level g/L	130-175	131.0 (118.0-140.0)	129.0 (120.0-140.5)	131.0 (115.0-140.0)	0.7728

Laboratory findings	Disease severity				
	Normal Range	Total (N=116)	Non-severe (n=61)	Severe (n=55)	PValue
Prothrombin time, s	10.0-14.0	12.2 (11.5-13.0)	11.9 (11.5-13.0)	12.4 (11.6-13.1)	0.3330
Activated partial thromboplastin time, s	23.5-39.1	32.3 (30.1-34.8)	32.3 (30.2-34.6)	32.6 (29.8-35.0)	0.8014
D-dimer, ng/mL	0-243	168.0 (92.0-393.5)	122.0 (75.0-254.0)	222.0 (132.5-529.0)	0.0004
≥243ng/mL, No./total No. (%)			15/61 (24.6)	25/55 (45.5)	0.0204
Creatinine, umol/L	45-110	69.5 (56.0-81.0)	68.5 (54.3-75.8)	72.0 (58.8-86.5)	0.0156
≥110umol/L, No./total No. (%)		8/116 (6.9)	0/61 (0)	8/55 (14.5)	0.0019
Blood urea nitrogen, mmol/L	2.5-6.3	4.4 (3.4-5.8)	3.9 (3.2-5.0)	4.9 (3.5-7.0)	0.0080
≥6.3mmol/L, No./total No. (%)		22/116 (19.0)	5/61 (8.2)	17/55 (30.9)	0.0021
Alanine aminotransferase, U/L	9-50	24.0 (17.3-37.8)	22.0 (15.5-37.5)	27.0 (20.0-39.0)	0.1227
≥50.0U/L, No./total No. (%)		23/116 (19.8)	11/61 (18.0)	12/55 (21.8)	0.6470
Aspartate aminotransferase, U/L	15-40	34.5 (25.0-52.8)	31.0 (22.5-49.5)	38.0 (26.0-59.0)	0.0810
≥40.0U/L, No./total No. (%)		46/116 (39.7)	21/61 (34.2)	25/55 (45.5)	0.2573
Total bilirubin, umol/L	4.42-20.52	11.0 (8.1-14.1)	11.0 (8.3-13.0)	10.5 (7.8-16.3)	0.9308
Creatine kinase-MB, U/L	0-24	17.0 (15.0-21.0)	17.0 (15.0-22.5)	18.0 (15.8-20.0)	0.6960
≥24.0U/L, No./total No. (%)		19/116 (16.4)	12/61 (19.7)	7/55 (12.7)	0.4518

Laboratory findings	Laboratory findings		Disease severity		PValue		
	Normal Range	Total (N=116)	Non-severe (n=61)	Severe (n=55)			
Lactate dehydrogenase, U/L	109-225	245.5 (194.3-319.8)	225.0 (184.0-303.5)	275.0 (212.0-378.0)	0.0041		
≥225.0U/L, No./total No. (%)		69/116 (59.5)	30/61 (49.2)	39/55 (70.9)	0.0230		
Hypersensitive troponin T, ng/mL	≤0.02	0.008 (0.005-0.013)	0.007 (0.004-0.01)	0.0115 (0.006-0.0175)	0.0015		
≥0.02ng/mL, No./total No. (%)		16/116 (13.8)	4/61 (6.6)	12/55 (21.8)	0.0288		
Myoglobin, ng/mL	28-72	39.4 (22.3-92.7)	26.1 (21.0-47.8)	54.1 (33.4-120.7)	0.0002		
≥72.0ng/mL, No./total No. (%)		34/116 (29.3)	12/61 (19.7)	22/55 (40.0)	0.0096		
NT-proBNP, pg/mL		71.5 (27.0-363.5)	37.0 (19.0-118.8)	233.0 (53.3-693.0)	< 0.0001		
elevated cases, No./total No. (%)		8/116 (6.9)	1/61 (1.6)	7/55 (12.8)	0.0260		
Procalcitonin, ng/mL	0.00-0.50	0.07 (0.03-0.16)	0.05 (0.03-0.09)	0.09 (0.06-0.23)	< 0.0001		
≥0.5ng/mL, No./total No. (%)		8/116 (6.9)	2/61 (3.3)	6/55 (10.9)	0.1473		
IL-6, pg/mL	0-7.0	18.7 (7.7-46.4)	11.8 (3.3-27.5)	33.0 (13.1-69.2)	< 0.0001		
C-reactive protein level, mg/L	0-10	15.1 (9.0-50.9)	10.0 (7.0-27.5)	40.0 (10.0-8.0)	0.0023		
Sodium,mmol/L	137.0-147.0		138.8 (137.1-141.0)	140.0 (138.2-141.7)	138.0 (134.8-139.3)	0.0001	
Potassium,mmol/L	3.5-5.3		4.0 (3.5-4.3)	3.9 (3.5-4.3)	4.1 (3.6-4.3)	0.4548	

Laboratory findings			Disease severity			
	Normal Range	Total (N=116)	Non-severe (n=61)	Severe (n=55)	PValue	
⊗5.3mmol/L, No./total No. (%)	1/116 (0.9)		0/61 (0)		1/55 (1.8)	0.4741
⊗3.5mmol/L, No./total No. (%)	24/116 (20.7)		12/61 (19.7)		12/55 (21.8)	0.8213
Chloride,mmol/L	99.0-110.0		103.3 (99.8-105.5)		104.1 (101.3-106.2)	102.2 (99.2-103.8) 0.0011
Calcium,mmol/L	2.02-2.6		2.10 (2.04-2.21)		2.14 (2.09-2.28)	2.07 (2.02-2.13) < 0.0001
Magnesium,mmol/L	0.6-1.2		0.73 (0.68-0.78)		0.76 (0.70-0.79)	0.70 (0.64-0.76) 0.0025

Complications, Main Interventions, And Outcomes

The organ dysfunction and treatment of the 116 patients were shown in **Table 4**. On April 13, 2020, 4 patients (3.4%) were still hospitalized. A total of 101 patients (90.5%) had been discharged, and 7 patients (6.0%) had died. Among the 116 patients, the most common complication was acute cardiac injury (23 [19.8%]), followed by acute heart failure (21 [18.1%]), ARDS (20 [17.2%]), shock (16 [13.8%]), or liver dysfunction (15 [12.9%]). Severe cases were more liable to having one of these complications than non- severe cases.

Table 4
Complications, treatment and outcomes of patients with COVID-19

Characteristics	Total (N=116)		Disease severity	PValue
			Non-severe (n=61)	Severe (n=55)
Complications, No./total No. (%)				
Shock	16/116 (13.8)	0/61 (0.0)	16/55 (29.1)	< 0.0001
Acute cardiac injury	23/116 (19.8)	4/61 (6.6)	19/55 (34.5)	0.001
Acute respiratory distress syndrome	20/116 (17.2)	0/61 (0.0)	20/55 (36.4)	< 0.0001
Liver dysfunction	15/116 (12.9)	2/61 (3.3)	13/55 (23.6)	0.0011
Acute kidney injury	3/116 (2.6)	0/61 (0.0)	3/55 (5.5)	0.0565
Acute heart failure	21/116 (18.1)	3/61 (4.9)	18/55 (32.7)	0.0002
Treatment, No./total No. (%)				
Administration of oseltamivir	91/116 (78.4)	46/61 (75.4)	45/55 (81.8)	0.4019
Administration of lopinave/litonawe(LPV/r)	74/116 (63.8)	42/61 (68.9)	32/55 (58.2)	0.2325
Administration of arbidolhydrochloride	27/116 (23.3)	15/61 (24.6)	12/55 (21.8)	0.7243
Administration of ribavirin injection	71/116 (61.2)	37/61 (60.7)	34/55 (61.8)	0.8979
Administration of antibiotics	113/116 (97.4)	58/61 (95.1)	55/55 (100.0)	0.0956
Use of antifungal medications	12/116 (10.3)	0/61 (0.0)	12/55 (21.8)	0.0001
Administration of systemic corticosteroids	55/116 (47.4)	14/61 (23.0)	41/55 (74.5)	< 0.0001
Oxygen inhalation	103/116 (88.8)	50/61 (82.0)	53/55 (96.4)	0.0141
Noninvasive ventilation	18/116 (15.5)	0/61 (0.0)	18/55 (32.7)	< 0.0001
Invasive mechanical ventilation	10/116 (8.6)	0/61 (0.0)	10/55 (18.2)	0.0005

Extracorporeal membrane oxygenation	3/116 (2.6)	0/61 (0.0)	3/55 (5.5)	0.0646
Use of intravenous immunoglobulin	74/116 (63.8)	25/61 (41.0)	49/55 (89.1)	< 0.0001
Use of thymopeptides injection	92/116 (79.3)	45/61 (73.8)	47/55 (85.5)	0.1209
Traditional Chinese medical herbal treatment	103/116 (88.8)	55/61 (90.2)	48/55 (87.3)	0.2955
Clinical outcomes, No./total No. (%)				
Discharge from hospital	105/116 (90.5)	61/61 (100.0)	44/55 (80.0)	0.0002
Death	7/116 (6.0)	0/61 (0.0)	7/55 (12.7)	0.0040
Staying in hospital	4/116 (3.4)	0/61 (0.0)	4/55 (7.3)	0.0321

All patients were treated in isolation. 112 (96.6%) patients received antiviral treatment, including oseltamivir phosphate (91, [78.4%]), lopinavir/ritonavir (74, [63.8%]), arbidol hydrochloride (27, [21.4%]), and ribavirin Injection (71, [61.2%]). Antibacterial therapy was performed for most patients (113, [97.4%]). Twelve (10.3%) patients adopting antifungal medications were all enrolled in severe group. Systemic corticosteroid was given to 47.4% of cases and more so in the severe patients than in the non-severe group (74.5% vs 23.0%, $P < 0.0001$). Obviously, more severe cases received mechanical ventilation (non-invasive: 32.7% vs. 0%, $P < 0.0001$; invasive: 18.2% vs. 0%, $P < 0.0001$) as compared with non-severe cases. Three severe patients were treated with extracorporeal membrane oxygenation. Traditional Chinese medical herbal treatment and immune support therapy were applied in most cases. Overall, all these interventions were initiated in significantly higher percentages of severe patients.

Dynamic profile of blood pressure and cardiac markers

To determine the cardiovascular implications during COVID-19 progression, the dynamic changes in heart rate, blood pressure, and 5 clinical laboratory parameters, including creatine kinase-MB, α -hydroxybutyric dehydrogenase, lactate dehydrogenase, cTnT, and NT-proBNP were tracked in survivors (Figure 1, 2). The results of heart rate and blood pressure were recorded every day at the same time. Laboratory parameters were examined once every 3 days. During hospitalization, higher levels of systolic blood pressure were observed in severe cases (Figure 1B). A total of 8 patients from non-severe group and 16 patients from severe group were diagnosed with new onset hypertension (Figure 1D). The morbidity rate of hypertension was significantly higher in severe cases (42 [76.4%] vs 27 [44.3%]; $P=0.0006$) (Figure 1E).

During hospitalization, severe cases exhibited higher levels of cardiac markers (Figure 2). The prevalence of cardiac complications, including acute cardiac injury and heart failure was significantly higher in

severe cases (**Table 4**). Increases of lactate dehydrogenase and α -hydroxybutyric dehydrogenase were also more likely to occur in severe patients (Figure 2B, 2C). Myocardial enzymes increased rapidly in the early stage of illness, but gradually decreased with the disease progression. In the end period of the 28-day time frame, abnormal results of cTnT and NT-proBNP were still common in some of the severe patients because of progression of COVID-19 ((Figure 2D, 2E).

Discussion

This retrospective study described the clinical characteristics and cardiovascular implications in hospitalized patients with COVID-19 in Wuhan. By April 13, 2020, of the 116 patients included in this study, 69% were male, 47% were severe cases, 90.5% were discharged, 6.0% (7) died, and 3.4% remained hospitalized. Most severe patients were older and had more underlying conditions. Common symptoms at onset of illness were fever, dry cough, fatigue, dyspnea, and chest discomfort. Local and/or bilateral patchy shadowing was a typical hallmark of CT imaging for COVID-19. Lymphopenia and elevated levels of neutrophil count, C-reactive protein, interleukin 6, D-dimer, creatinine, lactate dehydrogenase, cTnT, and NT-proBNP were more commonly seen in severe cases. During hospitalization, the prevalence of new onset hypertension, acute heart injury, and heart failure was significantly higher in severe patients.

In our cohort, 69% (80) of COVID-19 patient were male. Severe patients were older and had a greater number of comorbid conditions. Evidence from previous studies suggest that older, male patients are the most susceptible to SARS-CoV-2 infection [4, 5, 7, 13], which is supported by our data. It has been confirmed that increased age was associated with death in COVID-19 patients [14], and the coexistence of agedness and comorbidity could lead to an even higher risk of death [13]. Older age has been regarded as an important independent predictor of mortality in COVID-19.

Cardiovascular diseases have a high incidence rate in the middle aged and elderly population [8]. As previously reported [14], we observed that many COVID-19 patients had a comorbidity, with hypertension being the most common (45 [38.8%]), followed by diabetes (19 [16.4%]) or coronary heart disease (17 [14.7%]). The morbidity rates of coronary heart disease and cerebrovascular diseases were significantly higher in the severe group. Thus, older people with comorbidities, such as coronary heart disease and hypertension were thought to be more vulnerable to SARS-CoV-2 and result in more severe outcomes and elevated case-fatality rate[5, 9, 15]. In the present study, 4 of 7 dead patients had preexisting hypertension and coronary heart disease. Previously, coronary heart disease has also been found to be correlated with acute cardiac events and poor outcomes in influenza and other respiratory viral infections [16, 17], Multivariate logistic regression analysis demonstrated that coronary heart disease and heart injury were the independent risk factors for critical disease status in COVID-19 patients [18]. More intense clinical care is in need for COVID-19 patients with cardiac-related chronic diseases.

Incident cardiovascular complications including new or worsening heart failure, new or worsening arrhythmias, or myocardial infarction are common in patients with pneumonia and are associated with increased short-term mortality [19]. Acute pneumonia brings important effects on the status of cardiovascular system irrespective of severities of infection [16, 19]. Risk factors for cardiac

complications after pneumonia include older age, preexisting cardiovascular diseases, and greater severity of pneumonia [16, 19]. An analysis of 112 cardiovascular disease patients with COVID-19 found that, COVID-19 patients combined with cardiovascular disease were associated with a higher risk of mortality [15]. In this study, compared with non-severe patients with COVID-19, severe patients showed abnormalities in numerous cardiac markers. During hospitalization, the morbidity of new onset hypertension, acute heart injury, and heart failure was significantly higher in severe patients. Increased level of myocardial enzymes and cTnT was found in all 3 dead cases. As far as we know, this is the first study that reports the prevalence rate of new onset hypertension was significantly higher in hospitalized severe patients with COVID-19. These findings suggest a higher possibility of cardiovascular complications in severe patients with COVID-19. Outcomes of patients with COVID-19 may be improved by prevention of the development and progression of associated cardiac complications.

Angiotensin-converting enzyme 2 (ACE2) acts as a receptor for SARS-CoV-2 entry into cells and contributes to the pathogenesis of COVID-19 [20]. Meanwhile, ACE2 is widely expressed in myocytes and vascular endothelial cells. At least, there is theoretically a possibility of direct cardiovascular involvement induced by the virus. The only pathological result of heart biopsy in a fatal case with COVID-19 showed a few interstitial mononuclear inflammatory infiltration, but no other substantial damage in the heart tissue [21]. However, given that this patient had no clinical manifestations of myocardial injury during the whole course of this disease, it could not be concluded whether myocardium was involved in SARS-CoV-2 infection yet. We speculated that the potential pathogenesis of myocardial injury in COVID-19 may include several processes, SARS-CoV-2 may directly invade myocytes via ACE2 and cause viral myocarditis; the imbalance between supply and demand in oxygen further results in myocardial injury; and inflammatory cytokines storm. In order to further clarify the etiology of SARS-CoV-2 infection related myocardial injury, it is of great need to obtain pathological evidence from COVID-19 patients showing definite myocardial injury.

Recently, the safety of treatment applying angiotensin converting enzyme inhibitors (ACEI) or angiotensin receptor blockers (ARBs) in relation to COVID-19 has been concerned. An observational study containing 112 patients with cardiovascular diseases infected by COVID-19 reported that there was no significant difference in the proportion of ACEI/ARB medication between the critical group and the general group or between non-survivors and survivors [15]. Currently, there is a lack of any experimental or clinical evidence suggesting adverse or beneficial outcomes with background use of ACEI, ARBs or other RAAS antagonists in COVID-19 or among COVID-19 patients with a history of cardiovascular disease treated with such agents. Statements of ACC and ESC Council on Hypertension do not recommend to discontinue ACEI/ARB treatment in the lack of any evidence supporting adverse effect of ACEI and ARB in the context of the pandemic COVID-19 outbreak [22, 23]. Individualized treatment strategies should be approached according to each patient's hemodynamic status and clinical manifestations.

Our study has some limitations. First, only 116 patients with laboratory-confirmed COVID-19 were included. It would be better to include a large population of patients from other centers in Wuhan, and even in other cities in China to get a more comprehensive understanding of COVID-19. Second, not all

laboratory tests were dynamically performed in all patients, including the counts of lymphocyte subsets and inflammatory cytokines, therefore their role in the pathogenesis of COVID-19 might be underestimated. Third, due to the retrospective study design, echocardiography and electrocardiograph were only performed in some of the patients. The detailed information of ACEI/ARB medication was incomplete. Therefore, we could not further assess the changes of cardiac structure and function during the progression of COVID-19, and the possible effect of ACEI/ARB on SARS-CoV-2 infection. Last but not least, we were unable to obtain myocardial tissues from deceased COVID-19 patients with heart injury. The characteristics of myocardial damage should be further demonstrated by pathologists.

Conclusions

In this single-centered, retrospective, observational study, we found that the infection of SARS-CoV-2 were more likely to occur in older population with cardiovascular comorbidities. Cardiovascular complications, including new onset hypertension and heart injury were common in severe patients with COVID-19. More comprehensive and in-depth researches are in need to unveil the cardiovascular involvement in COVID-19 to further understand the disease.

Abbreviations

COVID-19

Corona Virus Disease 2019; SARS-CoV-2:2019 novel coronavirus; ARDS:acute respiratory distress syndrome; cTnT:hypersensitive cardiac troponin T; NT-proBNP:N-terminal B-type natriuretic peptide; ACC:American College of Cardiology; ESC:European Society of Cardiology; SEM:standard error; IQR:interquartile ranges.

Declarations

Ethics approval and consent to participate

The Ethics Commission of General Hospital of Central Theater Command approved this study ([2020]025-1).

Consent for publication

Not applicable.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This research was supported by the National Natural Science Foundation of China (31600942), Wuhan young and middle-aged medical backbone talent training program, and 2014 Major Projects of PLA Logistics Research Fund (AWS13C003).

Authors' contributions

SX was a major contributor in writing the manuscript and statistical analysis. LL analyzed and interpreted the patient data. All authors participated in collecting data, designing the analyses, and interpreting the results. SD had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis. All authors read and approved the final manuscript.

Acknowledgements

Not applicable.

Authors' information

¹Department of Cardiology, ²Department of integrated traditional Chinese and Western internal medicine, ³Division of medical management, the General Hospital of Central Theater Command, Wuhan 430000, China.

References

1. Liu W, Zhang R, Shu R, Yu J, Li H, Long H, Jin S, Li S, Hu Q, Yao F, et al. Study of the Relationship between Microbiome and Colorectal Cancer Susceptibility Using 16SrRNA Sequencing. *Biomed Res Int.* 2020;2020:7828392–2.
2. Wu F, Zhao S, Yu B, Chen YM, Wang W, Song ZG, Hu Y, Tao ZW, Tian JH, Pei YY, et al: **A new coronavirus associated with human respiratory disease in China.** *Nature* 2020.
3. Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, Zhao X, Huang B, Shi W, Lu R, et al. A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N Engl J Med.* 2020;382(8):727–33.
4. Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX, Liu L, Shan H, Lei CL, Hui DSC, et al: **Clinical Characteristics of Coronavirus Disease 2019 in China.** *The New England journal of medicine* 2020.
5. Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, Qiu Y, Wang J, Liu Y, Wei Y, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet.* 2020;395(10223):507–13.
6. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, Zhang L, Fan G, Xu J, Gu X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet.* 2020;395(10223):497–506.

7. Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, Wang B, Xiang H, Cheng Z, Xiong Y, et al: **Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China.** *Jama* 2020.
8. Joseph P, Leong D, McKee M, Anand SS, Schwalm JD, Teo K, Mente A, Yusuf S. Reducing the Global Burden of Cardiovascular Disease, Part 1: The Epidemiology and Risk Factors. *Circulation research*. 2017;121(6):677–94.
9. Wu Z, McGoogan JM: **Characteristics of and Important Lessons From the Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72314 Cases From the Chinese Center for Disease Control and Prevention.** *Jama* 2020.
10. WHO. **Clinical management of severe acute respiratory infection when novel coronavirus (nCoV) infection is suspected.** <https://>. 2020-01-11.
11. Kellum JA, Lameire N, Group KAGW. Diagnosis, evaluation, and management of acute kidney injury: a KDIGO summary (Part 1). *Critical care*. 2013;17(1):204.
12. Ponikowski P, Voors AA, Anker SD, Bueno H, Cleland JGF, Coats AJS, Falk V, González-Juanatey JR, Harjola V-P, Jankowska EA, et al. 2016 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: The Task Force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC) Developed with the special contribution of the Heart Failure Association (HFA) of the ESC. *Eur Heart J*. 2016;37(27):2129–200.
13. Yang X, Yu Y, Xu J, Shu H, Xia J, Liu H, Wu Y, Zhang L, Yu Z, Fang M, et al: **Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study.** *The Lancet Respiratory medicine* 2020.
14. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, Xiang J, Wang Y, Song B, Gu X, et al: **Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study.** *Lancet* 2020.
15. Peng YD, Meng K, Guan HQ, Leng L, Zhu RR, Wang BY, He MA, Cheng LX, Huang K, Zeng QT. [Clinical characteristics and outcomes of 112 cardiovascular disease patients infected by 2019-nCoV]. *Zhonghua xin xue guan bing za zhi*. 2020;48(0):E004.
16. Corrales-Medina VF, Musher DM, Shachkina S, Chirinos JA. Acute pneumonia and the cardiovascular system. *Lancet*. 2013;381(9865):496–505.
17. Bhatt DI Fau - Udell JA, Zawi R, Fau - Bhatt DL, Bhatt DI Fau - Keshtkar-Jahromi M, Keshtkar-Jahromi M, Fau - Gaughran F, Gaughran F, Fau - Phrommintikul A, Phrommintikul A, Fau - Ciszewski A, Ciszewski A, Fau - Vakili H, Vakili H. Fau - Hoffman EB, Hoffman Eb Fau - Farkouh ME, Farkouh Me Fau - Cannon CP et al: **Association between influenza vaccination and cardiovascular outcomes in high-risk patients: a meta-analysis.** (1538–3598 (Electronic)).
18. Chen C, Chen C, Yan JT, Zhou N, Zhao JP, Wang DW. [Analysis of myocardial injury in patients with COVID-19 and association between concomitant cardiovascular diseases and severity of COVID-19]. *Zhonghua xin xue guan bing za zhi*. 2020;48(0):E008.

19. Corrales-Medina VF, Musher Dm Fau - Wells GA, Wells Ga Fau - Chirinos JA, Chirinos Ja Fau - Chen L, Chen L Fau - Fine MJ, Fine MJ: **Cardiac complications in patients with community-acquired pneumonia: incidence, timing, risk factors, and association with short-term mortality.** (1524–4539 (Electronic)).
20. Xu X, Chen P, Wang J, Feng J, Zhou H, Li X, Zhong W, Hao P: **Evolution of the novel coronavirus from the ongoing Wuhan outbreak and modeling of its spike protein for risk of human transmission.** *Science China Life sciences* 2020.
21. Xu Z, Shi L, Wang Y, Zhang J, Huang L, Zhang C, Liu S, Zhao P, Liu H, Zhu L, et al: **Pathological findings of COVID-19 associated with acute respiratory distress syndrome.** *The Lancet Respiratory medicine* 2020.
22. American College of Cardiology. **ACC Clinical Bulletin: Cardiac Implications of Novel Wuhan Coronavirus (COVID-19).** 2020-02–13.
23. ESC Council on Hypertension: **Position Statement of the ESC Council on Hypertension on ACE-Inhibitors and Angiotensin Receptor Blockers.** 2020-03–13.

Figures

Figure 1




Figure 1

Dynamic monitoring heart rate and blood pressure of patients hospitalized with COVID-19 During hospitalization, heart rate and blood pressure were recorded every day at the same time by nurses. The dynamic monitoring results of heart rate (A), systolic blood pressure (B), and diastolic blood pressure (C) were compared between non-severe and severe cases. D, the counts of normal blood pressure, preexisting hypertension, and new onset hypertension cases in non-severe and severe groups. E, rate of preexisting hypertension, new onset hypertension, and the total hypertension were compared between non-severe and severe subgroups. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ for Non-severe vs Severe.

Figure 2




Figure 2

Temporal changes in laboratory markers in patients hospitalized with COVID-19 Figure shows temporal changes in creatine kinase-MB (A), α-hydroxybutyric dehydrogenase (B), lactate dehydrogenase (C), cTnT (D), and NT-proBNP (E) of non-severe and severe patients every other day after admission. The dotted lines in black show the lower limit of normal for each parameter. *P<0.05, **P<0.01 for Non-severe vs Severe.