

# Fully automatic three-dimensional MRI analysis for the cartilage and meniscus of the knee: Kanagawa Knee Study

**Hayato Aoki**

Tokyo Ika Shika Daigaku

**Nobutake Ozeki**

Tokyo Ika Shika Daigaku

**Hisako Katano**

Tokyo Ika Shika Daigaku

**Akinobu Hyodo**

Tokyo Ika Shika Daigaku

**Junpei Matsuda**

Tokyo Ika Shika Daigaku

**Kimiko Takanashi**

Tokyo Ika Shika Daigaku

**Kenji Suzuki**

Fujifilm Corporation

**Jun Masumoto**

Fujifilm Corporation

**Noriya Okanouchi**

Kanagawa Prefectural Government

**Takeo Fujiwara**

Tokyo Ika Shika Daigaku

**Ichiro Sekiya** (✉ [sekiya.arm@tmd.ac.jp](mailto:sekiya.arm@tmd.ac.jp))

Tokyo Ika Shika Daigaku <https://orcid.org/0000-0002-6331-722X>

---

## Research article

**Keywords:** Cartilage, Meniscus, 3D MRI, U-Net, CNN

**Posted Date:** May 7th, 2020

**DOI:** <https://doi.org/10.21203/rs.3.rs-26389/v1>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# Abstract

## Objective

We have developed a fully automatic three-dimensional MRI analysis software that measures the projected cartilage area ratio (PCAR) to allow for the quantification of the cartilage in the knee. Our objectives for this cross-sectional study were to verify our software's accuracy and to quantify cartilage and meniscus extrusion. We also examined which cartilage quantification was most affected by age and analyzed the relationship between PCAR and meniscus extrusion.

## Methods

108 subjects were selected for training, and Dice similarity coefficients were determined from 5 other subjects. This study included 561 subjects between 30–70 years of age. From their knee MRI data, we quantified cartilage thickness, cartilage volume, and PCAR (0.0–1.5 mm) in four regions, including the medial tibial (MT) cartilage. Furthermore, each region was divided into nine subregions. The medial central (mc) subregion was also analyzed. As a quantification for meniscus extrusion, the medial meniscus coverage ratio (MMCR) was also investigated.

## Results

Dice similarity coefficients were 0.911 and 0.892 for the femoral and tibial cartilage and 0.916 and 0.891 for the medial and lateral meniscus. Among 48 cartilage quantifications, the highest absolute value of the correlation coefficient with age was mcMT PCAR 1.0 mm in females and mcMT cartilage thickness in males. In females, mcMT PCAR 1.0 mm was correlated with MMCR, although MMCR was not correlated with age. In males, mcMT PCAR 0.0 mm was correlated with MMCR.

## Conclusions

Our software showed high segmentation accuracy and provided numerous quantifications of cartilage related to age and meniscus extrusion.

# Introduction

Three-dimensional magnetic resonance imaging (3D MRI) analysis is useful for the quantification of cartilage and the meniscus in osteoarthritis of the knee [1–4]. However, this promising method is not popular at present because segmentation of cartilage and the meniscus often requires manual operation or correction, which requires time and effort. To solve this problem, automatic segmentation techniques using deep neural networks have been developed [5–8]. We have also recently developed a system for automatic extraction of cartilage and the meniscus using deep neural networks. The first purpose of our study was to verify the accuracy of our software for automatic cartilage and meniscus segmentation on knee MRI.

For quantification of knee cartilage using 3D MRI, cartilage thickness and volume are used in many cases. As another cartilage measurement, we recently proposed “projected cartilage area ratio” (PCAR), which correlates the projected area ratio of the cartilage with intended thickness with the total area of the region of interest (ROI). The cartilage thickness used in assessing PCAR can be adjusted according to disease conditions so that even slight changes can be revealed in a relatively short period (9).

Furthermore, PCAR, unlike cartilage thickness and volume, may not be affected by body size and may have an advantage in cross-sectional studies when analyzing the effects of age on cartilage. However, in our past development, the 3D-reconstructed femoral cartilage was projected directly onto the 2D plane, so the apparent thickness in the 2D projection was greater than the actual thickness due to the slope of the cartilage. To overcome this problem, we have improved the 3D-reconstructed femoral cartilage to project radially onto the 2D plane. We also recently developed a PCAR evaluation for tibial cartilage. In addition, for more detailed analysis, we enabled PCAR evaluation in five regions (including the medial femoral condyle cartilage, lateral femoral condyle cartilage, trochlea femoral cartilage, medial tibial cartilage, and lateral tibial cartilage) and nine subregions.

The meniscus plays a critical role in shock absorption and is especially important in regulating load-bearing distribution. Meniscus extrusion is one of the strongest risk factors for the progression of osteoarthritis (10). For the index of meniscus extrusion, the deviation width of the meniscus in the coronal view of 2D MRI or ultrasound images is often used [9]. However, in meniscus extrusion, the meniscus does not always simply displace externally, and an evaluation on only one plane cannot explain the pathological condition of the meniscus extrusion in detail. On the other hand, 3D MRI analysis can provide the meniscus extrusion area, volume, and meniscus coverage ratio (MCR), which also covers the tibial cartilage area (12–14). These indicators can be more useful for analyzing the relationship between meniscus extrusion and osteoarthritis.

Extensive epidemiological studies have shown that osteoarthritis of the knee increases with age and that it affects females more frequently than males [10–12]. However, it remains unknown whether PCAR and the quantification of meniscus extrusion are affected by age or gender. Furthermore, analyzing the relationship between PCAR and the quantification of meniscus extrusion will be useful for continued elucidation of the pathology of osteoarthritis. To address these factors, we collected 3D MRI knee data from 561 volunteers, including more than 50 females and 50 males in their 30 s, 40 s, 50 s, 60 s, and 70 s. The second purpose of our study was to quantify cartilage (including PCAR) and meniscus extrusion using 3D MRI analysis to examine whether these quantifications were affected by gender, body size, and age, as well as to analyze the relationship between PCAR and meniscus extrusion.

## **Materials And Methods**

### **Magnetic resonance imaging (MRI)**

Images were collected with a 3.0-T MRI (Achieva 3.0T TX; Philips) using 16-channel coils. The sagittal plane of the knee joint was imaged using both a fat-suppressed spoiled gradient echo (SPGR) sequence

to extract cartilage and a proton density weighted imaging (PDWI) sequence to extract meniscus and bone. For both sequences, sagittal imaging was performed at an in-plane resolution of  $0.31 \times 0.31$  mm, a partition thickness of 0.36 mm (320 slices), and a field of view of  $150 \times 150$  mm.

## Automatic Segmentation Algorithm Of 3D MRI

This study was approved by the Medical Research Ethics Committee of Tokyo Medical and Dental University and written informed consent was obtained from all participants. The protocol was enrolled in a database of the National University Hospital Council of Japan (UMIN000031924) and disclosed.

We used 3D U-Net as the neural network model for our software [13]. The network used convolution with padding to improve the accuracy of edge extraction. As a loss function, we used Soft Dice loss [14] (Supplementary Fig. 1).

For neural network construction, we randomly chose 10 healthy volunteers and 103 patients with knee pain who had visited our hospital between July 7, 2012, and July 24, 2018. These data were manually segmented by two authors (H.A. and A.H.) who had both trained as orthopedic surgeons for six years and who each had experience in the manual correction of over 200 knees. Segmentation data were converted by professional engineers (K.S. and J.M.) to train the neural network. To train the network to construct a region of interest (ROI) of the femoral subchondral bone and the medial/lateral tibial plateau, we manually segmented the ROI by using a reconstructed 3D knee model.

To validate our algorithm 108 subjects, among 113, were randomly selected for training. After training with 108 subjects, we computed the segmentation accuracy of another 5 subjects as one validation test. We performed the validation test three times, which selected another 108 subjects for training and 5 subjects for the test each time. After completing the three validation tests, the software was trained by all data collected from 113 subjects and was then used for the cross-sectional research of this study.

## Kanagawa Knee Study

This study was approved by the Medical Research Ethics Committee of Tokyo Medical and Dental University and written informed consent was obtained from all participants. The protocol was enrolled in a database of the National University Hospital Council of Japan (UMIN000032826) and disclosed. We mainly recruited workers or retirees from the Kanagawa Prefectural Office and named this study the “Kanagawa Knee Study.” The subjects included more than 50 females and 50 males between 30–79 years of age. Those who had a history of lower limb surgery or who were diagnosed with rheumatoid arthritis or cancer were excluded. We announced recruitment of these subjects at the Kanagawa Prefectural Government between September 1, 2018, and August 30, 2019. Participants joined our study voluntarily. We collected their heights, weights, knee radiographs, and MRIs between November 3, 2018, and September 28, 2019, at the AIC Yaesu clinic of Tokyo.

# Nomenclature and quantitative measurements for cartilage and the meniscus

To quantify cartilage, femoral (F) cartilage was radially projected and divided into three regions inside of the ROI of the femoral subchondral bone, femoral trochlea (TrF: anterior cartilage from proximal to intercondylar notch), medial femoral condyle (MF: medial cartilage from intercondylar notch to posterior), and lateral femoral condyle (LF: lateral cartilage from intercondylar notch to posterior) (Fig.1A). The tibial cartilage was vertically projected and divided into two areas inside the ROI of the medial/lateral tibial plateau, medial tibia (MT), and lateral tibia (LT) (Fig.1B).

Each area was divided into 3×3 subregions according to the previous report. From medial to lateral, the subregions in the TrF were named medial (m), central (c), and lateral (l). From proximal to distal, they were named proximal (p), intermediate (i), and distal (d) (Fig. 1A). From anterior to posterior, subregions in the other four parts (MF, LF, MT, and LT) were named anterior (a), medial (m), and posterior (p). From internal to external, they were named internal (i), central (c), and external (e) (Figs. 1A, 1B) [15].

Our software automatically computed cartilage average thickness (ThC), cartilage volume (VC), and projected cartilage area ratio (PCAR) in each region and subregion. Our software could also display the cartilage thickness mapping (Fig. 1C).

In this study, PCAR represented the ratio between the projected cartilage area and the total area of the ROI. A PCAR = 1.0 meant that the entire ROI of the subchondral bone was covered by cartilage. A PCAR = 0.0 meant that no cartilage covered the ROI of the subchondral bone. For example, a schematic diagram of the projected cartilage area (cartilage thickness  $\geq 1.0$  mm) is shown in Fig. 1D. As practical examples, the projected cartilage area (cartilage thickness  $\geq 0.0$  mm) is shown in Fig. 1E, and the projected cartilage area (cartilage thickness  $\geq 1.0$  mm) is shown in Fig. 1F. We evaluated PCAR for the threshold of cartilage thicknesses at  $\geq 0.0$  mm,  $\geq 0.5$  mm,  $\geq 1.0$  mm, and  $\geq 1.5$  mm. PCARs for the threshold of cartilage thicknesses at each of these measurements were named PCAR0.0, PCAR0.5, PCAR1.0, and PCAR1.5, respectively.

To evaluate the meniscus, medial and lateral meniscus volume, meniscus extrusion (ME) volume, ME area, and meniscus coverage ratio (MCR) were automatically computed. MCR was defined as the ratio between the overlapping area of the meniscus and the area of the ROI for the tibial plateau (Fig. 1G).

## Statistical analysis

To evaluate the accuracy of automatic segmentation, we calculated the Dice similarity coefficient (DSC) between manual segmentation and automatic segmentation [16]. In each validation test, DSC was computed for five test subjects at the femoral bone, tibial bone, femoral cartilage, tibial cartilage, medial/lateral meniscus, femoral subchondral bone ROI, and medial/lateral tibia plateau ROI. After the three validation tests, we calculated the mean DSC of each test.

To compare gender differences in the quantification of cartilage and the meniscus, we used the Mann–Whitney U test. To evaluate the correlation between each quantitative value and age or other quantitative value, we used Spearman’s rank correlation test. All statistical analyses were done using JMP® 14 (SAS Institute Inc., Cary, NC, USA). P values < 0.05 were considered statistically significant.

## Results

### Segmentation accuracy

The mean Dice similarity coefficient (DSC) of all tests combined was 0.985 for the femoral bone, 0.980 for the tibial bone, 0.911 for the femoral cartilage, 0.892 for the tibial cartilage, 0.916 for the medial meniscus, 0.891 for the lateral meniscus, 0.905 for the ROI of the femoral subchondral bone, and 0.888 for the ROI of the medial/lateral tibia plateau (Table 1). Our algorithm provided high segmentation accuracy.

Table 1  
Automatic segmentation accuracy.

Subregions	DSC (average ± SD)			
	1st test	2nd test	3rd test	Total
Femoral bone	0.985 ± 0.004	0.985 ± 0.007	0.983 ± 0.004	0.985 ± 0.005
Tibial bone	0.981 ± 0.008	0.978 ± 0.013	0.983 ± 0.002	0.980 ± 0.009
Femoral cartilage	0.905 ± 0.024	0.913 ± 0.034	0.916 ± 0.013	0.911 ± 0.024
Tibial cartilage	0.893 ± 0.072	0.868 ± 0.119	0.914 ± 0.009	0.892 ± 0.072
Medial meniscus	0.921 ± 0.012	0.906 ± 0.020	0.922 ± 0.014	0.916 ± 0.016
Lateral meniscus	0.893 ± 0.038	0.905 ± 0.033	0.875 ± 0.053	0.891 ± 0.041
ROI of femoral subchondral bone	0.905 ± 0.016	0.904 ± 0.019	0.906 ± 0.017	0.905 ± 0.016
ROI of medial/lateral tibia plateau	0.889 ± 0.036	0.884 ± 0.055	0.892 ± 0.022	0.888 ± 0.037
*DSC averages and SDs were calculated for the first to third tests with 5 subject data, for a total of 15 subject data.				

### Characteristics of study subjects in the cross-sectional study

This study included 561 subjects: 276 females and 285 males (Table 2). The overall average age of the subjects was 53.7 ± 13.9 years for females and 55.2 ± 11.1 years for males. The body mass index was 22.5 ± 3.0 for females and 23.6 ± 3.3 for males. The rate of grade 3 or 4 on the Kellgren–Lawrence OA scale was 1% for both females and males.

Table 2  
Characteristics of the study subjects.

Characteristic	Female	Male
Total	n = 276	n = 285
Age	53.7 ± 13.9	55.2 ± 11.1
30–39 years	n = 56	n = 52
40–49 years	n = 61	n = 62
50–59 years	n = 52	n = 51
60–69 years	n = 55	n = 57
70–79 years	n = 52	n = 63
Height: means ± SD (cm)	158.6 ± 7.2	168.4 ± 6.3
Weight: means ± SD (kg)	56.7 ± 9.2	66.9 ± 11.2
Body mass index: means ± SD	22.5 ± 3.0	23.6 ± 3.3
Grade 3 or 4 on the Kellgren–Lawrence OA scale	n = 2 (1%)	n = 4 (1%)

## Gender Differences In Quantification Of Cartilage And The Meniscus

The cartilage volume and thickness were significantly higher in males than in females in the MF, MT, LF, and LT (Table 3). PCAR was higher in males than females except for MT PCAR1.0 and LF PCAR0.0, 0.5, 1.0. The volume and area of the medial/lateral meniscus were higher in males than in females.

Table 3  
Cartilage and meniscus quantification at medial compartment in both genders.

	Female	Male	p value
MF VC (mm <sup>3</sup> )	2503 ± 416	3349 ± 531	< 0.0001
MF ThC (mm)	1.83 ± 0.25	2.02 ± 0.22	< 0.0001
MF PCAR0.0	0.99 ± 0.05	0.99 ± 0.01	0.01
MF PCAR0.5	0.99 ± 0.05	0.99 ± 0.01	0.01
MF PCAR1.0	0.98 ± 0.06	0.99 ± 0.01	0.002
MF PCAR1.5	0.85 ± 0.10	0.88 ± 0.07	0.001
LF VC (mm <sup>3</sup> )	2274 ± 370	3297 ± 492	< 0.0001
LF ThC (mm)	1.72 ± 0.23	1.94 ± 0.20	< 0.0001
LF PCAR0.0	0.99 ± 0.03	0.99 ± 0.04	0.90
LF PCAR0.5	0.99 ± 0.03	0.99 ± 0.00	0.99
LF PCAR1.0	0.99 ± 0.03	0.99 ± 0.01	0.32
LF PCAR1.5	0.84 ± 0.12	0.88 ± 0.08	< 0.0001
MT VC (mm <sup>3</sup> )	1286 ± 227	1872 ± 334	< 0.0001
MT ThC (mm)	1.75 ± 0.24	1.93 ± 0.25	< 0.0001
MT PCAR0.0	0.99 ± 0.06	0.99 ± 0.01	< 0.0001
MT PCAR0.5	0.99 ± 0.05	0.99 ± 0.01	< 0.0001
MT PCAR1.0	0.95 ± 0.06	0.96 ± 0.03	0.53
MT PCAR1.5	0.71 ± 0.19	0.78 ± 0.13	< 0.0001
LT VC (mm <sup>3</sup> )	1537 ± 309	2203 ± 383	< 0.0001
LT ThC (mm)	2.20 ± 0.37	2.44 ± 0.32	< 0.0001
LT PCAR0.0	0.96 ± 0.05	0.97 ± 0.02	< 0.0001
LT PCAR0.5	0.96 ± 0.05	0.97 ± 0.03	< 0.0001
LT PCAR1.0	0.92 ± 0.06	0.94 ± 0.04	< 0.0001
LT PCAR1.5	0.79 ± 0.11	0.83 ± 0.08	< 0.0001
MM Volume (mm <sup>3</sup> )	2179 ± 491	2974 ± 567	< 0.0001

	Female	Male	p value
MM Area (mm <sup>2</sup> )	655 ± 88	834 ± 99	< 0.0001
MME Volume (mm <sup>3</sup> )	300 ± 286	351 ± 217	0.0004
MME Area (mm <sup>2</sup> )	75 ± 54	83 ± 39	0.001
MMCR	0.49 ± 0.09	0.52 ± 0.07	0.003
LM Volume (mm <sup>3</sup> )	2184 ± 480	2820 ± 475	< 0.0001
LM Area (mm <sup>2</sup> )	651 ± 82	818 ± 83	< 0.0001
LME Volume (mm <sup>3</sup> )	25 ± 60	14 ± 34	0.01
LME Area (mm <sup>2</sup> )	6.39 ± 13.3	3.49 ± 7.57	0.002
LMCR	0.66 ± 0.08	0.67 ± 0.06	0.28

## Correlation Between Cartilage Quantifications And Body Size

Thickness and volume of femoral and tibial cartilage in each region were correlated with height in both genders (Supplementary Table 1). On the other hand, the MF PCAR, LF PCAR, MT PCAR, and LT PCAR0.0-1.5 were not correlated with height. Even if they were correlated, their  $r_s$  were less than 0.2 in both genders. There were no correlations between cartilage quantifications (CV, ThC, and PCAR) and weight (data not shown).

## Correlation Between Cartilage Quantification And Age

Among the nine subregions in the MF, MT, LF, and LT, we focused on the medial central subregion because it was considered more representative than other subregions. In females, among 48 cartilage quantifications (including PCAR0.0, 0.5, 1.0, 1.5, VC, and ThC for the MF, LF, MT, LT, mcMF, mcLF, mcMT, and mcLT), the highest absolute value of the correlation coefficient with age was mcMT PCAR1.0 ( $r_s = -0.34$ ) (Table 4). In males, the highest absolute value of the correlation coefficient with age was mcMT ThC ( $r_s = -0.35$ ). The second highest correlation coefficient with age was mcMT PCAR1.5 ( $r_s = -0.33$ ).

Table 4  
Correlation of PCAR, VC, and ThC with age.

Regions & subregions	rs (p value)					
	PCAR0.0	PCAR0.5	PCAR1.0	PCAR1.5	VC	ThC
Female						
MF	-0.14 (0.019)	-0.14 (0.025)	-0.06 (0.34)	-0.07 (0.27)	-0.01 (0.84)	0.02 (0.70)
LF	-0.23 (0.0001)	-0.21 (0.0005)	-0.11 (0.072)	0.07 (0.26)	-0.02 (0.69)	0.02 (0.76)
MT	-0.06 (0.30)	-0.07 (0.242)	-0.11 (0.080)	-0.16 (0.041)	-0.03 (0.59)	-0.12 (0.042)
LT	0.15 (0.013)	0.15 (0.016)	0.06 (0.30)	-0.06 (0.34)	-0.15 (0.0012)	-0.19 (0.012)
mcMF	-0.21 (0.0004)	-0.21 (0.0004)	-0.06 (0.30)	-0.06 (0.36)	0.03 (0.66)	0.05 (0.42)
mcLF	0.15 (0.014)	0.15 (0.014)	0.06 (0.35)	-0.03 (0.58)	-0.12 (0.054)	-0.04 (0.52)
mcMT	-0.21 (0.0004)	-0.21 (0.0004)	-0.34 (< 0.0001)	-0.28 (< 0.0001)	-0.16 (0.0063)	-0.25 (< 0.0001)
mcLT	-0.12 (0.042)	-0.12 (0.042)	-0.13 (0.039)	-0.15 (0.016)	-0.12 (0.055)	-0.13 (0.038)
Male						
MF	-0.26 (< 0.0001)	-0.25 (< 0.0001)	-0.16 (0.0063)	-0.02 (0.71)	-0.09 (0.127)	-0.08 (0.206)
LF	-0.08 (0.19)	-0.08 (0.18)	-0.04 (0.50)	0.06 (0.31)	-0.10 (0.10)	-0.09 (0.13)
MT	-0.11 (0.066)	-0.11 (0.069)	-0.13 (0.028)	-0.17 (0.0034)	-0.15 (0.014)	-0.22 (0.0002)
LT	-0.02 (0.78)	-0.02 (0.76)	-0.02 (0.70)	-0.09 (0.12)	-0.14 (0.019)	-0.20 (0.0007)
mcMF	-0.10 (0.085)	-0.05 (0.38)	-0.05 (0.36)	-0.02 (0.73)	-0.10 (0.097)	-0.12 (0.045)
mcLF	-0.10 (0.094)	-0.10 (0.094)	-0.09 (0.15)	-0.09 (0.14)	-0.19 (0.0011)	-0.18 (0.0019)
mcMT	-0.13 (0.025)	-0.13 (0.025)	-0.18 (0.0018)	-0.33 (< 0.0001)	-0.25 (< 0.0001)	-0.35 (< 0.0001)

-: The correlation coefficient could not be measured because they were all the same and had no variance.

Regions &	rs (p value)					
mcLT	-	-	-	-0.06 (0.33)	-0.07 (0.22)	-0.10 (0.081)
-: The correlation coefficient could not be measured because they were all the same and had no variance.						

## Correlation Between The Quantification Of Meniscus Extrusion And Age

In females, medial meniscus extrusion (MME) area and MME volume were correlated with age. The MM coverage ratio was not correlated with age (Table 5). In males, there were no correlations between the quantification of meniscus extrusion (MME area, MME volume, and MMCR) and age.

Table 5  
Correlation of age and mcMT with quantification of MM extrusion.

	rs (p value)				
	Age	mcMT PCAR0.0	mcMT PCAR0.5	mcMT PCAR1.0	mcMT PCAR1.5
Female					
MME Area	0.15 (0.012)	-0.32 (< 0.0001)	-0.32 (< 0.0001)	-0.19 (0.0017)	-0.05 (0.38)
MME Volume	0.15 (0.013)	-0.32 (< 0.0001)	-0.32 (< 0.0001)	-0.18 (0.0022)	-0.02 (0.67)
MMCR	0.001 (0.987)	0.30 (< 0.0001)	0.30 (< 0.0001)	0.25 (< 0.0001)	0.28 (< 0.0001)
Male					
MME Area	-0.09 (0.066)	-0.18 (0.0029)	-0.17 (0.0036)	-0.17 (0.0036)	-0.01 (0.82)
MME Volume	-0.11 (0.12)	-0.18 (0.0029)	-0.18 (0.0022)	-0.18 (0.0029)	-0.01 (0.90)
MMCR	0.03 (0.62)	0.17 (0.0033)	0.11 (0.057)	0.13 (0.033)	0.04 (0.54)

### Correlation between projected cartilage area ratio (PCAR) and meniscus extrusion

In females, mcMT PCAR0.0, 0.5, and 1.0 were significantly correlated with MME area, MME volume, and MMCR (Table 5). In particular, the absolute value of rs with mcMT PCAR0.0 and 0.5 exceeded 0.3. In males, mcMT PCAR0.0, 0.5, and 1.0 were significantly correlated with MME area and MME volume.

mcMT PCAR0.0 and 1.0 were also significantly correlated with MMCR. However, the absolute value of  $r_s$  between them was less than 0.2.

## Discussion

Our software provided high automatic segmentation accuracy for cartilage and the meniscus. The cartilage and meniscus volumes, and most PCARs, were higher in males than in females. Thickness and volume of femoral and tibial cartilage in each region were correlated with height, while MF PCAR0.0-1.5 and MT PCAR0.0-1.5 were not correlated in both genders. Among the 48 cartilage quantifications, the highest absolute values of the correlation coefficient with age were mcMT PCAR1.0 in females and mcMT ThC in males. In females, mcMT PCAR0.0-1.0 was correlated with MMCR, although MMCR was not correlated with age. In males, there were no correlations between the quantification of meniscus extrusion and age.

For segmentation accuracy, DSCs for bone and cartilage were higher in our algorithm than in previous reports [5–8]. This was possibly because we used 3D U-Net, a learning algorithm specialized for 3D images. DSCs for medial and lateral menisci were also higher in our algorithm than in previous reports [7, 8]. This was probably due to the use of an additional PDWI sequence for meniscus segmentation. It is often difficult to determine meniscus segmentation using the SPGR sequence alone.

Among 561 volunteers in our cross-sectional study, 20 had visited hospitals for more than three months due to their knee symptoms for the right knee only, 15 had visited for more than three months for the left knee only, and 6 had visited for more than three months for both knees. This means that 25 volunteers (4.4%) had ever visited hospitals due to their knee symptoms. On the other hand, those who had radiographic osteoarthritis at grade 3 or 4 on the Kellgren–Lawrence scale included 6 volunteers (1.1%). These data indicate that most of the subjects in this cross-sectional study were healthy.

Gender differences related to cartilage thickness is still a controversial concept. In our current study, the MT ThC was  $1.75 \pm 0.24$  mm in females ( $n = 276$ ) and  $1.93 \pm 0.25$  mm in males ( $n = 285$ ), meaning the MT ThC was significantly higher in females than in males ( $p < 0.0001$ ). On the other hand, according to a 3D MRI study by Faber et al., the MT ThC was  $1.20 \pm 0.19$  mm in females ( $n = 8$ ) and  $1.36 \pm 0.15$  mm in males ( $n = 9$ ), meaning there was no significant difference between females and males [17]. The subjects in the study by Faber et al. were young, healthy volunteers (age  $22 \pm 2$  years) and MRIs were performed with a 1.5 T fat-suppressed 3D gradient-echo sequence. The significant gender difference in cartilage thickness in these studies depended on the ages of the subjects and the imaging sequences, but the difference in sample number will greatly be affected. In our study, as is true of most quantifications of cartilage, quantifications of the meniscus (including meniscus coverage ratio) were significantly higher in males than in females. For analysis of cartilage and meniscus pathology, we should take gender differences into consideration.

In this study, thickness and volume of femoral and tibial cartilage in each region were correlated with height. This means that to accurately determine the relationship between these quantifications and age, it

is necessary to correct these quantifications by considering the heights of each individual. On the other hand, MF PCAR0.0-1.5 and MT PCAR0.0-1.5 were not correlated with height. Therefore, PCARs are useful cartilage quantifications for examining the relationship to age since they do not need to be corrected for height.

Among the 48 cartilage quantifications in this study, the highest absolute value of the correlation coefficient with age was mcMT PCAR1.0 ( $r_s = -0.34$ ) in females. In other words, in females, mcMT PCAR1.0 was more sensitive than thickness and volume at each region and subregion for age-related cartilage quantifications. Also, among mcMT PCAR0.0, 0.5, 1.0, 1.5, PCAR1.0 was the best for detecting age-related cartilage quantifications in this population. In males, the highest absolute value of the correlation coefficient with age was mcMT ThC ( $r_s = -0.35$ ), and the second highest was mcMT PCAR1.5 ( $r_s = -0.33$ ). Though MT ThC was significantly correlated with height, mcMT ThC without correction for height was correlated with age. Among mcMT PCAR0.0, 0.5, 1.0, 1.5, PCAR1.0 was highest in females, but PCAR1.5 was highest in males. This is due to the fact that cartilage is thicker in males than in females.

We also showed that MM extrusion area and volume were correlated with age in females. Achtnich et al. reported, using ultrasound examination, that the width of MM extrusion increased with age [18], and this was similar to our results, though the subjects in the Achtnich et al. study included 17 males in addition to 58 females. In our study, MMCR, which is considered an index that reflects meniscus function more directly than the MME area and MME volume, was not correlated with age in both genders. On the other hand, MMCR was associated with a decrease in mcMT PCAR, which was associated with age in both genders. These data mean that decreases in MMCR possibly promote decreases in MT cartilage over normal aging. At present, MRI is not recommended for early detection of osteoarthritis due to a lack of validated consensus [19]. Our findings suggest that mcMT PCAR is a useful quantification for detecting early knee osteoarthritis and that MMCR is an important risk factor.

In this study, we propose three limitations. First, we only recruited subjects who were interested in participating. Therefore, the composition of the subjects did not match that of the general population, and this selection bias cannot be ignored. Second, though we obtained the cartilage quantification of nine subregions in the MF, MT, LF, and LT, we only focused on the medial central subregion because it was considered more representative than other subregions. However, this was not fully investigated. Third, we examined PCAR0.0, PCAR0.5, PCAR1.0, and PCAR1.5. Among these PCARs, the thickness of the cartilage was not fixed but varied according to several uses. As an advantage, PCAR can detect small differences by adjusting cartilage thickness, but this may reduce the reliability of this study.

## Conclusion

Our 3D U-Net algorithm showed high segmentation accuracy for knee cartilage and menisci. In our cross-sectional study, the absolute value of the correlation coefficient with age was high in females (mcMT PCAR1.0) and males (mcMT PCAR1.5). Though MMCR was not correlated with age, MMCR was

correlated with mcMT PCAR1.0. Our software provided numerous quantifications of cartilage related to age and meniscus extrusion.

## Abbreviations

3D MRI: Three-dimensional magnetic resonance imaging, PCAR: Projected cartilage area ratio, ThC: cartilage average thickness, VC: cartilage volume, ROI: region of interest, ME: meniscus extrusion, MME: medial meniscus extrusion, MCR: meniscus coverage ratio, MMCR: medial meniscus coverage ratio, SPGR: spoiled gradient echo, PDWI: proton density weighted imaging, TrF: femoral trochlea, MF: medial femoral condyle, LF: lateral femoral condyle, MT: medial tibia, LT: lateral tibia, DSC: Dice similarity coefficient, mc: medial central subregion

## Declarations

### Ethics approval and consent to participate

This study was approved by the Medical Research Ethics Committee of Tokyo Medical and Dental University and written informed consent was obtained from all participants. The protocol was enrolled in a database of the National University Hospital Council of Japan (UMIN000031924, UMIN000032826) and disclosed.

### Consent for publication

This manuscript does not contain any individual data.

### Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

### Competing interests

The authors declare that they have no competing interests

### Funding

This study was funded by Fujifim.

### Authors' contributions

Study conception and design: Aoki, Ozeki, Hyodo, Suzuki, Masumoto, and Sekiya. Acquisition of data: Aoki, Takanashi, Katano, Okanouchi, and Sekiya. Analysis and interpretation of data: Aoki, Ozeki, Matsuda, Fujiwara, and Sekiya.

## Acknowledgements

This research was supported by Fujifilm. This manuscript was proofread by Jason Stillings.

## References

1. Buck RJ, Wyman BT, Le Graverand MP, Hudelmaier M, Wirth W, Eckstein F, investigators A: **Osteoarthritis may not be a one-way-road of cartilage loss– comparison of spatial patterns of cartilage change between osteoarthritic and healthy knees.** *Osteoarthritis Cartilage* 2010, **18**(3):329–335.
2. Eckstein F, Peterfy C. A 20 years of progress and future of quantitative magnetic resonance imaging (qMRI) of cartilage and articular tissues-personal perspective. *Semin Arthritis Rheum.* 2016;**45**(6):639–47.
3. Okazaki Y, Furumatsu T, Yamaguchi T, Kodama Y, Kamatsuki Y, Masuda S, Hiranaka T, Zhang X, Ozaki T. **Medial meniscus posterior root tear causes swelling of the medial meniscus and expansion of the extruded meniscus: a comparative analysis between 2D and 3D MRI.** *Knee Surg Sports Traumatol Arthrosc* 2019.
4. Berthiaume MJ, Raynauld JP, Martel-Pelletier J, Labonté F, Beaudoin G, Bloch DA, Choquette D, Haraoui B, Altman RD, Hochberg M, et al. Meniscal tear and extrusion are strongly associated with progression of symptomatic knee osteoarthritis as assessed by quantitative magnetic resonance imaging. *Ann Rheum Dis.* 2005;**64**(4):556–63.
5. Ambellan F, Tack A, Ehlke M, Zachow S. Automated segmentation of knee bone and cartilage combining statistical shape knowledge and convolutional neural networks: Data from the Osteoarthritis Initiative. *Med Image Anal.* 2019;**52**:109–18.
6. Liu F, Zhou Z, Samsonov A, Blankenbaker D, Larison W, Kanarek A, Lian K, Kambhampati S, Kijowski R. Deep Learning Approach for Evaluating Knee MR Images: Achieving High Diagnostic Performance for Cartilage Lesion Detection. *Radiology.* 2018;**289**(1):160–9.
7. Norman B, Pedoia V, Majumdar S. Use of 2D U-Net Convolutional Neural Networks for Automated Cartilage and Meniscus Segmentation of Knee MR Imaging Data to Determine Relaxometry and Morphometry. *Radiology.* 2018;**288**(1):177–85.
8. Tack A, Mukhopadhyay A, Zachow S. Knee menisci segmentation using convolutional neural networks: data from the Osteoarthritis Initiative. *Osteoarthritis Cartilage.* 2018;**26**(5):680–8.
9. Costa CR, Morrison WB, Carrino JA. Medial meniscus extrusion on knee MRI: is extent associated with severity of degeneration or type of tear? *AJR Am J Roentgenol.* 2004;**183**(1):17–23.
10. Lawrence RC, Felson DT, Helmick CG, Arnold LM, Choi H, Deyo RA, Gabriel S, Hirsch R, Hochberg MC, Hunder GG, et al. Estimates of the prevalence of arthritis and other rheumatic conditions in the United States. Part II. *Arthritis Rheum.* 2008;**58**(1):26–35.
11. Jordan JM, Helmick CG, Renner JB, Luta G, Dragomir AD, Woodard J, Fang F, Schwartz TA, Abbate LM, Callahan LF, et al. Prevalence of knee symptoms and radiographic and symptomatic knee

- osteoarthritis in African Americans and Caucasians: the Johnston County Osteoarthritis Project. *J Rheumatol*. 2007;34(1):172–80.
12. Yoshimura N, Muraki S, Oka H, Mabuchi A, En-Yo Y, Yoshida M, Saika A, Yoshida H, Suzuki T, Yamamoto S, et al. Prevalence of knee osteoarthritis, lumbar spondylosis, and osteoporosis in Japanese men and women: the research on osteoarthritis/osteoporosis against disability study. *J Bone Miner Metab*. 2009;27(5):620–8.
  13. Ronneberger O, Fischer P, Brox T: **U-net: Convolutional networks for biomedical image segmentation**. In: *International Conference on Medical image computing and computer-assisted intervention: 2015*. Springer; 2015: 234–241.
  14. Anbeek P, Vincken KL, van Osch MJ, Bisschops RH, van der Grond J. Probabilistic segmentation of white matter lesions in MR imaging. *Neuroimage*. 2004;21(3):1037–44.
  15. Eckstein F, Ateshian G, Burgkart R, Burstein D, Cicuttini F, Dardzinski B, Gray M, Link TM, Majumdar S, Mosher T, et al. Proposal for a nomenclature for magnetic resonance imaging based measures of articular cartilage in osteoarthritis. *Osteoarthritis Cartilage*. 2006;14(10):974–83.
  16. Zou KH, Warfield SK, Bharatha A, Tempany CM, Kaus MR, Haker SJ, Wells WM, Jolesz FA, Kikinis R. Statistical validation of image segmentation quality based on a spatial overlap index. *Acad Radiol*. 2004;11(2):178–89.
  17. Faber SC, Eckstein F, Lukasz S, Mühlbauer R, Hohe J, Englmeier KH, Reiser M. Gender differences in knee joint cartilage thickness, volume and articular surface areas: assessment with quantitative three-dimensional MR imaging. *Skeletal Radiol*. 2001;30(3):144–50.
  18. Achtnich A, Petersen W, Willinger L, Sauter A, Rasper M, Wörtler K, Imhoff AB, Diermeier T. Medial meniscus extrusion increases with age and BMI and is depending on different loading conditions. *Knee Surg Sports Traumatol Arthrosc*. 2018;26(8):2282–8.
  19. Luyten FP, Bierma-Zeinstra S, Dell'Accio F, Kraus VB, Nakata K, Sekiya I, Arden NK, Lohmander LS. Toward classification criteria for early osteoarthritis of the knee. *Semin Arthritis Rheum*. 2018;47(4):457–63.

## Figures

Fig 1.




Figure 1

Projected cartilage area and meniscus coverage ratio. (A) Subregions of femoral cartilage. (B) Subregions of tibial cartilage. (C) Cartilage thickness mapping in the medial tibia. (D) Schematic diagram of the projected cartilage area (cartilage thickness  $\geq 1.0$  mm). (E) Practical example of the projected cartilage area (cartilage thickness  $\geq 0.0$  mm) in the medial tibia. (F) Practical example of the projected cartilage

area (cartilage thickness  $\geq 1.0$  mm) in the medial tibia. (G) Schematic diagram of the meniscus coverage ratio.

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [AokiKanagawasupplementarytable.pdf](#)
- [supplementaryfigure1.TIF](#)
- [EthicsapprovalAoki2.pdf](#)
- [EthicsapprovalAoki1.pdf](#)