ARABIDOPSIS POLLEN ABORTION MUTANT 2, Encoding an Unknown Protein with Three WD40-Repeats, Is Essential for Late Pollen Mature in *Arabidopsis thaliana*

Zhaoxia Ma

Kunming University

Xueqin Zhang

China Agricultural University

Liqun Chen

China Agricultural University

Guobin Zhang

Kunming University

Yue Liu

China Agricultural University

De Ye

China Agricultural University

Hongyu Yang (✉ yanghongyukm@126.com)

Kunming University <https://orcid.org/0000-0001-5778-5396>

Research article

Keywords: Arabidopsis thaliana, AtAPAM2, late pollen mature, WD40 protein

Posted Date: July 23rd, 2019

DOI: <https://doi.org/10.21203/rs.2.11828/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background In flowering plants, pollen formation is a very complex process. It is strictly regulated by various genetic factors. Some of these factors have been identified, but the regulatory mechanism concerning the process of late pollen maturing remains unknown. **Results** Previously we identified and reported an ARABIDOPSIS POLLEN ABORTION MUTANT 2 (APAM2) gene which was required for pollen formation in *Arabidopsis thaliana*. In this study, we further analysed the phenotype of *atapam2* and the subcellular localisation of AtAPAM2. Mutation in AtAPAM2 drastically affected male gametophytic function, impacted the late pollen mature process, and resulted in mature pollen grains from *atapam2/+* plants being shrivelled and dead. It was expressed among seedlings, roots, stems, leaves, siliques, flowers, and pollen grains. However, its expression levels in flowers and pollen grains were much higher than in other tissues. Gene AtAPAM2 encodes an unknown protein with three WD40-repeats with localisation in plasma membrane and nucleus, and homologous proteins exist in many species. **Conclusions** We determined that AtAPAM2 may play a role in pollen formation and other development processes by interacting with other proteins, but we did not identify the protein that interacts with it.

Background

The WD40 domain is one of the most abundant domains in eukaryotic genomes, and consists of 44–60 amino acid (aa) **residues** with a glycine–histidine di-peptide close to the N-terminal and a tryptophan–aspartic acid (WD) di-peptide at the C-terminal end [1]. When present in a protein, the WD40 motif is typically found as several (usually 4–10) tandem repeat units and they usually function as a protein–protein or protein–DNA/RNA interaction platform [2, 3, 4]. The first identified WD40 domain was the β -subunit of heterotrimeric G protein. The WD40 repeats form four-antiparallel β -sheets which in turn fold into a higher-level structure denoted the β -propeller. At least four repeats are believed to be required to form a β -propeller. In G β , which contains seven WD40-repeats, the first and last repeats participate in the same propeller blade, potentially reinforcing the structure [5, 6].

In the *Arabidopsis thaliana* genome, 269 proteins containing at least one copy of the WD40 motif were identified, and of these 237 contain four or more recognisable copies. These 237 proteins can be classified into 143 distinct families, and are components of basic cellular mechanisms [2].

The WD40 proteins are involved in a variety of cellular functions; for example, AtNEDD1, a WD40 repeat protein is associated with animal NEDD1/GCP-WD protein, which interacts with the γ -tubulin complex and participates in microtubule organisation during cell division [7]. The WD40 protein TRANSPARENT TESTA GLABRA1 (TTG1) regulates several developmental and biochemical pathways in *Arabidopsis*, including formation of hairs on leaves, stems, and roots, and production of seed mucilage and anthocyanin pigments [8]. The WD40 proteins also participate in the progress of gametophytic development, and several have been identified. Gene *SLOW WALKER1* (*SWA1*) encodes a protein with six WD40 repeats that localises in the nucleolus in interphase cells, which are involved in 18S ribosomal RNA biogenesis, and is essential for gametogenesis in *Arabidopsis* [9]. The *Arabidopsis* Yaozhe (YAO) is a nucleolar WD40

protein involved in embryogenesis and gametogenesis [10]. Rice Immature Pollen1RIP1), containing five WD40 repeats, regulates late pollen development in rice [11]. The *OsLIS-L1* gene encoding the lissencephaly type-1-like protein with WD40 repeats is involved in plant height and male gametophyte formation in rice [12]. The *LuWD40-1* gene encoding a WD40 repeat protein regulates growth and pollen viability in flax (*Linum usitatissimum* L.) [13]. The *JINGUBANG* gene encodes a putative hydrophilic protein containing seven WD40 repeats, which is a negative regulator of pollen germination and prevents pollination in moist environments [14].

In recent years, although a number of WD40 proteins have been identified, their functions are little known. Yang et al. isolated four gametophytic male-sterile mutants (*apam1–apam4*) in *Arabidopsis* [15]. Mutation of *AtAPAM2* (*ARABIDOPSIS POLLEN ABORTION MUTANT 2*) drastically affected male gametophytic function and the *Ds* element was inserted in the 18th intron of *At3g50590* [15]. In this study, we report the characterisation of *atapam2*. Gene *AtAPAM2* is constitutively expressed including in seedlings, roots, stems, leaves, siliques, flowers, and pollen grains. It encodes an unknown protein with three WD40-repeats that localises in plasma membrane and the nucleus. Homologous proteins of *AtAPAM2* exist in many species, and we speculate that this protein may play a role in pollen formation and other development processes by interacting with other proteins.

Results

The *atapam2/+* Mutant is Defective in Mature Pollen Grains

We crossed the *atapam2/+* plant with the *quartet1* (*qrt1*) mutant to obtain the *atapam2/+; qrt1/qrt1* plants for tetrad analysis. The *qrt1* mutant caused the pollen grains not to separate from the tetrad, but had little effect on growth of the pollen tube [16]. Therefore, the *atapam2/+; qrt1/qrt1* plants produced a tetrad consisting of two *qrt1* (representing wild type) pollen grains and two *atapam2* pollen grains. Scanning electron microscopy (SEM) showed that all four pollen grains in the tetrad from *qrt1* plants had normal morphology (Fig. 1a), yet the *atapam2/+; qrt1/qrt1* plants had only two normal pollen grains (Fig. 1b). Alexander staining (Fig. 1c and 1d) and 4',6-diamidino-2-phenylindole (DAPI) staining showed the same results (Fig. 1e–1h). These results showed that the *AtAPAM2* mutation affected male gametophytic function.

Because mature pollen grains from *atapam2/+* were shrivelled and had no vitality, we were interested in how the mutation influenced development of pollen grains. The DAPI staining was used to investigate at which developmental stage pollen became abnormal (Fig. 2). Before the 12th anther development stage occurred, the developing nuclei of *atapam2/+* pollen grains were of normal phenotype, as were wild-type pollen grains (Fig. 2a–2c and 2k–2m). However, when anthers developed to the 12th stage in which pollen grains had developed three nuclei, the pollen grains began to turn abnormal (Fig. 2d and 2n). The mutant grains were smaller than normal pollen grains in bright field image and their nuclei began to

disappear under UV-light (Fig. 2d and 2n). By the 13th stage, the nuclei of mutant pollen grains had disappeared completely (Fig. 2e and 2o).

To further determine the results of DAPI staining, the different development stages of buds were embedded in resin. Then we obtained transverse sections of anthers from wild-type and *atapam2/+* plants using the semi-thin section method (Fig. 3). The results showed that in the 11th stage of anther development or before, the appearance and size of pollen grains from mutants were the same as for wild type (Fig. 3a–3d). However, in the 12th stage of anther development, anthers were cracking and pollen grains were shed from anthers. Transverse sections of anthers showed that some pollen grains were of irregular appearance and were not as smooth as the wild type (Fig. 3e and 3f). Combining the results of DAPI staining with semi-thin sections, we conclude that pollen grains of *atapam2/+* mutant began to turn abnormal at the 12th anther development stage.

Next, we used transmission electron microscopy (TEM) to observe differences in pollen wall structure between wild-type and *atapam2* pollen (Fig. 4). Mature *atapam2* pollen had characteristic wrinkled intine (Fig. 4c–4e), which was smooth in wild type and closely connected to the inner side of nexine (Fig. 4a and 4b). Mutant pollen also had deformed cytoplasm (Fig. 4c–4e).

At the 11th stage of anther development, pollen grains from *atapam2/+* mutants were normal, with two sperm cell nuclei and a vegetative nucleus, suggesting that pollen grains had finished the second mitosis process. But at the 12th stage, pollen grains from *atapam2/+* mutant began to turn abnormal. During this stage, the three nuclei pollen grains completed the process of dehydration and the mutation may have affected this process.

The Phenotype of *atapam2/+* is Caused by a *Ds* Insertion in *AtAPAM2*

In a previous study, thermal asymmetric interlaced PCR was used to localise the mutation site in *atapam2*. Sequence analysis indicated that the *Ds* element was inserted in the 18th intron of *At3g50590* [15]. To confirm that the *atapam2/+* mutant phenotype was caused by the *Ds* insertion in this gene, we subcloned the full-length genomic DNA fragment of *At3g50590*, including the promoter and 5' and 3' UTRs into the pCambia1300 vector. Then, this construct was introduced into *atapam2/+* mutant plants. Transgenic lines were screened by MS culture medium containing kanamycin and hygromycin. The self-pollinated T₁ transgenic seeds were plated on MS culture medium containing kanamycin to determine the segregation ratio of *Ds*. The Kan^R:Kan^S segregation ratio was approximately 2:1 for most of the transformant lines (19 of 25). The complemented mutant plants produced offspring of *atapam2/+* mutation homozygotes in T₂ and subsequent generations. When pollen grains from the two independent complemented lines homozygous for the *atapam2/+* mutation were observed by SEM, Alexander staining assay, and DAPI staining assay, the configuration of surface, vitality, and nuclei development were

returned to the level of wild type (Fig. 5c, 5f, 5i, and 5l). At the same time, we [calculated](#) the [abnormal](#) rate of pollen grains from the two complemented lines, and found about 1.28% (8/624) and 1.42% (10/700) of pollen grains were abnormal, [respectively](#) (Fig. 5m). These results demonstrate that the defect in male gametophytic function in *atapam2/+* plants was fully complemented by *At3g50590* full-length genomic DNA.

To validate this result, *atapam2-2* (SAIL_1288_C09) was [obtained](#) from the seed stocks of the Arabidopsis Biological Resource Center (ABRC). The *atapam2-2* is a T-DNA insertion mutant of [Columbia](#) (Col) background. Using a T-DNA left-border primer Lba1 and a gene-specific primers (*atapam2-2-F* and *atapam2-2-R*) (Table 1) for PCR analysis, we confirmed that T-DNA was inserted in the 13th exon (Fig. 8a). The *atapam2-2* mutant carried a Basta^R-selective marker. The self-pollinated *atapam2-2/+* progeny exhibited a Basta^R:Basta^S segregation ratio of approximately 1:1 (1590:1500) (Table 2) rather than 3:1. We also used SEM to [observe](#) the [surface](#) of pollen grains from *atapam2-2/+*, and found about 50% (134/271) of pollen grains were [shrivelled](#) (data not shown). These findings are consistent with previous results concerning *atapam2/+* plants [15], and confirm that the defect in the male gametophyte in *atapam2/+* was caused by mutation in *At3g50590*.

***AtAPAM2* is Constitutively Expressed and Highly Expressed in Flowers and Pollen Grains**

To understand the function of *AtAPAM2*, we used RT-PCR to assess its expression pattern. The RNA was isolated from various wild-type plant tissues including seedlings, roots, stems, leaves, [inflorescences](#), and siliques. Then the RNA was [reverse-transcribed](#) to cDNA. The RT-PCR analyses showed that *AtAPAM2* was ubiquitously expressed in many tissues, was constitutively expressed, and highly expressed in [inflorescences](#) (Fig. 6a).

To further study the expression pattern of *AtAPAM2*, the promoter fragment of *AtAPAM2* was incorporated into *GUS* reporter gene and introduced into wild-type plants. In the T₂ transgenic plants harbouring this fusion, GUS staining signals were detected in seedlings, roots, leaves, inflorescences, flowers, siliques, and pollen grains (Fig. 6b–6h). This further demonstrated that *AtAPAM2* was expressed constitutively.

Subcellular Localisation of *AtAPAM2* Protein

To further study the function of *AtAPAM2* protein, the GFP–*AtAPAM2* fusion protein under control of the 35S promoter (p35S:GFP–*AtAPAM2*) was constructed. This construct was transformed into leaves of *Nicotiana benthamiana* for transient expression and *Arabidopsis* wild-type plants for stable expression. Both in epidermal cells of *N. benthamiana* (Fig. 7a–7c) and roots of *Arabidopsis* (Fig. 7d–7f), the GFP signals indicated *AtAPAM2* localisation in plasma membrane and nucleus.

***AtAPAM2* Encodes a Putative WD40-repeat Protein and is Evolutionarily Conserved**

The *AtAPAM2* mRNA encodes a putative WD40-repeat protein (Fig. 8b) of 1614 aa and contained three WD40-repeat domains (Fig. 8c). We used the entire aa sequence of *AtAPAM2* to carry out a basic local alignment search tool (BLAST) search from NCBI. The results showed several proteins with >60% aa sequence similarity to *AtAPAM2* in higher plants. Specifically, *AtAPAM2* had 98% identity with EOA23389.1 of *Capsella rubella*, 96% with XP_002876038.1 of *Arabidopsis lyrata* subsp. *lyrata*, 77% with XP_002522312.1 of *Ricinus communis*, 76% with EOX91354.1 of *Theobroma cacao*, 76% with XP_004161728.1 of *Cucumis sativus*, 75% with XP_004232045.1 of *Solanum lycopersicum*, and 68% with EEC84558.1 of *Oryza sativa* (Fig. 8d). The above are WD40-repeat proteins, but of unknown function.

The *AtAPAM2* has three WD40-repeats and in eukaryotes this domain usually functions as a protein–protein or protein–DNA/RNA interaction platform [1, 2]. In order to understand the function of this protein, we made use of the WD40 domain to carry out a yeast two-hybrid assay, with the aim of finding the protein that interacted with it; however, this was not successful.

Discussion

In flowering plants, the male gametophyte plays a crucial role in plant fertility and crop production by producing and transferring male gametes to embryos for double fertilisation. The male gametophyte is derived from the anther, which is the male sexual organ. Formation of the male gametophyte involves several important steps. First, reproductive cells divide and microspore mother cells form. Then, the microspore mother cells undergo meiosis and produce **tetrad** microspores surrounded by callose. Later, the callose is degraded by callose enzyme, releasing the haploid microspores. The individual microspores further form three-celled pollen grains, comprising a larger vegetative cell and two smaller sperm cells. The three-celled pollen grain further undergoes dehydration to form a mature pollen grain [17]. The dehydration of pollen grains is very **important** for maintaining maximum pollen viability so that pollen grains can tolerate various environmental stresses after release from the anther [18]. Defects in any of these steps will **affect** pollen formation or male gametophytic function [17, 19, 20]. In recent years, many mutants with altered processes of pollen formation have been identified in *Arabidopsis*, rice, and maize. Because of both male and female gametogenesis defects, loss-of-function of nucleoporin NUP1 in *Arabidopsis* caused fertility defects. The ovules are arrested during meiosis, and pollen grains are aborted at mitosis I [21]. Loss of *AP1G* function compromised dynamic vacuolar remodelling during pollen development and impaired vacuolar acidification of pollen, which resulted in male gametophytic lethality due to defective pollen development [22]. A conserved glycerol–3-phosphate acyltransferase gene *OsGPAT3* plays an important role in regulating anther wall degradation and pollen exocrine formation in rice [23]. Gene *OsMS1* regulates tapetal programmed cell death and pollen exine formation in rice; it functions as a transcriptional activator and interacts with known tapetal regulatory factors through its

plant homeodomain [24]. Loss-of-function of *OsTKPR1* delays degeneration of the tapetum, reduces the lipid level of the anther cuticle, disrupts the Ubisch body and pollen exine formation, and results in complete male sterility [25]. *Abnormal Pollen Vacuolation1 (APV1)*, a tapetum-specific gene, affects anther cuticle and pollen exine formation in maize. The *apv1* mutant was completely male sterile [26].

In the present study, we reported the characterisation of *atapam2*, which affects pollen maturation. It was widely expressed in seedlings, roots, stems, leaves, siliques, flowers, and pollen grains. Gene *AtAPAM2* encodes an unknown protein with three WD40-repeats that localises in plasma membrane and nucleus. Homologous proteins of *AtAPAM2* exist in many species, and we speculate that this protein may interact with other proteins and play a role in pollen formation and other developmental processes.

Conclusions

We determined that *AtAPAM2* protein may play a role in pollen formation and other development processes by interacting with other proteins, but we did not identify the protein that interacts with it.

Methods

Plant Materials and Mutant Isolation

The plants used were of Landsberg *erecta* (*Ler*) and Colombia (*Col*) backgrounds. The *Ler* and *Col* seeds were propagated and preserved in the laboratory of YeDe (China Agricultural University). The seeds were surface-sterilised and pre-germinated on MS-salt agar plates with or without 50 µg/mL of kanamycin (E004000, Sigma, USA) or 25 µg/mL hygromycin (Roche, Mannheim, Germany) or 10 µg/mL Basta (45520, Sigma, USA) at 22°C with a 16/8 h of light/dark cycle. The plants were grown in soil at 22°C under the same light cycle as for seed germination. The *Ds* insertion line was generated and mutants selected according to methods described previously [27]. The SALK T-DNA insertion lines were obtained from the ABRC (<http://www.arabidopsis.org/>).

Phenotypic Characterisation of Mutant

Morphological observations of pollen grains by SEM were carried out as previously described [28]. The DAPI staining was performed as previously described [29]. Pollen viability was assayed using Alexander's staining [30]. For semi-thin section analysis, developing anthers were fixed overnight in a fixation solution of 5% (v/v) glutaraldehyde and 4% (v/v) paraformaldehyde (pH 7.2) and then rinsed three times with 0.1 M phosphate buffer, post-fixed in 1% osmium tetroxide for 3–4 h, and rinsed three times with 0.1 M phosphate buffer again. The samples were dehydrated through 30 min of exposure to a series of ethanol/water mixtures (30, 50, 70, 80, 90, and 100% ethanol) and 100% epoxypropane for two times. Subsequently, samples were transferred to a 1:1 dry epoxypropane:resin mix [31] on a rotator (12 h) and

infiltrated in a 1:3 dry [epoxypropane](#):resin mix for an additional 12 h before transferring to freshly mixed resin. Following two fresh resin changes, the materials were polymerised in moulds (70°C for 8–10 h) [31]. Semi-thin sections (1.0 µm thick) were cut using a [glass cutter](#) on a Leica RM2265 microtome (Leica, Wetzlar, Germany). The sections were dyed with toluidine blue and viewed under an optical microscope (DM2500, Leica, Wetzlar, Germany). The TEM of pollen grains was performed as described previously [32].

Molecular Cloning of *AtAPAM2* and Complementation Experiments

The full-length *AtAPAM2* genomic DNA fragment was amplified by PCR with primers AtAPAM2-C-F and AtAPAM2-C-R (Table 1). For complementation experiments, the full-length *AtAPAM2* genomic DNA fragment was subcloned into pCAMBIA1300 vector (CAMBIA, Canberra, Australia) and introduced into *atapam2/+* heterozygous plants using the *Agrobacterium tumefaciens*-mediated infiltration method. The transformant plants were selected in MS medium containing 50 mg/L of kanamycin and 25 mg/L of hygromycin. The homozygous transformant plants were selected in T₂ generations and used for further analyses.

Analysis of Expression Pattern

The promoter fragments of *AtAPAM2* were amplified using *LA* Taq and *Pfu* DNA polymerase (Takara, Dalian, China) with the gene-specific primers AtAPAM2-P-F and AtAPAM2-P-R (Table 1). The resulting fragments were subcloned upstream of the *GUS* reporter gene in pCAMBIA1300 Ti-derived binary vector and introduced into wild-type plants. Transformation and GUS activity analysis were carried out as described previously [27, 33].

For the RT-PCR assay, total RNAs were extracted from roots, stems, leaves, inflorescences, and siliques of 5-week-old wild-type and 10-day-old transformant seedlings, respectively, using an RNAPrep Pure Plant Kit (Tiangen, Beijing, China) according to the manufacturer's instructions. First-strand cDNA was synthesised from 1 µg of total RNA using the M-MLV [reverse transcriptase](#) from Invitrogen (CA, USA) according to the manufacturer's instructions. The gene-specific primers AtAPAM2-RT-F and AtAPAM2-RT-R were used for PCR reactions. *TUBULIN8* expression level was applied as the internal control by including the specific primers TUBULIN-F and TUBULIN-R (Table 1).

Subcellular Localisation Assay of the *AtAPAM2* Protein

The *AtAPAM2* cDNA fragment was amplified by PCR using the primers AtAPAM2GFP-F and AtAPAM2GFP-R (Table 1) and then cloned into the modified pCAMBIA1300 vector to generate the fusion protein-expressing cassette p35S:GFP–*AtAPAM2*. The resulting construct was transformed into leaves of

N. benthamiana for transient expression and *Arabidopsis* wild-type plants for stable expression. The GFP signals were observed under a Leica DM2500 microscope as previously described [34, 35, 36].

Abbreviations

DAPI: 4',6-Diamidino-2-Phenylindole; *SEM*: [Scanning Electron Microscope](#);

TEM: [Transmission Electron Microscope](#); *GUS*: β -Glucuronidase;

RT-PCR: Reverse transcription-Polymerase Chain Reaction;

GFP: Green Fluorescent Protein; *BLAST*: Basic Local Alignment Search Tool.

Declarations

Ethics approval and consent participate

Not applicable.

Consent to publish

Not applicable.

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests

Funding

This work was supported by the National Natural Science Foundation of China (31760078 and 30860121). The funders did not play any roles in the design of the study, collection, analysis, and interpretation of the relevant data, and writing the manuscript.

Author Contributions

The work presented here was carried out in collaboration among all the authors. ZM, YL, and GZ performed the experiments, analysed data, and contributed to writing the manuscript. HY designed the project, analysed data, wrote the manuscript, and obtained funds to support the project. DY contributed to

performing the experiment and manuscript writing. XZ and LC helped to perform the experiments. All authors have read and approved the final manuscript.

Acknowledgements

Not applicable.

Authors' Information

Zhaoxia Ma, Email: zxma0402@126.com, Address: Yunnan Key Laboratory for Basic Research on Bone and Joint Diseases & Yunnan Stem Cell Translational Research Center, Kunming University, Kunming, Yunnan 650214, P. R. China.

Xueqin Zhang, Email: zhangxq711-1@163.com; Liquan Chen, Email: chenliquan@cau.edu.cn; De Ye, Email: yede@cau.edu.cn, Address: College of Biological Sciences, China Agricultural University, Beijing 100193, P. R. China.

Guobin Zhang, Email: zgb1982@126.com; Hongyu Yang, Email: yanghongyukm@126.com, Address: College of Agriculture and Life Sciences, Kunming University, Kunming, Yunnan 650214, P. R. China.

YueLiu, Email: liuyue984@126.com; Address: College of Life Sciences, Yunnan Normal University, Kunming, Yunnan 650500, P. R. China.

References

1. Neer EJ, Schmidt CJ, Nambudripad R, Smith TF. The ancient regulatory-protein family of WD-repeat proteins. *Nature*. 1994; 371: 297–300.
2. Van Nocker S, Ludwig P. The WD-repeat protein superfamily in Arabidopsis: conservation and divergence in structure and function. *BMC Genomics*. 2003; 4: 50.
3. Xu C, Min JR. Structure and function of WD40 domain proteins. *Protein Cell*. 2011; 2: 202–214.
4. Schapira M, Tyers M, Torrent M, Arrowsmith CH. WD-repeat domain proteins: a novel target class? *Nat Rev Drug Discov*. 2017; 16: 773–786.
5. Wall MA, Coleman DE, Lee E, Iniguez-Lluhi JA, Posner BA, Gilman AG, Sprang SR. The structure of the G protein heterotrimer Gi alpha 1 beta 1 gamma 2. *Cell*. 1995; 83: 1047–1058.
6. Lambright DG, Sondek J, Bohm A, Skiba NP, Hamm HE, Sigler PB. The 2.0 Å crystal structure of a heterotrimeric G protein. *Nature*. 1996; 379: 311–319.
7. Zeng CJ, Lee YR, Liu B. The WD40 repeat protein NEDD1 functions in microtubule organization during cell division in Arabidopsis thaliana. *Plant Cell*. 2009; 21: 1129–1140.
8. Walker AR, Davison PA, Bolognesi-Winfield AC, James CM, Srinivasan N, Blundell TL, Esch JJ, Marks MD, Graya JC. The TRANSPARENT TESTA GLABRA1 locus, which regulates trichome differentiation

- and anthocyanin biosynthesis in *Arabidopsis*, encodes a WD40 repeat protein. *Plant Cell*. 1999; 11: 1337–1349.
9. Shi DQ, Liu J, Xiang YH, Ye D, Sundaresan V, Yang WC. SLOW WALKER1, essential for gametogenesis in *Arabidopsis*, encodes a WD40 protein involved in 18S ribosomal RNA biogenesis. *Plant Cell*. 2005; 17: 2340–2354.
 10. Li HJ, Liu NY, Shi DQ, Liu J, Yang WC. YAO is a nucleolar WD40-repeat protein critical for embryogenesis and gametogenesis in *Arabidopsis*. *BMC Plant Biol*. 2010; 10: 169.
 11. Han MJ, Jung KH, Yi G, Lee DY, An G. Rice Immature Pollen 1 (RIP1) is a regulator of late pollen development. *Plant Cell Physiol*. 2006; 47: 1457–1472.
 12. Gao X, Chen Z, Zhang J, Li X, Chen G. OsLIS-L1 encoding a lissencephaly type-1-like protein with WD40 repeats is required for plant height and male gametophyte formation in rice. *Planta*. 2012; 235: 713–727.
 13. Kumar S, Jordan MC, Datla R Cloutier S. The LuWD40-1 gene encoding WD repeat protein regulates growth and pollen viability in Flax (*Linum Usitatissimum* L.). *PLoS One*. 2013; 8: e69124.
 14. Ju Y, Guo L, Cai Q, Ma F, Zhu QY, Zhang Q, Sodmergen. *Arabidopsis* JINGUBANG is a negative regulator of pollen germination that prevents pollination in moist environments. *Plant Cell*. 2016; 28: 2131–2146.
 15. Yang HY, Li WS, Chen SN. Isolation and characterization of 4 gametophytic male sterile mutants in *Arabidopsis thaliana*. *Chinese Sci Bull*. 2007; 52: 1949–1956.
 16. Preuss D, Rhee SY, Davis RW. Tetrad analysis possible in *Arabidopsis* with mutation of the QUARTET (QRT) genes. *Science*. 1994; 264: 1458–1460.
 17. McCormick S. Control of male gametophyte development. *Plant Cell*. 2004; 16: S142-S153.
 18. Swanson R, Edlund AF, Preuss D. Species specificity in pollen-pistil interactions. *Annu Rev Genet*. 2004; 38: 793–818.
 19. Scott R, Hodge R, Paul W, Draper J. The molecular biology of anther differentiation. *Plant Sci*. 1991; 80: 167–191.
 20. McCormick S. Male gametophyte development. *Plant Cell*. 1993; 5: 1265 –1275.
 21. Bao S, Shen G, Li G, Liu Z, Arif M, Wei Q, Men S. The *Arabidopsis* nucleoporin NUP1 is essential for megasporogenesis and early stages of pollen development. *Plant Cell Rep*. 2019; 38 (1): 59–74.
 22. Feng C, Wang JG, Liu HH, Li S, Zhang Y. *Arabidopsis* adaptor protein 1G is critical for pollen development. *J Integr Plant Biol*. 2017; 59 (9): 594–599.
 23. Sun LP, Xiang XJ, Yang ZF, Yu P, Wen XX, Wang H, Abbas A, Muhammad Khan R, Zhang YX, Cheng SH, Cao LY. OsGPAT3 plays a critical role in anther wall programmed cell death and pollen development in rice. *Int J Mol Sci*. 2018; 19 (12): 4017.
 24. Yang ZF, Liu L, Sun LP, Yu P, Zhang PP, Abbas A, Xiang XJ, Wu WX, Zhang YX, Cao YL, Cheng SH (2019) OsMS1 functions as a transcriptional activator to regulate programmed tapetum development and pollen exine formation in rice. *Plant Mol Biol*. 99 (1–2): 175–191.

25. Xu DW, Qu SY, Tucker MR, Zhang DB, Liang WQ, Shi JX. Ostkpr1 functions in anther cuticle development and pollen wall formation in rice. *BMC Plant Biol.* 2019; 19 (1):104.
26. Somaratne Y, Tian YH, Zhang H, Wang MM, Huo YQ, Cao FG, Zhao L, Chen HB. ABNORMAL POLLEN VACUOLATION1 (APV1) is required for male fertility by contributing to anther cuticle and pollen exine formation in maize. *Plant J.* 2017; 90 (1): 96–110.
27. Sundaresan V, Springer P, Volpe T, Haward S, Jones JD, Dean C, Ma H, Martienssen R. Patterns of gene action in plant development revealed by enhancer trap and gene trap transposable elements. *Genes Dev.* 1995; 9: 1797–1810.
28. Jiang LX, Yang SL, Xie LF, Puah CS, Zhang XQ, Yang WC, Sundaresan V, Ye D. VANGUARD1 encodes a pectin methylesterase that enhances pollen tube growth in the Arabidopsis style and transmitting tract. *Plant Cell.* 2005; 17: 584–596.
29. Jia DJ, Cao X, Wang W, Tan XY, Zhang XQ, Chen LQ, Ye D. GNOM-LIKE 2, encoding an adenosine diphosphate-ribosylation factor-guanine nucleotide exchange factor protein homologous to GNOM and GNL1, is essential for pollen germination in Arabidopsis. *J Integr Plant Biol.* 2009; 51: 762–773.
30. Alexander MP. Differential staining of aborted and nonaborted pollen. *Stain Technol.* 1969; 44: 117–122.
31. Spurr AR. A low-viscosity epoxy resin embedding medium for electron microscopy. *J Ultrastruct Res.* 1969; 26: 31–43.
32. Tan XY, Liu XL, Wang W, Jia DJ, Chen LQ, Zhang XQ, Ye D. Mutations in the Arabidopsis nuclear-encoded mitochondrial phage-type RNA polymerase gene RPOTm led to defects in pollen tube growth, female gametogenesis and embryogenesis. *Plant Cell Physiol*, 2010; 51: 635–649.
33. Yang WC, Ye D, Xu J, Sundaresan V. The SPOROCTELESS gene of Arabidopsis is required for initiation of sporogenesis and encodes a novel nuclear protein. *Genes Dev.* 1999; 13: 2108–2117.
34. Xia C, Wang YJ, Li WQ, Chen YR, Deng Y, Zhang XQ, Chen LQ, Ye D. The Arabidopsis eukaryotic translation initiation factor 3, subunit F (AtelF3f), is required for pollen germination and embryogenesis. *Plant J.* 2010; 63: 189–202.
35. Niu QK, Liang Y, Zhou JJ, Dou XY, Gao SC, Chen LQ, Zhang XQ, Ye D. Pollen-expressed transcription factor 2 encodes a novel plant-specific TFIIB-related protein that is required for pollen germination and embryogenesis in Arabidopsis. *Mol Plant.* 2013; 6: 1091–1108.
36. Zhou JJ, Liang Y, Niu QK, Chen LQ, Zhang XQ, Ye D. The Arabidopsis general transcription factor TFIIB1 (AtTFIIB1) is required for pollen tube growth and endosperm development. *J Exp Bot.* 2013; 64: 2205–2218.

Tables

Table 1. The primers used in this study

Primers	Sequences
AtAPAM2-C-F	5'-TACAAAGAAATGGACTAACTC-3'
AtAPAM2-C-R	5'-CTGTCAAGAACGATGAAC-3'
AtAPAM2-P-F	5'- gctgcagctcttaaagggaagatttgta-3'
AtAPAM2-P-R	5'-ctctagagaacttctgaaagacaaaatac-3'
AtAPAM2-RT-F	5'-GAGCCTGCTGCTATGGAATC-3'
AtAPAM2-RT-R	5'-TAGCTGACTCTGGCCCTTGT-3'
TUBULIN-F	5'-CTTCGTATTTGGTCAATCCGGTGC-3'
TUBULIN-R	5'-GAACATGGCTGAGGCTGTCAAGTA-3'
atapam2-2-F	5'-CCAGAGTCAGCTAACAATTTTC-3'
atapam2-2-R	5'-CTTCTGTCTCAAACATGAAT-3'
AtAPAM2GFP-F	5'-GGGGTACCGAGAGCAGAAGTTATGGAGTGGGCAACGGTGCAGCAT-3'
AtAPAM2GFP-R	5'-ACGCGTCGACTTAGATGTTGATTAACACCAACGGTGA-3'

Table 2 Genetic analysis of *atapam2-2/+* mutant

Crosses (Female×Male)	Basta ^R	Basta ^S	Basta ^R :Basta ^S
<i>atapam2-2/+</i> × <i>atapam2-2/+</i>	1590	1500	1.06:1
<i>atapam2-2/+</i> ×WT	851	900	0.95:1
WT× <i>atapam2-2/+</i>	0	2108	0:1

Basta^R: Basta-resistant; Basta^S: Basta-sensitive; WT: Wild-Type

Figures

Figure 1

The phenotype of *atapam2/+; qrt1/qrt1* pollen. a Scanning electron microscopy of pollen from *qrt1/qrt1*. b Scanning electron microscopy of pollen from *atapam2/+; qrt1/qrt1*. c Alexander staining of pollen from *qrt1/qrt1*. d Alexander staining of pollen from *atapam2/+; qrt1/qrt1*. e DAPI staining of pollen from *atapam2/+; qrt1/qrt1*. f DAPI staining of pollen from *qrt1/qrt1*. g and h Bright fields of e and f.

Figure 2

DAPI staining of different stages of pollen development. a, f, k, p Uninucleate pollen grains. b, g, l, q Binucleate pollen grains. c, h, m, r Early trinucleate pollen grains. d, i, n, s Middle trinucleate pollen grains. e, j, o, t Late trinucleate pollen grains. a–e DAPI staining of different stages of pollen grains from *atapam2/+*. f–j Bright fields of a–e. k–o DAPI staining of different stages of pollen grains from wild type. p–t Bright fields of k–o. Red arrowheads indicate abnormal pollen grains. Bars = 100 μ m.

Figure 3

Semi-thin section images of anthers in wild-type and atapam2/+ mutant. a, c, e Wild-type plants. b, d, f atapam2/+ plants. a, b Stage 10 of anther development. c, d Stage 11 of anther development. e, f Stage 12 of anther development. Red arrowheads indicate the atapam2 mutant pollen grains. Bars = 50 μm.

Figure 4

Transmission electron microscopy images of pollen grains from wild-type and *atpam2/+*. a, b wild-type. c–e *atpam2/+*. Bars = 1 μ m.

Figure 5

Characterisation of *atapam2/+* mutant pollen grains and complementation plants pollen grains. a–c Scanning electron microscopy image of pollen grains from wild-type, *atapam2/+* mutant, and complementation plants. d–f Alexander-stained pollen grains from wild type, *atapam2/+* mutant, and complementation plants. g–i DAPI-stained pollen grains from wild type, *atapam2/+* mutant, and complementation plants. j–l Light field of g–i. m Statistics of normal pollen percentage from wild type, *atapam2/+* mutant, and complementation plants. Bars = 100 μ m in a–f and 50 μ m in g–l.

Figure 6

Expression pattern of *AtAPAM2*. a RT-PCR assay comparing *AtAPAM2* mRNA levels in roots, stems, leaves, inflorescences, siliques, and seedlings. Expression level of *TUBULIN* was used as an internal control. b–h GUS activity in seedlings (b), mature leaves (c), stems (d), siliques (e), inflorescences (f), flowers (g), and anthers (h) from transgenic wild-type plants carrying the p*AtAPAM2*:GUS construct. Bars = 500 μm (b–f) and 100 μm (g, h).

Figure 7

Subcellular localisation of AtAPAM2 protein. a–c Transient expression of p35S:GFP–AtAPAM2 in epidermal cells of *Nicotiana benthamiana*, showing localisation of AtAPAM2 protein in plasma membrane and nucleus. a GFP image. b Bright field image of a. c Merged image of a and b. d–f Stable expression of p35S:GFP–AtAPAM2 in roots of *Arabidopsis*, showing localisation of AtAPAM2 protein in plasma membrane and nucleus. d GFP image. e Bright field image of d. f Merged image of d and e. Bars = 20 μm .

Figure 8

Molecular characterisation of the AtAPAM2 gene. a Schematic diagram of AtAPAM2 structure, showing the Ds insertion site in the *atpam2/+* mutant and T-DNA insertion site in the *atpam2-2* mutant. Black boxes indicate coding sequences and lines indicate introns. b Putative conserved domains of AtAPAM2 detected. c Predicted peptide sequence of AtAPAM2, showing the conserved domains. WD40 domains are shown by underline and red characters. d The sequence similarity analysis of AtAPAM2 and similar proteins from different species. AtAPAM2, Arabidopsis APAM2 protein; EOA23389.1, *Capsella rubella*; EEC84558.1, *Oryza*; XP_002522312.1, *Ricinus communis*; XP_004232045.1, *Solanum lycopersicum*;

XP_004161728.1, Cucumis sativus; EOX91354.1, Theobroma cacao; XP_002876038.1, Arabidopsis lyrata subsp. Lyrata.