

Comparison of Apical and Basolateral Cu Treatment for Iron-related Gene Regulation During Deferoxamine Induced Iron Deficiency Anemia

Ezgi Evcan

Izmir Institute of Technology: Izmir Yuksek Teknoloji Enstitusu

sukru gulec (✉ sukrugulec@iyte.edu.tr)

Izmir Institute of Technology <https://orcid.org/0000-0002-6789-1050>

Research

Keywords: Iron deficiency anemia, Copper, Iron, Intestine, Caco-2, Apical, Basolateral

Posted Date: March 6th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-267161/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Intestinal copper transporter (Atp7a) mutant brindled mice with systemic Cu deficiency had elevated Cu levels in enterocyte cells without any perturbation of iron regulating genes, suggesting that blood Cu level might be important for intestinal iron homeostasis during iron deficiency anemia (IDA). We hypothesized that the blood Cu level and polarization (apical and basolateral) of enterocyte cells might be important regulators for the compensatory response on the regulation of genes in enterocyte cells during IDA. **Methods:** We grew Caco-2 cells on a bicameral cell culture plate to mimic the human intestine system and on a regular tissue culture plate. IDA was induced by Deferoxamine (DFO). The cells were treated with Cu and Cu with Fe following mRNA expressions of *DMT1*, *FPN*, *TFR*, and *ANKRD37* were analyzed.

Results: Our main finding was that basolateral treatment of Cu significantly reduced mRNA expressions of iron-regulated genes, including *DMT1*, *FPN*, *TFR*, and *ANKRD37*, compared to DFO treated and DFO with apical Cu treated groups in both bicameral and regular tissue culture plates.

Conclusions: Cu level in the basolateral side of Caco-2 cells significantly influenced the intracellular gene regulation in DFO induced iron-deficient condition, and polarization of the cells might be important factor gene regulation in enterocyte cells.

Background

Iron is an essential and vital nutrient for life to sustain the functions of living organisms. In the human body, iron is essential to carry oxygen to tissues since it is the main component of hemoglobin. Moreover, it is necessary to control cell division and differentiation, mitochondrial energy production, DNA replication and repair, and immune response against pathogenic microorganisms^{1,2}. Iron levels in the human body are tightly controlled via intestinal absorption since mammals do not have a specific active iron excretory mechanism. Therefore, intestinal enterocyte cells maintain adequate levels of iron in the body. Inadequate dietary iron absorption through enterocyte cells of the intestine leads to iron deficiency anemia, one of the major global nutritional deficiencies, especially in women, infants and children. Iron deficiency causes elevated copper mineral levels in the intestinal mucosa³, liver, and serum⁴ in many mammalian species, including humans⁵. Previous investigations have implicated that copper homeostasis perturbations influence iron metabolism^{1,5-8}.

Divalent metal transporter (*DMT1*), the iron exporter ferroportin 1 (*FPN*), can be a potential connection between iron and copper metabolisms since there are studies that suggest iron amount is regulated by copper in intestine⁹⁻¹¹. Intestinal hypoxia-inducible factor 2 α (HIF-2 α) is essential for iron absorption during iron deficiency since it regulates apical and basolateral iron transporters¹². Copper influences the DNA-binding activity of HIF-2 α , illustrating another connection of copper influencing iron homeostasis. *DMT1* and *FPN* were shown as direct target genes related to HIF-2 α ¹³. mRNA regulation of Ankyrin repeat

domain 37 (*ANKRD37*), prolyl 4-hydroxylase (*P4ha1*), and HIF prolyl hydroxylase 3 (*EGLN3*) are the most well-known marker genes for the hypoxic signal under iron deficiency.

Recent studies have shown that the regulation of genes involved in iron and copper homeostasis were altered in brindled mice⁹. However, when intracellular Cu level increased during IDA, Cu did not affect the genes' regulation, which plays a role in enterocyte iron metabolism. In the same study, enterocyte cells of mice intestine were treated with high intracellular and low blood Cu levels. Thus, blood Cu level might be a more dominant regulator of iron metabolism-related genes during IDA. The apical side of the enterocyte is in contact with dietary nutrients, whereas the basolateral part is in contact with the blood. Considering this apical and basolateral polarization, different sides of enterocytes which in relation to different environmental conditions (diet or blood) may be necessary for controlling the molecular and genetic regulation of iron metabolism in enterocyte cells. However, to the best of our knowledge, no research exists related to the dependency of the apical versus basolateral Cu for enterocyte cell iron metabolism during IDA. Given this background, the current study investigated the effects of dietary and blood copper treatments separately on iron and iron-dependent hypoxic gene regulations of anemic enterocyte cells by *in vitro* modeling of the human intestine system.

Results

Modeling of Human Intestine System and TEER Measurements of Monolayer Caco-2 Cells

The apical and basolateral polarization of Caco-2 cells can behave like the human small intestine¹⁵. Therefore, Caco-2 cells were grown on the bicameral cell culture system for 21 days and then DFO and mineral treatments were performed. TEER was measured from all experimental groups at the end of 21 days and after treatments to control the stability of the monolayer integrity of the cells. As shown in the Fig. 1 we did not observe any significant change in TEER values between the experimental groups before and after Cu and Fe treatments. Furthermore, TEER values that are much higher than 250 ohm/cm² reflect the experimentally polarized cells (as apical and basolateral) in the human intestinal system.

Apical and Basolateral Treatments of Cu in DFO Induced Anemic Monolayer Caco-2 Cells

When the Caco-2 cells are grown on the bicameral cell culture system, it allows us to study the effects of apical and basolateral Cu treatment on regulation of iron metabolism-related genes in enterocyte cells during DFO induced IDA. DFO treatment induced *DMT1*, *TFR*, and *ANKRD37*, and *EGLN3* mRNA levels; however, *FPN* and *MT1a* were not affected (Figure-2a). When Cu was given to the apical side of the cells, it did not affect mRNA levels of genes, including *DMT1*, *TFR*, and *ANKRD37*, and *EGLN3* that were induced by DFO. Cu treatment of the basolateral side of cells significantly reduced *DMT1*, *FPN* and *TFR*, and *ANKRD37* mRNA levels under the IDA condition. Moreover, *DMT1*, *FPN*, and *TFR* gene expressions

were not changed by apical side treatments of the cells by Cu and Fe, whereas *ANKRD37* and *EGLN3* mRNA levels were slightly reduced. Furthermore, when both Cu and Fe were given to the basolateral side of the cells, mRNA levels of iron regulating genes (*DMT1*, *FPN*, and *TFR*), and hypoxia-related genes (*ANKRD37*, and *EGLN3*) were significantly lower than all treatment groups. *MT1a* mRNA levels were significantly increased by basolateral Cu treatment of cells, and this induction was higher when both Cu and Fe were given to the basolateral side of the cells (Fig. 2a and 2b).

The Effect of Cu on Iron Deficiency Anemia in Caco-2 Cells Grown on the 12-well Tissue Culture Plate

Copper and iron interaction in enterocyte cells has attracted interest since their homeostasis is controlled by intestinal absorption. In this study, we grew Caco-2 cells on 12-well tissue culture plates for 21 days and then treated by DFO to induce IDA. We treated the cells Cu with and without Fe. Then we analyzed the mRNA levels of iron regulating and iron-dependent hypoxia regulating genes to see the effects of Cu on iron metabolism during IDA (Fig. 3a and 3b). DFO treatment significantly induced *DMT1*, *TFR*, and *ANKRD37*, and *EGLN3* mRNA expression levels, whereas *FPN* and *MT1a* mRNA levels were not affected. The Cu treatment did not reduce DFO induced gene mRNA expressions and did not affect *FPN* and *MT1a* mRNA levels. However, Cu with Fe treatment significantly reduced *DMT1*, *TFR*, *ANKRD37*, and *EGLN3* mRNA levels. Moreover, we observed that Cu and Fe together significantly upregulated *MT1a* mRNA expression.

Discussion

Enterocyte cells are unique cell types in the intestine due to the interaction between diet and the internal circulatory system of the body. Thus, the dietary and blood-derived stimulus might play a role in enterocyte nutrient metabolism. Copper and iron have physiological interactions in enterocyte cells during IDA. In many mammalian species, iron deficiency leads to elevated Cu levels in enterocyte¹⁶, serum¹⁷, and liver¹⁸, indicating that copper might influence intestinal iron metabolism. Iron and copper interactions have been investigated in brindled mice (*Mo^{Br/y}*) and researchers observed no perturbation in iron absorption during anemia⁹. In this study, the enterocyte copper level of mutant mice was high due to nonfunctional ATP7A protein. In contrast, these mice were systemic copper-deficient; thus, enterocyte cells were exposed to two different copper levels. Furthermore, it was observed that dietary-induced iron deficiency upregulated mRNA expressions of iron-regulated genes, including *DMT1*, *TFR*, and *ATP7A*, but high enterocyte copper level did not affect the gene regulations suggesting that blood Cu level might be an important dietary factor for gene regulations in enterocyte cells during IDA. Thus, we grew Caco-2 cells on a bicameral cell culture plate to mimic the human intestine system to test whether dietary (apical side) or blood copper (basolateral side) are involved in iron-dependent genes regulation during IDA.

In the current study, we selected marker genes that were regulated by iron deficiency anemia including iron transporters (*DMT1*, *TFR* and *FPN*) and iron-dependent hypoxic genes (*ANKRD37* and *EGLN3*). Furthermore, *MT1A* mRNA level was used to control intracellular mineral uptake since *MT1A* is regulated by intracellular Cu and Zn, respectively^{19,20}. Our results showed that Cu and Cu with Fe in the basolateral

side of the cells reduced marker genes mRNA levels (*DMT1*, *FPN*, *TFR1*, and *ANKR37*) compared to the apical Cu and Cu with Fe treatment group during IDA. This suggests that Cu sensing of the basolateral side of the cells might be different from Cu sensing of the apical side. Cu regulates the *MT1A* mRNA levels and its level is correlated by elevated intracellular Cu²⁰. We observed that *MT1A* mRNA levels were significantly induced when Cu and Cu with Fe were given to the basolateral side of the cells suggesting that the level of Cu in the blood might be a compelling factor for the regulation of iron deficiency-related genes in the intestine. Furthermore, the basolateral side of the enterocyte cells may play a role in Cu sensing. The importance of the molecular and genetic interactions between basolateral and apical sides of polarized enterocyte cells has been indicated in different studies. It was shown that glucose treatment in the basolateral side of the Caco-2 cells induced apical cholesterol uptake and the mRNA levels of the cholesterol transporter gene²¹, indicating that basolateral signaling influences fatty acid metabolism through apical side of enterocyte cells. Moreover, Han et al. (2002)²² showed that both apical and basolateral Cu treatment increased Fe uptake in non-anemic Caco-2 cells. Moreover, Cu treatment of apical and basolateral sides also induced *DMT1*, *TFR*, and *FPN* mRNA levels compared to the control group. All together results suggest that basolateral and apical sides of enterocyte cells might have a different physiological response regarding the same nutritional stimulus.

Caco-2 cells can differentiate on the regular tissue culture plate when they are grown 21 post-confluent days. Differentiation of Caco-2 cells gives different genetic responses for a variety of cellular physiological pathways¹⁵. However, polarization of those cells on the insert system might be another factor that can influence genetic regulation. Thus, we also grew Caco-2 cells on regular tissue culture plates at the same time that we performed bicameral cell culture experiments. We observed similar results for marker genes regulation between apical side Cu treatment groups and Cu treatment on Caco-2 cells that were grown on tissue culture plates. However, basolateral Cu treatment affected regulations of *DMT1*, *FPN*, *MT1A*, *ANKRD37* genes compared to results from the cells that were grown on tissue culture plates. Our results suggest that the polarization of the Caco-2 cells might be an important factor for gene regulation in terms of Cu treatment in our study. It might be better to account polarization of the Caco-2 cells for gene regulation or whole-genome array studies.

Conclusion

The nutrient-dependent regulation of enterocyte cells is central to the intestinal nutrient-sensing mechanism. The basolateral and apical sides of the enterocyte cells are the primary targets to understand nutrient sensing in terms of nutrient overload or deficiency. Furthermore, polarization of Caco-2 cells also might influence gene regulation. Our results suggest that intracellular gene regulation was mainly affected by Cu treatment in the basolateral side of enterocyte cells during IDA, indicating that blood copper level might have ability to control the enterocyte iron metabolism at molecular and genetic levels during iron deficiency anemia. The blood Cu level might be an important regulator for intestinal iron metabolism during iron deficiency anemia.

Methods

Cell Culture

The human colorectal adenocarcinoma epithelial cell line, Caco-2, was purchased from the American Type Culture Collection (ATCC, HTB-37, Manassas, VA). Caco-2 cells were cultured in minimal essential medium (MEM) (Sigma, United Kingdom) supplemented with 20% fetal bovine serum (FBS) (Gibco, Cat. No. 10500), 1% penicillin and streptomycin (100 U/mL) and, 1% nonessential amino acid solution (Gibco, Cat. No. 11140). They were maintained in 75 cm² culture flasks at 37°C in a constant humidified incubator with an air atmosphere of 5% CO₂/95% O₂. When cultures reached 70–80% confluency, they were plated for either subsequent passage or treatment. Caco-2 cells used for the treatment experiments were between the 20th -30th passages.

Modeling of the Human Intestinal System and Cu Treatments of Apical and Basolateral Sides of the Cells During Iron Deficiency Anemia

To test the effect of the dietary and blood copper on iron deficiency anemia, we mimicked the human intestines. The Caco-2 cells were seeded into bicameral collagen-coated polytetrafluoroethylene membrane with 0.4- μ m pore size and 1.12 cm² diameter (12-well inserts) (Corning, Cat. No.: 3493). After three days of confluent culture, the cells were grown for an additional 21 days to form a monolayer and became polarized under standard conditions in the indicated MEM media. The experimental cell medium was changed every other day. Transepithelial electrical barrier resistance (TEER) was measured by an epithelial volt-ohm meter (EVOM; World Precision Instruments, Inc., FL, USA) to confirm the cell tight junction integrity. We only used cell monolayers with the TEER values above 250 Ω /cm² for the treatment experiments (Sambuy et al. 2005). Iron deficiency anemia was induced in the cells at 21-day of post-seeding using a chemical agent (Deferoxamine, DFO) at a concentration of 200 μ M. After incubation for 24 h, the apical and/or basolateral compartments of anemic Caco-2 cells were treated with the copper (copper (II) chloride, CuCl₂; 100 μ M) or iron (ferric ammonium citrate, FAC; 100 μ g/mL) and/or together at the same time for further 18h in the incubator.

Determination of the Effect of Copper on Iron Metabolism at the Molecular and Genetic Level

The Caco-2 cells were seeded at 1×10^5 cells/well into classical cell culture plates (12-well plates) (Costar, Cambridge, MA) to investigate the effects of Cu on molecular and genetic regulation of iron metabolism. Iron deficiency anemia was induced in the cells at 21-day of post-seeding 200 μ M DFO. After incubation for 24 h, anemic cells were treated with the copper and/or iron minerals at the same concentrations as previously described in the method section.

Quantitative Real-Time Polymerase Chain Reaction (qRT-PCR)

We evaluated the mRNA expression levels of the marker genes that regulate iron metabolism and hypoxic condition. Total RNA was isolated from cells with RNAzol reagent (MRC, Cat. No.: RN190) following the manufacturer's protocol. One microgram of each total RNA was converted to cDNA using the cDNA synthesis kit (Lifetech, Cat. No.: 4368814). qRT-PCR was performed on an ABI StepOnePlus instrument (Lifetech, CA, USA) by using gene-specific oligonucleotide primers (*CYP1A*: Forward-TACGGGTCCTGGCATCTTG, Reverse-CGAGTTGTCCACAGTCAGCA; *DMT1* Forward-TGCATCTTGCTGAAGTATGTCACC, Reverse-CTCCACCATCAGCCACAGGAT; *FPN*: Forward-GCAGGAGAAGACAGAAGCAAAC, Reverse-TCCTTCGAATTGTGGCATTTCAT; *MT1A*: Forward-ACTGGTGGCTCCTGCACCTGCACT, Reverse-ACAGCAGCTGCACTTCTCTGAT; *TFR*: Forward-TCAGAGCGTCGGGATGATATCGG, Reverse-CTTGATCCATCATCATTCTGAACTGCC; *ANKRD37*: Forward-AGCAGTCGCCTGTCCACTTAGC, Reverse-AGCAGGCTTAGGCACTCCAGG; *EGLN3*: Forward-GCAAATACTACGTCAAGGAGAGGTCTAA, Reverse-GGCATCCCAATTCTTGTTTCAGATAG) SYBR-Green mix (Lifetech, Cat. No.: 4367659). C_T (threshold cycle) levels were normalized according to *CYP1A* mRNA expression as a housekeeping gene. Mean fold changes in gene-specific mRNA levels from all experimental groups were calculated by the $2^{-\Delta\Delta C_t}$ analysis method¹⁴.

Statistical Analyses

The results were expressed as the mean values \pm standard deviation of three independent experiments with at least two parallels of each experimental group. Statistical analysis for more than two groups was carried out by one-way analysis of variance (ANOVA) with Tukey's *post hoc* test by using GraphPad Prism 6 (GraphPad Software Inc., CA, USA). Data were considered significant for $p \leq 0.05$.

List Of Abbreviations

Atp7a	Intestinal copper transporter
IDA	Iron deficiency anemia
Caco-2	Human colorectal adenocarcinoma epithelial cell line
DFO	Deferoxamine
DMT1	Divalent metal transporter
FPN1	Ferroportin
TFR	Transferrin receptor
ANKRD37	Ankyrin repeat domain 37
HIF-2 α	Intestinal hypoxia-inducible factor
P4ha1	Prolyl 4-hydroxylase

EGLN3	HIF prolyl hydroxylase 3
MT1a	Metallothionein-1A
CYPA	Cyclophilin A
ATCC	American Type Culture Collection
MEM	Minimal essential medium
FBS	Fetal bovine serum
TEER	Transepithelial electrical barrier resistance
EVOM	Epithelial volt-ohm meter
FAC	Ferric ammonium citrate
qRT-PCR	Quantitative real-time polymerase chain reaction
CT	Threshold cycle

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interest.

Funding

This work was supported by the Scientific and Technological Research Council of Turkey (grant number:215Z041).

Author Contributions

SG had the original project idea. EE and SG designed the investigation, wrote the manuscript, and prepared figures; SG and EE performed the experiments. All authors approved the final manuscript.

References

1. Gulec S, Collins JF. Molecular mediators governing iron-copper interactions. *Annu Rev Nutr.* 2014;34:95-116.
2. Verma C, Tapadia K, Soni AB. Determination of iron (III) in food, biological and environmental samples. *Food Chem.* 2016;221:1415-1420.
3. El-Shobaki FA, Rummel W. Binding of copper to mucosal transferrin and inhibition of intestinal iron absorption in rats. *Res Exp Med (Berl).* 1979;174(2):187-95.
4. Ece A, Uyanik BS, Iscan A, et al. Increased serum copper and decreased serum zinc levels in children with iron deficiency anemia. *Biol Trace Elem Res.* 1997;59(1-3):31-9
5. Fox PL. The copper-iron chronicles: the story of an intimate relationship. *Biometals.* 2003;16(1):9-40.
6. Collins JF, Prohaska JR, Knutson MD. Metabolic crossroads of iron and copper. *Nutr Rev.* 2010;68(3):133-147.
7. Ha JH, Doguer C, Collins JF. Consumption of a high-iron diet disrupts homeostatic regulation of intestinal copper absorption in adolescent mice. *Am J Physiol Gastrointest Liver Physiol.* 2017;313:G353-G360.
8. Wang T, Xiang P, Ha JH, et al. Copper supplementation reverses dietary iron overload-induced pathologies in mice. *J Nutr Biochem.* 2018;59:56-63.
9. Gulec S, Collins JF. Investigation of iron metabolism in mice expressing a mutant Menke's copper transporting ATPase (Atp7a) protein with diminished activity (Brindled; Mo^(Br/y)). *PLoS One.* 2013;8(6):e66010.
10. Matak P, Zumerle S, Mastrogiannaki M, et al. Copper deficiency leads to anemia, duodenal hypoxia, upregulation of HIF-2 α and altered expression of iron absorption genes in mice. *PLoS One.* 2013;8(3):e59538.
11. Wang X, Flores SRI, Ha JH, et al. Intestinal DMT1 is essential for optimal assimilation of dietary copper in male and female mice with iron-deficiency anemia. *J Nutr.* 2018;148(8):1244-1252.
12. Schwartz AJ, Das NK, Ramakrishnan SK, et al. Hepatic hepcidin/intestinal HIF-2 α axis maintains iron absorption during iron deficiency and overload. *J Clin Invest.* 2019;129(1):336-348.
13. Anderson ER, Taylor M, Xue X, et al. Intestinal HIF2 α promotes tissue-iron accumulation in disorders of iron overload with anemia. *Proc Natl Acad Sci.* 2013;110(50):E4922-4930.
14. Hu Z, Gulec S, Collins JF. Cross-species comparison of genomewide gene expression profiles reveals induction of hypoxia-inducible factor-responsive genes in iron-deprived intestinal epithelial cells. *Am J Physiol Cell Physiol.* 2010;299(5):C930-938.
15. Sambuy Y, De Angelis I, Ranaldi G, et al. The Caco-2 cell line as a model of the intestinal barrier: influence of cell and culture-related factors on Caco-2 cell functional characteristics. *Cell Biol Toxicol.*

2005;21(1):1-26.

16. Ravia JJ, Stephen RM, Ghishan FK, et al. Menkes Copper ATPase (Atp7a) is a novel metal-responsive gene in rat duodenum, and immunoreactive protein is present on brush-border and basolateral membrane domains. *J Biol Chem.* 2005;280:36221-36227.
17. Yokoi K, Kimura M, Itokawa Y. Effect of dietary iron deficiency on mineral levels in tissues of rats. *Biol Trace Elem Res.* 1991;29:257-265.
18. Sherman AR, Moran PE. Copper metabolism in iron-deficient maternal and neonatal rats. *J Nutr.* 1984;114:298-306.
19. Gefeller EM, Bondzio A, Aschenbach JR, et al. Regulation of intracellular Zn homeostasis in two intestinal epithelial cell models at various maturation time points. *J Physiol Sci.* 2015;65:317-328.
20. Gulec S, Collins JF. Silencing the Menkes copper-transporting ATPase (Atp7a) gene in rat intestinal epithelial (IEC-6) cells increases iron flux via transcriptional induction of ferroportin 1 (Fpn1). *J Nutr.* 2014;144(1):12-9.
21. Ravid Z, Bendayan M, Delvin E, et al. Modulation of intestinal cholesterol absorption by high glucose levels: impact on cholesterol transporters, regulatory enzymes, and transcription factors. *Am J Physiol Gastrointest Liver Physiol.* 2008; 295(5):G873-885.
22. Han O, Wessling-Resnick M. Copper repletion enhances apical iron uptake and transepithelial iron transport by Caco-2 cells. *Am J Physiol Gastrointest Liver Physiol.* 2002;282(3):G527-533.

Figures




Figure 1

Trans epithelial electrical resistance (TEER) measurements. The Caco-2 cells were grown on the bicameral insert for 21 days. TEER measurements controlled monolayer cell polarization and tight junction integrity of the Caco-2 cells.




Figure 2

The effect of trace minerals on iron and hypoxia-regulated gene mRNA expression levels in polarized Caco-2 cells under the DFO treatment condition. Caco-2 cells were grown on the bicameral cell culture system for 21 days. Iron deficiency anemia was induced by 200 μ M for 24h, and then copper (CuCl₂; 100 μ M) and iron (FAC; 100 μ g/mL) were given to apical and basolateral sides of the cell culture system for 18h. The effects of Cu and Cu with Fe on iron (a) and hypoxia (b) regulating gene mRNA expression were analyzed.




Figure 3

The effect of minerals on iron-regulated gene mRNA expression levels in Caco-2 cells under the DFO treatment condition. Caco-2 cells were grown on the 12-well tissue culture plates for 21 days. The cells were treated by 200 μ M for 24h to induce iron deficiency anemia. The copper mineral (CuCl₂; 100 μ M) and iron (FAC; 100 μ g/mL) were given to Caco-2 cells for 18h to test the effect of Cu and Cu with Fe on iron (a) and hypoxia (b) regulated genes mRNA expressions.