

Lotus Leaf Extract Inhibits the Cell Migration and Metastasis of ER- Breast Cancer

Yuelin Tong

shandong university

Zhongwei Li

jiangnan university

Yikuan Wu

jiangnan university

Shenglong Zhu

jiangnan university

Keke Lu

shandong provincial hospital

Zhao He (✉ zhaoh7711@qq.com)

Research

Keywords: Lotus leaf, cell migration, breast cancer, metastasis, TGF- β 1 signaling pathway

Posted Date: May 15th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-26733/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on February 18th, 2021. See the published version at <https://doi.org/10.1186/s12986-021-00549-0>.

Abstract

Background: Patients with estrogen receptor negative (ER-) breast cancer have poor prognosis because of their high rates of metastasis. However, there is no effective treatment and drugs for ER⁻ breast cancer metastasis. Our purpose of this study was to evaluate the effect and mechanism of lotus leaf alcohol extract (LAE) on the cell migration and metastasis of ER- breast cancer.

Methods: The anti-migratory effect and mechanism of LAE were analysed in ER- breast cancer cells including SK-BR-3, MDA-MB-231 and HCC1806. Cell viability assay, wound-healing assay, RNA-sequence analysis and immunoblotting assay were applied in examining the cytotoxicity, anti-migratory effect and its possible pathways of LAE. To further investigate the inhibitory effect of LAE on metastasis *in vivo*, subcutaneous xenograft nude mice model and intravenous injection nude mice model were established. Lung and liver tissues were analysed by the Hematoxylin & eosin staining and immunoblotting assay.

Results: We found that lotus leaf alcohol extract (LAE), not nuciferine, inhibited cell migration significantly in SK-BR-3, MDA-MB-231 and HCC1806 breast cancer cells, and did not change viability of breast cancer cells. The anti-migratory effect of LAE was dependent on TGF- β 1 signaling, while independent of Wnt signaling and autophagy influx. Intracellular H₂O₂ participated in the TGF- β 1-related inhibition of cell migration. LAE inhibited significantly the breast cancer cells metastasis in mice models. RNA-sequence analysis showed that extracellular matrix signaling pathways are associated with LAE-suppressed cell migration.

Conclusions: Our findings demonstrated that lotus leaf alcohol extract inhibits the cell migration and metastasis of ER- breast cancer via TGF- β 1/Erk1/2 and TGF- β 1/SMAD3 signaling, which provides a potential therapeutic strategy for ER- breast cancer.

1. Introduction

Breast cancer is usually classified into three subtypes according to the expression of hormone receptors: estrogen receptor positive/progesterone receptor positive (ER+/PR+), human epidermal growth factor receptor positive (HER2+) and triple negative (TNBC) [1]. Among the three subtypes of breast cancer cells, ER- breast cancer such as HER2 + and Triple negative breast cancer cells have a higher migration capacity than that of ER+/PR + breast cancer cells. In breast cancer patients, HER2 + subtypes of breast cancer accounts for 20–30%, which is an invasive tumor with shorter metastatic time and poor prognosis compared to other two subtypes of breast cancer patients [2]. In addition, Triple negative breast cancer (TNBC) is considered as the most severe subtypes of breast cancer because of high rate of recurrence and metastasis [3]. However, there is no effective treatment and drugs for ER⁻ breast cancer metastasis.

About 90% death rates of breast cancer are resulted from metastasis. It's a complicated process associated with NF- κ B, TGF- β 1 and Wnt/ β -catenin signaling pathways [4–6]. TGF- β 1 signaling pathway is a pivotal signal for cell migration in breast cancer. TGF- β enhances ROS levels by enhancing production and reducing antioxidative/scavenging systems activity [7–8]. And increased ROS levels in

turn may promote TGF- β expression and stimulate the release of TGF- β from the secreted complex in cell [9]. Moreover, several studies have demonstrated that ROS is associated with carcinogenesis, abnormal growth, and angiogenesis, especially metastasis [10–14]. Thus, inhibition of breast cancer cell metastasis by TGF- β as well as ROS pathway may become a promising strategy for the treatment of ER⁻ breast cancer patients.

Lotus leaf (*Nelumbo nucifera* Gaertn) is a traditional Chinese medicine, also called He-Ye, which has a long history of usages against oxidation, diabetes, obesity and immunomodulatory effects in China [15]. Sometimes lotus leaf is also used as a seasoning in food due to its unique aroma. Considerable literatures have demonstrated that the main compounds contained in the lotus leaf are nuciferine, quercetin, quercetin-3-O-glycoside, kampherol-3-O-glycoside, and myricetin-3-O-glucoside [16]. These compounds from the extracts of lotus leaves, seed and rhizome display cytoprotective, anti-bacterial, anti-obesity especially anti-oxidant pharmacologic activities partly through the extracellular signal-regulated protein kinases like Erk1/2 and PKC [17–21]. However, the effect of lotus leaf extract on cancer metastasis is still remained unclear. Given that the important role of Erk1/2 and ROS in TGF- β -related cancer metastasis [22–23], we tested the effect of LAE on ER⁻ breast cancer metastasis.

In this study, our data showed that LAE significantly inhibits the ER⁻ breast cancer cell migration and metastasis via the SMAD3 and Erk1/2, accompanied by reduced intracellular H₂O₂ level in ER⁻ breast cancer cells. RNA-sequence analysis also showed that extracellular matrix signaling pathways and FAK might be another possible pathways to inhibit the cell migration induced by LAE. These results suggested that LAE has the potential to screen out novel drugs for the treatment of ER⁻ breast cancer clinically.

2. Materials And Methods

2.1 Cells culture

Breast cancer cell lines SK-BR-3, HCC1806 and MDA-MB-231 were purchased from cell bank of China. Cells were maintained in RPMI 1640 (HCC1806) and DMEM (SK-BR-3 and MDA-MB-231) media (Gibco, USA) with 10% FBS (Gibco, USA) and 100 μ g/ml penicillin/streptomycin (Gibco, USA) in a humidified atmosphere containing 5% CO₂ at 37 °C.

2.2 LAE treatment

The dry lotus leaf (*Nelumbo nucifera* Gaertn, <http://www.theplantlist.org>) powder was purchased from Tong Ren Tang Group in Wuxi, China, in November 2017 (Latitude: 31°33'17.44"N; Longitude: 120°19'15.24"E, organization code 101461180). The company is approved by the competent department, ensuring the good production and quality according to the approved content. All production lines had passed the national GMP recognition. Then, the lotus leaf were identified in National Functional Food Engineering Technology Research Center in Jiangnan University (China).

Lotus leaf alcohol extract: The lotus leaf powder and the 75% aqueous ethanol solution were uniformly mixed at a ratio of 1:20, and shaken at 37°C for 12 hours. Then separated by filtration, and the obtained residue was repeatedly extracted twice. All the collected filtrate was concentrated to dryness by evaporation on a rotary evaporator, and then dissolved in DMSO and filtered by a membrane filter with 0.22 µm pore size (Millipore, USA). Finally LAE was diluted into several concentrations (10, 25, 50, 100, 250 µg/ml) for cell treatments and stored at 4°C.

2.3 Human breast cancer xenograft model

This study was approved by the ethics committee of Jiangnan University with protocol number SYXK2016-0045. Female BALB/c nude mice, 4 weeks of age, were purchased from Lingchang Biotechnology Co., Ltd. (Shanghai, China). All the experiments were in accordance with national institution guidelines. MDA-MB-231 cells (1×10^6) suspending in medium were subcutaneously injected in the right flank of mice. After one week of injection, the mice bearing tumor were randomly subdivided into 2 groups and treated with either vehicle (model group) or LAE (0.5% w/w). Tumor dimensions were measured with vernier caliper and tumor volumes were estimated by the formula: length \times width² \times 0.5. Body weights and tumor growth were recorded twice a week. The mice were sacrificed after 56 days, and visible metastatic tumors were counted and subjected to further analysis.

2.4 Tail vein injection model

This study was approved by the ethics committee of Jiangnan University with protocol number SYXK2016-0045. Female BALB/c nude mice, 4 weeks of age, were purchased from Lingchang Biotechnology Co., Ltd. (Shanghai, China). All the experiments were in accordance with institution guidelines. MDA-MB-231 cells (5×10^5) suspending in PBS were injected through the tail vein of mice. Then, all the mice were randomly subdivided into 2 groups (10 mice/group) and treated with either vehicle (model group) or LAE (0.5% w/w). Body weights were scaled once a week. The mice were sacrificed after 41 days, and the metastatic tumors and the organs with visible metastatic tumors were analyzed.

2.5 Cell viability assay

Cell Counting Kit 8 (CCK8) was purchased from Dojindo Molecular Technology (Tokyo, Japan). Cells were cultured in 96-well plates at a density of 5000 cells per well in 100 µl medium. LAE were added into the wells and incubated for 48 h. Then, cells were added 10 µl CCK8 substrate and incubated for another 2 h at 37 °C. The optical density was measured at 450 nm on a microplate reader Multiskan GO (Thermo Scientific, USA).

2.6 Wound-healing assay

Cells were seeded in dish to form a confluent monolayer. Then 200 µl pipette tip was used to make a scratch in the middle of plate, and wash twice with PBS. The monolayer was maintained with the FBS-free media with LAE supplement. Wound closure was monitored after 0 and 36 h. The percent of wound closed are calculated by image J. TGF-β1 was purchased from Peprotech (USA, # 100-21-10) and

dissolved in 10 mM citric acid, pH3.0 at 0.1–1.0 mg/ml. Chloroquine diphosphate (CQ) was purchased from Medchem Express (Shanghai, China, cat# HY-17589) and dissolved in H₂O to 10 mg/ml. LiCl was purchased from Sigma-Aldrich (Sigma, St. Louis, MO, USA, # 213233) and dissolved in cell culture medium to 20 mM. Nuciferine was purchased from Medchem Express (Shanghai, China, cat# HY-N0049) and dissolved in DMSO to 1.6925 mM.

2.7 H₂O₂ content

The content of H₂O₂ was measured by the Hydrogen Peroxide assay kit (Beyotime, # S0038) according to the manufacturer protocols. In brief, test tubes containing 50 µl test solutions were placed at room temperature for 30 min and measured immediately with a spectrophotometer at a wavelength of 560 nm. Absorbance values were calibrated with a standard curve generated with known concentration of H₂O₂.

2.8 Cell signaling

Cells were lysed by lysis buffer (RIPA buffer contains protease inhibitors and phosphatase inhibitors). Protein concentrations were determined by using a BCA Protein Assay Kit. Equal amounts of protein were electrophoretically separated in 10% SDS–polyacrylamide gels, and then transferred onto PVDF membranes (Millipore, Beijing, China). The membranes were blocked with 5% fat free milk for 1 h at room temperature, further incubated with primary antibodies (1:1,000) and then probed with secondary peroxidase-labeled antibody. The signal was detected by Plus-enhanced chemiluminescence using FluorChem FC3 (ProteinSimple, USA). The following primary antibodies were used: Erk1/2 (# 9102), p-Erk1/2 (# 4370), SMAD3 (# 9523S), TGF-β1 (# 3711S), Snail (# 3879T), PTEN (# 9552S), β-catenin (# 8480S), p-AKT (# 4060), AKT (# 4685), p-JNK (# 9251S), JNK (# 9252T), p-p38 (# 4511), p38 (# 8690), p62 (# 23214S), LC3 (# 3868T), Catalase (# 12980T) were purchased from cell signaling technology; NOX4 (# ab133303), p-SMAD3 (# ab52903) and NOX2 (# ab129068) were from Abcam company; β-Actin (# sc-130656) and SOD1 (# sc-101523) were purchased from Santa Cruz Biotechnology.

2.9 Hematoxylin and eosin staining

For histopathology analysis, lung and liver tissues were fixed by paraformaldehyde and then were dehydrated, embedded in paraffin and cut into serial sections at 5 µm. Sections were stained with hematoxylin and eosin (H&E) solution and observed them under an optical microscope (DP73, OLUMPUS, Japan).

2.10 Statistical Analysis

All experiments were performed at least three times and data were presented as mean ± SEM. One-way ANOVA with Dunnett's post-test was performed (*P < 0.05; **P < 0.01; ***P < 0.001).

3. Results

3.1 LAE inhibits ER⁻ breast cancer cells migration independent of nuciferine

To investigate LAE effects on cell migration, metastatic SK-BR-3, MDA-MB-231 and HCC1806 ER⁻ breast cancer cell lines were used in the study. Previous studies have shown that cell proliferation interferes with cell migration. To address this issue, the proliferation of breast cancer cells was examined by CCK8 assay. The viability of SK-BR-3, MDA-MB-231 and HCC1806 cells was similarly unchanged between with low dose (50 and 100 µg/ml) of LAE supplement and without LAE (Fig. 1a). All of SK-BR-3, MDA-MB-231 and HCC1806 cells exhibited no cytotoxicity in morphological observation with 100 µg/ml or less LAE supplement (Fig. 1b). However, cell number was significantly reduced by high dose (250 µg/ml) of LAE supplement for 48 hours (Fig. 1a). These results indicated that low dose LAE did not alter breast cancer cell proliferation.

Next, we examined the LAE role in cell migration by wound healing assays with 50 and 250 µg/ml LAE supplement for 36 hours. Previous studies found that FBS promote cell proliferation to alter cell migration. To rule out the effects of FBS, all experiments were performed in an FBS-free medium. Cell wound healing distance analysis showed that LAE supplement significantly reduced the wound healing distance in three subtypes cells, suggesting cell migration was suppressed (Fig. 1c). Interestingly, 50 or 250 µg/ml LAE supplement showed similarly inhibitory effect on SK-BR-3 cells migration, while LAE supplement inhibited cell migration of MDA-MB-231 and HCC1806 cells in a dose dependent manner.

Nuciferine, a major bioactive component of lotus, inhibits the growth of cancer cells and breast cancer-associated bone loss [24–25]. To verify if nuciferine is involved in the inhibition of cell migration, SK-BR-3, MDA-MB-231 and HCC1806 cells were treated with nuciferine supplement. Cell viability (Supplementary Fig. 1a) and migration (Supplementary Fig. 1b) were unchanged in the presence of high dose of nuciferine, indicating that nuciferine was not associated with LAE-inhibited migration. Together, these results indicate that LAE supplement suppresses ER⁻ breast cancer cells migration independent of nuciferine.

3.2 LAE-inhibited cell migration is independent of autophagy and Wnt signaling

Autophagic process is recognized to suppress cancer metastasis through decreasing EMT [26]. To determine if LAE-suppressed cell migration was regulated by autophagy and EMT-related Wnt/β-catenin signaling pathways directly, we examined β-catenin, snail, p62 and LC3 proteins levels by immunoblotting. Elevated LC3 protein level was detected in cells with high dose (100, 250 µg/ml) of LAE supplement. However, p62 protein level was similarly in cells with low and high dose of LAE supplement (Supplementary Fig. 1a). To address the controversial results, we used autophagy/lysosome inhibitor chloroquine (CQ) to inhibit autophagy flux. Inhibition of autophagy flux did not alter cell migration, revealed by wound healing distance (Supplementary Fig. 1b). Unchanged β-catenin and snail protein levels were detected in three subtypes of breast cancer cells with low or high dose of LAE supplement

(Supplementary Fig. 1c). To further determine the role of Wnt signaling on the effect of LAE-inhibited cell migration, LiCl was used to activate the Wnt signaling. Wound healing assays revealed that the activation of Wnt signaling did not reverse the cell migration inhibited by LAE (Supplementary Fig. 1d), suggesting LAE-suppressed cell migration was independent of Wnt signaling. Thus, LAE-suppressed cell migration is independent of autophagy and Wnt signaling.

3.3 TGF- β 1 signaling is associated with LAE-suppressed cell migration

Considerable literatures showed that TGF- β 1 is recognized as a major regulator of EMT to regulate tumor metastasis [27–29]. Then we tested the protein level of TGF- β 1 after treating with LAE. The results showed that TGF- β 1 was obviously reduced in SK-BR-3 and HCC1806 cells at both low (even 10 μ g/ml) and high concentrations (Fig. 2a). To further determine whether TGF- β 1 signal was required for LAE-inhibited cell migration, 10 ng/ml TGF- β 1 growth factor with 100 μ g/ml LAE was used to treat SK-BR-3, MDA-MB-231 and HCC1806 cells. LAE-inhibited cell migration was significantly restored by the addition of TGF- β 1 growth factor, as revealed by wound healing assays (Fig. 2b). Consistent with literature, TGF- β 1 growth factor changed cells morphology from epithelial to mesenchymal transition (EMT), which was suppressed by LAE supplement (Fig. 2c). Thus, these observations demonstrate that TGF- β 1 signal is involved in LAE-inhibited cell migration via inhibition of EMT.

3.4 SMAD3 and Erk1/2 phosphorylation are inhibited

To explore the signaling pathway involved in LAE-inhibited cell migration, we examined the canonical TGF- β 1/SMAD3 signaling pathway and the non-canonical TGF- β 1/PTEN/AKT and MAPKs signaling pathways [30–31]. Immunoblotting analysis showed that AKT phosphorylation and PTEN levels were unchanged in three subtypes of breast cancer cells with LAE supplement (Fig. 3a). Similar p38 and JNK phosphorylation levels were detected in cells with and without LAE supplement (Fig. 3a). However, the phosphorylated Erk1/2 level was significantly decreased in SK-BR-3 and HCC1806 cells even at a low LAE supplement (10 μ g/ml), and mildly reduced in MDA-MB-231 cells (Fig. 3b). Interestingly, with LAE supplement, the phosphorylated SMAD3 level was significantly reduced in SK-BR-3 and MDA-MB-231 cells, but not in HCC1806 cells (Fig. 3c). Moreover, TGF- β 1 supplement significantly restored Erk1/2 and SMAD3 phosphorylation levels reduced by LAE supplement in three subtypes of breast cancer cells (Fig. 3d).

Since mutant p53 mediates the TGF- β 1 signaling pathway via Erk1/2 as well as SMAD3 signals, and the p53 in all of the three subtypes of breast cancer cells are mutant [32–35]. To address this issue, we examined the mutant p53 protein level in three subtypes of breast cancer cells with LAE supplement. Although p53 level was unchanged in SK-BR-3 and HCC1806 cells with LAE supplement compared to cells without LAE supplement, immunoblotting analysis showed that p53 level was significantly decreased in MDA-MB-231 cells with LAE supplement in a dose dependent manner (Fig. 3e). It provides another explanation for the effect of LAE on the inhibition of the metastasis in MDA-MB-231 cells via

Erk1/2 as well as SMAD3 signals. Above all, these results demonstrate that LAE inhibited the cell migration of ER- breast cancer cells via suppressing the phosphorylation of TGF- β 1/Erk1/2 as well as TGF- β 1/SMAD3 signals.

3.5 LAE inhibits TGF- β 1-related cell migration via downregulating hydrogen peroxide

Reactive oxygen species (ROS), especially hydrogen peroxide (H_2O_2) are recognized as associated with tumor metastasis [36–38]. TGF- β -related signaling promotes hydrogen peroxide production in several types of cells [39–41]. Simultaneously, H_2O_2 enhances TGF- β -mediated EMT via SMAD and MEK/ERK signaling [42]. To determine if intracellular H_2O_2 participate in the TGF- β 1-related inhibition of cell migration, we first measured H_2O_2 level in cells with LAE supplement for 48 hours. With LAE supplement, intracellular H_2O_2 level was significantly downregulated in MDA-MB-231 and HCC1806 cells, but not in SK-BR-3 (Fig. 4a). Consistently, H_2O_2 scavenger catalase protein level was significantly increased even at a low concentration LAE supplement (10 μ g/ml) (Fig. 4b). However, the proteins levels of SOD1, NOX2 and NOX4, responsible for the production of large amounts of ROS, were similarly in cells with or without LAE supplement. Thus, LAE supplement reduced H_2O_2 production in MDA-MB-231 and HCC1806 cells, likely via enhancing the catalase level.

To determine if downregulated H_2O_2 level caused LAE-induced inhibition of breast cancer cell migration, we used 25 nM H_2O_2 with 100 μ g/ml LAE supplement to treat SK-BR-3, MDA-MB-231 and HCC1806 cells. Wound healing distance analysis showed that addition of H_2O_2 significantly restored LAE-inhibited cell migration in MDA-MB-231 and HCC1806 cells (Fig. 4c). However, SK-BR-3 cells displayed similarly cell migration with or without H_2O_2 supplement, suggesting other signals involved in LAE-inhibited cell migration. Interestingly, elevated intracellular H_2O_2 level was significantly detected in MDA-MB-231 and HCC1806 cells with 10 ng/ml TGF- β 1 and 100 μ g/ml LAE compared to cells with single LAE supplement, but not in SK-BR-3 cells (Fig. 4d). These findings indicate that LAE inhibits TGF- β 1-related cell migration via downregulating hydrogen peroxide in MDA-MB-231 and HCC1806.

3.6 LAE inhibits tumor metastasis

To investigate the inhibitory effects of LAE on metastasis *in vivo*, we used subcutaneous xenograft nude mice model to examine tumor metastasis. The schedule of experiments was showed in Figure. 5a. To exclude the toxicity of LAE *in vivo*, xenograft nude female mice were fed with 0.5% LAE for 56 days and body weight of mice was monitored. Body weight in mice with and without LAE supplement was similar (Fig. 5b), indicating that LAE supplement did not cause significant systemic toxicity. Consistent with the above results, LAE supplement did not affect the growth of in-situ tumor compared to controls (Fig. 5c). However, the number and weight of new metastatic tumors derived from the primary tumor were significantly decreased in mice with LAE supplement compared to controls (Fig. 5d). Immunoblotting assays showed that LAE did not inhibit the phosphorylation of Erk1/2 and SMAD3 in liver. However, LAE supplement significantly downregulated the phosphorylation of Erk1/2 but not the phosphorylation of

SMAD3 in lung (Fig. 5e). These data indicate that the effect of LAE on SMAD3 and Erk1/2 is dependent on tissue environment or extracellular signal stimulation.

Next, we used intravenous injection nude mice model to investigate the anti-metastatic role of LAE. The experiments schedule was showed in Fig. 6a. Strikingly, LAE supplement significantly suppressed mice body weight loss induced by tumor compared to control mice (Fig. 6b), which indicated that LAE may have the effect of improving the weight loss in patients with advanced breast cancer. And the number and weight of new formed metastatic tumors were significantly decreased in mice with LAE supplement compared to control animals (Fig. 6c). Interestingly, we found less lung congestion in mice with LAE supplement than that of control animals (Fig. 6d). The number and size of metastatic tumor nodules in lung and liver was decreased significantly. H&E staining revealed inflammatory infiltration and vascular invasion in lung of mice, while LAE supplement improved inflammatory infiltration and vascular invasion in lung of mice (Fig. 6e). In liver, LAE supplement significantly decreased the number of large metastatic tumors. Mice with LAE supplement showed significantly downregulated the phosphorylation of SMAD3 in lung and liver, but not the phosphorylation of Erk1/2 (Fig. 6f). Taken together, these data suggest that LAE inhibited tumor metastasis via suppressing the phosphorylation of SMAD3 and Erk1/2, which is dependent on different tissue environment and extracellular signal.

3.7 Extracellular matrix signaling are associated with LAE-suppressed cell migration

To further explore the role of LAE in inhibiting cell migration, we performed RNA-sequence to analysis genes change in SK-BR-3, HCC1806 and MDA-MB-231. After LAE treatment, all the differentially expressed genes were analyzed by (Gene Ontology) GO method in DAVID with the standard of $P < 0.001$ and $FDR < 1$. Results showed that extracellular matrix (ECM) organization, oxidation-reduction process, negative regulation of cell migration and cell adhesion were significantly enriched by LAE supplement in MDA-MB-231 (Supplementary Fig. 2a), and oxidation-reduction process, extracellular matrix organization, cell-cell adhesion and cell migration were significantly enhanced by LAE supplement in SK-BR-3 (Supplementary Fig. 2b), while extracellular region part and extracellular matrix part were significantly changed by LAE supplement in HCC1806 (Supplementary Fig. 2c). Furthermore, Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway analysis found that pathways involved in ECM-receptor interaction and focal adhesion as well as their related genes were significantly enriched (Supplementary Fig. 2d-f) in SK-BR-3, HCC1806 and MDA-MB-231. Notably, the ECM components and cell adhesion could be induced by the TGF- β 1 and FAK signaling pathway [43–46]. Taken together, our results demonstrate that LAE-suppressed cell migration are associated with multiple different pathways in different cell lines, likely via extracellular matrix signaling.

4. Discussion

Breast cancer, especially the ER- breast cancer, does harm to the health of women because of its metastasis. It's recognized that more than 90% of breast cancer deaths are the results of tumor

metastasis [47], primarily to the bone, lung, liver, brain and lymph nodes [48]. Tumor metastasis is the dissemination of cancer cells from the initial site to distant organs, including epithelial-mesenchymal transition (EMT), cell migration, local tissue invasion, intravasation, extravasation, and metastatic niche formation. EMT endues cells not only migration ability but invasive properties and stem-like cells functions [49–51]. In the present study, we found that LAE inhibited EMT and cell migration via Erk1/2 and SMAD3 and reducing H₂O₂ production in ER- breast cancer even at a low concentration of 10 µg/ml. Our findings provided a new possible therapeutic approach for ER- breast cancer.

Although the anti-bacterial, anti-oxidant, and anti-obesity effects of lotus and its extracts have been known for many years [24, 52], little is known about the activity of LAE on cancer cell migration. We demonstrated that LAE had the potential inhibitory activity for cell migration and EMT *in vitro* and *in vivo*. However, few components of lotus leaf with the effects of anticancer have revealed so far. Previous studies showed that nuciferine, a major bioactive component of lotus, inhibits the growth of cancer cells and breast cancer-associated bone loss [25, 53]. However, our results showed that nuciferine did not participate in LAE-inhibited migration. Thus, further experiments are required for identification of bioactive compound in LAE of inhibiting cell migration and metastasis.

Many signaling pathways are associated with the migration of tumor cells. TGF-β1 signaling pathway has been recognized to be the main inducer. TGF-β1 modulates the EMT process by the canonical TGF-β1/SMAD3 signaling pathway and the non-canonical TGF-β1/PTEN/AKT and MAPKs signaling pathways. Our results showed that LAE inhibited TGF-β1 signaling pathway via Erk1/2 and SMAD3 in SK-BR-3, MDA-MB-231 and HCC1806 breast cancer cells. Furthermore, several studies reported that SMAD3 increases the adhesive ability of hepatocellular carcinoma (HCC) cells by the form of exosome, which promotes the lung metastasis [54]. These might explain the reason why LAE treatment significantly changed the ECM and cell-cell adhesive signaling pathway, which were mediated by the TGF-β1 and FAK signaling pathway according to the results of GSEA. Therefore, the different effect of LAE on SMAD3 and Erk1/2 in mice model could be possibly explained by the different extracellular matrix compounds and environment in liver and lung (Fig. 5e, Fig. 6e-f). Thus, ECM and FAK might be another targets to inhibit the cell migration and metastasis induced by LAE.

Literatures have demonstrated that activated Erk1/2 stimulates the metastasis in cancers by targeting Snail, Slug and matrix metalloproteinases (MMPs) [55]. However, in the present work, we found that LAE suppressed the activation of Erk1/2 and SMAD3 as well as reduced the intracellular H₂O₂ levels, which were regulated by TGF-β1 signaling pathway in MDA-MB-231 and HCC1806 cells, but not in SK-BR-3 cells. Exogenous H₂O₂ significantly restored LAE-inhibited cell migration in MDA-MB-231 and HCC1806 cells, but not SK-BR-3, revealed that H₂O₂ was not involved in LAE-inhibited cell migration in SK-BR-3 cells. Interestingly, although LAE treatment significantly reduced the phosphorylation levels of Erk1/2 and SMAD3, the expression of TGF-β1 protein which regulated them both did not change significantly in MDA-MB-231 cells. This might be explained by the significantly decreased expression of mutant p53 in MDA-MB-231 cells because the essential effects of mutant p53 in mediating the TGF-β1 signaling pathway [32]. Taken together, our findings strongly suggested that LAE inhibit EMT by primarily targeting the TGF-

β 1 pathway, and the underlying mechanism of LAE-suppressed cell migration is differently dependent on cell lines.

Lotus leaf has been used as “Chinese medicine” for a long history [56]. In the present study, we demonstrated that LAE inhibited the cell migration and metastasis of ER- breast cancer without significant cytotoxicity. Further identification and clinical studies are required to make LAE as a novel anticancer agent in clinical therapy, and our findings strongly contribute to achieve this goal.

Conclusion

In this study, our data showed that LAE significantly inhibits the ER⁻ breast cancer cell migration and metastasis via the TGF- β 1/Erk1/2 and TGF- β 1/SMAD3 signaling, accompanied by reduced intracellular H₂O₂ level in ER- breast cancer cells. RNA-sequence analysis also showed that extracellular matrix signaling pathways and FAK might be another possible pathways to inhibit the cell migration induced by LAE. These results suggested that LAE has the potential to screen out novel drugs for the treatment of ER⁻ breast cancer clinically.

Abbreviations

TGF- β 1: transforming growth factor- β 1; ER+/PR+: estrogen receptor positive/progesterone receptor positive; HER2+: human epidermal growth factor receptor positive; TNBC: triple negative; LAE: Lotus leaf alcoholic extract; CQ: Chloroquine diphosphate; H&E: hematoxylin and eosin; ROS: Reactive oxygen species; ECM: extracellular matrix.

Declarations

Consent for publication Funding

Not applicable.

Acknowledgements

Not applicable.

Funding

This research was supported by NSFC grants 31471321 (Z.H.), National Young 1000 Talents Plan (Z.H.), Jiangsu Province Recruitment Plan for High-level, Innovative and Entrepreneurial Talents (Z.H.) and Shandong University 2018TB019.

Ethics approval and consent to participate

This study was approved by the ethics committee of Jiangnan University. All the experiments were in accordance with institution guidelines.

Availability of data and materials

The datasets from the present study are available from the corresponding author upon request.

Competing interests

The authors declare to have no conflict of interest.

Authors' contributions

Yuelin Tong (1946331910@qq.com): Designed and performed most part of experimental work and wrote the scientific paper, Zhongwei Li (1072315826@qq.com): Searched the literatures and participated in animal experiment, Yikuan Wu (873334081@qq.com): Experimental work, Shenglong Zhu (490149835@qq.com): Study design, Keke Lu (3545978755@qq.com): Manuscript revision, Zhao He (zhaohe7711@qq.com): Study design and manuscript revision.

References

1. Lin, G. , Zhu, S. , Wu, Y. , Song, C. , & He, Z. ω -3 free fatty acids and all-trans retinoic acid synergistically induce growth inhibition of three subtypes of breast cancer cell lines. *Scientific Reports*.2017;7:2929.
2. Ji, M. , Qingli, Z. , & Jing, L. Effect of aspirin on invasion and migration of her-2-positive breast cancer skbr3 cells. *Chinese Journal of Breast Disease(Electronic Edition)*.2018; 1.
3. Xue, R. , Zhi, L. , & Emergency, D. O. Recurrence and metastasis risk of breast cancer in different molecular types. *China Continuing Medical Education*.2018;14.
4. Lin, K. , Baritaki, S. , Militello, L. , Malaponte, G. , Bevelacqua, Y. , & Bonavida, B. The role of b-raf mutations in melanoma and the induction of emt via dysregulation of the nf-kb/snail/rkip/pten circuit. *Genes & Cancer*.2010;1: 409-420.
5. Kasai, H. , Allen, J. T. , Mason, R. M. , Kamimura, T. , & Zhang, Z. Tgf- β 1 induces human alveolar epithelial to mesenchymal cell transition (emt). *Respiratory Research*.2005;6: 56.
6. Jin, Z. , Hongqing, C. , Lixin, S. , Panpan, Z. , Meng, C. , & Feng, Z. , et al. Lgr5, a novel functional glioma stem cell marker, promotes emt by activating the wnt/ β -catenin pathway and predicts poor survival of glioma patients. *Journal of Experimental & Clinical Cancer Research*. 2018;37: 225.

7. Liu, R. M. , & Pravia, K. A. G. Oxidative stress and glutathione in tgf- β -mediated fibrogenesis. *Free Radical Biology & Medicine*.2010;48: 1-15.
8. Fumihito, Ishikawa, Emi, Kaneko, Tadashi, & Sugimoto, et al. A mitochondrial thioredoxin-sensitive mechanism regulates tgf- β -mediated gene expression associated with epithelial–mesenchymal transition. *Biochemical and Biophysical Research Communications*.2014; 443:821-827.
9. Annes, & J., P. Making sense of latent tgf β activation. *Journal of Cell Science*. 2003;116:217-224.
10. Son, Y. O. , Wang, L. , Poyil, P. , Budhraj, A. , Hitron, J. A. , & Zhang, Z. , et al. Cadmium induces carcinogenesis in beas-2b cells through ros-dependent activation of pi3k/akt/gsk-3 β / β -catenin signaling. *Toxicology and Applied Pharmacology*.2012;264: 153-160.
11. Wang, J. , Luo, B. , Li, X. , Lu, W. , Yang, J. , & Hu, Y. , et al. Inhibition of cancer growth in vitro and in vivo by a novel ros-modulating agent with ability to eliminate stem-like cancer cells. *Cell Death and Disease*.2017;8:e2887.
12. Nishikawa, M. Reactive oxygen species in tumor metastasis. *Cancer Letters*.2008;266: 0-59.
13. Glasauer, A., & Chandel, N. Targeting antioxidants for cancer therapy. *Biochemical Pharmacology*.2014;92: 90-101.
14. WULF DRÖ` GE. Free radicals in the physiological control of cell function. *Physiol Rev*.2002;82:47-95.
15. Huang, C. F. , Chen, Y. W. , Yang, C. Y. , Lin, H. Y. , Way, T. D. , & Chiang, W. , et al. Extract of lotus leaf (*nelumbo nucifera*) and its active constituent catechin with insulin secretagogue activity. *Journal of Agricultural and Food Chemistry*.2011;59:1087-1094.
16. Lin, H. Y. , Kuo, Y. H. , Lin, Y. L. & Chiang, W. Antioxidative effect and active components from leaves of lotus (*nelumbo nucifera*). *Journal of Agricultural and Food Chemistry*.2009;57: 6623-6629.
17. Mingzhi, Z. , Ting, L. , & Mingquan, G.. Current advances in the metabolomics study on lotus seeds. *Frontiers in Plant Science*.2016;7:891.
18. Du, H. , You, J. S. , Zhao, X. , Park, J. Y. , & Chang, K. J. Antiobesity and hypolipidemic effects of lotus leaf hot water extract with taurine supplementation in rats fed a high fat diet. *Journal of Biomedical Science*.2010;17 :*Suppl 1*(Suppl 1)-S42.
19. Ha, J. Y. , Lee, K. , Park, J. , Dong, A. , & Shin, H. S. Cytoprotective activity of lotus (*nelumbo nucifera gaertner*) leaf extracts on the mouse embryonic fibroblast cell. *Food science and biotechnology*.2010;19:1171-1176.
20. Li, M. , & Xu, Z. Quercetin in a lotus leaves extract may be responsible for antibacterial activity. *Archives of Pharmacal Research*.2008;31: 640-644.
21. Ham, Y. K. , Hwang, K. E. , Song, D. H. , Kim, Y. J. , Shin, D. J. , & Kim, K. I. , et al. Lotus (*nelumbo nucifera*) rhizome as an antioxidant dietary fiber in cooked sausage: effects on physicochemical and sensory characteristics. *Korean Journal for Food Science of Animal Resources*.2017;37: 219-227.
22. Gomes, L. R. , Letícia F Terra, Rosângela Am Wailemann, Labriola, L. , & Sogayar, M. C. Tgf-1 modulates the homeostasis between mmps and mmp inhibitors through p38 mapk and erk1/2 in highly invasive breast cancer cells. *BMC Cancer*.2012;12:26.

23. Hsieh, H. L. , Wang, H. H. , Wu, W. B. , Chu, P. J. , & Yang, C. M. Transforming growth factor- β 1 induces matrix metalloproteinase-9 and cell migration in astrocytes: roles of ros-dependent erk- and jnk-nf-kb pathways. *Journal of Neuroinflammation*.2010;7:88.
24. Ji, K. E. , Kyoung, L. S. , Kwang-Kyun, P. , Hwa, S. S. , Rim, K. K. , & Won-Yoon, C. Liensinine and nuciferine, bioactive components of *nelumbo nucifera*, inhibit the growth of breast cancer cells and breast cancer-associated bone loss. *Evidence-Based Complementary and Alternative Medicine*.2017;2017:1-12.
25. Qi, Q. , Li, R. , Li, H. Y. , Cao, Y. B. , Bai, M. , & Fan, X. J. , et al. Identification of the anti-tumor activity and mechanisms of nuciferine through a network pharmacology approach. *Acta Pharmacologica Sinica*.2016; 37: 963–972.
26. L. Poillet-Perez, G. Despouy, R. Delage-Mourroux, M. (2015). Interplay between ros and autophagy in cancer cells, from tumor initiation to cancer therapy. *Redox Biology, Transforming growth factor-beta and oxidative stress interplay: implications in tumorigenesis and cancer progression*.2016;4: 184-192.
27. Jelena, K. , Drenka, T. , Slavko, M. , & Santibanez, J. F. Transforming growth factor-beta and oxidative stress interplay: implications in tumorigenesis and cancer progression. *Oxidative Medicine and Cellular Longevity*.2015;2015: 1-15.
28. Xu, Zhang, Peng, Meng, Shao, & Xueyan, et al. Sall4 activates tgf- β /smad signaling pathway to induce emt and promote gastric cancer metastasis. *Cancer management and research*.2018.
29. Y.P. Wang, G.R. Yu, M.J. Lee, S.Y. Lee, I.S. Chu, S.H. Leem, D.G. Kim. Lipocalin-2 negatively modulates the epithelial-to-mesenchymal transition in hepatocellular carcinoma through the epidermal growth factor (tgf-beta1)/lcn2/twist1 pathway. *Hepatolog*.2013;58.
30. Wrighton, K. H. , Lin, X. , & Feng, X. H. Phospho-control of tgf-beta superfamily signaling. *Cell Research*.2009;19: 8.
31. Boldbaatar, A. , Lee, S. , Han, S. , Jeong, A. , Ka, H. , & Buyanravjikh, S. , et al. Eupatolide inhibits the tgf- β 1-induced migration of breast cancer cells via downregulation of smad3 phosphorylation and transcriptional repression of alk5. *Oncology Letters*.2017; 14:6031-6039.
32. Lin, S. , Yu, L. , Yang, J. , Liu, Z. , Karia, B. , & Bishop, A. J. R. , et al. Mutant p53 disrupts role of shca protein in balancing smad protein-dependent and -independent signaling activity of transforming growth factor- β (tgf- β) *. *Journal of Biological Chemistry*.2011; 286:44023-44034.
33. Marzec, K. A. , Lin, M. Z. , Martin, J. L. , & Baxter, R. C. Involvement of p53 in insulin-like growth factor binding protein-3 regulation in the breast cancer cell response to dna damage. *Oncotarget*.2015;6: 26583-26598.
34. Chander, H. , Brien, C. D. , Truesdell, P. , Watt, K. , Meens, J. , & Schick, C. , et al. Toca-1 is suppressed by p53 to limit breast cancer cell invasion and tumor metastasis. *Breast Cancer Research*.2014 16 3413.
35. Hennel, R. , Brix, N. , Seidl, K. , Ernst, A. , Scheithauer, H. , & Belka, C. , et al. Release of monocyte migration signals by breast cancer cell lines after ablative and fractionated γ -irradiation. *Radiation*

- Oncology.2014;9:85.
36. Liu, Z. , Li, S. , Cai, Y. , Wang, A. , He, Q. , & Zheng, C. , et al. Manganese superoxide dismutase induces migration and invasion of tongue squamous cell carcinoma via h₂O₂-dependent snail signaling. *Free Radical Biology & Medicine*.2012;53: 44-50.
 37. Wei, L. , Lun, Z. , Xin, C. , Zhengdong, J. , Liang, Z. , & Qingyong, M. Hyperglycemia promotes the epithelial-mesenchymal transition of pancreatic cancer via hydrogen peroxide. *Oxidative Medicine and Cellular Longevity*.2016; 1-9.
 38. Satooka, H. , & Hara-Chikuma, M. Aquaporin-3 controls breast cancer cell migration by regulating hydrogen peroxide transport and its downstream cell signaling. *Molecular and Cellular Biology*.2016; 36:1206-1218.
 39. Thannickal, V. J. , & Fanburg, B. L. Activation of an h₂O₂-generating nadh oxidase in human lung fibroblasts by transforming growth factor- β 1. *Journal of Biological Chemistry*.1995;270:30334-30338.
 40. Huang, L. S. , Jiang, P. , Feghalibostwick, C. , Reddy, S. P. , Jgn, G. , & Natarajan, V. Lysocardiolipin acyltransferase regulates tgf- β mediated lung fibroblast differentiation. *Free Radical Biology & Medicine*.2017:112.
 41. Hongqiao, Zhang, Henry, Jay, & Forman. S1-4 - tgf β 1 rapidly activates src through a non-canonical redox mechanism. *Free Radical Biology and Medicine*.2014:75.
 42. Iizuka, D. , Sasatani, M. , Barcellos-Hoff, M. H. , & Kamiya, K. Hydrogen peroxide enhances tgf β -mediated epithelial-to-mesenchymal transition in human mammary epithelial mcf-10a cells. *Anticancer research*.2017;37:987-995.
 43. Yu, L. , Liu, X. , Cui, K. , Di, Y. , Xin, L. , & Sun, X. , et al. Snd1 acts downstream of tgf- β 1 and upstream of smurf1 to promote breast cancer metastasis. *Cancer Research*.2015;75: 1275-1286.
 44. Sahoo, S. S. , Quah, M. Y. , Nielsen, S. , Atkins, J. , & Tanwar, P. S. Inhibition of extracellular matrix mediated tgf- β signalling suppresses endometrial cancer metastasis. *Oncotarget*.2017;8: 71400-71417.
 45. Nakashio, T. , Narita, T. , Akiyama, S. , Kasai, Y. , & Kannagi, R. Adhesion molecules and tgf-beta1 are involved in the peritoneal dissemination of nUGC-4 human gastric cancer cells. *International Journal of Cancer*.1997;70: 612-618.
 46. Maschler, S. , Grunert, S. , Danielopol, A. , Beug, H. , & Wirtl, G. Enhanced tenascin-c expression and matrix deposition during ras/tgf- β -induced progression of mammary tumor cells. *Oncogene*.2004; 23: 3622-3633.
 47. Han, D. , Wu, G. , Chang, C. , Zhu, F. , & Zhang, L. Disulfiram inhibits tgf- β -induced epithelial-mesenchymal transition and stem-like features in breast cancer via erk/nf- κ b/snail pathway. *Oncotarget*.2015;6: 40907-40919.
 48. Z.J. Liu, G.L. Semenza, H.F. Zhang. Hypoxia-inducible factor 1 and breast cancer metastasis, *J Zhejiang Univ Sci B*. 2015;16: 32-43.

49. Mani, S. A. , Guo, W. , Liao, M. J. , Eaton, E. N. , Ayyanan, A. , & Zhou, A. Y. , et al. The epithelial-mesenchymal transition generates cells with properties of stem cells. *Cell*.2008;133:704-715.
50. Ota, I. , Masui, T. , Kurihara, M. , Yook, J. I. , Mikami, S. , & Kimura, T. , et al. Snail-induced emt promotes cancer stem cell-like properties in head and neck cancer cells. *Oncology Reports*. 2016;35:261-266.
51. Wen-Tai, C. , Yu-Fang, H. , Huei-Yu, T. , Chien-Chin, C. , Chang-Hwa, C. , & Soon-Cen, H. , et al. Foxm1 confers to epithelial-mesenchymal transition, stemness and chemoresistance in epithelial ovarian carcinoma cells. *Oncotarget*.2015;6: 2349-2365.
52. Shukla, S. , Park, J. , Park, J. H. , Lee, J. S. , & Kim, M. . Development of novel meju starter culture using plant extracts with reduced bacillus cereus counts and enhanced functional properties. *Scientific Reports*.2017;7:11409.
53. Lee, D. B. , Kim, D. H. , & Je, J. Y. Antioxidant and cytoprotective effects of lotus (nelumbo nucifera) leaves phenolic fraction. *Preventive Nutrition and Food Science*.2015;20: 22-28.
54. Qihan, F. , Qi, Z. , Yu, L. , Jiaqi, Y. , Gang, N. , & Qi, C. , et al. Primary tumor-derived exosomes facilitate metastasis by regulating adhesion of circulating tumor cells via smad3 in liver cancer. *Oncogene*.2018;37: 6105–6118.
55. Wu, Z. , Wang, T. , Fang, M. , Huang, W. , & Yan, W. Mfap5 promotes tumor progression and bone metastasis by regulating erk/mmp signaling pathways in breast cancer. *Biochemical and Biophysical Research Communications*.2018;498:495-501.
56. Y. Zhang, X. Lu, S. Zeng, X. Huang, Z. Guo, Y. Zheng, Y. Tian, B. Zheng. Nutritional composition, physiological functions and processing of lotus (nelumbo nuciferagaertn.) seeds: a review. *Phytochemistry Review*.2015;14:321-334.

Figures




Figure 1

LAE effect on ER- breast cancer cells migration. SK-BR-3, MDA-MB-231 and HCC1806 breast cancer cells were treated with different concentrations LAE for 48h. a The viability of LAE-treated cells was determined by CCK8 assays. b The cell morphology after treating with LAE. c Wound healing assays after treating with LAE for 36h (x100 magnification).




Figure 2

TGF-β1 signaling with LAE supplement. a TGF-β1 was determined by immunoblotting under different concentrations of LAE (0, 10, 25, 50, 100, 250 μg/ml). b Wound healing assays after treating with 100 μg/ml LAE alone or combined with 10 ng/ml TGF-β1 for 36h (x100 magnification). c The morphology of SK-BR-3, MDA-MB-231 and HCC1806 breast cancer cells after exposure to 100 μg/ml LAE alone or combined with TGF-β1 (10 ng/ml) for 48h (x 400 magnification).




Figure 3

SMAD3 and Erk1/2 phosphorylation levels. a PTEN, p-Akt, Akt, p-p38, p38, p-JNK and JNK were determined by immunoblotting under different concentrations of LAE (0, 10, 25, 50, 100, 250 $\mu\text{g/ml}$). b p-Erk1/2 and Erk1/2 were examined by immunoblotting; c p-SMAD3 and SMAD3 were examined by immunoblotting; d p-SMAD3 and p-Erk1/2 proteins derived from cells treated with 100 $\mu\text{g/ml}$ alone or combined with TGF- β 1 (10ng/ml). e p53 protein level was examined by immunoblotting.




Figure 4

Hydrogen peroxide level in LAE-inhibited cell migration. a The content of intracellular H₂O₂ was determined after treating with 100 µg/ml LAE for 48h in SK-BR-3, MDA-MB-231 and HCC1806. b NOX4, NOX2, catalase and SOD1 proteins levels were examined by immunoblotting with different concentrations of LAE (0, 10, 25, 50, 100, 250 µg/ml). c Wound healing assays after treating with LAE alone or combined with 25nM H₂O₂ for 36h (x100 magnification). d The content of intracellular H₂O₂

was determined after treating with 100 μ g/ml LAE alone or combined with TGF- β 1 (10ng/ml) for 48h in SK-BR-3, MDA-MB-231 and HCC1806. Values represent the mean \pm SD (n = 3).




Figure 5

LAE effect on tumor metastasis in xenograft model. a Flowchart of animal experiments. MDA-MB-231 (5 x10⁷) were subcutaneously injected in the right flank of mice. After one week of injection, the tumor bearing mice were randomly subdivided into 2 groups: Control; 0.5% LAE. b Growth curves of body weight in xenograft model. c Upper panel: Representative tumors and weight of tumors. Lower panel: Growth curves of tumor volume in xenograft model. d The numbers and weights of new metastatic tumors derived from the primary tumor. e p-SMAD3 and p-Erk1/2 proteins levels were examined by immunoblotting in lung and liver tissues (xenograft model).




Figure 6

LAE effect on tumor metastasis in tail vein injection model. a Flowchart of animal experiments. MDA-MB-231 (5×10^7) were intravenously injected in nude mice. After one week of injection, the tumor-bearing mice were randomly subdivided into 2 groups: Control; 0.5% LAE. b Growth curves of body weight in intravenous injection nude mouse model. c Left panel: Representative new metastatic tumors which formed after injecting the MDA-MB-231 breast cancer cells into the vein of mice; Middle and right panels:

The numbers and weights of new metastatic tumors. d Representative lung and liver tissues of intravenous injection nude mice. White arrows represent the metastatic tumor nodules. e Metastatic tumor nodules in the lung and liver tissues were examined by H&E staining (White arrow: Cancer cells invade the lumen of a large blood vessel. Black arrow: Metastatic tumor nodules substitute a segment of the vascular wall or exist in the liver tissues. Red arrow: inflammatory infiltration in blood vessel). f p-SMAD3 and p-Erk1/2 levels were examined by immunoblotting in lung and liver tissues (tail vein injection model). Values represent the mean \pm SD (n = 3).

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryFigures.doc](#)