

Quality of sleep and associated factors among people living with HIV/AIDS on follow up at Ethiopian Zewditu Memorial Hospital,2018

Nebiyu Mengistu

Dilla University

Telake Azale

University of Gondar

Solomon Yimer (✉ sopsycha@gmail.com)

Dilla University <https://orcid.org/0000-0003-0110-3792>

Mahilet Fikreyesus

Amanuel Mental Specialized Hospital

Elsa Melaku

Amanuel Mental Specialized Hospital

Seid Shumye

Dilla university

Research

Keywords: HIV/AIDS, Sleep habits, AIDS, Sleep hygiene, Prevalence Studies

Posted Date: October 30th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-26745/v3>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on January 4th, 2021. See the published version at <https://doi.org/10.1186/s41606-020-00056-w>.

Abstract

Background: Sleep disturbance is a common complaint in people living with HIV/AIDS. Despite the influence of sleep disturbance on treatment adherence, quality of life, work productivity, risk of chronic illness, it remains under-recognized and under-studied in Ethiopia. Therefore it is necessary to produce scientific evidence to fill the knowledge gap and areas of management. The current study aimed to assess sleep quality and its associated factors among people living with HIV/AIDS.

Methods: An institution-based cross-sectional study was utilized among 408 participants who were selected by a systematic random sampling technique at Zewditu memorial hospital from April to May 2018. The Pittsburgh Sleep Quality Index questionnaire was used to measure sleep quality. Ethical clearance was obtained from the joint ethics committee of the University of Gondar and Amanuel Mental Specialized Hospital. Oral informed consent was obtained from each participant. Binary and multivariable logistic regression models were fitted. Odds ratios (OR) with the corresponding 95% confidence interval (95%CI) was computed.

Results: The magnitude of poor sleep quality was 55.6%. Being female [AOR=3.40, 95% CI: (1.80, 6.41)], depression [AOR =3.52, 95% CI: (1.95, 6.32)], CD₄count \leq 200 cells/mm³ [AOR=3.18,95%CI: (1.65,6.13)], duration of HIV/AIDS diagnosis [AOR=3.43,95% CI: (1.61,7.29)], current use of tobacco [AOR=5.69, 95% CI: (2.04,15.9)] and chat or caffeinated drinks [AOR=2.65, 95% CI: (1.06,6.64)] and poor sleep hygiene [AOR=3.55, 95% CI: (1.85, 6.78)] were significantly associated with poor sleep quality.

Conclusions: More than half of the study participants were found to have poor sleep quality. A range factors influence quality of sleep of people with HIV/AIDS. Routine screening of sleep condition among people living with HIV/AIDS and early intervention based on the findings is suggested.

Background

Sleep is a natural process that the brain requires to keep proper functioning and maintain the health of the human body. Sleep occupies one third of human life and its deprivation causes negative consequences that may be physical, cognitive, or emotional. Suffering from inadequate sleep has been associated with chronic medical illness like diabetes, stroke, heart disease and Human immunodeficiency virus or Acquired immune deficiency syndrome (HIV/AIDS) [1, 2].

HIV/AIDS is a chronic, potentially life threatening condition caused by the HIV, which affects mostly the immune system and nervous system. HIV/AIDS is among the most overwhelming health problems throughout the world and especially in developing countries [3]. An approximated 36.7 million people are living with HIV/AIDS in the world. In Ethiopia, the national adult HIV/AIDS prevalence has been reported to be 1.14%. It has been estimated that 542,121 adults and 178,500 children require antiretroviral treatment in Ethiopia [4]. Sleep disturbance is one of the most prevalent symptoms in individuals infected with HIV/AIDS where 40% to 70% of individuals infected with HIV/AIDS are reporting significant sleep disturbances, including difficulty falling asleep, awakenings during the night, and reduced sleep time [5-7].

Sleep disturbance among people living with HIV/AIDS occurs throughout the stages of the infection, but more prevalent in the advanced stage. Despite this fact, still the condition receives little attention[8]. This may be, because sleep disturbance is considered as a normal consequence of the disease and its treatment, or considered to be insignificant in comparison with other complications of HIV infection [9].

But sleep deprivation among people living with HIV/AIDS leads to various effects, such as impaired immune system, hampered physical performance, affected cell growth and repair, deteriorated neuronal connections and neuronal malfunctions [10]. In addition, it may also increase the risk of cardiovascular morbidity and mortality, and the degree of suffering with psychiatric disorders [11].

According to study results, sleep disturbance is not only prevalent, but also there are more intense and distressful symptoms experienced by the HIV population. As a result, such individuals are less likely to adhere to their antiretroviral therapy regimens probably as a result of depression [6, 12-14].

Cross-sectional surveys conducted in China and France showed that prevalence of sleep disturbances and Poor sleep quality was 43.1% and 47% respectively. Prevalence of sleep disturbances differed significantly between those who suffered anxiety and depression and those who did not. Moreover, being male, a smoker, living single, being unemployed or moderately or seriously depressed were significantly associated with poor sleep quality [15, 16].

According to a survey conducted in Latin America countries Mexico and South East Brazil 58.6% and 46.7% had poor sleep quality respectively. Depressive symptoms, illicit drug use, a CD4 count < 200 cells/ μ l, and longer duration of HIV/AIDS diagnosis was positively associated with poor sleep quality[17, 18].

Similarly, studies conducted in Nigeria among HIV-positive outpatients reported that 46.2% in University of Calabar Teaching Hospital and 59.3% in Lagos State University Teaching Hospital had sleep disorders and poor sleep quality respectively. Elevated systolic blood pressure, shorter duration of HIV diagnosis and highly active antiretroviral therapy (HAART) type were associated with sleep disorders and poor sleep quality [19, 20].

Despite this significant health problems and higher prevalence, sleep quality among people with HIV/AIDS remains under-studied with no published study in Ethiopia. Beside that the current study addressed key correlated factors that could be managed by health care providers for better treatment outcome and control of psychological problems among people with HIV/AIDS. Therefore, the aim of this study was to assess the magnitude of sleep quality and its associated factors among people living with HIV/AIDS in Zewditu Memorial Hospital, Ethiopia.

Methods

Study design, period, setting, and population

An institutional-based cross-sectional study was conducted from April to May 2018 in Zewditu Memorial Hospital which is found in Addis Ababa, Ethiopia. Center for Disease Control and prevention (CDC-

Ethiopia) helped the launch of Ethiopia's first antiretroviral therapy (ART) program at Zewditu Memorial Hospital in July 2003, and in March 2005. The program received technical assistance from Johns Hopkins University's TSEHAI Program. Zewditu Memorial Hospital became the largest HIV clinic in Ethiopia and a leading hospital in the treatment of antiretroviral therapy patients. Currently, the hospital treats over 7299 patients each month. There were 17,857 HIV-positive patients having HIV care follow-up in the hospital in 2018. The study population was patients who were attending at ART clinics during the data collection period. All adults who were seriously ill and unable to communicate were not included in the study.

Sample size and sampling technique

Sample size (n) was calculated based on single population proportion formula, by assuming 95% confidence level, the prevalence of poor sleep quality among HIV/AIDS patients which was found to be 59.3% in Nigeria [20] and a precision of 5% between the sample and the parameter was taken. $\alpha = 0.05(95\%) = 1.96$

By considering a 10% non-response rate the final sample size was 408. We used a systematic sampling technique to select the four hundred eight (408) HIV/AIDS patients who were included in our survey. We determined the sampling interval by dividing the total study population who had to follow up during the average 1-month data collection period (3264) by total sample size (408). Hence, the sample interval is eight. We selected the first study participant by lottery method and the next study participants were chosen every 8th interval for interview.

Data collection tools and procedures

Data were collected by trained nurses by face-to-face interviewing of the participant attending ART service using paper and pencil. The questionnaire was pre-tested by taking 5% of the calculated sample size and modified accordingly to easily understand by the study participants. The questionnaire contained socio-demographic characteristics (age, income, education, occupation, marital status, and others), questions to collect data on clinical factors and standard tools to address other independent variables of the study.

Data on the components of sleep quality was collected by using the Pittsburgh Sleep Quality Index (PSQI), a self-report measure instrument composed of 19 items evaluating seven components of sleep. Each part was scored (range: 0-3; higher scores indicating worse sleep). A total global PSQI was derived by summing the seven components (range: 0 to 21; higher scores indicating poor sleep quality). A global PSQI score >5 yielded a diagnostic sensitivity of 89% and specificity of 86.5% ($\kappa=0.75$, $p \leq 0.001$) in distinguishing "good" from "poor" sleepers. "Good sleep" was defined as global PSQI scores of 0-5 and "poor sleep" was global PSQI scores of 6-12[21]. The tool was validated in Ethiopia among community dwellers having Cronbach's alpha of 0.59, sensitivity of 82%, and specificity of 56.2% [22].

Hospital anxiety and depression scale (HADS) were used to assess anxiety and depression. The tool has anxiety subscale (HADS-A) and the depression subscale (HADS-D). It has a cutoff point ≥ 8 for each subscale [22]. HADS had an internal consistency of 0.78 for the anxiety, 0.76 for depression subscales and 0.87 for the full scale in a validation study conducted among HIV Infected Patients in Ethiopia [23]. HIV/AIDS related stigma scale was used to assess stigma. It is a 12-item screening tool developed by Annelies Van Rie, Sohini Sengupta. The tool has 4-point Likert response. Each items were scored with 0 (strongly disagree) and 3 (strongly agree). Participants who scored above the mean score were considered as stigmatized[24, 25]. The scale demonstrated acceptable internal consistency (Cronbach's alpha=0.73) in a study conducted in Uganda[26].

Sleep hygiene index (SHI), a 13-item self-report measure designed to assess the practice of sleep hygiene behaviors. Each item is rated on a five-point scale ranging from 0 (never) to 4 (always). Total scores range from 0 to 52 with a higher score representing poor sleep hygiene[27]. Oslo Social Support Scale (OSS) was used to assess respondents support system which is a 3 items measure of support with internal consistency (Cronbach's $\alpha = 0.50$) [28].

Data quality control issues

Training was given to the data collectors and supervisors on the data collection tool and sampling techniques. Supervision was held regularly during the data collection period both by the researcher, co-investigators and supervisors to check on a daily basis for completeness and consistency.

Analysis

Data were analyzed using SPSS version 20. Description statistics (frequencies, proportions, means, and standard deviations) were used to present the sociodemographic characteristics and the prevalence of sleep quality. Both bivariate and multivariate logistic regression analysis were carried out to see the association of each independent variable with the outcome variable. A p-value of less than 0.05 was considered statistically significant, and an adjusted odds ratio with 95% CI was calculated to determine the association.

Ethical Considerations

Ethical clearance was obtained from the joint ethics committee of the University of Gondar and Amanuel Mental Specialized Hospital and Addis Ababa health bureau based upon Declaration of Helsinki (DoH). The data collectors clearly explained the purpose and importance of the study to each study participant before they proceed into actual activities. Information was collected after obtaining written consent from each participant. Written informed consent was obtained from the study participants and comparison of study participants after oral information about the study, including an assurance that they could withdraw from the study at any time. Confidentiality was maintained by anonymous questionnaire.

Results

A total of 396 study participants were interviewed, giving a response rate of 97.1%. The mean age of the respondents was 38.6 years with SD (± 10.8). The proportion of male to female participants was almost equal (50.3% to 49.7%). More than half of the clients ($n=276$, 69.7%) were Orthodox by religion and most of the respondents were married ($n=208$, 52.5%) and regarding the education level of the respondents 176(44.4%) had attended secondary school (table 1)

HIV related clinical characteristics of the participants

Among the study participants, the majority ($n= 294$, 74.2%) was on clinical stage I, 219 (55.3%) had greater than 200 cells/mm³ of CD₄ count, 296 (74.7%) on the first line regimen of ART drugs and 272 (68.7%) had duration of HIV/AIDS diagnosis greater than 10 years (table 2).

Psycho-social characteristics of the participants

Regarding to psycho-social characteristics, more than half (51.5%) were in high depression severity, 109 (27.5%) of the participants had general anxiety disorder symptoms, 194(49.0%) of them were stigmatized and 164 (41.4%) had poor social support. Among one hundred ninety seven female participants 117 (59.4%) and of one hundred ninety nine male participants 87 (43.7 %) had depression symptoms.

Substance use by participants

Two hundred six (52.0%) the study participants were current substances users. Of them, majority ($n=183$, 46.2%) use alcohol followed by ($n=109$, 27.5%) use tobacco (fig 1).

Poor sleep quality and Factors associated among people with HIV/AIDS

The magnitude of poor sleep quality among people living with HIV/AIDS was found to be 55.6%. Multivariable logistic regression revealed female sex, depression, CD4 less than 200 cells/mm³ count, duration of HIV/AIDS diagnosis, current use of tobacco, current use of chat or caffeinated drinks and poor sleep hygiene were significantly associated with poor sleep quality.

The magnitude of poor sleep quality was found to be more than 3 times higher in females compared to males [AOR=3.40, 95% CI: (1.80, 6.41)]. The odds of having poor sleep quality among depressed respondents were 3.52 times higher as compared to non-depressed respondents [AOR =3.52, 95% CI: (1.95, 6.32)]. The odds of experiencing poor sleep quality among Participants with shorter duration of HIV diagnosis were more than three times as compared to those who had longer duration of diagnosis [AOR=3.43,95% CI : (1.61,7.29)]. Similarly, Individuals whose CD₄ count were less than 200cells/mm³ were 3.18 times more likely to have poor sleep quality than individuals with CD₄ counts greater than 200cells/mm³ [AOR=3.18, 95% CI: (1.65,6.13)]. Regarding substances, those who smoke cigarette currently had 5.69 times more likely to have poor sleep quality as compared with non-smokers [AOR=5.69, 95% CI: (2.04, 15.9)]. In addition, those taking chat or caffeinated drinks were 2.65 times higher to have poor sleep quality as compared to those who didn't use [AOR=2.65,95%CI: (1.06,6.64)]. Finally, The odds

of poor sleep quality was 3.55 times higher among participants with poor sleep hygiene as compared to those with good sleep hygiene [(AOR=3.55, 95%CI: (1.85, 6.78)] (Table 3).

Discussion

This study was a first attempt to ascertain the magnitude of sleep quality and its possible association with various variables among people living with HIV/AIDS in Ethiopia. The results from the current survey revealed that a remarkable proportion of people living with HIV/AIDS had experienced poor sleep quality. More than half of the people living with HIV/AIDS (55.6 %) experienced poor sleep quality. The magnitude of this study is in line with the studies conducted in Nigeria (59.3%), South Africa (55.8%), Indonesia (52%) and Mexico (58.6%) [17, 20-22].

Contrarily, the magnitude of poor sleep quality in this study was higher than study results in Brazil (46.7%) [18], Iran (47.5%) [23], China (43.1%) [15], Romania (42%) [24], Spain (40.9%) [25], France (47%) [16] and USA (26%) [26]. Furthermore the current finding is lower than the study conducted in USA 66.7% [27]. The possible reason for the observed variation might be the difference in socio-cultural conditions, in methodological designs and difference in hospital setting and caring modality.

In this study, poor sleep quality was significantly associated with female gender among people with HIV/AIDS. This could be due to the fact that females are more prone to stress due to the burden of excessive household responsibilities and changes in hormonal level. This result is supported by a study carried out in Nigeria [20].

The current study also found that depressed respondents were significantly associated with poor sleep quality as compared with non-depressed respondents. This may be due to depressed individuals have decreased serotonin neurotransmitters that results in diminished cognitive performance affects normal sleep pattern. This finding is in agreement with studies done in China [15], Europe and USA [16, 29-32]. Similarly different studies supported that poor emotion regulation modifies the point and prospective relationship between sleep quality and depression symptom [33-35].

CD₄ count less than 200cells/mm³ were a significant factor to develop poor sleep quality among people with HIV/AIDS. Probably, immune system is directly linked to the brain by a complex network of nerves, hormones, and neuropeptides. This network of specific physiological pathways is the primary determinant of neuropathology to have a direct impact on health including sleep. The finding of this study coincides with a study done in Nigeria [19, 20] and Mexico [29].

Similarly, shorter duration of HIV diagnosis were significantly associated with poor sleep quality as compared longer duration of diagnosis. The possible reasoning might be HIV-positive patients feel stigmatized and this may contribute to frequent psychological stress and emotional disturbances. The result is supported by research conducted in Nigeria [20].

Furthermore, use of substances within the current three months (i.e. tobacco, chat or caffeinated drinks) were significantly associated with poor sleep quality. The possible reasons behind these is biological effect of the substances on the brain results in disturbance of circadian rhythms which is associated with symptoms of poor sleep quality. This finding agrees with a study conducted in France[16], US geographic areas [36, 37] and studies conducted in Ethiopia at community level[38, 39].

Finally, in the present study there is a strong relationship between poor sleep hygiene and poor sleep quality. Even if, the finding of this study has no corresponding literature among the same population, it's supported by study undertaken in India to identify the relationship between sleep hygiene practices and sleep quality among cancer patients [40]. The current study implicates that health care providers must give attention to the sleep conditions of people with HIV/AIDs which have an enormous impact on treatment adherence and risk of psychological problem development.

Limitation of the study

The current study was limited to assess chronic insomnia having a duration of six months due to the nature of the assessment tool used and missed participants who might experience more sleep disturbances (i.e. participants with severe mental illness and acute physical illness) due to the exclusion criteria set. Hence, future studies focused on casual relations are of great significance.

Conclusion

In summary, this study revealed that more than half of people living with HIV/AIDS were found to have poor sleep quality. Being female, depression symptoms, lower CD4 count, shorter duration of HIV/AIDS diagnosis, current use of tobacco, chat or caffeinated drinks, and poor sleep hygiene were factors significantly associated with poor sleep quality. The findings suggest that routine screening of sleep disturbances among people living with HIV/AIDS is highly recommended. An intervention targeted the factors influencing sleep quality is also suggested.

Abbreviations

AOR: Adjusted odd ratio; CI: Confidence interval; PSQI: Pittsburgh Sleep Quality Index

Declarations

Ethics approval and consent to participation

Ethical approval was obtained from joint Ethical Review Committee of University of Gondar and Amanual Mental Specialized Hospital (No Committee reference number). Written consent was taken from the participants for their voluntary participation. Confidentiality was maintained throughout the study process.

Consent to publication: N/A

Availability of data and materials.

All data generated or analyzed during this study are included in this published article. The data sets of the current study is available from [Nebiyu Mengistu, email: nebiyumen@gmail.com ; Mobile: +251931333504, Dilla university, Dilla] upon reasonable request

Competing interests

We declare that there is no any financial or non-financial conflict of interest

Funding

The sources of funding for the current research is from Amanuel Mental specialized Hospital. The funding organization did not have any role in the design of the study, collection, analysis, interpretation of data and in writing the manuscript.

Authors' contributions

Nebiyu Mengistu (nebiyumen@gmail.com) organized the original investigation; coordinated the collection of data; analysis and writing report. Telake Azale (atelake07@gmail.com), Mahlet Fikreyesus (mahifyf@gmail.com) and Elsa Melaku (elsa_mlk@yahoo.com) equally contributed to the design of the study, performed the statistical analyses, Solomon Yimer (sopsycha@gmail.com) and Seid Shumye (Seidshumye22@gmail.com) contributed to the statistical analyses, drafted the manuscript and revision of the Manuscript.

Acknowledgements

We would like to thank Amanuel Mental Specialized Hospital, University of Gondar and Zewditu Memorial Hospital staffs for their support and assistance throughout the study process.

References

1. Barlow DH: **Clinical handbook of psychological disorders: A step-by-step treatment manual**: Guilford publications; 2014.
2. Association AP: **Diagnostic and statistical manual of mental disorders (DSM-5®)**: American Psychiatric Pub; 2013.
3. Johnston L, O'Malley P, Bachman J, Schulenberg J, Patrick M, Miech R: **HIV/AIDS: Risk & protective behaviors among adults ages 21 to 40 in the US, 2004-2016**. 2017.
4. Azagew AW, Woreta HK, Tilahun AD, Anlay DZ: **High prevalence of pain among adult HIV-infected patients at University of Gondar Hospital, Northwest Ethiopia**. *Journal of pain research* 2017, **10**:2461.

5. Taibi DM: **Sleep disturbances in persons living with HIV.** *Journal of the Association of Nurses in AIDS Care* 2013, **24**(1):S72-S85.
6. Reid S, Dwyer J: **Insomnia in HIV infection: a systematic review of prevalence, correlates, and management.** *Psychosomatic Medicine* 2005, **67**(2):260-269.
7. Rubinstein ML, Selwyn PA: **High prevalence of insomnia in an outpatient population with HIV infection.** *Journal of Acquired Immune Deficiency Syndromes and Human Retrovirology* 1998, **19**:260-265.
8. Junqueira P, Bellucci S, Rossini S, Reimão R: **Women living with HIV/AIDS: sleep impairment, anxiety and depression symptoms.** *Arquivos de neuro-psiquiatria* 2008, **66**(4):817-820.
9. McGrath L, Reid S: **Sleep and Quality of Life in HIV and AIDS.** In: *Sleep and Quality of Life in Clinical Medicine.* edn.: Springer; 2008: 505-514.
10. Qaseem A, Kansagara D, Forcica MA, Cooke M, Denberg TD: **Management of chronic insomnia disorder in adults: a clinical practice guideline from the American College of Physicians.** *Annals of internal medicine* 2016, **165**(2):125-133.
11. Gamaldo CE, Gamaldo A, Creighton J, Salas RE, Selnes OA, David PM, Mbeo G, Parker BS, Brown A, McArthur JC: **Sleep and cognition in an HIV+ cohort: a multi-method approach.** *Journal of acquired immune deficiency syndromes (1999)* 2013, **63**(5).
12. Saberi P, Neilands TB, Johnson MO: **Quality of sleep: associations with antiretroviral nonadherence.** *AIDS patient care and STDs* 2011, **25**(9):517-524.
13. Babson KA, Heinz AJ, Bonn-Miller MO: **HIV medication adherence and HIV symptom severity: the roles of sleep quality and memory.** *AIDS patient care and STDs* 2013, **27**(10):544-552.
14. Hudson A, Kirksey K, Holzemer W: **The influence of symptoms on quality of life among HIV-infected women.** *Western Journal of Nursing Research* 2004, **26**(1):9-23.
15. Huang X, Li H, Meyers K, Xia W, Meng Z, Li C, Bai J, He S, Cai W, Huang C: **Burden of sleep disturbances and associated risk factors: a cross-sectional survey among HIV-infected persons on antiretroviral therapy across China.** *Scientific Reports* 2017, **7**(1):3657.
16. Allavena C, Guimard T, Billaud E, De la Tullaye S, Reliquet V, Pineau S, Hùe H, Supiot C, Chennebault J-M, Michau C: **Prevalence and risk factors of sleep disturbance in a large HIV-infected adult population.** *AIDS and Behavior* 2016, **20**(2):339-344.
17. Ferreira LTK, Ceolim MF: **Sleep quality in HIV-positive outpatients.** *Revista da Escola de Enfermagem da USP* 2012.
18. Rodríguez-Estrada E, Iglesias-Chiesa MC, Fresán-Orellana A, Reyes-Terán G: **Factors associated with poor sleep quality among HIV-positive individuals in Mexico City.** *Salud mental* 2018, **41**(3):123-129.
19. Bisong E: **Predictors of Sleep Disorders among HIV Out-Patients in a Tertiary Hospital.** *Recent Advances in Biology and Medicine* 2017, **3**(2017):2747.
20. Oshinaike O, Akinbami A, Ojelabi O, Dada A, Dosunmu A, John Olabode S: **Quality of sleep in an HIV population on antiretroviral therapy at an urban tertiary centre in Lagos, Nigeria.** *Neurology research*

international 2014, **2014**.

21. Buysse DJ, Reynolds CF, Monk TH, Berman SR, Kupfer DJ: **The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research.** *Psychiatry res* 1989, **28(2)**:193-213.
22. Salahuddin M, Maru TT, Kumalo A, Pandi-Perumal SR, Bahammam AS, Manzar MD: **Validation of the Pittsburgh sleep quality index in community dwelling Ethiopian adults.** *Health and quality of life outcomes* 2017, **15(1)**:58.
23. Reda AA: **Reliability and validity of the Ethiopian version of the hospital anxiety and depression scale (HADS) in HIV infected patients.** *PLoS One* 2011, **6(1)**:e16049.
24. Reda AA: **Reliability and validity of the Ethiopian version of the hospital anxiety and depression scale (HADS) in HIV infected patients.** *PLoS One* 2011, **6(1)**.
25. Van Rie A, Sengupta S, Pungrassami P, Balthip Q, Choonuan S, Kasetjaroen Y, Strauss RP, Chongsuvivatwong V: **Measuring stigma associated with tuberculosis and HIV/AIDS in southern Thailand: exploratory and confirmatory factor analyses of two new scales.** *Tropical medicine & international health* 2008, **13(1)**:21-30.
26. Tsai AC, Weiser SD, Steward WT, Mukiibi NF, Kawuma A, Kembabazi A, Muzoora C, Hunt PW, Martin JN, Bangsberg DR: **Evidence for the reliability and validity of the internalized AIDS-related stigma scale in rural Uganda.** *AIDS and Behavior* 2013, **17(1)**:427-433.
27. Cho S, Kim G-S, Lee J-H: **Psychometric evaluation of the sleep hygiene index: a sample of patients with chronic pain.** *Health and quality of life outcomes* 2013, **11(1)**:213.
28. Dalgard O: **Social support-Consequences for individual and society.** *EUPHIX, Euphact Bilthoven: RIVM,* < <http://www.euphix.org> > *Euphact\Determinants of health\Environment\Social support* 2009, **16**.
29. Rodríguez-Estrada E, Iglesias-Chiesa M, Fresán-Orellana A, Reyes-Terán G: **Factors associated with poor sleep quality among HIV-positive individuals in Mexico City.** *Salud Mental* 2018, **41(3)**:123-129.
30. Redman K: **Sleep quality and immune changes in HIV positive people in the first six months of starting highly active antiretroviral therapy (HAART).**
31. Dabaghzadeh F, Khalili H, Ghaeli P, Alimadadi A: **Sleep quality and its correlates in HIV positive patients who are candidates for initiation of antiretroviral therapy.** *Iranian journal of psychiatry* 2013, **8(4)**:160.
32. Arbune M, Dumitru I-M, Cretu-Stuparu M: **Characteristics of sleep disorders in Romanian adults infected with human immunodeficiency virus.** *ARS Medica Tomitana* 2017, **23(3)**:126-131.
33. O'Leary K, Bylsma LM, Rottenberg J: **Why might poor sleep quality lead to depression? A role for emotion regulation.** *Cognition and Emotion* 2017, **31(8)**:1698-1706.
34. Pillai V, Kalmbach DA, Ciesla JA: **A meta-analysis of electroencephalographic sleep in depression: evidence for genetic biomarkers.** *Biological psychiatry* 2011, **70(10)**:912-919.
35. Zhai L, Zhang H, Zhang D: **Sleep duration and depression among adults: A meta-analysis of prospective studies.** *Depression and anxiety* 2015, **32(9)**:664-670.

36. Crum-Cianflone NF, Roediger MP, Moore DJ, Hale B, Weintrob A, Ganesan A, Eberly LE, Johnson E, Agan BK, Letendre S: **Prevalence and factors associated with sleep disturbances among early-treated HIV-infected persons.** *Clinical Infectious Diseases* 2012, **54**(10):1485-1494.
37. Ramamoorthy V, Campa A, Rubens M, Martinez SS, Fleetwood C, Stewart T, Liuzzi JP, George F, Khan H, Li Y: **Caffeine and Insomnia in People Living With HIV From the Miami Adult Studies on HIV (MASH) Cohort.** *Journal of the Association of Nurses in AIDS Care* 2017, **28**(6):897-906.
38. Manzar MD, Salahuddin M, Maru TT, Dadi TL, Abiche MG, Abateneh DD, Pandi-Perumal SR, Bahammam AS: **Sleep correlates of substance use in community-dwelling Ethiopian adults.** *Sleep and Breathing* 2017, **21**(4):1005-1011.
39. Berhanu H, Mossie A, Tadesse S, Geleta D: **Prevalence and Associated Factors of Sleep Quality among Adults in Jimma Town, Southwest Ethiopia: A Community-Based Cross-Sectional Study.** *Sleep disorders* 2018, **2018**.
40. Roselin V. Rhenius¹ JA: **Relationship of Sleep Hygiene Practices and Sleep Quality among Cancer Patients.** *IOSR Journal of Nursing and Health Science (IOSR-JNHS)* (Jan.- Feb .2018),, **Volume 7**(Issue 1 Ver. II.):2320–1940.

Tables

Table 1: Socio demographic characteristics of people living with HIV/AIDS on follow-up at Zewditu Memorial Hospital, Ethiopia, 2018. (n=396)

Variable	Category	Frequency	M(SD) or %
Age(mean and SD)			38.57±10.76
Sex	Male	199	50.3
	Female	197	49.7
Religion	Orthodox	265	66.9
	Muslim	24	6.1
	Protestant	84	21.2
	Others*	23	5.8
Marital status	Single	166	41.9
	Married	208	52.5
	Others**	22	5.6
Educational status	Can't write and read	20	5.1
	Primary	88	22.2
	Secondary	176	44.4
	Tertiary and above	112	28.3
Occupational status	Employed	154	38.8
	Private business	150	37.9
	Students	66	16.7
	Others***	26	6.6
Living arrangement	with family	244	61.6
	alone	133	33.6
	Others****	19	4.8

N.B. * Catholic, Wakifeta ** Separated, Divorced, widowed *** Daily laborer, House wife, Farmer **** with relatives,
Adopted

Table 2. HIV related clinical characteristics of people living with HIV/AIDS on follow up at Zewditu Memorial Hospital, Addis Ababa, Ethiopia, 2018.(n=396)

Variable	Category	Frequency(396)	Percentage (%)
WHO clinical stage	stage I	294	74.2
	stage II	59	14.9
	stage III	35	8.8
	stage IV	8	2.0
CD ₄ count	≤ 200 cells/mm ³	177	44.7
	> 200 cells/mm ³	219	55.3
Duration of HIV/AIDS diagnosis	≤ 10 years	124	31.3
	> 10 years	272	68.7
ART type	first line regimen	296	74.7
	Second line regimen	100	25.3
Co-morbid chronic medical illness	No	255	64.4
	Yes	141	35.6
Types of chronic medical conditions	Hypertension	47	33.4
	Diabetes mellitus	41	29.0
	Tuberculosis	35	24.8
	Others	18	12.7

Table 3. Factors associated with poor sleep quality among people living with HIV/AIDS at Zewditu Memorial Hospital (Bivariate and multivariate analysis) (n=396), Addis Ababa, Ethiopia, 2018.

Variables	Categories	sleep quality		COR(95%CI)	AOR(95%CI)
		Poor	Good		
Age				0.98(0.96,1.00)	0.98(0.95,1.01)
Sex	Male	75	124	1.00	1.00
	Female	145	52	4.61(3.00,7.06)	3.40(1.80, 6.41)**
Depression	No	72	120	1.00	1.00
	Yes	148	56	4.40(2.88,6.73)	3.52(1.95, 6.32)**
Anxiety	No	153	134	1.00	1.00
	Yes	67	42	1.39(0.89,2.19)	0.92(0.48,1.76)
Perceived stigma	No	78	124	1.00	1.00
	Yes	142	52	4.34(2.83,6.64)	0.82(0.39,1.76)
Social support	Poor	128	36	7.11(3.51,14.3)	2.48(0.91,6.75)
	Moderate	76	108	1.40(0.72,2.74)	1.20(0.50,2.88)
	Strong	16	32	1.00	1.00
CD ₄ count	>200 cells/mm ³	79	140	1.00	1.00
	≤200 cells/mm ³	141	36	6.94(4.38,10.9)	3.18(1.65,6.13) **
Duration HIV diagnosis	>10 years	115	157	1.00	1.00
	≤10 years	105	19	7.54(4.37,13.0)	3.43(1.61,7.29) **
Co-morbid medical illness	No	111	144	1.00	1.00
	Yes	109	32	4.41(2.77,7.03)	0.67(0.28,1.62)
Current use of tobacco	No	123	164	1.00	1.00
	Yes	97	12	10.7(5.66,20.5)	5.69(2.04,15.9)**
Current use of alcohol	No	84	129	1.00	1.00
	Yes	136	47	4.44(2.88,6.83)	0.94(0.43,2.05)
Current use of chat or caffeinated drinks	No	140	166	1.00	1.00
	Yes	80	10	9.48(4.73,19.0)	2.65(1.06,6.64) *
Sleep hygiene	Good	84	109	1.00	1.00
	Poor	136	67	2.64(1.75,3.96)	3.55(1.85,6.78)**

Figures




Figure 1

Current use of substances among people living with HIV/AIDS on follow up at Zewditu Memorial, Addis Ababa, Ethiopia, 2018(n=396).