Laccase immobilization via cross-linked aggregate preparation: Characterization, thermodynamic/kinetic properties and application in removal of bisphenol A from solutions

Adedeji Ademakinwa (✉ adedeji.ademakinwa@elizadeuniversity.edu.ng)

EU: Elizade University <https://orcid.org/0000-0001-7736-0459>

Research Article

Keywords: CLEA, laccase, RSM, biodegradation, Bisphenol A

Posted Date: March 5th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-267600/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

**Laccase immobilization via cross-linked aggregate preparation: Characterization,
thermodynamic/kinetic properties and application in removal of bisphenol A from
solutions**

Adedeji Nelson Ademakinwa

Department of Physical and Chemical Sciences, Elizade University, Ilara-Mokin, Ondo-State.

E-mail: adedeji.ademakinwa@elizadeuniversity.edu.ng

24 **Abstract**

25 Fungal laccase from *Aureobasidium pullulans* was immobilized via cross-linked enzyme
26 aggregate (CLEA) preparation under statistically optimized conditions. The stability of the
27 CLEA to heat inactivation was studied via investigation of its thermodynamic and kinetic
28 parameters. The immobilized enzyme was then deployed in the biodegradation of a bisphenol-A
29 (BPA). The optimum conditions for CLEA preparation resulting in the highest immobilization
30 yield were ammonium sulphate (60% v/v), glutaraldehyde (30 mM), pH (4.5), time (6 h) and
31 temperature (55°C). The CLEA retained about 51% of its activity after eight catalytic cycles. The
32 optimum pH and temperature of the laccase CLEA were 5.5 and 60°C respectively. The SEM
33 indicated that the laccase CLEA was type II (unstructured). The data obtained from the heat
34 inactivation kinetics and thermodynamic characterization indicated that the CLEA was stable
35 to heat denaturation than the free enzyme. The kinetic parameters obtained for the CLEA with
36 ABTS as substrate were 101.3 μ M, 2.94 μ mol $\cdot$ s $^{-1}$ mg $^{-1}$ and 0.03 dm 3 s $^{-1}$ mg $^{-1}$ for the K_m , K_{cat}
37 and K_{cat}/K_m respectively. The optimum conditions for BPA biodegradation using the CLEA
38 were temperature (55°C), time (2 h), CLEA (1.0 mg) and BPA concentration (40 mg/L). After the
39 7th cycle, laccase CLEA retained about 63 $\pm$ 2.3% biodegradation efficiency. A heat-resistant
40 laccase CLEA was able to remove BPA from solutions under statistically optimized conditions.
41 The laccase CLEA has properties for other futuristic applications.

42 **Keywords:** CLEA, laccase, RSM, biodegradation, Bisphenol A

43

44

45

46

47 **Introduction**

48 Laccases (multicopper oxidases) are useful enzymes mostly in the industry as they play roles
49 such as in xenobiotic biodegradation, pulp and paper bleaching etc. They can perform these
50 functions due to their low substrate specificity and potentials in the oxidation of a large number
51 of substrates [1]. Substrate oxidation is often accompanied by the release of oxygen to water.
52 Moreso, laccases are often regarded as being environmentally friendly, hence, its application in
53 numerous field is quite understandable [2]. Laccases have been reported to be involved in the
54 removal of pollutants such as endocrine-disrupting chemicals and dyestuff (EDC) [3, 4]. These
55 EDC's are present in wastewater bodies and are known to cause deleterious effects on human
56 and animal health [5].

57 Major hindrances to the application of laccases (in an aqueous form) industrially especially in
58 xenobiotics biodegradation is the low yields and very high costs, loss of enzyme activity and poor
59 reusability [6]. These problems are often solved via immobilizing the enzymes because the use
60 of immobilized enzymes provides options for cost reduction, enzyme stability and enhanced
61 reusability [7]. Several techniques of laccase immobilization for BPA biodegradation exists with
62 the most prominent one being immobilization on a solid support [6]. This technique comes with
63 usual problems such as low enzyme: solid support. This brings about the technique of deploying
64 cross-linked enzyme aggregates (CLEA) [8, 9]. This technique is devoid of solid support as it
65 involves precipitating the enzyme and then cross-linking it with a cross-linker such as
66 glutaraldehyde. The production of laccase CLEA's has been reported in the literature. Laccase
67 CLEA's are known to be thermostable and resistant to denaturants.

68 The preparation of laccase CLEA from the fungus, *Aureobasidium pullulans* with a focus on its
69 application in the biodegradation of endocrine-disrupting compounds was the main objective of

70 this study. There are reports of production of extracellular laccase by this fungus [10, 11]. The
71 utilization of laccase and mycelia from this fungus in the biodegradation of textile dyes has been
72 reported [12]. This study is the first-ever for the immobilization of laccase via CLEA from this
73 fungus. Despite the numerous advantages posed by using CLEA, very few reports exist in
74 literature where laccase CLEA was produced for applications in BPA biodegradation. Only the
75 report of Cabana et al. [3] established the application of CLEA in the removal of BPA using a
76 fluidized bed reactor. Most recent studies on CLEA and its application focused on dye
77 biodegradation [13]. Hence, this study provides needed information on the application of laccase
78 CLEA in the removal of EDC's.

79 In this study, the use of response surface methodology via central composite rotatable design
80 (CCRD) was utilized in both the preparation of the CLEA and its application in the
81 biodegradation of BPA. During optimization of most bioprocesses, CCRD provides unique
82 advantages such as easy monitoring of the several interactive variable effects [14].

83 **Methods**

84 **Fungal growth and enzyme isolation**

85 *Aureobasidium pullulans* was maintained on malt extract agar and sub-cultured every 4 days
86 at 23±1°C. The fungus was inoculated into a laccase-producing medium containing (in g/L):
87 CuSO₄·5H₂O (4) Tyrosine (3.0), glucose (10), ZnSO₄ (0.1), FeSO₄ (0.05), MnSO₄ (0.5), MgSO₄
88 (0.5) yeast extract (5). Growth was monitored for 5 days and the mycelia was harvested after
89 centrifugation at 6000 x g for 10 min. The mycelia (10 g) was homogenized using acid-washed
90 sand (in a sterile and pre-chilled mortar/pestle) in 1:1 ratio [15] Thereafter, three-time volume
91 of 10 mM citrate buffer pH 4.0 buffer (chilled at 4°C) was added to the homogenized mycelia and

it was stirred. All operations were carried out on ice at 4°C. The homogenate was centrifuged at 6000xg for 30 min at 4°C and the resulting supernatant served as the source of enzyme.

Laccase assay and protein concentration determination

Laccases was assayed using 13 mM ABTS as substrate [10]. ABTS oxidation was monitored at 460 nm. One unit of laccase activity was defined as the amount of enzyme that oxidizes one micromole of substrate to form the product ABTS⁺ in one min. Protein concentration was determined using bovine serum albumin as standard via methods described by Bradford [16].

Preparation of CLEA

Selection of precipitant

The choice of precipitant was determined over a concentration range of 0-70% v/v of each chilled precipitant (acetone, ethanol, ammonium sulphate, PEG-2000, PEG-5000). The precipitants were added to the enzyme solution in a 10 ml centrifuge tube and incubated for 8 h at 4°C. The enzymatic activity of the aggregates were determined with ABTS as substrate.

CLEA synthesis via optimization

In preliminary studies, the CLEA was synthesized by using the methods described by Cabana et al.³ and Ademakinwa et al.[17]. Briefly, 2 ml of 800 mU/mL of laccase was added to 4 ml of ammonium sulphate (50% v/v). After a 16 h incubation, the reaction mixture was centrifuged at 3500 x g for 10 min at 4°C. The pellet was collected and re-dissolved in minute amount of buffer (10 mM citrate-phosphate pH 4.0). Thereafter, 2 ml of glutaraldehyde (25 mM) was added to the re-dissolved pellet and it was allowed to cross-link for 2 h at 4°C. The resulting pellet was collected by centrifugation at 3500 x g for 10 min at 4°C. The pellet was washed in 10 mM citrate-phosphate buffer, pH 4.0 and stored at 4C. The immobilization yield was determined using the equation below

$$\text{Immobilization yield(\%)} = \frac{\text{Specific activity of immobilized enzyme}}{\text{Specific activity of free enzyme}} \quad [1]$$

To optimize CLEA preparation, response surface methodology via central composite design was adopted. Five independent variables (at five unique levels) were considered such as precipitant concentration, precipitation time, glutaraldehyde concentration, cross-linking time and temperature. The experimental design was generated using Design-Expert version 6.0.

The high and low levels of the process variables adopted for the RSM optimization were determined from results of preliminary experiments (not shown). The precipitation concentration 10 and 50% (v/v), precipitation time 4 and 16 h, glutaraldehyde concentration 2 and 30% v/v; cross-linking time 3 and 12 h and temperature 30 and 70°C. The CCRD is shown in Supplementary S1 and it illustrates the five variables studied at the five different levels, the observed and residual activity of the CLEA. The five different levels for the individual experiment were coded thus: $-\alpha$, -1 , 0 , 1 , $+\alpha$.

$$\alpha = 2^{N/4} \text{ where } N \text{ is the number of variable} \quad [2]$$

Data analysis

Equation 1 illustrates the analysis of the experimental data for each independent variable by fitting of the second-order polynomial equation to the experimental data of each dependent variable.

$$X_i = \mu_0 + \sum_{i=1}^n \mu_i x_i + \sum_{i=1}^{n-1} \sum_{j=i+1}^n \mu_{ij} x_j + \sum_{i=1}^n \mu_{ii} x_i^2 \quad [3]$$

X_i is the response (%Immobilization yield); x_i represents the independent variables considered in this study, i.e. x_1 : precipitant concentration [% v/v], x_2 : precipitation time [h], x_3 : glutaraldehyde concentration [% v/v]; x_4 : cross-linking time [h] and x_5 : temperature [C]. μ_0 represent the value of the fitted response at the central points. The liner, interactive and quadratic terms are designated by μ_i , μ_{ij} and μ_{ii} respectively.

138 **CLEA characterization**

139 **Effect of pH and temperature**

140 The effect of pH on the free enzyme and CLEA was determined using several buffers: 10 mM
141 citrate-phosphate buffer pH 3.0-5.0, phosphate buffer 6.0-8.0, Tris-HCl buffer 9.0-11.0. The
142 effect of temperature on CLEA was determined at temperatures ranging from 20 to 90°C with
143 ABTS as substrate.

144 **pH and heat stability studies**

145 CLEA was incubated at optimum pH for 8 h and the residual activity determined every 1 h with
146 ABTS as the substrate. For the heat stability studies, the CLEA was incubated at 40 to 80°C for
147 6 h and the residual activity was estimated.

148 **Heat inactivation kinetics of the free enzyme and CLEA**

149 The irreversible heat inactivation process is often kinetically described by a first-order reaction,
150 see equation (4)

151
$$v = k_d E \quad [2], \quad [4]$$

152 where v is defined as the rate of inactivation of the enzyme and E is the concentration of the
153 active form of the enzyme. The k_d is obtained from the slopes of the plot of the activity coefficient,
154 ρ , against time at different temperature range (40-80°C) i.e. $\ln \rho$ vs t . From the Arrhenius plot,
155 $\ln k_d$ vs $1/T$, the energy of thermal denaturation, E_d , was estimated from the slopes of the
156 straight lines. Other thermodynamic parameters were estimated from the equations stated
157 below as described by Melikoglu et al. [18].

158
$$\Delta H_d^* = E_d - RT \quad [5]$$

159
$$\Delta G_d^* = -RT \ln\left(\frac{k_d h}{k_b T}\right) \quad [6]$$

160
$$\Delta S_d^* = \frac{\Delta H^* - \Delta G^*}{T} \quad [7]$$

161 ΔH_d^* : *Change in enthalpy*
 162 ΔG_d^* : *Gibb's free energy*
 163 ΔS_d^* : *Change in entropy*
 164 R: universal gas constant (= 8.314 J/mol/K)
 165 h: Planck's constant (= 6.63 x 10⁻³⁴)
 166 k_b : Boltzmann constant (= 1.38 x 10⁻²³)
 167 The half-life of the enzyme, $t_{1/2}$, was determined according to the equation described below.
 168
$$t_{1/2} = 0.693/k_d$$
 [8]
 169 **Determination of kinetic parameters**
 170 The kinetic parameters for the immobilized laccase were determined using ABTS (25-1500 μ M)
 171 as substrate. The kinetic parameters (K_m , K_{cat} and V_{max}) were determined from the Michealis-
 172 Menten plot (v against S) using the GraphPad Prism software version 7.0.
 173 **SEM and FTIR studies**
 174 The surface structure of the CLEA was determined by FTIR spectroscopy in a transmittance
 175 mode while the structural features of the CLEA was investigated using SEM. For the SEM
 176 studies, the dried CLEA was coated with gold, viewed under 2500 X magnification under 10 kV
 177 using scanning electron microscope model TESCAN. For the FTIR studies, the dried CLEA was
 178 ground in a mortar containing KBr (in 1:1) and screwed to pellets [17]. Scans were recorded
 179 from 4000-400 cm in transmittance mode using a Shimadzu FTIR-8400S spectrophotometer.
 180 **Biodegradation efficiency of BPA by CLEA**
 181 The potential of the CLEA to biodegrade BPA was investigated under response surface
 182 methodology via CCRD (described earlier) only that in this case, the response is %
 183 biodegradation efficiency and the independent variables are concentration of the BPA (mg/L),

time of incubation (h), temperature (C) and CLEA amount (mg). Supplementary S2 provides the information on the CCRD experimental design and it illustrates the four variables studied at five different levels. As seen in equation 3,

X_i is the response (% Biodegradation efficiency); x_i represents the independent variables considered in this study, i.e. x1: BPA concentration, [mg/L], x2: time of incubation [h], x3: CLEA amount [mg]; x4: temperature [C].

μ_0 represent the value of the fitted response at the central points. The liner, interactive and quadratic terms are designated by μ_i , μ_{ij} and μ_{ii} respectively.

$$\% \text{ Biodegradation Efficiency} = \left(1 - \frac{[BPA]_t}{[BPA]_0}\right) \times 100 \quad [9]$$

Where $[BPA]_t$ and $[BPA]_0$ represent the BPA concentration at a specified time after treatment with CLEA and the initial BPA concentration respectively.

2.4.1 Model validation

To validate the model, the optimum conditions obtained in the experimental runs were repeated at least three times and the biodegradation efficiency determined.

2.4.2. Reusability of CLEA in BPA biodegradation

The CLEA was used in the biodegradation of BPA at the optimum conditions for several catalytic cycles. The experiment was repeated thrice and the biodegradation efficiency determined.

Result and discussion

Selection of best precipitating agent for laccase prior to cross-linking

It was observed that acetone had the least recovery at all concentrations studied (0-75% v/v). The reduced recovery might be due to the acetone inducing certain changes in structural properties of the laccase, which resulted in a loss of enzymatic activity. Ammonium sulphate had the highest recovery of about 80% at 65% (v/v). The choice of precipitant for laccase is

necessary because of the recovery obtained after precipitation and the change in structural conformation/enzymatic features afterwards [8]. The conclusion that ammonium sulphate was the best precipitating agent prior to crosslinking was also reported by Yang et al. [19].

Statistical optimization of the preparation of laccase CLEA

From Supplementary S1, it was observed that the optimum conditions resulting in the highest yield (73%) for the laccase CLEA were ammonium sulphate (60% v/v), glutaraldehyde (30 mM), pH (4.5), time (6 h) and temperature (55°C). The second order RSM obtained via ANOVA of the immobilization yield is provided in Table 1. The model F value of 36.31 indicated that the model was significant which implied that there was a 0.01% chance that this model could occur due to noise. The standard deviation, coefficient of variation, adequate precision and coefficient of determination values were 1.27, 1.92, 23.80 and 0.96 respectively. The p-value obtained was less than 0.0001 and it indicated that the model was significant. Other significant parameters included from the viewpoint of the p-value were X_1 , X_2, X_3, X_4, X_5 , $X_1X_2, X_1X_3, X_1X_4, X_1X_5, X_2X_3, X_2X_5$ and X_4X_5 . The implications of the significance of these variables is that they provide information that these variables affect the immobilization yield. Other parameters were not significant ($p > 0.05$) indicating that those variables had no profound effect on the immobilization yield of the laccase CLEA.

A second-order polynomial equation was obtained that described the link between the variables studied and the response (immobilization yield). The equation is described as:

$$\begin{aligned} \text{Immobilization Yield (\%)} = & 70.91 - 3.96X_1 + 0.57X_2 + 1.29X_3 - 0.83X_4 + 0.46X_5 - 1.57X_1^2 - \\ & 0.37X_2^2 - 1.57X_3^2 - 1.39X_4^2 - 0.69X_5^2 + 0.46X_1X_2 - 0.27X_1X_3 - 0.85X_1X_4 + 0.38X_1X_5 + 0.26X_2X_3 - \\ & 0.52X_2X_4 - 0.060X_2X_5 - 0.31X_3X_4 + 0.39X_3X_5 + 0.24X_4X_5 \end{aligned}$$

The variables were coded such that, x1: precipitant concentration [% v/v], x2: precipitation time [h], x3: glutaraldehyde concentration [% v/v]; x4: cross-linking time [h] and x5: temperature [°C].

To critically understand the interactions between the variables considered in this study and how it influences the response (immobilization yield), response surface plots (Supplementary S3)

were obtained for two variables while the others are kept at a constant (central levels). For example, the effects of pH and time (while precipitant and glutaraldehyde concentration and temperature were kept constant at their central levels) on the CLEA immobilization yield is provided in Fig. 1. It was observed that the immobilization yield gradually increased as both the time and pH values were increased until the optimum was reached. Further increased above the optimum values led to decreases in the response.

Model Validation

The immobilization yield obtained after repeating the experiments under the optimum conditions predicted by the model was $75.3 \pm 2.8\%$ which was very close to that predicted by the model. The immobilization yield obtained for laccase CLEA was higher than the values obtained for laccase CLEA from *Cerrena sp.* [19] and *Coriolopsis polyzona* [3] which had values of 68.1% and 60.2% respectively.

Reusability of CLEA

Supplementary S4 shows the potential of the laccase CLEA to catalyze ABTS oxidation over several cycles of catalysis. After eight reaction cycles, the resulting immobilization yield was 51%. A gradual decrease in the rate of catalysis was observed after the fifth cycle and this might be due to substrate mass-transfer limitations that result in a decreased rate of ABTS oxidation by the CLEA.

Laccase CLEA characterization

Effect of pH and temperature

The optimum pH obtained for the free laccase and CLEA were 4.5 and 5.5 respectively (Fig.2). The optimum pH of most fungal laccases falls within the acidic region [20]. The pH of the CLEA increased by a unit when compared with the free enzyme. This increase could be due to the

changes in the microenvironment of the immobilized enzyme. This observation was in agreement with the reports of several authors [21, 22]. The optimum temperature for the free laccase and CLEA were 40 and 60°C respectively (Fig. 3). Most immobilized laccases have an optimum temperature that falls within 40 and 60°C [20].

pH and temperature stability

The pH stability is shown in Supplementary S5. It was noted that after 4 h incubation the free laccase retained about 100% of its initial activity at pH 4.5. This decreased after 5 h incubation with a loss of about 5%. Upon the increase of the incubation time to 8 h, the free laccase retained about 85% of its residual activity. The results obtained in this study was in agreement with what Lassaoune et al. [21] obtained for free laccase where the enzyme retained about 85.2% after incubation. Meanwhile, the CLEA retained more than 95% of its initial enzymatic activity after 8 h incubation at pH 5.5

The thermostability of the free laccase is shown in Supplementary S6. It was observed that after 8 h incubation at 40°C, the free laccase retained about 80% of its initial activity. It was observed that at very high temperatures (60-80°C) more than 80% of the initial activity was lost. Supplementary S8 illustrates the thermostability of the laccase CLEA. It was noted that after 8 h incubation, at 40-50°C, the laccase CLEA retained more than 90% of its initial activity. Increase of the temperature from 50-60°C still showed that the immobilized enzyme retained more than 75% of its initial activity after 8 h incubation. A drastic drop to about 55% of its initial activity was observed after the incubating temperature was increased to 80°C after 8 h incubation. It was concluded that the laccase CLEA was more thermostable than the free enzyme. The thermostability of CLEA or other immobilized enzymes over their free enzyme has been reported in recent literature [17, 21, 23-25]

Heat inactivation studies

Table 2 shows the summary of the kinetic and thermodynamic parameters for both the free and immobilized laccase. Both enzyme form's enzymatic activity decayed in a typical first-order heat inactivation pattern. This inactivation pattern is similar to other enzymes produced by this fungus [15, 23-25].

The values of the first-order thermal denaturation constant, k_d , for the CLEA (0.0113-0.0791 h⁻¹) and the free laccase (0.0317-0.1995 h⁻¹) is indicated in Table 2. These k_d values were initially obtained from the plot of $\ln \left(\frac{E}{E_0} \right)$ vs time (see supplementary S7 and S9) and they increased as the temperature was gradually raised. At any temperature range considered, the k_d values obtained for the CLEA lower than the free enzyme. This also implies that the immobilized enzyme displayed a greater degree of thermostability at each temperature increase than the free enzyme. Several authors have reported that immobilized forms of the enzyme tend to be more thermally stable than the free enzyme [23, 24].

From Table 2, the half-life ($t_{1/2}$) values for both free laccase and laccase CLEA decreased progressively while the heat inactivation constant increased as the temperature was gradually increased from 40 to 80°C. A higher $t_{1/2}$ value is required if the enzyme is to applied industrially [26]. The laccase CLEA prepared in this study had very high $t_{1/2}$ even at higher temperatures and hence could be useful in numerous industrial applications.

The values energy of thermal inactivation, E_d , for both enzyme forms were obtained from the plot indicated in Fig. 4. E_d values obtained for both free and immobilized laccase were 39.9 ($R^2 = 0.86$) and 42.4 ($R^2 = 0.96$) kJ/mol/K for free laccase and CLEA respectively. It was observed that the E_d values for the CLEA was higher than the free enzyme. This further attests to the fact

301 that the free enzyme was relatively less stable to thermal inactivation compared to the laccase
 302 CLEA.

303 The E_d is directly related to the enthalpy of thermal inactivation, ΔH_d . The ΔH_d values provide
 304 information on the overall energies necessary to cause heat inactivation of proteins. Therefore,
 305 it can be suggested that a high value of ΔH_d and E_d indicates increased resistance to heat [27].

306 The laccase CLEA had higher values of ΔH_d compared to the free enzyme (Table 2).

307 The degree of heat-mediated inactivation of enzymes is quite dependent on the entropy of
 308 thermal inactivation, ΔS_d . ΔS_d values indicate the energies required for the transition from a
 309 native (N) to a denatured state (U) ($N \leftrightarrow I \leftrightarrow U$). Every thermal inactivation process leads to
 310 changes in the enzyme conformation leading to increased orderliness and randomness (or
 311 positive values for ΔS_d) [28]. In this study, the negative values obtained for ΔS_d for both free
 312 laccase (-162.3 to -155.0 J/mol/K) and CLEA (-155.6 to -155.5 J/mol/K) is suggestive of even more
 313 increased resistance to disorderliness. This means that both enzyme forms were resistant to
 314 heat inactivation. A comparative look into the ΔS_d values obtained for both free and laccase
 315 CLEA, it can be concluded that the process of CLEA preparation (precipitating with ammonium
 316 sulphate and crosslinking with glutaraldehyde) allowed for more orderliness which
 317 concomitantly increased its thermal stability. This conclusion was also reached for protease [28],
 318 rhodanese [24], cellulase [23].

319 The values for Gibb's free energy (ΔG_d) obtained for the free laccase (88.1 to 91.7 kJ/mol) and
 320 laccase CLEA (88.5 to 94.4 kJ/mol) is provided in Table 3. The ΔG_d values obtained for both the
 321 free and laccase CLEA were quite very close. This is indicative of thermostability.

When all thermodynamic and kinetic parameters were taken into consideration, it was confidently asserted that the CLEA was very stable to thermal denaturation even at very high temperatures (50-60°C) compared to the free laccase.

3.4.4 Kinetic Parameters for laccase CLEA

The K_M for free laccase (106.2 μM) and CLEA (101.3 μM) was obtained from the Michaelis-Menten plot (v against S) with ABTS as substrate. These values were about the same. This observation was quite similar to the report of Cabana et al. [3] for free laccase and laccase CLEA. The K_{cat} obtained in this study for the free laccase and the laccase CLEA were 1.94 and 2.94 ($\mu\text{mol/s/mg}$) respectively. The rate of catalysis by the CLEA was approximately twice that of the free enzyme as observed from the K_{cat} values. The values obtained in this study is quite lower than what was obtained for laccase CLEA from *C. polyzona* [3] (which was six times higher than the free enzyme). Another kinetic parameter of great importance is the catalytic efficiency, K_{cat}/K_M . From the K_{cat}/K_M values obtained for the free ($0.04 \text{ dm}^3\text{s}^{-1}\text{mg}^{-1}$) and immobilized laccase ($0.06 \text{ dm}^3\text{s}^{-1}\text{mg}^{-1}$), it can be concluded that the catalytic efficiency in the CLEA is approximately twice that of the free enzyme. This suggests the laccase CLEA had a better overall catalytic potential than the free enzyme.

3.4.5 SEM and FTIR studies on CLEA

The FTIR spectra of the laccase CLEA is shown in supplementary S10. Several characteristic peaks were noted such as the peaks at 1647 cm^{-1} which is indicative of the amide I (carbonyl stretching $\text{C}=\text{O}$). Another peak that represents the bending and stretching of both $\text{C}-\text{N}$ and $\text{N}-\text{H}$ [17]. The SEM of the CLEA is indicated in Fig. 5. Typically, CLEA structure from SEM falls into two categories, the ball type (type 1) and unstructured (type 2) [29]. From the SEM, it was evident that the laccase CLEA falls into type 2.

3.5 RSM-modelled biodegradation of BPA using laccase CLEA

The optimum conditions predicted by the mathematical model were BPA (40 mg/L), time (2 h), temperature (55°C) and CLEA amount (1.0 mg) (Supplementary S2). The optimum conditions resulted in a biodegradation efficiency of 99.9%. Asagdol et al. [30] reported the removal of BPA with an efficiency of 88.3% by the free laccase which was lower than that obtained for the immobilized enzyme used in this study or reported elsewhere [21]. The results obtained in this study was quite similar to the findings of Lassaoune et al. [21] where it was reported that laccase immobilized by cross-linking prior to entrapment in alginate had a biodegradation efficiency of 99%.

From Table 3, the R², standard deviation and, CV were 0.97, 5.91 and 11.39 respectively. The model-F value obtained in this study was 34.56 and the model was deemed significant (p<0.0001). A model-F value that is as large as the one obtained in this study has a 0.01% chance it occurred due to noise. Significant model terms (based on the values of “Prob>F” less than 0.05) where A, C, D, A², B², D² and AD. The final equation in terms of the coded factor was:

$$\begin{aligned} \text{\%Biodegradation Efficiency} = & +65.68 - 21.92A - 1.77B + 8.50C - 5.87D - 3.63A^2 - 4.01B^2 - \\ & 1.25C^2 - 8.41D^2 - 1.31AB - 2.47AC + 3.74AD + 1.48BC + 0.51BD - 1.37CD \end{aligned}$$

The response surface plots are shown in supplementary S11. S11 (a) examines the effects of time and temperature on the BE. It was observed that increasing the temperature increased the BE by the CLEA. Meanwhile, when the optimum time was reached for maximum BE by the CLEA, a gradual decrease in the BE was noted. S11 (b) illustrates the effects of time and BPA concentration on the BE of the CLEA. It was noted that a steady decline in the BE was observed as the BPA concentration was increased above the optimum value. It can be deduced that

perhaps, the CLEA enzymatic activity had been inhibited either by the substrate (BPA) or the products formed.

3.5.1 Model validation of BPA biodegradation by laccase CLEA

The optimum conditions predicted by the model were validated under the following experimental conditions: temperature (55°C), time (2 h), CLEA (1.0 mg) and BPA concentration (40 mg/L). The biodegradation efficiency obtained after the experiment was conducted thrice was 98.4 ± 1.2 .

3.6. Reusability of CLEA in BPA biodegradation

Laccase CLEA under the optimum conditions removed BPA from solutions while retaining about $63 \pm 2.3\%$ biodegradation efficiency after the 7th cycle (Supplementary S12). The gradual loss of enzymatic activity by the CLEA might be associated with mass-transfer limitations

4.0 Conclusions

Response surface methodology provided a unique mathematical/statistical model for both the preparation of laccase CLEA and its application in the removal of BPA. From this study, the optimum conditions for CLEA preparation resulting in the highest immobilization yield were ammonium sulphate (60% v/v), glutaraldehyde (30 mM), pH (4.5), time (6 h) and temperature (55°C) while the optimum conditions for BPA biodegradation using the CLEA were temperature (55°C), time (2 h), CLEA (1.0 mg) and BPA concentration (40 mg/L). The thermodynamic and kinetic properties of the CLEA indicated the biocatalyst was thermally stable and resistant to heat inactivation. This property is necessary for biocatalysts of industrial importance. The laccase CLEA retained about $63 \pm 2.3\%$ of BPA biodegradation efficiency after seven rounds of catalysis. The CLEA could be deployed in possible large-scale BPA removal and other futuristic biotechnological applications.

390 Ethical Approval: Ethics approval not required for this study.

391

392 -Consent to Participate: Not applicable

393

394 -Consent to Publish: Not applicable

395

396 -Authors Contributions: Ademakinwa Adedeji Nelson carried out the Investigation, Writing,

397 Conceptualization, Methodology, Software, Data curation, Writing- Original draft preparation.

398

399 -Funding: The author has no relevant financial or non-financial interests to disclose.

400

401 -Competing Interests: The author declares that there are no conflicts of interests

402

403 -Availability of data and materials: Yes

References

- [1] Giardina, P., Faraco, V., Pezzella, C., Piscitelli, A., Vanhulle, S. and Sannia, G. (2009). Laccases: a never-ending story. *Cellular and Molecular Life Sciences*, 67(3), 369–385. doi:10.1007/s00018-009-0169-1
- [2] Viswanath, B., Rajesh, B., Janardhan, A., Kumar, A.P. and Narasimha, G. (2014). Fungal Laccases and Their Applications in Bioremediation. *Enzyme Research*, 2014, 1–21. doi:10.1155/2014/163242
- [3] Cabana, H., Jones, J.P. and Agathos, S.N. (2007). Preparation and characterization of cross-linked laccase aggregates and their application to the elimination of endocrine disrupting chemicals. *Journal of Biotechnology*, 132(1), 23–31. doi:10.1016/j.jbiotec.2007.07.948
- [4] Bosso, L. and Cristinzio, G. A. (2014). A comprehensive overview of bacteria and fungi used for pentachlorophenol biodegradation. *Reviews in Environmental Science and Bio/Technology*, 13(4), 387–427. doi:10.1007/s11157-014-9342-6
- [5] Dekant, W. and Völkel, W. (2008). Human exposure to bisphenol A by biomonitoring: Methods, results and assessment of environmental exposures. *Toxicology and Applied Pharmacology*, 228(1), 114–134. doi:10.1016/j.taap.2007.12.008
- [6] Bilal, M., Iqbal, H.M.N. and Barceló, D. (2019). Mitigation of bisphenol A using an array of laccase-based robust bio-catalytic cues – A review. *Science of The Total Environment*. doi:10.1016/j.scitotenv.2019.06.403
- [7] Durán, N., Rosa, M.A., D’Annibale, A. and Gianfreda, L. (2002). Applications of laccases and tyrosinases (phenoloxidases) immobilized on different supports: a review. *Enzyme and Microbial Technology*, 31(7), 907–931. doi:10.1016/s0141-0229(02)00214-4
- [8] Sheldon, R.A. (2007). Cross-linked enzyme aggregates (CLEA®s): stable and recyclable biocatalysts. *Biochemical Society Transactions*, 35(6), 1583–1587. doi:10.1042/bst0351583
- [9] Cao, L., van Rantwijk, F. and Sheldon, R.A. (2000). Cross-Linked Enzyme Aggregates: A Simple and Effective Method for the Immobilization of Penicillin Acylase. *Organic Letters*, 2(10), 1361–1364. doi:10.1021/ol005593x

- [10] Rich, J.O., Leathers, T.D., Anderson, A.M., Bischoff, K.M. and Manitchotpisit P. (2013). Laccases from *Aureobasidium pullulans*. *Enzyme and Microbial Technology*, 53(1), 33–37. doi:10.1016/j.enzmictec.2013.03.015
- [11] Ademakinwa, A.N. and Agboola, F.K. (2016). Biochemical characterization and kinetic studies on a purified yellow laccase from newly isolated *Aureobasidium pullulans* NAC8 obtained from soil containing decayed plant matter. *Journal of Genetic Engineering and Biotechnology*, 14(1), 143–151. doi:10.1016/j.jgeb.2016.05.004
- [12] Ademakinwa, A.N. and Agboola, F.K. (2015). Bioremediation of textile dye solutions, textile dye mixtures and textile effluents by laccase from *Aureobasidium pullulans* (de Bary) G. Arnaud (1918)(Fungi: Ascomycota). *Brazilian Journal of Biological Sciences*, 2(4), 253-262.
- [13] Vršanská, M., Voběrková, S., Jiménez Jiménez, A., Strmiska, V. and Adam, V. (2017). Preparation and Optimisation of Cross-Linked Enzyme Aggregates Using Native Isolate White Rot Fungi *Trametes versicolor* and *Fomes fomentarius* for the Decolourisation of Synthetic Dyes. *International Journal of Environmental Research and Public Health*, 15(1), 23. doi:10.3390/ijerph15010023
- [14] Ferreira, S., Duarte, A.P., Ribeiro, M.H.L., Queiroz, J.A. and Domingues, F.C. (2009). Response surface optimization of enzymatic hydrolysis of *Cistus ladanifer* and *Cytisus striatus* for bioethanol production. *Biochemical Engineering Journal*, 45(3), 192–200. doi:10.1016/j.bej.2009.03.012
- [15] Ademakinwa, A.N. and Agboola, F.K. (2020). Some biochemical, catalytic, thermodynamic and kinetic properties of purified fructosyltransferase from wild and improved mutant-type *Aureobasidium pullulans* NAC8. *Biocatalysis and Biotransformation*, 1–12. doi:10.1080/10242422.2019.1671376
- [16] Bradford, M.M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry*, 72(1-2), 248–254. doi:10.1016/0003-2697(76)90527-3
- [17] Ademakinwa, A.N., Ayinla, Z.A., Omitogun, O.G. and Agboola, F.K. (2018) Preparation, characterization and optimization of cross-linked fructosyltransferase aggregates for the production of prebiotic fructooligosaccharides. *BioTechnologia*, 99(4), 417-434. <https://doi.org/10.5114/bta.2018.79972>

- [18] Melikoglu, M., Lin, C.S.K. and Webb, C. (2013). Kinetic studies on the multi-enzyme solution produced via solid state fermentation of waste bread by *Aspergillus awamori*. *Biochem. Eng. J.* 80 76–82. doi:10.1016/j.bej.2013.09.016
- [19] Yang, J., Xu, X., Yang, X., Ye, X. and Lin, J. (2016). Cross-linked enzyme aggregates of *Cerrena laccase*: Preparation, enhanced NaCl tolerance and decolorization of Remazol Brilliant Blue Reactive. *Journal of the Taiwan Institute of Chemical Engineers*, 65, 1–7. doi:10.1016/j.jtice.2016.04.025
- [20] Ramírez-Cavazos, L.I., Junghanns, C., Ornelas-Soto, N. (2014). Purification and Characterization of two thermostable laccases from *Pycnoporus sanguineus* and potential role in degradation of endocrine disrupting chemicals. *Journal of Molecular Catalysis B: Enzymatic*, 104, 32–42.
- [21] Lassouane, F., Aït-Amar, H., Amrani, S. and Rodriguez-Couto, A. (2019). A promising laccase immobilization approach for Bisphenol A removal from aqueous solutions. *Bioresour. Technol.* doi:10.1016/j.biortech.2018.09.129
- [22] Teerapatsakul, C., Parra, R., Keshavarz, T. and Chitradon, L. (2017). Repeated batch for dye degradation in an airlift bioreactor by laccase entrapped in copper alginate. *International Biodeterioration and Biodegradation*, 120, 52–57. doi:10.1016/j.ibiod.2017.02.001
- [23] Ademakinwa, A.N. and Agboola, F.K. (2019). Kinetic and thermodynamic investigations of cell-wall degrading enzymes produced by *Aureobasidium pullulans* via induction with orange peels: application in lycopene extraction. *Prep. Biochem. Biotech.* 49, 949-960. <https://doi.org/10.1080/10826068.2019.1650375>
- [24] Ademakinwa, A.N. and Agunbiade, M.O. and Fagbohun, O. (2020). Biodegradation of cyanide in cassava wastewater using a novel thermodynamically-stable immobilized rhodanese. *Preparative Biochemistry & Biotechnology*, 1–11. doi:10.1080/10826068.2020.1846053
- [25] Ademakinwa, A.N., Agunbiade, M.O., Ayinla, Z.A. and Agboola, F.K. (2019). Optimization of aqueous two-phase partitioning of *Aureobasidium pullulans* α -amylase via response surface methodology and investigation of its thermodynamic and kinetic properties. *International Journal of Biological Macromolecules*. doi:10.1016/j.ijbiomac.2019.08.159
- [26] Souza, P.M., Aliakbarian, B., Filho, E.X.F., Magalhães, P.O., Junior, A.P., Converti A. and Perego P. (2015). Kinetic and thermodynamic studies of a novel acid protease from

Aspergillus foetidus. International Journal of Biological Macromolecules, 81, 17–21 (2015). doi:10.1016/j.ijbiomac.2015.07.043

[27] Saqib, A.A.N., Hassan, M., Khan, N.F, Baig S. (2010). Thermostability of crude endoglucanase from Aspergillus fumigatus grown under solid state fermentation (SSF) and submerged fermentation (SmF). Process Biochemistry, 45(5), 641–646 (2010). doi:10.1016/j.procbio.2009.12.011

[28] Gohel, S.D. and Singh, S.P. (2013). Characteristics and thermodynamics of a thermostable protease from a salt-tolerant alkaliphilic actinomycete. International Journal of Biological Macromolecules, 56, 20–27. doi:10.1016/j.ijbiomac.2013.01.028

[29] Schoevaart, R., Wolbers, M.W., Golubovic, M., Ottens, M., Kieboom, A.P.G., van Rantwijk, F. and Sheldon, R.A. (2004). Preparation, optimization, and structures of cross-linked enzyme aggregates (CLEAs). Biotechnology and Bioengineering, 87(6), 754–762. doi:10.1002/bit.20184

[30] Asadgol, Z., Forootanfar, H. and Rezaei, S. (2014). Removal of phenol and bisphenol-A catalyzed by laccase in aqueous solution. J Environ Health Sci Engineer 12, 93. <https://doi.org/10.1186/2052-336X-12-93>

546 **Fig.1.** Response surface plots for the preparation of CLEA. Effect of time and pH (A). The Figures
547 indicating the effects of glutaraldehyde and precipitant (B) precipitant and pH (C) temperature and
548 time (D) and temperature and pH on the immobilization yield of the laccase CLEA are shown in the
549 supplementary section.

550

551

552 **Fig. 2.** Effect of pH on free an immobilized laccase

553

Fig. 3. Effect of temperature on free and immobilized laccase

Fig. 4. Arrhenius-type plot to estimate the activation energy of thermal denaturation (E_d^*) for free and immobilized laccase

Fig. 5. Scanning electron microscopy of the laccase CLEA

586 Table 1: Regression analysis and analysis of variance (ANOVA) table for the immobilization yield (%) of
 587 laccase CLEA preparation

Source	Sum of squares	df	Mean square	F-value	p-value prob > F	
Model	1168.20	20	58.41	36.31	<0.0001	Significant
X_1	680.29	1	680.29	422.85	<0.0001	Significant
X_2	14.02	1	14.02	8.71	0.0062	Significant
X_3	71.66	1	71.66	44.54	0.0002	Significant
X_4	29.74	1	29.74	18.48	0.0024	Significant
X_5	9.36	1	9.36	5.82	<0.0001	Significant
X_1X_2	136.11	1	136.11	84.60	0.043	Significant
X_1X_3	7.42	1	7.42	4.61	<0.0001	Significant
X_1X_4	137.32	1	137.32	85.36	<0.0001	Significant
X_1X_5	107.19	1	107.19	66.63	0.0003	Significant
X_2X_3	26.79	1	26.79	16.65	0.004	Significant
X_2X_4	6.83	1	6.83	4.25	0.2499	Not Significant
X_2X_5	2.27	1	2.27	1.41	0.0007	Significant
X_3X_4	23.15	1	23.15	14.39	0.1036	Not Significant
X_3X_5	4.54	1	4.54	2.82	0.2527	Not Significant
X_4X_5	2.19	1	2.19	1.36	0.0276	Significant
X_1^2	8.66	1	8.66	5.38	0.7915	Not Significant
X_2^2	0.11	1	0.11	0.071	0.1775	Not Significant
X_3^2	3.07	1	3.07	1.91	0.0930	Not Significant
X_4^2	4.91	1	4.91	3.05	0.2859	Not Significant
X_5^2	1.90	1	1.90	1.18		
Residual	46.66	29	1.61			
Lack of fit	46.66	22	2.12			
Pure error	0.00	7	0.00			
Cor Total	1214.86	49				

588 *Std. Dev* = 1.27; R^2 = 0.96; *Adj R*² = 0.94; *C.V.* = 1.92; *Adeq Precision* = 23.79

589

590

591

592

593

594

595

596

597

598

599 Table 2. Summary of the kinetic and thermodynamic parameters for free and immobilized laccase

T (K)	k_d (h^{-1})	$t_{1/2}$ (h)	E_d (kJ/mol/K)	ΔH_d^* (kJ/mol)	ΔG_d^* (kJ/mol)	ΔS_d^* J/mol/K
40	#0.0317 $R^2 = 0.96$ *0.0113 $R^2 = 0.94$	#21.8 *61.1	#39.9 $R^2 = 0.86$ *42.4 $R^2 = 0.96$	#37.3 *39.8	#88.1 *88.5	#-162.3 *-155.6
50	#0.0645 $R^2 = 0.98$ *0.0128 $R^2 = 0.95$	#10.8 *54.1		#37.2 *39.7	#88.6 *91.0	#-159.1 *-158.8
60	#0.1718 $R^2 = 0.98$ *0.026 $R^2 = 0.94$	#4.0 *26.3		#37.1 *39.6	#86.7 *91.9	#-149.0 *-157.1
70	#0.1822 $R^2 = 0.99$ *0.0401 $R^2 = 0.94$	#3.8 *17.3		#37.0 *39.6	#89.3 *93.6	#-152.5 *-157.4
80	#0.1955 $R^2 = 0.89$ *0.0791 $R^2 = 0.97$	#3.5 *8.8		#37.0 *39.5	#91.7 *94.4	#-155.0 *-155.5

600 #- Free enzyme

601 *- CLEA

602
603
604
605
606
607
608
609
610

611 Table 3: Regression analysis and analysis of variance (ANOVA) table for the immobilization yield (%) of
612 laccase CLEA preparation

Source	Sum of squares	df	Mean square	F-value	p-value	
--------	----------------	----	-------------	---------	---------	--

					prob > F	
Model	16877.51	14	1205.54	34.56	< 0.0001	Significant
A-BPA	11530.10	1	11530.10	330.56	< 0.0001	Significant
B-Time	75.57	1	75.57	2.17	0.1617	
C-Temp	1735.02	1	1735.02	49.74	< 0.0001	Significant
D-CLEA	826.26	1	826.26	23.69	0.0002	Significant
A2	360.85	1	360.85	10.35	0.0058	Significant
B2	439.98	1	439.98	12.61	0.0029	Significant
C2	42.93	1	42.93	1.23	0.2847	
D2	1940.95	1	1940.95	55.65	< 0.0001	Significant
AB	27.28	1	27.28	0.78	0.3905	
AC	97.71	1	97.71	2.80	0.1149	
AD	224.37	1	224.37	6.43	0.0228	
BC	35.18	1	35.18	1.01	0.3312	
BD	4.17	1	4.17	0.12	0.7342	
CD	30.24	1	30.24	0.87	0.3665	
Residual	523.20	15	34.88			
Lack of fit	523.20	10	52.32			
Pure error	0.000	5				
Cor Total	17400.71	29				

Std. Dev = 5.91; R^2 = 0.97; *Adj R*² = 0.94; *C.V.* = 11.39; *Adeq Precision* = 20.99

630

S1 Central composite rotatable experimental design variables for the optimization of immobilization yield

631

of laccase CLEA preparation

	Precipitant	Glutaraldehyde	pH	Time	Temperature	Immobilization Yield	
	Conc. (v/v)	(mM)		(h)	°C	(%)	
	(x_1)	(x_2)	(x_3)	(x_4)	(x_5)	Experimental	Observed
1	-1(50)	-1(30)	-1(3.5)	-1(4)	-1(30)	67.87	67.52
2	1(65)	-1(30)	-1(3.5)	-1(4)	-1(30)	60.72	60.15
3	-1(50)	1(70)	-1(3.5)	-1(4)	-1(30)	68.58	68.37
4	1(65)	1(70)	-1(3.5)	-1(4)	-1(30)	62.15	62.85
5	-1(50)	-1(30)	1(5.5)	-1(4)	-1(30)	54.23	57.34
6	1(65)	-1(30)	1(5.5)	-1(4)	-1(30)	62.15	61.50
7	-1(50)	1(70)	1(5.5)	-1(4)	-1(30)	70.73	71.84
8	1(65)	1(70)	1(5.5)	-1(4)	-1(30)	67.87	65.25
9	-1(50)	-1(30)	-1(3.5)	1(8)	-1(30)	67.87	68.54
10	1(65)	-1(30)	-1(3.5)	1(8)	-1(30)	59.30	57.96
11	-1(50)	1(70)	-1(3.5)	1(8)	-1(30)	68.30	67.51
12	1(65)	1(70)	-1(3.5)	1(8)	-1(30)	59.30	58.58
13	-1(50)	-1(30)	1(5.5)	1(8)	-1(30)	70.94	69.92
14	1(65)	-1(30)	1(5.5)	1(8)	-1(30)	57.15	58.08
15	-1(50)	1(70)	1(5.5)	1(8)	-1(30)	70.94	69.73
16	1(65)	1(70)	1(5.5)	1(8)	-1(30)	57.15	59.74
17	-1(50)	-1(30)	-1(3.5)	-1(4)	1(55)	67.87	66.55
18	1(65)	-1(30)	-1(3.5)	-1(4)	1(55)	60.72	60.68
19	-1(50)	1(70)	-1(3.5)	-1(4)	1(55)	68.58	67.16
20	1(65)	1(70)	-1(3.5)	-1(4)	1(55)	62.15	63.14
21	-1(50)	-1(30)	1(5.5)	-1(4)	1(55)	71.01	70.53
22	1(65)	-1(30)	1(5.5)	-1(4)	1(55)	62.15	63.30
23	-1(50)	1(70)	1(5.5)	-1(4)	1(55)	70.95	72.19

24	1(65)	1(70)	1(5.5)	-1(4)	1(55)	67.87	67.11
25	-1(50)	-1(30)	-1(3.5)	1(8)	1(55)	67.94	68.74
26	1(65)	-1(30)	-1(3.5)	1(8)	1(55)	59.30	59.47
27	-1(50)	1(70)	-1(3.5)	1(8)	1(55)	66.08	67.27
28	1(65)	1(70)	-1(3.5)	1(8)	1(55)	59.30	59.45
29	-1(50)	-1(30)	1(5.5)	1(8)	1(55)	70.95	71.48
30	1(65)	-1(30)	1(5.5)	1(8)	1(55)	62.15	61.15
31	-1(50)	1(70)	1(5.5)	1(8)	1(55)	70.95	71.06
32	1(65)	1(70)	1(5.5)	1(8)	1(55)	62.30	62.58
33	-2.38(40)	0(30)	0(4.5)	0(6)	0(45)	70.88	71.48
34	2.38(70)	0(30)	0(4.5)	0(6)	0(45)	52.87	52.63
35	0(60)	-2.38	0(4.5)	0(6)	0(45)	66.44	67.49
36	0(60)	2.38	0(4.5)	0(6)	0(45)	70.88	70.19
37	0(60)	0(30)	-2.38(3.0)	0(6)	0(45)	58.58	58.95
38	0(60)	0(30)	2.38(7.0)	0(6)	0(45)	65.08	65.07
39	0(60)	0(30)	0(4.5)	-2.38(2)	0(45)	63.58	65.02
40	0(60)	0(30)	0(4.5)	2.38(10)	0(45)	62.15	61.08
41	0(60)	0(30)	0(4.5)	0(6)	-2.38(20)	64.30	65.87
42	0(60)	0(30)	0(4.5)	0(6)	2.38(70)	69.30	68.08
43	0(60)	0(30)	0(4.5)	0(6)	0(45)	73.21	72.89
44	0(60)	0(30)	0(4.5)	0(6)	0(45)	72.98	73.12
45	0(60)	0(30)	0(4.5)	0(6)	0(45)	73.66	72.99
46	0(60)	0(30)	0(4.5)	0(6)	0(45)	74.21	73.87
47	0(60)	0(30)	0(4.5)	0(6)	0(45)	73.00	72.67
48	0(60)	0(30)	0(4.5)	0(6)	0(45)	72.86	73.10
49	0(60)	0(30)	0(4.5)	0(6)	0(45)	72.87	73.03
50	0(60)	0(30)	0(4.5)	0(6)	0(45)	73.00	72.65

632

633

634 S2: Central composite rotatable experimental design variables for the optimization of biodegradation of
635 BPA by laccase CLEA

	BPA	Time	Temp	CLEA	Biodegradation	
	(mg/L)	(h)	(C)	(mg)	(%)	
	(x_1)	(x_2)	(x_3)	(x_4)	Experimental	Observed
1	1(40)	-1(2)	1(50)	-1(1.0)	73.8	70.0
2	-2(25)	0(3)	0(40)	0(1.5)	31.5	26.2
3	1(40)	1(4)	1(50)	1(2.0)	64.4	65.1
4	0(35)	2(5)	-1(30)	0(1.5)	18.1	16.1
5	-1(30)	1(4)	0(40)	-1(1.0)	87.0	91.7
6	1(40)	-1(2)	-1(30)	1(2.0)	35.7	38.1
7	1(40)	-1(2)	-1(30)	-1(1.0)	99.9	92.7
8	-1(30)	-1(2)	-1(30)	-1(1.0)	26.5	33.8
9	2(45)	0(3)	1(50)	0(1.5)	56.9	52.5
10	0(35)	0(3)	0(40)	0(1.5)	20.8	23.7
11	0(35)	0(3)	0(40)	0(1.5)	56.2	49.5
12	0(35)	0(3)	0(40)	0(1.5)	17.5	15.3
13	0(35)	0(3)	0(40)	-2(0.5)	70.9	68.6
14	1(40)	-1(2)	0(40)	1(2.0)	27.8	30.8
15	0(35)	-2(1)	1(50)	0(1.5)	63.6	71.1
16	-1(30)	-1(2)	0(40)	1(2.0)	28.4	27.1
17	0(35)	0(3)	1(50)	2(2.5)	90.3	95.0
18	0(35)	0(3)	0(40)	0(1.5)	10.5	7.3
19	0(35)	0(3)	2(60)	0(1.5)	52.1	53.0
20	1(40)	1(4)	0(40)	-1(1.0)	45.7	46.1
21	-1(30)	-1(2)	1(50)	1(2.0)	34.0	43.7
22	1(40)	1(4)	-1(30)	1(2.0)	85.8	77.8
23	-1(30)	1(4)	1(50)	1(2.0)	42.8	43.6

24	1(40)	1(4)	-1(30)	-1(1.0)	19.8	20.9
25	-1(30)	1(4)	1(50)	-1(1.0)	65.6	65.8
26	0(35)	0(3)	0(40)	0(1.5)	65.6	65.8
27	-1(30)	1(4)	1(50)	1(2.0)	65.6	65.8
28	0(35)	0(3)	-2(20)	0(1.5)	65.6	65.8
29	0(35)	0(3)	0(40)	0(1.5)	65.6	65.6
30	-1(30)	-1(2)	-1(30)	-1(1.0)	65.6	65.6

636

637

638

639

640

641

642

643

644

645

646

647

648

663 S3: Response surface plots for the preparation of CLEA. Effect of time and pH (A) glutaraldehyde and
664 precipitant (B) precipitant and pH (C) temperature and time (D) and temperature and pH on the
665 immobilization yield of the laccase CLEA

666
667 S4: Reusability of CLEA

668
669 S5: pH stability of free and immobilized laccase

S6: Thermostability of free laccase

S7: Plot of $\ln (E/E_o)$ against $1/T$ for free laccase

682

683

684

685

686 S8: Thermostability of laccase CLEA

687

688 S9: Plot of ln (E/Eo) against 1/T for immobilized laccase

689

S10: Laccase CLEA FTIR

703

704

705

706

707

708

709

710

S11: Effect of (A) temperature x time (B) time x [BPA] (C) CLEA amount x [BPA] (D) temperature x [BPA] (E) CLEA amount x time and (F) CLEA amount x temperature on the biodegradation efficiency of CLEA on BPA.

711

712

713 S12: Reusability of CLEA in BPA biodegradation

714

715

Figures

Figure 1

Response surface plots for the preparation of CLEA. Effect of time and pH (A). The Figures indicating the effects of glutaraldehyde and precipitant (B) precipitant and pH (C) temperature and time (D) and temperature and pH on the immobilization yield of the laccase CLEA are shown in the supplementary section.

Figure 2

Effect of pH on free and immobilized laccase

Figure 3

Effect of temperature on free and immobilized laccase

Figure 4

Arrhenius-type plot to estimate the activation energy of thermal denaturation (E_d^{**}) for free and immobilized laccase

Figure 5

Scanning electron microscopy of the laccase CLEA

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SiFiles.docx](#)