

Ghost cell glaucoma after intravitreal injection of ranibizumab in proliferative diabetic retinopathy

Jun XU

Beijing Tongren Hospital

Meng ZHAO (✉ zhaomeng_jasmine@sina.com)

Beijing Tongren eye center <https://orcid.org/0000-0002-3705-3056>

Jipeng LI

Beijing Tongren Hospital

Ningpu LIU

Beijing Tongren Hospital

Research article

Keywords: intravitreal injection, vitrectomy, proliferative diabetic retinopathy, ghost cell glaucoma

Posted Date: February 7th, 2020

DOI: <https://doi.org/10.21203/rs.2.11864/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Ophthalmology on April 15th, 2020.

See the published version at <https://doi.org/10.1186/s12886-020-01422-z>.

Abstract

BACKGROUND: The development of ghost cell glaucoma in patients with PDR after intravitreal injection was rare. Here we reported a series of patients with PDR who received Intravitreal Ranibizumab (IVR) and developed ghost cell glaucoma and analyzed the potential factors that might be related to the development of ghost cell glaucoma.

METHODS: Retrospective case series study. The medical records of 71 consecutive eyes of 68 PDR patients who received vitrectomy after IVR from January 2015 to January 2017 were reviewed. The development of ghost cell glaucoma after IVR was recorded. Characteristics of enrolled patients were retrieved from their medical charts. Factors associated with ghost cell glaucoma were compared between eyes with the development of ghost cell glaucoma and eyes without the development of ghost cell glaucoma. Variables were further enrolled in a binary backward stepwise logistic regression model, and the model that had the lowest AIC was chosen.

RESULTS: There were 8 out of 71 eyes of the PDR patients developed ghost cell glaucoma after they received IVR. The interval between detection of elevation of intraocular pressure (IOP) and intravitreal injection ranged from 0 to 2 days. Among them, After IVR, there were two eyes had IOP greater than 30mmHg within 30 minutes, four eyes showed normal IOP at 30 min, and then developed ghost cell glaucoma within 1 day, two eyes developed ghost cell glaucoma between 24 and 48 hours. The mean maximum IOP was 46.5 ± 8.0 mmHg. All patients gained normal IOP after vitrectomy without medicine for lowering IOP. The presence of ghost cell glaucoma was associated with tractional retinal detachment (RR= 4.60 [2.02~8.48], p= 0.004) and fibrovascular membrane involving disk (RR=-3.57 [-7.59~-0.92], p=0.03) (AIC= 39.23 AUC=0.88) in a logistic regression model.

CONCLUSION: Attention to postoperative IOP should be paid to patients with PDR undergoing vitrectomy who receive a preoperative intravitreal injection of anti-VEGF agents. PDR patients with tractional retinal detachment or fibrovascular membrane involving optic disc are more likely to develop ghost cell glaucoma after IVR.

Background

Ghost cell glaucoma can occur in patients with long-standing vitreous hemorrhage[1, 2] and can be refractive to lower intraocular pressure medications. It can cause irreversible visual impairment [3]. It has been reported that ghost cell glaucoma generally occurs where there are vitreous hemorrhage and disruption of the anterior hyaloid surface following surgery or trauma[3-6].

Severe vision impairments caused by proliferative diabetic retinopathy (PDR) often result from complications such as neovascularization and fibrovascular proliferation[1]. Non-clearing vitreous hemorrhage, tractional retinal detachment (TRD), extensive fibrovascular proliferation are common indications of vitrectomy in PDR patients [2, 3]. Intravitreal injection of anti-vascular endothelial growth factor (VEGF) agents can be used as an adjunctive method to vitrectomy in eyes with vitreous

hemorrhage due to PDR [4-8]. It has been proved to be a safe and effective method to shorten the overall surgery time, lower the rate of intraoperative complications, and reduce the occurrence of postoperative hemorrhage[9, 10]. It is reported that the development of ghost cell glaucoma in PDR patients who received IV is rare [7-10].

At present, IV of anti-VEGF agents has also been proved to be safe and effective in the treatment of age-related macular degeneration[11], cystoid macular edema due to retinal vein occlusion [12]in large clinical trials. The sustained intraocular pressure elevation was reported to be related to the total number of injections, a greater frequency of injection, and preexisting glaucoma[13]. The occurrence of ghost cell glaucoma after IV was rarely reported in patients with either age-related macular degeneration or retinal vein occlusion who need IV[11, 12].

Here we reported a series of patients with PDR who developed ghost cell glaucoma after pre-vitrectomy intravitreal injection and analyzed the potential factors that might be related to the development of ghost cell glaucoma.

Methods

This study was a retrospective case series of consecutive patients undergoing intravitreal injection of ranibizumab with a diagnosis of PDR who were planned to take vitrectomy after IV. Records of 126 patients were retrospectively reviewed in the study from January 2015 to January 2017 in the Beijing Tongren Eye center. This study was approved by the Ethics Committee of Beijing Tongren Hospital and adhered to the tenets of the Declaration of Helsinki.

Inclusion criteria: 1) Patients diagnosed with PDR, 2) patients took an intravitreal injection of ranibizumab before vitrectomy, 3)Records with intraocular pressure (IOP) values measured before and after intravitreal injection of ranibizumab (IVR). Exclusion criteria: 1) patients failed to finish at least 1-month follow-up after vitrectomy; 2) patients with history of preexisted open-angle glaucoma, 3) patients with preexisted narrow/closed angle, 4) patients received an intravitreal or subtenon injection of corticosteroids or steroid eye drops within the latest six months, 5) uncontrolled neovascular glaucoma by at least 3 kinds of antiglaucoma medicines.

All patients underwent comprehensive ophthalmological examinations, including best-corrected visual acuity (BCVA) testing using a decimal VA chart, slit-lamp biomicroscopy, IOP measurement, dilated fundus examination with indirect ophthalmoscopy, color fundus photograph, optical biometry, optic coherent tomography (OCT) B scan. Gonioscopy was considered when iris neovascularization was found. BCVA, the axial length, the presence of posterior vitreous detachment (PVD, was defined as the presence of a Weiss ring and visible posterior vitreous cortex under the slit-lamp biocular biomicroscopy examination by the same surgical doctor or by B scan[14], PVD was confirmed by findings in triamcinolone acetonide-assisted vitrectomy), history of diabetes mellitus, history of visual acuity decrease, use of insulin, history of retinal photocoagulation for diabetic retinopathy, sex, age, refraction, presence of iris neovascularization, intraocular lens (IOL), dense vitreous hemorrhage that obscured the

view of optic disc and details of fundus, tractional retinal detachment that threatened the central vision or caused repeated vitreous hemorrhage, fibrovascular membrane involving the disk, presence of macular edema were recorded as the baseline data.

An intravitreal injection of ranibizumab 0.5 mg was performed 1-10 days before vitrectomy. All injections were performed by one surgeon. All patients underwent a 3-port pars plana 23-gauge vitrectomy under general anesthesia. Phacoemulsification surgery was performed before vitrectomy in case of necessary determined by the surgeon. The presence of PVD, a tractional retinal detachment that threatened central vision, fibrovascular membrane involving the disk were confirmed in the vitrectomy after removal of dense vitreous hemorrhage and recorded. The silicon oil tamponade and laser points during vitrectomy were recorded.

All patients underwent a complete series of IOP measurements with an air tonometer (Nidek, Tonoref 3). IOP was measured before IVR, 30min, 2h, 1d, 2d, 3d after intravitreal injection. If elevated IOP occurred, IOP was measured twice a day until the IOP was controlled. The ghost cell glaucoma was defined as the presence of both high IOP and ghost cells in the anterior chamber^[12]. Follow-up visits were scheduled at 1,2,7,14 and 30 days after the initial surgery. The examination included BCVA, IOP, slitlamp dilated fundus examination.

Statistical analysis was performed using R version 3.20 (<http://www.R-project.org>). Patient characteristics were retrieved from their medical charts and recorded in Epidata EntryClientversion2.0.3.15 (<http://epidata.dk>). BCVA results were converted to a logMAR value for statistical analysis. Mean and standard deviation (SD) were calculated for continuous variables with a normal distribution. Median with quartiles was calculated for continuous variables with a non-normal distribution. The t-test or Mann-Whitney U test was carried out for continuous variables. The Chi-square test or Fisher's exact test was carried out for discrete data. To explore the potential factors that might influence the occurrence of ghost cell glaucoma, we divided the patients into two groups, patients with ghost cell glaucoma and patients without ghost cell glaucoma after IVR. Several factors including duration of diabetes mellitus, onset of decrease vision, use of insulin, pre-existence of pan-retinal photocoagulation (PRP) for DR, refraction error, axial length, sex, age, presence of PVD, presence of iris neovascularization (NVI), tractional retinal detachment that threatened the central vision or caused repeated vitreous hemorrhage, IOL, fibrovascular membrane involving the disk, presence of clinical significant macular edema were compared between two groups (Table1). Variables ($p \leq 0.4$) were further enrolled in a binary backward stepwise logistic regression model. One variable was included or excluded from the model each time by comparing the Akaike information criterion (AIC) value, and the model that had the lowest AIC was chosen. The model was assessed by the receiver operating characteristic curve (ROC curve).

Results

A total of 71 eyes of 68 patients were included. Among them, 3 patients received an intravitreal injection in both eyes. The patients' baseline characteristics are presented in Table 1.

There were 8 out of 71 eyes of the PDR patients developed ghost cell glaucoma after they received IVR. The basic characteristics of the patients developed ghost cell glaucoma were listed in Table 1. The interval between the onset of visual symptom and intravitreal injection ranged from 0.2 m to 24 m, with a median of 4 m [1st quartile was 2, 3rd quartile was 7]. There were 7 out of 8 patients who failed to complete pan-retinal photocoagulation due to dense vitreous hemorrhage. Three patients were presented with tiny iris neovascularization on the pupil margin, and they had normal IOP before intravitreal injection. Three patients were presented with dense vitreous hemorrhage and sight threatened tractional retinal detachment. The interval between detection of elevation of IOP and intravitreal injection ranged from 0 to 2 days. Among them, there were 2 eyes had IOP higher than 30mmHg within 30 minutes after IVR, 4 eyes showed normal IOP at 30 min after IVR, and then developed ghost cell glaucoma within 1 day after IVR, 2 eyes developed ghost cell glaucoma between 24-48 hour after the IVR. The mean maximum IOP was 46.5 ± 8.0 mmHg. Five patients required methazolamide, brimonidine tartrate, and carteolol hydrochloride to control IOP, and three patients required additional paracentesis to control IOP. Three patients had persistent ghost cells in the anterior chamber after vitrectomy. All patients gained normal IOP without medication for lowering IOP after vitrectomy. All patients did not show vitrectomy related complications during follow-up.

To confirm the potential factors that might be related to the development of ghost cell glaucoma, the data were divided into two groups by the presence of post-IVR ghost cell glaucoma. Variables in Table 1 with a p-value < 0.4 in the two independent sample comparison were selected in the initial logistic regression model, including age, onset of vision decrease, history of retinal photocoagulation, presence of iris neovascular, presence of tractional retinal detachment, presence of fibrovascular membrane involving disk, and pseudophakic eye. Variables were further enrolled in a binary backward stepwise logistic regression model. One variable was included or excluded from the model each time by comparing the Akaike information criterion (AIC) value, and the model that had the lowest AIC was chosen. The presence of ghost cell glaucoma was associated with tractional retinal detachment (RR= 4.60 [2.02~8.48], p= 0.004) and fibrovascular membranes involving disk (RR=-3.57 [-7.59~-0.92], p=0.03) (AIC= 39.23, AUC=0.88). The logistic regression ROC curve was shown in figure 2.

Discussion

Here we report a group of patients with PDR who developed ghost cell glaucoma after intravitreal injection of ranibizumab and showed some potential factors that may be related to the development of ghost cell glaucoma.

The ghost cell glaucoma is rare [7, 15] after IVR in PDR patients, as reported. It has been said that sustained IOP elevation after IVR caused by ghost cell glaucoma only occurs in 3% PDR eyes with

vitreous hemorrhage [15], but do not occur in eyes with diabetic macular edema or PDR without vitreous hemorrhage [16, 17]. In our series of patients with PDR, the incidence of ghost cell glaucoma after IVR (8/71) is higher than previously reported, which was published as 0.7-2.2%[7, 15, 18], but is much lower than eyes with PDR with post-vitrectomy vitreous hemorrhage [8]. In L. Liu et al.'s study[8], the incidence of ghost cell glaucoma in eyes with PDR after intravitreal bevacizumab for post-vitrectomy vitreous hemorrhage is 3 out of 8 eyes. Ghost cell glaucoma generally occurs where there are vitreous hemorrhage and disruption of the anterior hyaloid surface following surgery or trauma[3-6]. A higher incidence of ghost cell glaucoma after vitrectomy in L. Liu et al.'s study might suggest that the ghost cell could gain entrance to the anterior chamber more easily in the condition of the removal of vitreous by vitrectomy. In our series, the PDR patients had a long-standing or recurrent vitreous hemorrhage that required vitrectomy intervention. Although the anterior hyaloid surface of the vitreous body was relatively intact, but a large amount of pre-existing ghost cell in the vitreous, the disturbance of fluid in the vitreous and transient changes of intraocular pressure during IVR may push the ghost cell into the anterior chamber and contribute to the higher occurrence of ghost cell glaucoma in our study. Moreover, transient elevated pressure in vitreous may damage the anterior hyaloid surface of the vitreous body during IVR. More attention should be paid to a slower injection and maintain a relatively stable IOP during IVR to lessen the development of ghost cell glaucoma.

The onset of ghost cell glaucoma after IVR in our series is different from eyes without IVR. It has been reported that the ghost cell can occur in the vitreous and enter the anterior chamber in phakic eyes spontaneously between 18 months and 4 years after vitreous hemorrhage[1, 2]. The development of ghost cell glaucoma within 1 week after intravitreal injection of bevacizumab in eyes with postoperative vitreous hemorrhage after vitrectomy for PDR has also been reported[8]. In our study, the development of ghost cell glaucoma might be found right after IVR or 1-2 days delay. The development of ghost cell glaucoma was more rapidly in eyes with IVR compared with eyes without intravitreal injection [1]. It is suggested that IVR was a risk factor for the development of ghost cell glaucoma in eyes with vitreous hemorrhage. More attention should be paid on postoperative IOP on the first 2 days after IVR in patients with PDR and vitreous hemorrhage.

We tried to identify the potentially related factors to the development of ghost cell glaucoma in PDR patients who received IVR before vitrectomy. Among the patients who developed ghost cell glaucoma, we found most of the patients failed to receive any PRP treatment, and some of them had iris neovascularization, which may indicate long-standing history and poorly controlled and severe ischemic retinal condition. We also found that tractional retinal detachment (RR= 4.60 [2.02~8.48], p= 0.004) and fibrovascular membranes involved optic disc (RR=-3.57 [-7.59~-0.92], p=0.03) are factors related to the development of ghost cell glaucoma. We failed to identify either NVI or lacking PRP as potential factors related to the development of ghost cell glaucoma in the logistics regression model. It should be provided by a further large sample study. The late complications of PDR result from the development of posterior vitreous detachment and contraction of the fibrovascular membranes [19, 20]. Both tractional retinal detachment and fibrovascular membrane on optic disk may result from adherent of vitreous to the retina, which provides a scaffold for retinal neovascularization to grow into vitreous and can cause repeated

vitreous hemorrhage [16].The latter one may be the source of a large amount of ghost cell in vitreous[1, 8] in our series. The higher incidence of ghost cell glaucoma in our series of PDR patients may also due to the high amount of ghost cells in the vitreous body caused by long-standing vitreous hemorrhage or repeated vitreous hemorrhage.

Vitrectomy is effective as a treatment to lowering IOP in the condition of ghost cell glaucoma by cleaning both the vitreous hemorrhage and reservoir of ghost cells in the vitreous as previously reported[3, 21].In our study, All patients gained normal IOP after vitrectomy without further medications for lowering IOP.

We failed to show the presence of vitreous hemorrhage is associated with the development of ghost cell glaucoma due to a small sample size of patients with vitreous hemorrhage without tractional retinal detachment. Further study with a larger sample is required to confirm whether vitreous hemorrhage caused by neither tractional retinal detachment nor fibrovascular membrane on the optic disc was related to the development of ghost cell glaucoma after intravitreal injection.

The limitation of this study was due to it was a retrospective case series. We failed to include enough samples with pseudophakic eye or vitreous hemorrhage without tractional retinal detachment. The selected bias was also presented due to the patients enrolled in this study was in a tertiary hospital. More complicated cases with longer duration may be enrolled in this study, which may contribute to the high occurrence of ghost cell glaucoma after intravitreal injection.

Conclusion

In summary, this study reports a series of patients with PDR who developed ghost cell glaucoma after pre-vitrectomy intravitreal injection of ranibizumab. The development of ghost cell glaucoma varies from 0 to 2 days and is most anticipated within 1 day after intravitreal injection. The presence of tractional retinal detachment and fibrovascular membrane on the optic disc are factors may be related to the development of ghost cell glaucoma after intravitreal injection. Attention on postoperative IOP should be paid to patients with PDR undergoing vitrectomy who receive a preoperative intravitreal injection of anti-VEGF agents, especially in patients with severe PDR.

Declarations

Ethics approval and consent to participate:

Ethical approval was given by the medical ethics committee of Beijing Tongren Hospital. It does not have reference number as a retrospective case series study.

Consent for publication: Not applicable

Availability of data and material

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

Not applicable.

Authors' contributions

All authors read and approved the final manuscript. MZ collected and analyzed the data, she was the one major contributor in writing the manuscript. XJ interpreted the data and performed the vitrectomy surgeries, he was one major contributor in writing and reviewing the manuscript. JPL performed the vitrectomy surgeries and reviewed the manuscript. NPL reviewed the manuscript and contribute in writing the manuscript.

Acknowledgments

We acknowledge the residents who worked as assistants.

Abbreviations

Akaike information criterion AIC

best-corrected visual acuity BCVA

fluorescence angiography FA

intraocular pressure IOP

intravitreal injection of ranibizumab IVR

intraocular lens IOL

iris neovascularization NVI

optical coherence tomography OCT

posterior vitreous detachment PVD

receiver operating characteristic curve ROC curve

standard deviation SD

References

1. Frazer DG, Kidd MN, Johnston PB: **Ghost cell glaucoma in phakic eyes.** *Int Ophthalmol* 1987, **11**(1):51-54.
2. Brooks AM, Gillies WE: **Haemolytic glaucoma occurring in phakic eyes.** *Br J Ophthalmol* 1986, **70**(8):603-606.
3. Montenegro MH, Simmons RJ: **Ghost cell glaucoma.** *Int Ophthalmol Clin* 1995, **35**(1):111-115.
4. Campbell DG, Simmons RJ, Grant WM: **Ghost cells as a cause of glaucoma.** *American journal of ophthalmology* 1976, **81**(4):441-450.
5. Campbell DG, Simmons RJ, Tolentino FI, McMeel JW: **Glaucoma occurring after closed vitrectomy.** *American journal of ophthalmology* 1977, **83**(1):63-69.
6. Campbell DG: **Ghost cell glaucoma following trauma.** *Ophthalmology* 1981, **88**(11):1151-1158.
7. Arevalo JF, Wu L, Sanchez JG, Maia M, Saravia MJ, Fernandez CF, Evans T: **Intravitreal bevacizumab (Avastin) for proliferative diabetic retinopathy: 6-months follow-up.** *Eye (London, England)* 2009, **23**(1):117-123.
8. Liu L, Wu WC, Yeung L, Wang NK, Kuo YH, Chao AN, Chen KJ, Chen TL, Lai CC, Hwang YS *et al*: **Ghost cell glaucoma after intravitreal bevacizumab for postoperative vitreous hemorrhage following vitrectomy for proliferative diabetic retinopathy.** *Ophthalmic surgery, lasers & imaging : the official journal of the International Society for Imaging in the Eye* 2010, **41**(1):72-77.
9. Dong F, Yu C, Ding H, Shen L, Lou D: **Evaluation of Intravitreal Ranibizumab on the Surgical Outcome for Diabetic Retinopathy With Tractional Retinal Detachment.** *Medicine* 2016, **95**(8):e2731.
10. Smith JM, Steel DH: **Anti-vascular endothelial growth factor for prevention of postoperative vitreous cavity haemorrhage after vitrectomy for proliferative diabetic retinopathy.** *Cochrane Database Syst Rev* 2015(8):CD008214.
11. Modi YS, Tanchon C, Ehlers JP: **Comparative safety and tolerability of anti-VEGF therapy in age-related macular degeneration.** *Drug safety* 2015, **38**(3):279-293.
12. Sangroongruangsri S, Ratanapakorn T, Wu O, Anothaisintawee T, Chaikledkaew U: **Comparative efficacy of bevacizumab, ranibizumab, and aflibercept for treatment of macular edema secondary to retinal vein occlusion: a systematic review and network meta-analysis.** *Expert review of clinical pharmacology* 2018, **11**(9):903-916.
13. Bracha P, Moore NA, Ciulla TA, WuDunn D, Cantor LB: **The acute and chronic effects of intravitreal anti-vascular endothelial growth factor injections on intraocular pressure: A review.** *Survey of ophthalmology* 2018, **63**(3):281-295.
14. Kicova N, Bertelmann T, Irle S, Sekundo W, Mennel S: **Evaluation of a posterior vitreous detachment: a comparison of biomicroscopy, B-scan ultrasonography and optical coherence tomography to surgical findings with chromodissection.** *Acta ophthalmologica* 2012, **90**(4):e264-268.
15. Diabetic Retinopathy Clinical Research N: **Randomized clinical trial evaluating intravitreal ranibizumab or saline for vitreous hemorrhage from proliferative diabetic retinopathy.** *JAMA*

ophthalmology 2013, **131**(3):283-293.

16. Bressler SB, Beaulieu WT, Glassman AR, Gross JG, Jampol LM, Melia M, Peters MA, Rauser ME: **Factors Associated with Worsening Proliferative Diabetic Retinopathy in Eyes Treated with Panretinal Photocoagulation or Ranibizumab.** *Ophthalmology* 2017, **124**(4):431-439.
17. Writing Committee for the Diabetic Retinopathy Clinical Research N, Gross JG, Glassman AR, Jampol LM, Inusah S, Aiello LP, Antoszyk AN, Baker CW, Berger BB, Bressler NM *et al*: **Panretinal Photocoagulation vs Intravitreal Ranibizumab for Proliferative Diabetic Retinopathy: A Randomized Clinical Trial.** *JAMA* 2015, **314**(20):2137-2146.
18. Arevalo JF, Lasave AF, Wu L, Maia M, Diaz-Llopis M, Alezzandrini AA, Brito M, Pan-American Collaborative Retina Study G: **INTRAVITREAL BEVACIZUMAB FOR PROLIFERATIVE DIABETIC RETINOPATHY: Results From the Pan-American Collaborative Retina Study Group (PACORES) at 24 Months of Follow-up.** *Retina (Philadelphia, Pa)* 2017, **37**(2):334-343.
19. Newman DK: **Surgical management of the late complications of proliferative diabetic retinopathy.** *Eye (London, England)* 2010, **24**(3):441-449.
20. Farahvash MS, Majidi AR, Roohipour R, Ghassemi F: **Preoperative injection of intravitreal bevacizumab in dense diabetic vitreous hemorrhage.** *Retina (Philadelphia, Pa)* 2011, **31**(7):1254-1260.
21. Abu el-Asrar AM, al-Obeidan SA: **Pars plana vitrectomy in the management of ghost cell glaucoma.** *Int Ophthalmol* 1995, **19**(2):121-124.

Figures




Figure 1

The receiver operating characteristic curve for the model of logistic regression analysis for potential factors associated with development of ghost cell glaucoma

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [table1supp.docx](#)