

Adsorption and pH values determine the distribution of cadmium in terrestrial and marine soils in the Nansha area, Pearl River Delta

Fangting Wang (✉ ftwang1991@cug.edu.cn)

China University of Geosciences School of Environmental Studies

Ke Bao

CWRC: Changjiang Water Resources Commission

Changsheng Huang

China Geological Survey

Wenjing Han

China University of Geosciences

Zhibin Yin

China University of Geosciences

Zhijia Chen

China University of Geosciences

Xinwen Zhao

China Geological Survey

Research Article

Keywords: Cadmium contamination, Soil pH, Isothermal adsorption experiment, Adsorption kinetic experiment, longitudinal migration, the Pearl River Delta

Posted Date: March 11th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-270015/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at International Journal of Environmental Research and Public Health on January 11th, 2022. See the published version at <https://doi.org/10.3390/ijerph19020793>.

Abstract

Cadmium is a toxic element with a half-life of several decades, which can accumulate in the human body by entering the food chain and seriously harm health. Coastal zones are located in the transitional zone between land and sea, and large amounts of terrigenous material input have important environmental effects on this ecosystem. The pH, Eh, hydrodynamic conditions, soil organic matter(SOM) and other factors defining the sea-land interaction within the sedimentary environment are significantly different from those defining land facies. It was found that the adsorption constant K_f and the Cd^{2+} adsorption capacity of marine soils were higher than those of terrestrial soils. However, the saturation adsorption of cadmium in terrestrial sediments was higher than in marine sediments. In order to study the key factors affecting cadmium adsorption in soils at the sea-land interface in the Nansha area of the Pearl River Delta, a test was conducted on a column of undisturbed soil. The results showed that soil pH was an important factor affecting cadmium adsorption capacity in both terrestrial and marine sediments. Neutral and alkaline topsoil conditions inhibited the vertical migration of cadmium, while the acidic environment favored it. The higher the clay and SOM were, the stronger the Cd^{2+} adsorption capacity of the soil was. These findings suggest that the distribution of cadmium in marine and continental sedimentary soils is not only related to adsorption, but also to the physical and chemical processes occurring in different sedimentary environments.

1. Introduction

Cadmium (Cd) is a carcinogen that can be accumulated in the body by entering the food chain and seriously endanger human health(Godt et al. 2006, Peng et al. 2016). Excessive importation of this heavy metal into the soil has attracted worldwide attention because of its high toxicity, persistence and bio-enrichment capacity(Dong et al. 2018, Li et al. 2017). Survey results show that 13.86% of China's grain production is affected by inorganic pollutants present in soils, among which cadmium ranks as the most polluting metal, accounting for 7.75%(Zhang et al. 2015). Multi-target regional geochemical surveys conducted in China show that cadmium pollution in land areas around the Pearl River Delta is increasing. Major factors contributing to the increase are: wastewater irrigation, atmospheric precipitation, stacking and leaching of solid waste in industrial and mining enterprises, and the use of agricultural supplies(Lai et al. 2005). Overall, the environmental quality of the soil in the Pearl River Delta is poor. Most of the soils in the area, covering an area of 9298.3km², are classified as grade III or inferior to grade III. The main toxic elements detected are Cd, Hg, As, Cu and F(Du et al. 2011). Due to its stable chemical properties, a small part of the cadmium entering the soil is absorbed by the soil medium, while the rest is transferred and transformed in the soil environment, resulting in soil cadmium pollution. Plants absorb cadmium from the contaminated soil and accumulate large amounts of this element, which then enters the food chain once the edible parts of plants are ingested by various animal species(Zhang & Reynolds 2019). Cadmium-contaminated soils will eventually produce cadmium-contaminated grains, posing a serious threat to human health(Feng et al. 2020). Therefore, finding viable solutions to reduce the health risk posed by cadmium pollution has become a research priority in recent years.

The cadmium adsorption and desorption processes occurring in the soil are key processes that control cadmium concentration in soil solutions, which directly affect the migration, transformation, bioavailability and ecotoxicity of this element in soil-plant systems(Elbana & Selim 2010), ultimately affecting the quality of agricultural products and of the environment. Therefore, it is of great practical significance to study the adsorption and desorption characteristics of cadmium in order to understand and control its behavior in the soil-plant system and to predict its environmental effects(Li et al. 2016). The cadmium adsorption behavior in the soil varies with soil type and is affected by soil properties such as pH(Ardestani & van Gestel 2016, Shen et al. 2020), soil organic matter (SOM)(Hamid et al. 2020), ion exchange capacity and clay content(Bai et al. 2017). Cadmium exists in nature as a divalent form, with stable chemical properties, and it is easily soluble in acid solutions. As the surface soil becomes polluted with heavy metals, cadmium is precipitated out under the leaching action of acid rainfall and it gradually sinks and migrates, which may pollute phreatic water aquifers in serious cases. The Pearl River Delta region is characterized by permanent agricultural activities and abundant rainfall(Zhang et al. 2018), which is a distinctive feature of the subtropical monsoon climate(Huang et al. 2018). Acid rain is a widespread phenomenon in the area, with rainfall pH values ranging between 4.0 and 5.6; surface soil pH values are usually low, thus affecting cadmium migration, transformation and bioavailability. On the other hand, coastal zones represent an area where important interactions occur between land, ocean and atmosphere. These zones are vulnerable to environmental change, and they are highly economically developed. At the land-sea interface, various physical, chemical and biological factors are subjected to dramatic variations, making this area particularly fragile. It is very sensitive to the fluctuations caused by various natural processes and by reactions in response to human activities(R & Vvss 2015). Moreover, the large amounts of land-originated terrigenous input have important environmental effects on the entire coastal ecosystem(Liu 2019). Geomorphology in the Pearl River Delta is determined by the sea-land interaction, which also affects the geochemical characteristics of the quaternary soils present in the area(Bai et al. 2012, Li et al. 2014). The pH, Eh, hydrodynamic conditions, SOM and other factors pertaining the sedimentary environments of the sea-land interface are significantly different from those of continental facies, which usually constitute geochemical barriers to the elements that migrate stably in them, ultimately leading to the precipitation and enrichment of elements in the sea-land transition zone(Tang 2017).

Several studies have investigated the effects of soil properties on the adsorption-desorption, transport and transformation of heavy metals(Li et al. 2013, Zeng et al. 2019, Zhou et al. 2018, Zhu et al. 2016). However, these studies primarily focused on surface soils of cultivated lands, and often ignored the soil properties of different sedimentary environments, both terrestrial and marine. In order to prevent and reduce the environmental damage caused by cadmium in the sea-land interaction zone of the Pearl River Delta, it is urgent to study the mechanisms of cadmium migration and transformation in soils within different sedimentary environments. Various studies have investigated the long-term equilibrium effect of metals in soil, but have not paid enough attention to the influence of important soil properties, such as pH(Diagboya et al. 2015, Furuya et al. 2016, Huang et al. 2015, Huang et al. 2013). Consequently, the

interaction between the soil's depositional environment and soil properties during metal adsorption is poorly understood.

This paper is one of a series of studies on the effects of soil properties on heavy metal adsorption. In Wang et al.(2019) (Wang et al. 2019) multivariate statistical and spatial analysis methods were used to analyze the factors affecting the accumulation of cadmium in sedimentary soils at the sea-land interaction zone in the Pearl River Delta. Here, we further extend the model of cadmium adsorption in soils within marine and continental inter-deposition environments considering important additional environmental parameters. In terms of global geography, China's coastal areas lie between the largest continent and the largest ocean, and China's marginal sea is a natural laboratory for studying the interaction between sea and land (Yao 2014). In this study, the soil of both terrestrial and marine sedimentary environments in the Nansha area of the Pearl River Delta and its cadmium adsorption characteristics were studied using isothermal adsorption methods. Through the measurement of external environmental parameters (pH value, temperature, time, etc.), the cadmium adsorption behavior was analyzed in terms of isothermal and kinetic adsorption; and the adsorption characteristics and influencing factors at play within the sea-land transition zone were explored.

2. Material And Methods

2.1 Study Area

The Nansha district connects the Pearl River basin to the ocean, and covers a total area of approximately 803 km². The region is characterized by a southern subtropical monsoon climate, the annual average temperature is 22.2°C, the annual average rainfall is 1646.9 mm. This region is highly affected by thunderstorms which occur 78.3 days per year. The landforms in Nansha are hilly, marine-terrigenous plains, and tidal flats. The low hills are mainly distributed around Huangshanlu Mountain and the highest point in the region is 295.3 m above sea level. The marine-terrigenous plains are distributed in Huangge and Hengli; the upper part is silt or silty soil (sand) and the lower part is river-deposited sand. The tidal flats are mainly distributed in the southeastern area of Wanqingsha, Longxue Island, and the Xinken coast. The terrain is zonal, parallel to the coast, broad, and gentle.

2.2 Soil Sampling and Chemical Analysis

Two types of soils were collected in Wanqingsha Town, Nansha District, to be used in laboratory adsorption experiments to study the adsorption characteristics of soils with different origins. Soil sample 1 belonged to the continental sediment layer; it was identified as silted clay and was collected at a depth ranging from 0 to 0.5 m. Soil sample 2 belonged to the marine sediment layer, it was identified as sandy silt and was collected at a depth ranging from 1 to 1.5 m. All soil samples were air-dried, mashed with a 2 mm screen and sealed separately in airtight polyethylene bags.

A total of 18 groups of deep soil profiles (0~200 cm) were collected in the Nansha District, Guangzhou (Fig.1C). The sampling sites were mainly located in the thicker soil layer of gardens. Care was taken to avoid sections with obvious point source pollution, newly transported and accumulated soil, soil occupied by garbage and ridge fields; sampling was conducted at least 100 m away from main highways and railways. The deep soil was sampled by hand drill, and the soil samples were collected in each section at a depth between 0 and 2 m. They were divided into five groups (0 ~ 20, 20 ~ 50, 50 ~ 90, 90 ~ 140 and 140 ~ 200 cm), A total of 90 groups of soil samples were sealed in airtight polyethylene bags.

All the soil samples were taken back to the laboratory and the physical and chemical indexes were determined. Soil pH was determined by a pH meter with an accuracy of 0.01(2016). The SOM content and cation exchange capacity (CEC) were determined using reference standards(1121.6-2006) (2016). The total cadmium content in soil samples was determined by an inductively-coupled plasma mass spectrometer with a detection limit of 0.02 μ g/g (14506.30-2010). According to the International Classification of Soil Texture, soil particles are classified into the following types(1121.3-2006): clayey (<0.002mm), silty (0.002~0.02mm) and sandy (0.02~2mm).

2.3 Batch experiments

The main reagents used in the experiments were pH standard buffer solution (4.01, 6.86, 9.18), hydrochloric acid and the national standard cadmium solution (product number: GSBG62040-90, concentration: 1000 μ g/mL, medium: 10%HCl). The chemical reagents used were high-purity analytical reagents. Deionized water (18.2 M Ω ·cm) was prepared with a Heal Force NW ultrapure water system and used in all experiments.

2.3.1. Isothermal adsorption experiment

A series of standard solutions containing cadmium concentrations of 0.1, 0.5, 1, 5, 10, 50, 100, 150, 200, 250, 300, 350, 400 and 450 mg/L were prepared, with 0.01 mol/L NaCl solution as supporting electrolyte solution. The basis for setting the soil-liquid ratio was as follows: the volume mass (bulk density) of the experimental soil was about 1.5 g/cm³, and considering that the 0~20 cm topsoil layer near was the main layer affected by acid rain, the weight of soil column per 1 cm² unit area was 30 g. According to meteorological data, the annual rainfall in Guangzhou is about 1646.9mm. Therefore, the soil-liquid ratio used in the experiment was 30g: (1g/cm³×1646.9mm×1cm²)≈1:5.5. A quantity of 8 ± 0.001 g of soil sample, and 44 mL of a series of standard cadmium solutions were added to a 50 mL plastic centrifuge tube. The mixtures were evenly mixed and shaken at a constant temperature of 25±1°C for 24h in a thermostatic oscillator to obtain full adsorption (two sets of parallel samples for each cadmium solution, and a total of 28 groups of mixed samples were prepared), then they were centrifuged at 4030 r/min for 20 min. The supernatant was removed to analyze pH \square PHS-3C with calibrated every day \square and conductivity (DDS-307A \square . The concentration of cadmium in the supernatant was measured with an atomic absorption spectrophotometer (ZCA-1000SF8).

The adsorption effect is usually defined by the adsorption capacity and adsorption rate: the adsorption capacity S and adsorption rate η of soil to cadmium are calculated as follows:

$$S = (C_0 - C_e) V / m \quad (1)$$

$$\eta (\%) = (C_0 - C_e) \times 100 / C_0 \quad (2)$$

where, S is the adsorption capacity of soil to cadmium, mg/kg; η is the adsorption rate of soil to cadmium, %; C_0 is the initial concentration of cadmium in the solution, mg/L; C_e is the concentration of cadmium in the equilibrium solution, mg/L; V is the volume of the equilibrium solution, mL; m is the mass of the soil, g.

2.3.2 Adsorption kinetic experiment

Using 0.01 mol/L NaCl as the supporting electrolyte solution and 0.1 mol/L HCl to adjust pH to 3.5, 4.5, 5.5 and 9.5, a standard solution with a cadmium concentration of 100 mg/L at different pH values was prepared. A quantity of 8 ± 0.001 g of soil sample and 44 mL of standard cadmium solution were added to a 50 mL centrifuge tube in that order, based on the soil-liquid ratio of 1:5.5. The mixtures were evenly mixed and shaken at a constant temperature of 25 ± 1 °C in a thermostatic oscillator for 2 min, 5 min, 10 min, 15 min, 20 min, 25 min, 30 min, 1 h, 2.5 h, 4 h, 6 h, 8 h, 10 h, 12 h and 24 h respectively (two sets of parallel samples for each time interval, and a total of 150 groups of mixed samples were taken). Each set of tubes was placed in a 5030 r/min centrifuge for 5 min immediately after the oscillation phase. The supernatant was removed to analyze pH (PHS-3C with calibrated every day) and conductivity (DDS-307A). The concentration of cadmium in the supernatant was measured with an atomic absorption spectrophotometer. Finally, the adsorption capacity S of soil to cadmium was calculated based on the initial equilibrium concentration of cadmium in the solution.

2.4 Analysis

SPSS 22.0 software was used for correlation analysis, and Origin 2018 software for experimental data sorting and analysis. The parameters of the isothermal adsorption and adsorption kinetics models were fitted using the results of the respective experiments.

The results of heavy metal adsorption analyses are usually expressed by isothermal adsorption curves. For different reactions, there are specific fitting equations that can describe adsorption behavior. Common isothermal adsorption equations include the Henry, Langmuir and Freundlich equations, which are described as follows:

$$\text{Henry equation: } S = K_d C_e \quad (3)$$

$$\text{Freundlich equation: } S = K_F C_e^{1/n} \quad (4)$$

Langmiur equation: $S = K_L S_m C_e / (1 + K_L C_e)$ (5)

where, S is the adsorption capacity, mg/kg; K_d is the solid/liquid distribution coefficient, L·kg⁻¹; C_e is the concentration of equilibrium solution, mg/L; K_F and $1/n$ are isothermal adsorption constants, K_L is the value of soil adsorption affinity, and S_m is the saturation adsorption, mg/kg.

Adsorption kinetics models are mainly used to describe adsorption rates, reactions at a stable temperature, relationships between adsorption substances and time, all of which collectively describe the mechanisms of heavy metal adsorption by soils. Common dynamics model equations include the Elvoich equation, the double constant equation, and the pseudo-first-order dynamics equation. These are described as follows:

Elvoich equation: $S_t = m + n \ln t$ (6)

Biconstant equation: $\ln S_t = m + n \ln t$ (7)

Quasi first order kinetic equation: $\ln(S_m - S_t) = \ln S_m - (K_1 t) / 2.303$ (8)

where, S_t is the solute adsorption capacity of soil in time t , mg/kg; m and n are constants; K_1 is the kinetic parameter; and S_m is the saturation adsorption, mg/kg.

3. Results And Discussions

3.1 Physical and chemical properties of soils in two sedimentary environments

The land morphology of the sampling site is delta plain, and this land is primarily used for banana plantations. The soil in the first terrestrial sedimentary environment (Soil 1) was a yellowish silty clay with a 35% clay ratio, pH value of 6.6, weak acidity, 1.8% SOM content, and 0.52 mg/kg cadmium content. The soil in the marine sedimentary environment (Soil 2) was gray-brown sandy silt, moist, with a soft to fluid plasticity, 64% of sand proportion, a 8.08 pH value (alkaline), 1.58% SOM content, and 0.48mg/kg cadmium content (Table 1). The cadmium background concentration in Guangdong Province soils was 0.056 mg/kg(Center 1990), while it was double in the Pearl River Delta, reaching 0.11mg/kg(DB44/T1415-2014). The cadmium content measured in the two sedimentary soils in the study area was significantly higher than the background cadmium value of both Guangdong Province and Pearl River Delta.

Table 1 Basic physical and chemical properties of soils in two sedimentary environments

	Soil mechanical composition ratio (%)			SOM	CEC	pH	Cd
	D<0.002 mm	0.002 mm <D<0.02 mm	0.02 mm <D<2 mm	g/kg	cmol/kg		mg/kg
Soil 1	35	36.8	28.2	18.0	15.2	6.40	0.52
Soil 2	18.6	17.4	64	15.8	10.5	8.08	0.48

3.2 Effects of initial concentrations on cadmium adsorption in the two soils

During the isothermal adsorption experiment, 8 g of each soil sample was evenly mixed with 44 mL of cadmium solutions with different concentrations, and the effects of the soil on the adsorption capacity of cadmium were studied while vibration and a constant temperature of $25\pm 1^\circ\text{C}$ were maintained for 24 h. As shown in Fig. 2A, the adsorption capacity of Cd^{2+} increased in all samples with the increase of the initial cadmium concentration, and the variation can be divided into three stages (Fig. 2B): (1) when the initial concentration was between 0 and 50 mg/L, the adsorption ratio of Cd^{2+} on soil increased significantly, (2) when it was between 50 and 200 mg/L, the adsorption ratio of Cd^{2+} was high, still greater than 95%, and (3) when it was more than 200 mg/L, the adsorption ratio decreased gradually.

In Figures 2C, 2D and 2E, the Henry, Freundlich and Langmuir models are shown, respectively. These models were used to fit the adsorption of cadmium by the two soils, and to establish the isotherm adsorption equation. According to the correlation coefficient analysis, the Freundlich equation has a better fitting for the cadmium adsorption process. The adsorption parameters of the two soils obtained through the three isothermal adsorption models are shown in Table 2.

Table 2 Isothermal adsorption parameters of cadmium in the two soils

Adsorption model	Parameter	Unit	Soil 1	Soil 2
Henry equation	K_d	L/kg	42.81	51.11
Freundlich equation	K_F	-	363.04	558.21
	$1/n$	-	0.43	0.34
Langmuir equation	K_L	-	0.072	0.162
	S_m	mg/kg	2419.83	2228.25

Houng et al.(1998) (Houng &Lee 1998) observed that the K_F constant in the Freundlich model is an important parameter to evaluate the adsorption capacity of heavy metals during soil adsorption processes, and the value of K_F is proportional to the capacity of the soil to adsorb heavy metal ions, but independent of saturation adsorption. Table 2 shows that the cadmium isothermal adsorption constant K_F of soil 2 in the study area is significantly higher than that of soil 1, measuring 558.21 and 363.04, respectively. The higher the K_F value is, the stronger the soil adsorption capacity for Cd^{2+} (Arabi &Sohrabi 2014)). Therefore, the adsorption capacity of soil 2 is stronger than that of soil 1. This is because soil 1 is a neutral soil while soil 2 is alkaline, and the amount of negative charge on the colloidal surface of alkaline soils is generally greater, resulting in an enhanced electrostatic action of the soil. This, in turn leads to the enhancement of the obligate adsorption capacity of the colloidal surface to metal ions(Barrow 1984). Furthermore, the SOM content of soil 1 is higher than that of soil 2, and the soil particles with more SOM content combine with inorganic colloids to form new complex colloids, thus weakening the adsorption capacity of soil particles to heavy metals. Nevertheless, the saturation adsorption of cadmium in soil 2 is smaller than that in soil 1, measuring 2228.25 mg/kg and 2419.83 mg/kg, respectively. This may be due to the clay content present in the soil. The specific surface area of clay particles is large, it is usually negatively charged and has a high cation exchange capacity. Under the action of static force, positively charged cations will adsorb on the surface of clay minerals(Huang et al. 2012). Therefore, it is believed that the higher the content of physical clay particles in the soil, the stronger the soil's ability to adsorb Cd^{2+} . The clay particle content in terrestrial sediments in the study area is high, accounting for 35% of the total, the cation exchange capacity is 15.2 cmol/kg, and the soil has a strong cadmium adsorption capacity. Conversely, the marine sediment soil has a high sand content (64%), a cation exchange capacity of 10.5 cmol/kg, and a weak soil adsorption capacity, indicating that clay minerals play an important role during the process of cadmium adsorption.

The solid-liquid phase equilibrium controls the mobility and toxicity of metals in soils. The higher the K_d value is, the easier it is for Cd^{2+} to be fixed in the soil solution through adsorption; the lower the K_d value is, the more Cd^{2+} exists in the soil solution(Duan et al. , Sauve et al. 2003). The variation of the solid-liquid distribution coefficient K_d with the initial solution concentration is shown in Fig. 2F. The variation trend of K_d seen in soils from different sedimentary environments is basically the same, showing a first increase, then a decrease and finally a stabilization with the increase of the initial concentration. Specifically, in this study, when the initial cadmium concentration of soil 1 was 0-10 mg/L, the K_d value increased, reaching the maximum peak. At a concentration of 10-100 mg/L, the value decreased rapidly until it stabilized. When the initial concentration of soil 2 was 0-50 mg/L, the K_d value increased reaching the maximum peak. At a concentration of 50-150 mg/L, the value decreased rapidly until it stabilized, indicating the soil's ability to hold Cd^{2+} by reaching adsorption equilibrium. Considering that the stability value of the soil's K_d does not change significantly in different sedimentary environments – but the peak value of K_d changes significantly – the latter was chosen to characterize the soil's ability to retain Cd^{2+} . By comparing the K_d peak values of the two soil types, it emerged that the K_d peak values of soil 2 were

significantly higher than those of soil 1, indicating that Cd^{2+} in the soil 2 solution was more easily fixated by adsorption, and its adsorption capacity was stronger. This process depends on the amount of cation exchange between the two soils. Generally, a higher total metal concentration results in a lower proportion of metals being adsorbed because potential adsorption sites are filled in order of decreasing affinity. The more a soil becomes “saturated” with cations, the lower the metal affinity of the remaining sites. We would then expect the resulting K_d values to decrease with the increasing of total metal levels (Sauve et al. 2003).

3.3 Effect of adsorption time on cadmium adsorption in the two soils

Adsorption kinetics express the relationship between solution concentration and reaction duration under fixed temperature conditions. In this study, 8g of each soil sample was evenly mixed with 44 ml Cd^{2+} solution with a concentration of 100 mg/L. By controlling the temperature ($t = 25 \pm 1^\circ\text{C}$), it was possible to analyze the cadmium adsorption capacity of the soil at different times. As shown in Fig. 3A, at the beginning of the adsorption process, the soil adsorbed Cd^{2+} combined with soil colloids by simple Coulomb force through the double potential layer (non-mandatory adsorption) (Lin & Xu 2008). The adsorption rate was fast and close to saturation, and the adsorption capacity reached the peak. Desorption occurred between one and 10 minutes, resulting in the increase of equilibrium concentration, and the soil adsorption capacity of cadmium decreased significantly from the maximum to the minimum value. After 10 minutes, the adsorption of Cd^{2+} changed into mandatory adsorption, and the adsorption rate was slow. With the increase of oscillation time, the equilibrium concentration gradually decreased, and the cadmium adsorption capacity on the soil gradually increased and tended to stabilize, reaching the equilibrium adsorption. By comparing the relationship curves of soil adsorption capacity and oscillation time in the two different sedimentary environments, it is observed that when the concentration of Cd^{2+} in aqueous solution was 100 mg/L, the initial maximum adsorption capacity of soil 1 and soil 2 ($t = 1\text{min}$) was 529.65 and 539.33 mg/kg, respectively, and the equilibrium adsorption capacity ($t = 24\text{h}$) was about 538 and 547 mg/kg, respectively. The initial maximum adsorption capacity and equilibrium adsorption capacity of soil 2 are both greater than those of soil 1.

The relationship between the adsorption capacity and time is characterized by the adsorption rate, which refers to the capacity of cadmium to be adsorbed by soil per unit mass in unit time (Wang et al. 2012), i.e., $V = S/t$. The relationship between the adsorption rate and adsorption time of marine and terrestrial soils is shown in Fig. 3B. At the beginning of the shock experiment ($t = 1\text{min}$), the adsorption rate was the fastest, then, with the increase in adsorption time, the soil adsorption rate slowed. At 2.5 h, the adsorption rate in the two soils was 3.6 mg/(kg·min), and after 24 h the rate was the slowest, at 0.4 mg/(kg·min), and in a state of equilibrium adsorption.

The biconstant equation, Elovich equation and a first order quasi-linear equation fitting were used to simulate the dynamic process of cadmium adsorption by terrestrial and marine soils in the Nansha area. The adsorption kinetic model was established based on the relationship between the amount of cadmium adsorbed and the adsorption time of the two soils. Both the Elovich equation and the biconstant equation can adequately describe the adsorption process, while the first order quasi-linear equation fitting is poor. However, a similar exponential asymptote equation can be used to fit the dynamic process of cadmium adsorption, one which provides better results than the other two equations. The dynamics fitting degree of the two soils is similar. Based on the exponential asymptote equation, when the ratio of soil to water was 5.5:1 and the concentration of Cd^{2+} was 100mg/L, the saturation adsorption of soils 1 and 2 was 537.03 mg/kg and 547.49 mg/kg, respectively.

3.4 Effect of solution pH value on cadmium adsorption in two soils

The adsorption behavior of cadmium in the two soils with different solution pH values is shown in Fig.4 (cadmium solution concentration is 100mg/L, water and soil ratio is 5.5:1). In general, the adsorption capacity increased as the pH became more alkaline. When the pH of the cadmium solution was 3.5, the cadmium adsorption capacity was significantly lower compared to the other three pH values; while at a pH value of 9.5, the adsorption capacity was significantly higher. When the pH of the cadmium solution was 4.5 and 5.5, the cadmium adsorption capacity was similar. The dynamic exponential asymptote equation was adopted to fit the adsorption kinetics of cadmium, as shown in Table 3: when the reaction time t tends to infinity, the saturation adsorption of S_m in the two soils increases as the overall pH of the solution increases (the saturation adsorption of soils 1 and 2 was 536.27 ~ 540.7 mg/kg and 545.32 ~ 547.48 mg/kg, respectively). Under the same pH conditions, the saturation adsorption S_m of soil 2 was greater than that of soil 1. In addition, when the reaction time was 24 h, the adsorption ratio of soils with different pH values was different (Fig. 4C).

These results indicate that pH value may affect cadmium adsorption mechanisms in soils. Davis and Leckie (1978) (Davis & Leckie 1978) compared the intrinsic surface complexation constants of free metal ions and hydroxy-oxygen complexes on different metal oxides, and found that hydroxy-oxygen complexes are more stable. Firstly, under neutral and weakly alkaline conditions, the exchange affinity of $\text{Cd}(\text{OH})^+$ generated by the hydrolysis of heavy metal ions is greater than that of metal free Cd^{2+} ions, resulting in an increase in the adsorption ratio. Secondly, the increase of soil pH increases the negative charge capacity on the soil colloidal surface, which leads to the enhancement of the soil electrostatic action and, consequently, to the enhancement of the obligate adsorption capacity of the soil colloidal surface to metal ions (Barrow 1984). The mobility of cadmium is very low under neutral and basic pH conditions due to its precipitation as carbonate (otavite: CdCO_3) and silicate (CdSiO_3) minerals, as well as to its adsorption of ferric hydroxide/oxide and clay minerals (Tabelin et al. 2018). However, the lower the

pH value, the more H^+ is present in the solution, leading to a greater competition between Cd^{2+} and H^+ for adsorption sites, which results in a decrease of cadmium adsorption capacity in low pH soils.

Table 3 Dynamic fitting of the exponential asymptote equation for the two soils under different solution pH values.

Soil	pH	Expression	COD (R^2)	S_m (mg/kg)
Soil 1	pH=3.5	$S = 536.27 - 12.13 * 0.968^t$	0.899	536.27
	pH=4.5	$S = 538.85 - 11.11 * 0.959^t$	0.948	538.85
	pH=5.5	$S = 537.18 - 9.69 * 0.852^t$	0.864	537.18
	pH=9.5	$S = 540.70 - 5.64 * 0.977^t$	0.906	540.70
Soil 2	pH=3.5	$S = 545.32 - 6.84 * 0.933^t$	0.883	545.32
	pH=4.5	$S = 546.27 - 5.83 * 0.933^t$	0.875	546.27
	pH=5.5	$S = 546.14 - 5.24 * 0.994^t$	0.801	546.14
	pH=9.5	$S = 547.48 - 4.22 * 0.994^t$	0.915	547.48

3.5 Factors affecting cadmium adsorption in undisturbed soil columns

According to previous studies, the cadmium adsorption capacity of marine sedimentary soils is greater than that of terrestrial sedimentary soils, and the adsorption rate is also faster. In an environment where intense interaction occurs between land and sea, the physical and chemical conditions of environmental media change dramatically, and the activity of heavy metals in sediments also changes correspondingly. Dou et al (2015) (Dou et al. 2015) summarized the characteristics of elements in marine and terrestrial sediments based on multi-objective geochemical surveys, and found that the content of most elements was higher in the marine sediment than in the continental one.

In this paper, an analysis of the influence of soil pH, SOM and soil mechanical composition on cadmium adsorption was conducted on 18 groups of deep soil samples in order to reveal the key factors affecting cadmium adsorption in undisturbed soil columns.

3.5.1 Effect of soil pH on cadmium migration in undisturbed soil columns

The above-mentioned studies on the effect of solution pH on cadmium adsorption in different soils suggest that pH is an important factor affecting cadmium adsorption capacity in both the terrestrial and marine sediments of the Nansha Area, and an increase in pH can enhance Cd^{2+} adsorption capacity. Figure 5A plots the variation of cadmium content and pH in the deep soil profile of the Nansha Area. The results show that, with increasing soil depth, pH tends to increase overall, while cadmium content gradually decreases, which is exactly the opposite of what has been observed in the previous analysis. What is the reason for such a different result ?

Previous studies(Wang et al. 2019) have shown that the surface soil in the Nansha area of the Pearl River Delta is to some extent polluted by cadmium, mainly because of anthropogenic inputs. Cadmium is mainly concentrated in the surface and analysis of pH and cadmium content in the 0-20 cm surface layer – which consists of silty clay – showed that soil pH was positively correlated with cadmium content ($P < 0.01$, correlation coefficient $r = 0.554$). The influence of soil pH on cadmium adsorption-desorption processes is not only reflected in the surface soil, but also in the longitudinal migration and transformation of this element. Cadmium content significantly decreases with depth, and its variation can be subdivided into three categories (Fig. 6), based on pH differences: (1) Between 0 ~ 20 cm of soil with a pH of 6.5 ~ 7.5, the soil is neutral and reaches the maximum average cadmium content. Between 20 ~ 50 cm of soil with pH > 7.5 (an increasing), cadmium content is significantly reduced, until reaching stability. Fig. 6A shows that the surface neutral soil inhibits the longitudinal migration of cadmium along the soil column; (2) Between 0 ~ 20 cm of soil with a pH value of 5.5 ~ 6.5, the soil is weakly acidic and the cadmium content reaches its maximum. Between 20 ~ 50 cm and 50 ~ 90 cm the soil is neutral, and at a depth of more than 90 cm it becomes alkaline. The cadmium content at 20 ~ 50 cm slowly decreases with increasing depth, showing initially a slower rate of curve variation compared to the first soil category, and then a faster rate, as depth continues to increase. Fig. 6B shows that the degree of vertical migration of cadmium is slightly higher than in the first soil category; (3) Between 0 ~ 20 cm of soil with a pH value of 4.5 ~5.5, the soil is acidic and the cadmium content is low. Between 20 ~ 50 cm and 50 ~ 90 cm the soil is weakly acidic: the average cadmium content in the first 20-50 cm layer reaches its maximum, while the average cadmium content in the 50-90 cm layer decreases, though remaining still higher than that in the surface soil. The soil at a depth > 90 cm is neutral, and the content of cadmium continues to decrease with increasing depth (Fig. 6C), indicating that the adsorption capacity in the surface acidic soil is poor, and that the longitudinal migration capacity of cadmium increases significantly with depth.

In general, the surface soil is neutral and alkaline, which can inhibit the longitudinal migration of cadmium. This is because in a weakly alkaline environment with a high pH, an increase in soil pH will determine the transition of cadmium into its carbonate state and Fe-Mn oxidation state, forming

precipitation complexes, increasing the negative charge, and therefore reducing the mobility of all heavy metals (Chen et al. 2020). In addition, the pH increase causes the OH⁻ in the solution to react with the groups on the surface of soil particles, resulting in the accumulation of negative charges; and the smaller the competition of Cd²⁺ for the adsorption sites, the stronger the Cd²⁺ adsorption capacity of the soil, and the faster the adsorption rate (Wang et al. 2006). In contrast, in a weakly acidic environment the cadmium adsorption in surface soil is limited, because the existing competitive adsorption of H⁺ and Cd²⁺ favors H⁺, as H⁺ ions can replace Cd²⁺ ions on the surface of the clay. This is due to the strong combination of H⁺ ions with oxygen atoms at the edge of the crystal (Inyang et al. 2016). In an acidic environment, Cd²⁺ could be effectively desorbed, and the lower the pH value is, the higher the reversibility of the adsorption process, as it is more difficult for Cd²⁺ to be adsorbed on the surface of soil particles (Cerqueira et al. 2011). However, this will increase the extent of the longitudinal migration, allowing cadmium ions to migrate into deep soil layers and pollute phreatic aquifers.

3.5.2 Soil mechanical composition

Based on the soil type, the deep profile soil in the Nansha area can be divided into two categories: (1) Continental sediment, at depths of 0 ~ 20 cm, 20 ~ 50 cm, 50 ~ 90 cm, where the soil type is silty clay; here the clay content in the soil is higher, accounting for 30% ~ 44.2%, and the soil adsorption capacity is strong. The clay content decreases slightly from top to bottom, and cadmium content at each depth is 0.39, 0.28, 0.22 mg/kg, respectively; (2) Marine sediment, at depths of 90-140 and 140-200 cm, where the soil type is mainly silty sand, with low clay content, weak soil adsorption capacity, and cadmium content at each depth ranging between 0.23-0.52 and 0.2-0.45 mg/kg, respectively (Fig. 7). There was no significant positive correlation between clay content and cadmium adsorption capacity in both the terrestrial and marine sedimentary soils. However, in general, the longitudinal variation of cadmium content correlates well with the change of sediment grain size. In particular, the cadmium content in coarse-grained sediments prevalent in sandy soils is low, while the element is enriched in fine-grained sediments which mainly characterize cohesive soils. This conclusion is consistent with the research by Covelo et al. (2004) (Covelo et al. 2004), which showed that soils with the highest heavy metal adsorption capacity also contained the highest contents of SOM, oxides and clays. The specific surface area of soil clay particles is large, and the surface is usually negatively charged, resulting in a high cation exchange capacity. The positively charged cations will adsorb on the surface of clay minerals under the action of electrostatic force. Therefore, it is believed that the higher the content of physical clay particles in the soil, the stronger the ability to adsorb Cd²⁺.

Basic chemical properties of the deep soil measured in the study area, such as pH, CEC, SOM and others, were correlated with the content of clay, silt and sand in soil mechanical composition analysis. The proportion of clay was negatively correlated with soil pH (correlation coefficient = -0.684), and was strongly correlated with CEC and SOM (correlation coefficients were 0.893 and 0.776, respectively). The

silt proportion was negatively correlated with soil pH (correlation coefficient = -0.647), and was strongly correlated with CEC and SOM (correlation coefficients were 0.867 and 0.861 , respectively). The mechanical composition of soil at the sea-land transition zone affects its basic chemical properties, and consequently it affects the migration and transformation of cadmium.

3.5.3 Soil organic matter

SOM is an important component of soil. The SOM content at $0\text{--}20$, $20\text{--}50$ and $50\text{--}90$ cm depths was similar, with mean values of 19.2 , 18.4 and 19.3 mg/kg, respectively. Similarly, the mean SOM content value at $90\text{--}140$ and $140\text{--}200$ cm depths was 15.9 and 13.6 mg/kg, respectively. As shown in Fig. 5B, cadmium concentrations generally declined with the decrease of SOM content. Ren et al.(2020) (Ren et al. 2020) pointed out that after the removal of SOM, the cadmium adsorption capacity generally decreased. SOM has a large surface area and abundant oxygen-containing functional groups (carboxyl, phenolic, hydroxyl, etc.), which produce organic ligands through decomposition (organic acids and humus, etc.). A variety of organic acids (oxalic, citric, formic, acetic, malic, succinic, lactic, and fumaric acids) were observed to complexate with Cd^{2+} in the soil through ion exchange mechanisms, and to form a complex with a stronger stability than the exchange reaction(Xiao et al. 2015), which may also reduce the mobility of heavy metals through precipitation(Bashir et al. 2019). As a result, the content of exchangeable and soluble cadmium in the soil can be effectively reduced, thus limiting the bioavailability of this element and the uptake potential of plants(Vranova et al. 2013). Therefore, the higher the SOM content is, the greater the adsorption capacity of the soil(Bashir et al. 2019).

4. Conclusions

Soils in the Nansha area of the Pearl River Delta are seriously polluted, and cadmium concentration in the two sedimentary environments analyzed is significantly higher than in Guangdong Province (0.056mg/kg) and the greater Pearl River Delta area (0.11mg/kg). The marine sedimentary soil is alkaline and it has a greater colloidal surface negative charge, which enhances its electrostatic action, resulting in the enhancement of specific metal ion adsorption. SOM content in the terrestrial sedimentary soil is high, so organic and inorganic colloids combine to form new colloid compounds during the adsorption process, which weakens the adsorption capacity of soil particles to heavy metals. The adsorption characteristics of cadmium in the two soils can be described by the Henry, Langmuir and Freundlich isotherm models. According to the fitting square variance, the Freundlich equation shows a better fitting performance. The fitting results show that the isothermal cadmium adsorption constant K_F in the marine sedimentary soil is significantly higher than that in the terrestrial sedimentary soil, measuring 558.21 and 363.04 , respectively, and indicating that the Cd^{2+} adsorption capacity of the marine soil is stronger. The continental sedimentary soil, however, presents a high content of clay—the clay specific surface area is large—and is usually negatively charged on the surface. This type of soil has the very high cation exchange capacity and, under the action of electrostatic force, positively charged cations can determine adsorption to the surface of clay minerals. As a result, the saturation adsorption of cadmium was higher

in the terrestrial (2419.83 mg/kg) than in the marine sedimentary soil (2228.25 mg/kg). In addition, it was observed that when the surface soil in the study area was neutral and alkaline, the vertical migration of cadmium was inhibited; when it was weakly acidic, the cadmium adsorption capacity was limited and when the topsoil was acidic, Cd^{2+} was effectively desorbed. In summary, the lower the pH value is, the higher the reversibility of the adsorption process, and the more difficult it is for Cd^{2+} to be adsorbed to the surface of soil particles. This favors the longitudinal migration of cadmium through the soil column allowing cadmium ions to reach the deep soil and pollute phreatic aquifers. Overall, the results of this study show that the distribution of cadmium in terrestrial and marine soils in the Nansha area of the Pearl River Delta is affected by adsorption and pH values.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request. All data generated or analysed during this study are included in this published article [and its supplementary information files].

Competing interests

The authors declare that they have no competing interests.

Funding

The work was supported by the Guangzhou Multi-factor Urban Geological Survey Project of Wuhan Geological Survey Center of China Geological Survey (DD20190291), National Key Research & Development Program of China (2018YFC1800800), Natural Science Foundation of China (No.51979252).

Authors' contributions

Fangting Wang: Roles/Writing - original draft, Investigation, Formal analysis, Project administration. Ke Bao: Investigation, Roles/Writing - original draft. Changsheng Huang: Conceptualization, Funding acquisition, Writing - review & editing. Wenjing Han: Data curation, Visualization, Validation. Zhibin Yin: Software, Methodology, Validation. Zhihua Chen: Conceptualization, Supervision, Resources. Xinwen Zhao: Funding acquisition, Project administration, Supervision. All authors read and approved the final manuscript.

References

1121.3-2006 NT Soil testing part 3: method for determination of soil mechanical composition

1121.6-2006 NT Soil testing part 6: method for determination of soil organic matter

2016 DT- Analysis methods for regional geochemical sample-part 1: determination of 24 components including aluminum oxide etc. by pressed power pellets-X-ray fluorescence spectrometry

14506.30-2010 GT Methods for chemical analysis of silicate rocks - Part 30 : Determination of 44 elements

Arabi S, Sohrabi MR (2014): Removal of methylene blue, a basic dye, from aqueous solutions using nano-zerovalent iron. *Water Sci Technol* 70, 24-31

Ardestani MM, van Gestel CAM (2016): Sorption and pH determine the long-term partitioning of cadmium in natural soils. *Environmental Science and Pollution Research* 23, 18492-18501

Bai JH, Xiao R, Zhang KJ, Gao HF (2012): Arsenic and heavy metal pollution in wetland soils from tidal freshwater and salt marshes before and after the flow-sediment regulation regime in the Yellow River Delta, China. *J Hydrol* 450, 244-253

Bai JH, Ye XF, Jia J, Zhang GL, Zhao QQ, Cui BS, Liu XH (2017): Phosphorus sorption-desorption and effects of temperature, pH and salinity on phosphorus sorption in marsh soils from coastal wetlands with different flooding conditions. *Chemosphere* 188, 677-688

Barrow N, J. (1984): Modeling the Effects Of Ph on Phosphate Sorption by Soils. *J Soil Sci* 35, 283-297

Bashir S, Adeel M, Gulshan A, B., Iqbal J, Khan S, Rehman M, Azeem M (2019): Effects of Organic and Inorganic Passivators on the Immobilization of Cadmium in Contaminated Soils: A Review. *Environ Eng Sci*

Center CNEM (1990): Chinese Soil Element Background Value Beijing

Cerqueira B, Covelo EF, Andrade L, Vega FA (2011): The influence of soil properties on the individual and competitive sorption and desorption of Cu and Cd. *Geoderma* 162, 20-26

Chen D, Wang XB, Wang XL, Feng K, Su JC, Dong JN (2020): The mechanism of cadmium sorption by sulphur-modified wheat straw biochar and its application cadmium-contaminated soil. *Sci Total Environ* 714

Covelo EF, Andrade ML, Vega FA (2004): Heavy metal adsorption by humic umbrisols: selectivity sequences and competitive sorption kinetics. *J Colloid Interf Sci* 280, 1-8

Davis J, A., Leckie J, O. (1978): Surface ionization and complexation at the oxide/water interface II. Surface properties of amorphous iron oxyhydroxide and adsorption of metal ions. *J Colloid Interf Sci* 67, 90-107

DB44/T1415-2014 Risk Screening Values for soil heavy metal The Pearl River Delta Area

Diagboya PN, Olu-Owolabi BI, Adebowale KO (2015): Effects of time, soil organic matter, and iron oxides on the relative retention and redistribution of lead, cadmium, and copper on soils. *Environmental Science And Pollution Research* 22, 10331-10339

Dong R, Z., Jia Z, M., Li S, Y. (2018): Risk assessment and sources identification of soil heavy metals in a typical county of Chongqing Municipality, Southwest China. *Process Saf Environ* 113, 275-281

Dou L, Du HY, Huang YH, Lai QH, You YH (2015): Main research achievements of agro-geological and eco-geochemical research in pearl river delta economic zone, Guangdong province, China. *Geological Survey of China* 2, 47-55

Du HY, Lai QH, Lin JP (2011): *Multi-objective Geochemical Atlas of the People's Republic of China (Pearl River Delta Economic Zone, Guangdong Province)*. Geological Publishing House, Beijing

Duan Y, Wang BG, Wang HM, Jiao TL, Qin YX, Ci RZM (2020): Adsorption characteristics of Cd in alluvial and lacustrine soils - a case study in Dangtu county, Anhui. *Earth Science*, 1-21

Elbana TA, Selim HM (2010): Cadmium Transport in Alkaline and Acidic Soils: Miscible Displacement Experiments. *Soil Sci Soc Am J* 74, 1956-1966

Feng L, Yan HL, Dai CH, Xu WX, Gu FY, Zhang F, Li T, Xian JY, He XQ, Yu YJ, Ma M, Wang F, He ZY (2020): The systematic exploration of cadmium-accumulation characteristics of maize kernel in acidic soil with different pollution levels in China. *Sci Total Environ* 729

Furuya M, Hashimoto Y, Yamaguchi N (2016): Time-Course Changes in Speciation and Solubility of Cadmium in Reduced and Oxidized Paddy Soils. *Soil Sci Soc Am J* 80, 870-877

- Godt J, Scheidig F, Grosse-Siestrup C, Esche V, Brandenburg P, Reich A, Groneberg DA (2006): The toxicity of cadmium and resulting hazards for human health. *J Occup Med Toxicol* 1
- Hamid Y, Tang L, Hussain B, Usman M, Lin Q, Rashid MS, He ZL, Yang XE (2020): Organic soil additives for the remediation of cadmium contaminated soils and their impact on the soil-plant system: A review. *Sci Total Environ* 707
- Houng KH, Lee DY (1998): Comparisons of linear and nonlinear Langmuir and Freundlich curve-fit in the study of Cu, Cd, and Pb adsorption on Taiwan soils. *Soil Sci* 163, 115-121
- Huang B, Li ZW, Huang JQ, Chen GQ, Nie XD, Ma WM, Yao HB, Zhen JM, Zeng GM (2015): Aging effect on the leaching behavior of heavy metals (Cu, Zn, and Cd) in red paddy soil. *Environmental Science and Pollution Research* 22, 11467-11477
- Huang JH, Wang SL, Lin JH, Chen YM, Wang MK (2013): Dynamics of cadmium concentration in contaminated rice paddy soils with submerging time. *Paddy Water Environ* 11, 483-491
- Huang MJ, Sun HR, Liu HT, Wang XM, Wang BM, Zheng D (2018): Atmospheric Arsenic Deposition in the Pearl River Delta Region, South China: Influencing Factors and Speciation. *Environ Sci Technol* 52, 2506-2516
- Huang S, Zhang RD, Zhang JY, Pan R (2012): Effects of physical and chemical characteristics of soils on the adsorption of Cd. *Journal of Irrigation and Drainage* 31, 19-22
- Inyang HI, Onwawoma A, Bae S (2016): The Elovich equation as a predictor of lead and cadmium sorption rates on contaminant barrier minerals. *Soil Till Res* 155, 124-132
- Lai QH, Du HY, Fang JW, Shao LB, Xia B (2005): Source and cause of regional Cd enrichment in alluvial soils in the pearl river delta plain. *Journal of Agro- environment Science*, 746-750
- Li P, Lang M, Wang XX, Zhang TL (2016): Sorption and Desorption of Copper and Cadmium in a Contaminated Soil Affected by Soil Amendments. *Clean-Soil Air Water* 44, 1547-1556
- Li QQ, Wang CQ, Dai TF, Shi WJ, Zhang X, Xiao Y, Song WP, Li B, Wang YD (2017): Prediction of soil cadmium distribution across a typical area of Chengdu Plain, China. *Sci Rep-Uk* 7
- Li TQ, Jiang H, Yang X, He ZL (2013): Competitive sorption and desorption of cadmium and lead in paddy soils of eastern China. *Environ Earth Sci* 68, 1599-1607
- Li Y, Zhang HB, Chen XB, Tu C, Luo YM, Christie P (2014): Distribution of heavy metals in soils of the Yellow River Delta: concentrations in different soil horizons and source identification. *J Soil Sediment* 14, 1158-1168
- Lin Q, Xu SH (2008): A review on competitive adsorption of heavy metals in soils. *Soils*, 706-711

- Liu JA 2019: Evaluating submarine groundwater discharge and its effects on eco-environment in estuarine and coastal waters by radium isotopes 镭 Thesis, East China Normal University
- Peng C, Wang M, Chen WP (2016): Modelling cadmium contamination in paddy soils under long-term remediation measures: Model development and stochastic simulations. *Environ Pollut* 216, 146-155
- R R, Vvss S (2015): Submarine groundwater discharge and nutrient addition to the coastal zone of the Godavari estuary. *Mar Chem* 172, 57-69
- Ren BZ, Wu Y, Deng DP, Tang XF, Li HT (2020): Effect of multiple factors on the adsorption of Cd in an alluvial soil from Xiba, China. *J Contam Hydrol* 232
- Sauve S, Manna S, Turmel MC, Roy AG, Courchesne F (2003): Solid-solution partitioning of Cd, Cu, Ni, Pb, and Zn in the organic horizons of a forest soil. *Environ Sci Technol* 37, 5191-5196
- Shen BB, Wang XM, Zhang Y, Zhang M, Wang K, Xie P, Ji HB (2020): The optimum pH and Eh for simultaneously minimizing bioavailable cadmium and arsenic contents in soils under the organic fertilizer application. *Sci Total Environ* 711
- Tabelin C, B., Igarashi T, Villacorte-Tabelin M, Park I, Opiso E, M., Ito M, Hiroyoshi N (2018): Arsenic, selenium, boron, lead, cadmium, copper, and zinc in naturally contaminated rocks: A review of their sources, modes of enrichment, mechanisms of release, and mitigation strategies. *Sci Total Environ* 645, 1522-1553
- Tang ZM 2017: Effect of Pearl River system on environmental quality of alluvial plain soil 镭 Thesis, China University of Geosciences
- Vranova V, Rejsek K, Formanek P (2013): Aliphatic, Cyclic, and Aromatic Organic Acids, Vitamins, and Carbohydrates in Soil: A Review. *Sci World J*
- Wang FT, Huang CS, Chen ZH, Bao K (2019): Distribution, Ecological Risk Assessment, and Bioavailability of Cadmium in Soil from Nansha, Pearl River Delta, China. *Int J Env Res Pub He* 16
- Wang YJ, Zhou DM, Sun RJ, Hao XZ, Cang L (2006): Competitive adsorption kinetics of copper and lead ions in soils. *Zhongguo Huanjing Kexue/China Environmental Science* 26, 555-559
- Wang Z, Zhang L, Dang XL, Zhang YL (2012): Effect of the freezing-thawing on kinetics of adsorption-desorption of the soil cadmium. *Acta Scientiae Circumstantiae* 32, 721-725
- Xiao H, Bottcher J, Utermann J (2015): Evaluation of field-scale variability of heavy metal sorption in soils by scale factors - Scaling approach and statistical analysis. *Geoderma* 241, 115-125
- Yao Q 2014: Climatic indicators and sedimentary environment studies inferred from transgressive and regressive sediments of core LZ908-south Bohai Sea 镭 Thesis, The institute of oceanology, Chinese

academy of sciences

Zeng HY, Chen LH, Zhou XH, Zeng QR (2019): Cadmium accumulation in winter crops and the assessment of paddy soil phytoremediation in southern China. *Environmental Science And Pollution Research* 26, 17173-17182

Zhang H, Reynolds M (2019): Cadmium exposure in living organisms: A short review. *Sci Total Environ* 678, 761-767

Zhang XY, Zhong TY, Liu L, Ouyang XY (2015): Impact of Soil Heavy Metal Pollution on Food Safety in China. *Plos One* 10

Zhang Y, Liu LP, Wen H, Wu C, Zhang YH (2018): Evaluation of the Polarimetric-Radar Quantitative Precipitation Estimates of an Extremely Heavy Rainfall Event and Nine Common Rainfall Events in Guangzhou. *Atmosphere-Basel* 9

Zhou T, Wu LH, Luo YM, Christie P (2018): Effects of organic matter fraction and compositional changes on distribution of cadmium and zinc in long-term polluted paddy soils. *Environ Pollut* 232, 514-522

Zhu HH, Chen C, Xu C, Zhu QH, Huang DY (2016): Effects of soil acidification and liming on the phytoavailability of cadmium in paddy soils of central subtropical China. *Environ Pollut* 219, 99-106

Figures




Figure 1

The location (A), terrain (B), and deep profile soil sampling maps (C) of the study area. Note: The designations employed and the presentation of the material on this map do not imply the expression of any opinion whatsoever on the part of Research Square concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. This map has been provided by the authors.




Figure 2

Changes in adsorption capacity S (A) and adsorption rate η (B) with the initial concentration of solution C_0 in the two soils (the cadmium adsorption isothermal curves for the two soils are shown in C, Henry equation; D, Freundlich equation; and E, Langmiur equation) and relationship between the solid/liquid distribution coefficient K_d and the initial concentration of cadmium C_0 in two soils.




Figure 3

The oscillation time t affected the cadmium adsorption capacity S and adsorption rate V of the two soils. Elvich equation fitting (C), biconstant equation fitting (D) and similar exponential asymptote equation fitting (E) for cadmium adsorption in the two soil types.





Figure 4

Relationship between oscillation time and cadmium adsorption capacity in soil 1 and soil 2 under different pH values.






Figure 6

Comparison of three trends of cadmium content and pH in the deep soil profile (the error bar represents the standard deviation)




Figure 7

Changes in soil mechanical composition and cadmium content with depth in the deep profile soil (0-200 cm)