Immunological repertoire linked to PSTPIP1-associated myeloid-related inflammatory (PAMI) syndrome

Leonardo Oliveira Mendonca (✉ leonardo.oliveira.mendonca@gmail.com)

Universidade de Sao Paulo <https://orcid.org/0000-0001-7533-1937>

Maria Teresa Terreri

Federal University of Sao Paulo: Universidade Federal de Sao Paulo

Fabiane Mitie Osaku

Hospital Infantil Rosa de Gusmão

Samar Freschi Barros

University of Sao Paulo: Universidade de Sao Paulo

Karen Francine Köhler

University of Sao Paulo: Universidade de Sao Paulo

Alex Isidoro Prado

University of Sao Paulo: Universidade de Sao Paulo

Myrthes Toledo Barros

University of Sao Paulo: Universidade de Sao Paulo

Jorge Kalil

University of Sao Paulo: Universidade de Sao Paulo

Fabio Fernandes Morato Castro

University of Sao Paulo: Universidade de Sao Paulo

Research Article

Keywords: PSTPIP1, PAPA syndrome, PAMI syndrome, E250K mutation

Posted Date: March 11th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-270780/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Pediatric Rheumatology on August 16th, 2021. See the published version at <https://doi.org/10.1186/s12969-021-00617-y>.

Abstract

Background: Mutations along PSTPIP1 gene are associated to two specific conditions, PAPA syndrome and PAMI syndrome, both autoinflammatory disorders associated to disturbances in cytoskeleton formation. Immunological aspects of PAMI syndrome has not yet been reported neither the clinical impact on therapeutical decisions.

Methods: Clinical data of patients records were retrospectively accessed. Genomic DNA were extracted and sequenced following standard procedures. Peripheral lymphocytes were quantified in T, B e FOXP3 phenotypes.

Results: We describe two related patients with PAMI syndrome harboring the usual E250K mutation. Anti-IL1 therapy could partially control the disease in the index patient. A broad spectrum of immunological effects as well as an aberrant expression of FOXP3 could be observed.

Conclusions. Here we report two related brazilian patients with PAMI syndromes harboring the E250K mutation in PSTPIP1, their immunological aspects and the therapeutical response to canakinumab.

Introduction

Proline-serine-threonine phosphatase-interacting protein 1 (PSTPIP1, also known as CD2BP1) is a cytoskeleton-associated adaptor protein that modulates T-cell activation, cytoskeletal organization and IL-1 β release (1). Mutations along the PSTPIP1 gene have been classically associated with PAPA syndrome (OMIM #604410), an autosomal dominant disorder (2). However, specific heterozygous mutations (E250k and E257K) in this same gene were recently linked to another condition, namely PSTPIP1-associated myeloid-related inflammatory (PAMI) syndrome (3).

In addition to its known involvement in cytoskeletal organization, interaction between PSTPIP1 and the WASP protein lead to defective podosome formation and interfere with macrophage migration (4). Additionally, gene MEFV, in which mutations have been associated with Familial Mediterranean fever (OMIM #249100), encodes the protein pyrin, which can alter the distribution of PSTPIP1 throughout the filamentous network (5). Accordingly, PSTPIP1-related diseases are considered cytoskeleton-associated disorders.

PAMI syndrome is classified as a rare autoinflammatory condition clinically characterized by the early onset of multisystemic inflammation. Affected patients present with recurrent episodes of fever, osteoarticular symptoms, skin lesions, diffuse reticula-endothelial enlargement, anemia and thrombocytopenia. These symptoms are usually associated with high levels of acute reactant markers (CRP, ESR and SAA); hyperzincemia is also a consistent finding. At the time of this publication, just 20 cases had been reported in the literature (6).

Here we report two related Brazilian patients with PAMI syndrome, both harboring the E250K mutation, one of whom was treated with canakinumab (Ilaris). We further discuss our observations of wide-ranging immunological aspects regarding T, B and regulatory T cells.

Methods

Clinical Data and Genomic Sequencing

Clinical data were retrieved from patient records after parents provided a signed term of consent. Genomic DNA was extracted from blood samples using a QIAamp® DNA Blood Maxi Kit (Qiagen®, Valencia, CA, USA) and peripheral blood mononuclear cells (PBMC) were obtained by density gradient centrifugation ($d=1.077$ g/ml), then cryopreserved for later use. Exome sequencing was performed using a commercially available kit (Medelics Genomiks) and primers were designed to directly target exon 11 of the PSTPIP1 gene. Sanger sequencing was performed for genetic confirmation and familial segregation following standard procedures.

Quantification of lymphocytic B, T and FOXP3 phenotypes by flow cytometry

Peripheral blood mononuclear cells (PBMC) were stained with titrated mouse anti-human monoclonal antibodies (mAbs) (all from BD Biosciences), as following: anti-CD3 FITC or PerCp, anti- CD4 AmCyan or Pacific Blue, anti-CD8 Pacific Blue or APCCy7 or PE-Cy7, anti-CD45RA PE-Cy7 or APC, anti-CCR7 (CD197) PerCp-Cy5.5, anti-CD27 APC or FITC, anti- CD16 PE, anti- CD56 PE, anti- CD19 PerCp, anti- CD20 PE-Cy7, anti- HLA-DR APC, anti- CD45 AmCyan, anti-CD62L PE, anti- TCR $\alpha\beta$ FITC, anti- TCR $\gamma\delta$ PE and anti-BB220 APC, according to manufacturer instructions. For Treg/CD28null panel, we stained the cell surface with anti-CD3 FITC, anti-CD4 AmCyan, anti-CD8 Pacific Blue, anti-CD25 PE-Cy7, anti-CD127 PE and CD28 APC. For intracellular staining of FOXP3 (PerCp-Cy5.5) cells were washed, fixed and permeabilized with Transcription Factor Buffer Set (BD Biosciences) immediately after surface staining, in accordance with the manufacturer's instructions. Fluorescence minus one (FMO) control were set up for CD45RA, CD127 and Foxp3 markers.

Flow cytometry was performed in FACSCanto II (BD Biosciences) and the analyses were made in FlowJo 9.9.5 software (TreeStar Inc, San Carlos, CA, USA). After exclusion of cell doublets, sequential gating of PBMC was performed in the lymphocyte region. For T lymphocytes, after gate of CD3⁺ T cells, followed by discrimination of CD8⁺ and CD4⁺ markers, we analyzed the T lymphocyte naïve/memory subpopulations through boolean gates: T naïve (CD45⁺CCR7⁺CD27⁺), T central memory (TCM) (CD45⁻CCR7⁺CD27⁺), T effector memory (TEM) (CD45⁻CCR7⁻CD27⁺) and T effector memory with RA re-expression (TEMRA) (CD45⁺CCR7⁻CD27⁻). To analyze the phenotypes CD3⁺ TCR $\alpha\beta$ CD4⁻ CD8⁻, CD3⁺ HLA-DR⁺ and CD3⁺ B220⁺, the cells were gated on CD3⁺ region, after exclusion of doublets and death cells. The phenotypes CD27⁺ was analyzed in B lymphocyte region (CD20⁺ cells). Lymphocyte subsets absolute counts were calculated using the percentages obtained in flow cytometry. The subset percentages analyzed were referred to total lymphocyte counts for T and B cells.

For Treg panel, we analyzed the following subpopulations within the CD4+ subgroup: TREG (CD25^{high}, CD127-, FOXP3+), TCD4/CD25+ (CD25+, CD127-, FOXP3+), TCD4 FOXP3+, TCD4 FOXP3+ CD28^{null} and TCD4 CD28^{null}. Within the CD8+ subgroup, we analyzed TCD8 FOXP3+, TCD8 FOXP3+ CD28^{null} and TCD8 CD28^{null}. The T cell subsets analyzed are expressed as percentages, referring to the total number of viable TCD4 and TCD8 lymphocytes.

T and B cell population results were compared to reference ranges from Brazilian subjects published in the literature. Treg cell values were compared to a patient harboring a TRNT1 mutation.

Results

Case Report

A 4-year-old girl in southern Brazil, born to non-consanguineous parents of German-Italian descent, came to our attention due to the suspicion of an inborn error of immunity. Her parents reported high fever in association with diarrhea, paleness and mucosal bleeding in her first month of life. She was first hospitalized at that time due to a severe and progressive deterioration in her clinical picture. The patient appeared severely dehydrated, with profoundly low levels of hemoglobin (6.6 mg/dL - reference range 11.5 - 15.5 g/dL), neutropenia (548 cells - reference range 2,500 - 7,500/mm³), thrombocytopenia (20,000 platelets - reference range 150,000-400,000) and high levels of CRP (57.6 mg/dL - reference range <0.5 mg/dL). The patient presented high cytomegalovirus viraemia in peripheral blood. Clinical remission was observed following prolonged intravenous administration of ganciclovir at standard doses.

Monthly episodes of fever, lasting from 2-3 days, were noted in association with intermittent and non-specific macular skin rash. Persistent splenomegaly and severe and refractory anemia, thrombocytopenia and neutropenia, as well as high levels of acute reactant markers (ESR, CRP and SAA) were consistent findings upon flares. Several hospitalizations ensued, entailing blood transfusions and the intravenous administration of broad-spectrum antibiotics to treat undefined systemic infections. During this period, a failure to thrive could also be observed. At the age of two years, an empiric dose of corticosteroids (1 mg/kg) was introduced, resulting in partial resolution of clinical manifestations. Steroids resulted in less frequent hospitalization, extended intervals between febrile episodes, and the amelioration of diarrhea, skin rash and arthritis. However, the patient presented failure to no less than two steroid sparing agents: IVIG and colchicine. Consequently, steroids were administered on demand during flares at a dose of 1mg/kg for 5-7 days. On two occasions serum zinc levels were evaluated in the index patient, returning results within normal range. Of note, the patient's mother also complained of recurrent fever and thrombocytopenia since early in life, yet remained asymptomatic throughout adulthood. Exome sequencing revealed a previously reported pathogenic mutation in PSTPIP1, c.748G>A, p.Glu250Lys, confirmed by Sanger sequencing. Segregation analysis confirmed that the mother harbored the same mutation (Figure 1). Therefore, anti-IL1 (canakinumab) at a dose of 2 mg/kg/month was initiated in the index patient and progressively titrated until reaching a dose of 4 mg/kg/month. Following the onset of anti-IL1 therapy, persistent clinical amelioration was observed in addition to the normalization of platelet

and hemoglobin levels, yet neutrophil counts remained low. While systemic inflammatory flares were still noted following anti-IL1B therapy, these were less severe and some resolved spontaneously (Figure 2). Her mother was not treated and is currently being regularly followed on an outpatient basis.

Peripheral lymphocyte repertoire

We observed high peripheral expression of T CD8 cells, low peripheral expression of T CD4 and normal B (CD19) cells, with normal levels of immunoglobulins A, G and M (Figure 1, Table 1). However, very low levels of CD8 and CD4 naïve and TCM cells were noted, together with normal levels of CD4 and CD8 TEM cells. In the B cell compartment, an expansion of B memory cells was observed with very low levels of B naïve cells. High levels of double-negative T cells and B220 cells were also noted (Table 1). Compared to another patient with SIFD syndrome, we identified higher FOXP3 expression in T CD4 cells, especially in cells dependent on CD127 (cytokine IL-7). We further observed high CD28^{null} expression in T-CD4 and T-CD8 cells, as well as FOXP3 expression dependent on CD28^{null} only in T-CD4 cells (Table 2). Conversely, high levels of FOXP3 in T-CD8 cells had been observed in the patient harboring the TRNT1 mutation.

Discussion

Here we describe a case of two related patients harboring the typical E250K mutation in PSTIIP1 who present classical signs of PAMI syndrome (1). While other patients typically present hyperzincemia and hyperprolactinemia, these were not seen in the young patient. This finding should be considered highly relevant, as patients lacking these characteristics must also be considered eligible for PSTIIP1 genetic screening (3,5).

As indicated by previous reports, no effective therapy exists for affected patients. Anakinra, an anti-IL1 receptor agonist, was demonstrated to partially control disease activity. The present report describes the first use of canakinumab, an anti-IL1B monoclonal antibody, in the treatment of this syndrome. Despite its significant control over the frequency and intensity of disease flares, canakinumab does not seem to be as thoroughly effective as anakinra in patients with PAMI syndrome. Bone marrow transplantation has been reported in just one patient refractory to a broad spectrum of immunosuppressants (5).

The immunological aspects of PAMI syndrome have not yet been reported. As F-actin is essential for immunological synapse (IS) formation (4,7), it follows that the strongly diminished actin formation in patients harboring pathogenic mutations in PSTIIP1 may lead to a wide range of T cell-related immune defects, as observed in the index patient (Table 1). Patients harboring other mutations along PSTIIP1 (e.g. R228C and T274M) who present a clinical phenotype of common variable immune deficiency without autoinflammation also exhibit similar T cell defects as those observed in our patient (7). Since PSTIIP1 is highly expressed in T cells rather than B cells, this may serve to explain the normal levels and functioning of B cells observed in our patient. Signs associated with T cell-related “immunodeficiency” may also explain the severe clinical course of CMV infection observed in the early life of our patient.

A subset of CD4⁺ T cells, Tregs are capable of limiting effector CD4⁺ T cells and suppressing immune-mediated inflammations. PSTPIP1 induces the activation of the so-called “pyroptosome,” which leads to cell death and the release of cytokines, such as IL-1 β , via the NLRP3 inflammasome that negatively regulates Treg cell differentiation. (1,8,9). Therefore, it is possible that the elevated FOXP3 expression (Table 1) in association with the high levels of double-negative T cells (Table 1) observed herein may be the result of non-canonical activation driven by PSTPIP1. Curiously, previous reports have described seemingly normal Treg (FOXP3) expression in patients with TRNT1 mutations (10). Our findings may shed light on a new pathway of FOXP3 activation mediated by PSTPIP1 in PAMI syndrome, which could entail therapeutic implications.

Conclusions

PAMI syndrome is a recently categorized autoinflammatory disorder clinically characterized by hematological involvement and recurrent episodes of autoinflammation. Here we describe two related patients with PAMI syndrome, harboring the same typical E205k mutation. We report for the first time a severe CMV infection as the first manifestation of PAMI syndrome. Our observations of T cell defects may help elucidate host predisposition to particular infections, such as cytomegalovirus here reported. The role of TReg cells in PAMI syndrome, as well the participation of PSTPIP1 in Treg cell activation and differentiation require further evaluation.

List Of Abbreviations

PSTPIP1 Proline-Serine-Threonine Phosphatase Interacting Protein 1

CD2BP1 CD2-binding protein 1

PAPA Pyogenic Arthritis, pyoderma gangrenous and acne

WASP Wiskott–Aldrich Syndrome **protein**

MEFV Mediterranean Fever

CRP C-reactive Protein

ESR erythrocyte sedimentation rate

SAA Serum Amyloid

FOXP3 Forkhead Box 3

CD Cluster Differentiation

TREG T-regulatory cells

TRNT1 tRNA-nucleotidyltransferase 1

IVIG Intravenous Immunoglobulin

Declarations

Ethical Approval and Consent for publication: Written informed consent for publication was obtained from all included in the study.

Availability of data and materials: The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Competing interests: LOM receives fees from Novartis Brazil as speaker

Funding: There was no funding for supporting this article

Acknowledgements We thank the family for allowing data collection.

Author's Contribution: Leonardo Oliveira Mendonça article draft and genetic analysis; Maria Teresa Terreri article revision and patient care ; Fabiane Mitie Osaku article revision and patient care ; Samar Freschi Barros genetic sequencing and article revision; Karen Francine Köhler lymphocytes quantification and article revision ; Alex Isidoro Prado article revision and patient care ; Myrthes Toledo Barros senior supervision and article revision ; Jorge Kalil senior supervision and critical article revision ; Fabio Fernandes Morato Castro senior supervision and critical article revision

Ethical Approval: The study is approved at the Hospital das Clínicas, University of São Paulo, School of Medicine. The number of study approval is CAAE 74305817.0.0000.0068.

References

- 1 - Janssen WJM, Grobarova V, Leleux J, Jongeneel L, van Gijn M, van Montfrans JM, Boes M. Proline-serine-threonine phosphatase interacting protein 1 (PSTPIP1) controls immune synapse stability in human T cells. *J Allergy Clin Immunol*. 2018 Dec;142(6):1947-1955. doi: 10.1016/j.jaci.2018.01.030. Epub 2018 Feb 9. PMID: 29432774.
- 2 - Almeida de Jesus A, Goldbach-Mansky R. Monogenic autoinflammatory diseases: concept and clinical manifestations. *Clin Immunol*. 2013 Jun;147(3):155-74. doi: 10.1016/j.clim.2013.03.016. Epub 2013 Apr 9. PMID: 23711932; PMCID: PMC3769788.
- 3 - Boursier G, Piram M, Rittore C, Sarabay G, Touitou I. Phenotypic Associations of PSTPIP1 Sequence Variants in PSTPIP1-Associated Autoinflammatory Diseases. *J Invest Dermatol*. 2020 Nov 17:S0022-202X(20)32168-0. doi: 10.1016/j.jid.2020.08.028. Epub ahead of print. PMID: 33218716.

4 - Cortesio CL, Wernimont SA, Kastner DL, Cooper KM, Huttenlocher A. Impaired podosome formation and invasive migration of macrophages from patients with a PSTPIP1 mutation and PAPA syndrome. *Arthritis Rheum.* 2010 Aug;62(8):2556-8. doi: 10.1002/art.27521. PMID: 20506269; PMCID: PMC2921034.

5 - Shoham NG, Centola M, Mansfield E, Hull KM, Wood G, Wise CA, Kastner DL. Pyrin binds the PSTPIP1/CD2BP1 protein, defining familial Mediterranean fever and PAPA syndrome as disorders in the same pathway. *Proc Natl Acad Sci U S A.* 2003 Nov 11;100(23):13501-6. doi: 10.1073/pnas.2135380100. Epub 2003 Oct 31. PMID: 14595024; PMCID: PMC263843.

6 - Mejbri M, Theodoropoulou K, Hofer M FRI0564 PSTPIP1- ASSOCIATED MYELOID-RELATED PROTEINEMIA INFLAMMATORY SYNDROME/PAMI SYNDROME: CASE REPORT AND REVIEW OF THE LITERATURE *Annals of the Rheumatic Diseases* 2019;78:977-978. doi.org/10.1136/annrheumdis-2019-eular.8133

7 - Janssen WJM, Grobarova V, Leleux J, Jongeneel L, van Gijn M, van Montfrans JM, Boes M. Proline-serine-threonine phosphatase interacting protein 1 (PSTPIP1) controls immune synapse stability in human T cells. *J Allergy Clin Immunol.* 2018 Dec;142(6):1947-1955. doi: 10.1016/j.jaci.2018.01.030. Epub 2018 Feb 9. PMID: 29432774.

8 - Lu L, Barbi J, Pan F. The regulation of immune tolerance by FOXP3. *Nat Rev Immunol.* 2017 Nov;17(11):703-717. doi: 10.1038/nri.2017.75. Epub 2017 Jul 31. PMID: 28757603; PMCID: PMC5793224.

9 - Park SH, Ham S, Lee A, Möller A, Kim TS. NLRP3 negatively regulates Treg differentiation through Kpna2-mediated nuclear translocation. *J Biol Chem.* 2019 Nov 22;294(47):17951-17961. doi: 10.1074/jbc.RA119.010545. Epub 2019 Oct 9. PMID: 31597697; PMCID: PMC6879348

10 - Yang L, Xue X, Zeng T, Chen X, Zhao Q, Tang X, Yang J, An Y, Zhao X. Novel biallelic *TRNT1* mutations lead to atypical SIFD and multiple immune defects. *Genes Dis.* 2020 Jan 23;7(1):128-137. doi: 10.1016/j.gendis.2020.01.005. Erratum in: *Genes Dis.* 2020 Apr 09;7(3):485. PMID: 32181284; PMCID: PMC7063413.

Tables

Due to technical limitations, Table 1 and Table 2 are only available as a download in the supplementary files section.

Figures

Figure 1 - Genetic sequencing and baseline laboratory analysis of patients harboring the E250K mutation in PSTIPI1 gene. In A) Family pedigree and electropherograms demonstrating the autosomal dominant fashion of inheritance. In B) Baseline laboratory analysis of the index patient and mother, evidencing anemia, neutropenia in both patients and thrombocytopenia in the index patient.

Figure 1

Genetic sequencing and baseline laboratory analysis of patients harboring the E250K mutation in gene PSTIPI1. A) Family pedigree and electropherograms demonstrating an autosomal dominant pattern of inheritance. B) Baseline laboratory analysis of the index patient and her mother, evidencing anemia, neutropenia in both patients and thrombocytopenia in the index patient.

Figure 2 - Laboratory analysis observed over 4 years in the index patient and response to anti-IL1 therapy . Systemic inflammation associated to marked hematological involvement can be observed with high levels of C-reactive protein (CRP), anemia, neutropenia and thrombocytopenia. Dashed lines indicates the moment anti-IL1 was started.

Figure 2

Four-year follow-up of laboratory parameters in the index patient, including response to anti-IL1B therapy. Systemic inflammation associated with marked hematological involvement is observed, with high levels of C-reactive protein (CRP), anemia, neutropenia and thrombocytopenia. Dashed lines indicate the onset of anti-IL1B (canakinumab).

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [TablestosubmissionJCI.001.jpeg](#)
- [TablestosubmissionJCI.002.jpeg](#)