

# Fluid Overload is Associated with Underestimation of Serum Creatinine in Cardiac Surgery Patients: A Retrospective Analysis

**Jifu Jin**

Zhongshan Hospital, Fudan University

**Jiarui Xu**

Zhongshan Hospital, Fudan University

**Sujuan Xu**

Zhongshan Hospital, Fudan University

**Jiachang Hu**

Zhongshan Hospital, Fudan University

**Wuhua Jiang**

Zhongshan Hospital, Fudan University

**Bo Shen**

Zhongshan Hospital, Fudan University

**Chunsheng Wang**

Zhongshan Hospital, Fudan University

**Jie Teng**

Zhongshan Hospital, Fudan University

**Xiaoqiang Ding** (✉ [dingxiaoqiang2018@163.com](mailto:dingxiaoqiang2018@163.com))

Zhongshan Hospital <https://orcid.org/0000-0002-0169-2729>

---

## Research article

**Keywords:** Acute kidney injury, Fluid balance, Cardiac surgery, Diagnosis, Prognosis

**Posted Date:** May 11th, 2020

**DOI:** <https://doi.org/10.21203/rs.3.rs-27199/v1>

**License:**   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

**Version of Record:** A version of this preprint was published on January 31st, 2021. See the published version at <https://doi.org/10.1186/s12872-021-01879-w>.

# Abstract

**Background:** Fluid overload is related to the development and prognosis of cardiac surgery-associated acute kidney injury (CSA-AKI). The study is to investigate the influence of serum creatinine (SCr) corrected by fluid balance on the prognosis of patients with cardiac surgery.

**Methods:** A retrospective study was conducted in 1334 patients who underwent elective cardiac surgery from January 1 to December 31, 2015. Kidney Disease: Improving Global Outcomes (KDIGO) criteria for AKI were applied to identify CSA-AKI. SCr was measured every 24 hours during ICU period and was accordingly adjusted for cumulative fluid balance. Change in SCr, defined as  $\Delta\text{Crea}$ , was determined by difference between before and after adjustment for cumulative fluid balance. All patients were divided as three groups: underestimation group ( $\Delta\text{Crea} \geq P 75$ ), normal group ( $P 25 < \Delta\text{Crea} < P 75$ ) and overestimation group ( $\Delta\text{Crea} \leq P 25$ ).

**Results:** The incidence of AKI increased from 29.5% to 31.8% after adjustment for fluid balance. Patients in underestimation group showed prolonged length of ICU stay compared with normal group and overestimation group (3.2[1.0-3.0] vs 2.1[1.0-3.0] d,  $P < 0.001$ ; 3.2[1.0-3.0] vs 2.3[1.0-3.0] d,  $P < 0.001$ ). Length of hospital stay and mechanical ventilation dependent days in underestimation group were significantly longer than normal group ( $P < 0.001$ ). Multivariate analysis showed age, baseline SCr and left ventricular ejection fraction were independently associated with underestimation of creatinine.

**Conclusions:** Cumulative fluid balance after cardiac surgery disturbs accurate measurement of serum creatinine. Patients with underestimation of SCr were associated with poor prognosis.

## Background

Acute kidney injury (AKI) develops in 17–30% of patients after cardiac surgery and is associated with an excessive mortality rate up to 60%-80% [1–4]. As the first-line treatment of critically ill patients, fluid resuscitation may cause positive fluid balance during treatment, which frequently results in a relative increase in body weight of 10%-15% in a short time [5, 6]. However, recent studies have illustrated that positive fluid balance was associated with worse outcome in critically ill patients with AKI [7–10]. Furthermore, positive fluid balance was associated with recognition, staging and outcome of AKI in patients with acute respiratory distress syndrome or undergoing cardiac surgery [11–14]. Fluid overload can dilute serum creatinine level, which may induce delayed diagnosis of AKI and incorrect estimation of glomerular filtration rate (GFR) [15]. Therefore, aim of our study is to investigate the influence of discrepancy of serum creatinine on the prognosis of patients with cardiac surgery, and moreover, to explore underlying risk factors for underestimation of serum creatinine.

## Methods

### 1. Patients

Patients who underwent cardiac surgery between January 1 and December 31, 2015 at the Department of Cardiovascular Surgery, Zhongshan Hospital, Fudan University were consecutively included in our study. Inclusion criteria were adult patients ( $\geq 18$  years) who received elective cardiac surgery with or without cardiopulmonary bypass (CPB). Exclusion criteria were patients who had preexisting renal dysfunction requiring renal replacement therapy or had a baseline creatinine  $\geq 4$  mg/dl; patients who died within 24 hours after surgery as well as patients who received cardiac transplant or aortic aneurysm surgery. The Institutional Ethics Committee of the Zhongshan Hospital (B2018-175) granted permission for data collection and informed consent was waived due to the retrospective design of the study.

## 2. Data collection

Acute Physiology and Chronic Health Evaluation II (APACHE II) score was used to assess severity of illness at intensive care unit (ICU) admission [16]. Intraoperative parameters, such as CPB and cross-clamp duration, type of surgery and ultrafiltration during extracorporeal circulation were recorded as well. Fluid input and output were obtained at least 24 hours since ICU admission or until discharge from the ICU, whichever occurred first. Insensitive loss of fluid was not taken into account in our study.

## 3. Definitions

Kidney Disease: Improving Global Outcomes (KDIGO) definition was utilized to identify cardiac surgery-associated acute kidney injury (CSA-AKI) [17]. The dilutional effect on the value of serum creatinine used to diagnose and staging AKI according to the KDIGO criteria was assessed.

Cumulative fluid balance was calculated based on total fluid input and output in every 24 hours. Patients' admission weights were utilized to estimate baseline total body water (TBW).  $TBW = 60\%$  of baseline weight in kilogram at admission [15].

Baseline serum creatinine was obtained immediately prior to surgery. Maximum serum creatinine was the highest value within 48 hours after cardiac surgery. During the postoperative ICU period, serum creatinine was measured at least once every 24 hours. The adjusted serum creatinine was calculated using formula utilized in previous studies [11, 15, 18].  $\text{Adjusted creatinine} = \text{serum creatinine} \times [1 + (\text{cumulative fluid balance in L/admission weight in kg} \times 0.6)]$ . The difference of adjusted and measured serum creatinine was defined as  $\Delta\text{Crea}$ .

## 4. Groups

Given that the normal distribution of  $\Delta\text{Crea}$  (Fig. 1), patients were divided as three groups: underestimation group ( $\Delta\text{Crea} \geq P_{75}$ ), normal group ( $P_{25} < \Delta\text{Crea} < P_{75}$ ) and overestimation group ( $\Delta\text{Crea} \leq P_{25}$ ).  $P_{25}$  and  $P_{75}$  are used to indicate the 25<sup>th</sup> and the 75<sup>th</sup> percentile value of the  $\Delta\text{Crea}$ , respectively. The primary outcome is in-hospital mortality rate and its relationship to the  $\Delta\text{Crea}$ . The

secondary outcome is length of ICU stay, total length of hospital stays as well as mechanical ventilation dependent days across these groups.

## 5. Statistical analysis

Continuous variables were expressed as means (standard deviations) or medians (interquartile range, IQR) and comparisons across three groups were made using analysis of variance (ANOVA) for normally distributed variables, Kruskal-Wallis test for non-normally distributed variables. Categorical variables were expressed as counts with proportions and compared using chi-square test or Fisher's exact test where appropriate. Difference in recognition of AKI before and after adjustment for cumulative fluid balance was evaluated with McNemar's test. Consistency of AKI diagnosis and staging were assessed in Cohen's weighted kappa coefficient. Univariate analysis was performed regarding underestimation of serum creatinine as an outcome variable. Risk factors of significance in univariate analysis were further included in multivariate analysis to confirm independent risk factors for underestimation of serum creatinine. A two-sided p value of  $< 0.05$  was considered to be statistically significant. All analyses were performed using SPSS 11.0 software (ver. 18.0, SPSS Inc., US).

## Results

### 1. Basic characteristics

A total of 1334 patients (776 men) with a mean age of 56 years (range, 18 to 82) were investigated. Patients were divided into three groups based on  $\Delta$ Crea level (Table 1). More patients had diabetes mellitus, hypertension, previous contrast exposure, surgery without CPB in underestimation group. Man sex, weight, and history of chronic kidney disease, acute coronary syndrome as well as stroke were similar across  $\Delta$ Crea groups. Differences were noted across groups with respect to age, left ventricular ejection fraction, baseline creatinine, APACHE II score, CPB and cross-clamp duration.

Table 1

Basic characteristics of the patients according to the difference between adjusted and unadjusted postoperative serum creatinine

	<b>Underestimation</b> <b>n = 353</b>	<b>Normal</b> <b>n = 641</b>	<b>Overestimation</b> <b>n = 340</b>	<b>P value</b>
Demographic data				
Male [n (%)]	228 (64.6)	338 (52.7)	210 (61.8)	0.687
Age (years)	61 ± 11	56 ± 12	52 ± 13	< 0.001
BMI (kg/m <sup>2</sup> )	23.2 ± 3.0	23.3 ± 3.3	22.9 ± 3.4	0.314
Comorbid conditions [n (%)]				
Hypertension	163 (46.2)	218 (34.0)	97 (28.5)	< 0.001
DM	73 (20.7)	66 (10.3)	20 (5.9)	< 0.001
CKD	6 (1.7)	4 (0.6)	4 (1.2)	0.272
Cancer	3 (0.8)	10 (1.6)	7 (2.1)	0.418
ACS	8 (2.3)	13 (2.0)	1 (0.3)	0.073
Stroke	8 (2.3)	19 (3.0)	6 (1.8)	0.494
Atrial fibrillation	12 (3.4)	25 (3.9)	13 (3.8)	0.921
Contrast exposure [n (%)]	210 (59.5)	356 (55.5)	168 (49.4)	0.027
NYHA III-IV [n (%)]	211 (59.8)	470 (73.3)	266 (78.2)	< 0.001
LVEF (%)	59.7 ± 9.6	61.9 ± 8.6	60.6 ± 9.3	0.006
Preoperative SCr (µmol/L)	87.4 ± 31.8	76.7 ± 17.1	81.9 ± 24.1	< 0.001
APACHE II at ICU admission	13.1 ± 3.4	12.0 ± 3.5	12.7 ± 3.9	< 0.001
Type of surgery [n (%)]				
Off-pump	154 (43.6)	119 (18.6)	5 (1.8)	< 0.001
On-pump	199 (56.4)	522 (81.4)	335 (98.5)	< 0.001
On pump-surgery variables				
CPB time (min)	105.3 ± 45.2	91.2 ± 32.3	104.4 ± 72.2	< 0.001
Cross-clamp (min)	63.9 ± 28.6	56.1 ± 30.6	60.1 ± 26.8	0.006
Ultrafiltration (ml)	1311 ± 1417	1727 ± 1036	1891 ± 1268	< 0.001
The data in the table are expressed as mean ± standard deviation or number (%). P value is for the comparison among groups.				

## 2. Incidence of CSA-AKI before and after adjustment

The incidence of AKI based on the KDIGO criteria was 29.5% (with 24.4%, 3.0% and 2.1% in AKI stage 1–3, respectively). The in-hospital mortality rate was 1.05% (14/1334) and incidence of requirement for continuous renal replacement therapy (CRRT) during 48 hours after ICU admission was 0.37% ( $n = 5$ ). After adjustment for cumulative fluid balance, the incidence of AKI elevated to 31.8% (increased from 24.4 to 26.5% in stage 1, decreased from 3.0 to 2.8% in stage 2 and increased from 2.1 to 2.5% in stage 3 respectively). AKI stage only increased by one stage in the 3 groups after adjustment for cumulative fluid balance. An increase was found in only 3.7% of those originally in stage 0, 0.4% in those originally in stage 1 and 0.5% in those originally in stage 2. The percentage of agreement for AKI diagnosis was 94.9% with a kappa of 0.86 (95% confidence interval [CI], 0.83–0.89), whereas percentage of agreement for AKI staging was 93.8% with a kappa of 0.86 (95% CI, 0.83–0.89) after adjustment for cumulative fluid balance (Table 2).

Table 2  
Diagnosis and staging of AKI before and after adjustment for fluid balance in patients after cardiac surgery

Unadjusted AKI stage	Adjusted AKI stage				Total, N (%)
	0	1	2	3	
0	890 (66.7)	50 (3.7)	0 (0)	0 (0)	940 (70.5)
1	19 (1.4)	301 (22.6)	6 (0.4)	0 (0)	326 (24.4)
2	0 (0)	2 (0.1)	31 (2.3)	7 (0.5)	40 (3.0)
3	0 (0)	0 (0)	1 (0.07)	27 (2.0)	28 (2.1)
Total, N (%)	909 (68.1)	353 (26.5)	38 (2.8)	34 (2.5)	1334
Kappa = 0.86 (95% CI 0.83–0.89) and percentage agreement = 94.9% for AKI diagnosis					
Kappa = 0.86 (95% CI 0.83–0.89) and percentage agreement = 93.8% for AKI staging					

## 3. Change in serum creatinine ( $\Delta$ Crea) and outcomes

Patients in underestimation group showed prolonged length of ICU stay compared to the normal group and overestimation group (3.2 [1.0–3.0] vs 2.1 [1.0–3.0] d,  $P < 0.001$ ; 3.2 [1.0–3.0] vs 2.3 [1.0–3.0] d,  $P < 0.001$ , respectively). Both length of hospital stays and ventilation dependent days in underestimation group were significantly higher than normal group (13.8 [10.0–15.0] vs 12.6 [10.0–14.0] d,  $P < 0.05$ ; 1.7 [1.0–1.5] vs 1.2 [0.5–1.5] d,  $P < 0.001$ , respectively) (Table 3). There were no statistical difference in the in-hospital mortality rate and the incidence of CRRT across these groups.

Table 3

Outcomes of the patients according to the difference between adjusted and unadjusted postoperative serum creatinine

	<b>Underestimation</b> <b>n = 353</b>	<b>Normal</b> <b>n = 641</b>	<b>Overestimation</b> <b>n = 340</b>	<b><i>P</i> value</b>
Outcome variables				
Ventilation dependent days (days)	1.7 [1.0-1.5]	1.2 [0.5–1.5]	1.0 [0.5–1.5]	< 0.001
ICU stay (days)	3.2 [1.0–3.0]	2.1 [1.0–3.0]	2.0 [1.0–3.0]	< 0.001
Hospital stay (days)	13.8 [10–15]	12.0 [10–14]	12.0 [10–14]	0.012
CRRT [n (%)]	0 (0.0)	2 (0.3)	3 (0.9)	0.591
In-hospital mortality [n (%)]	4 (1.1)	5 (0.8)	5 (1.5)	0.154
The data in the table are expressed as median ± interquartile range or number (%). <i>P</i> value is for the comparison among groups.				

## 4. Risk factors associated with underestimation of serum creatinine

Multivariate analysis indicated that factors independently associated with underestimation of serum creatinine due to the cumulative fluid balance were older age ( $P = 0.035$ ), higher baseline serum creatinine ( $P < 0.001$ ), lower left ventricular ejection fraction ( $P = 0.001$ ), and extent of cumulative fluid balance ( $P < 0.001$ ) during ICU stay after cardiac surgery (Table 4).

Table 4

Univariate and multivariate analysis with underestimation of serum creatinine as the outcome variable

Variable	Univariate Analysis		Multivariate Analysis	
	HR (95% CI)	<i>P</i> Value	HR (95% CI)	<i>P</i> Value
Sex, Male	1.44 (1.12–1.85)	0.004		
BMI (per 1 kg/m <sup>2</sup> increase)	1.01 (0.96–1.06)	0.723		
Age (per 1 unit increase)	1.05 (1.04–1.60)	< 0.001	1.05 (1.01–1.10)	0.035
Hypertension (present)	1.81 (1.41–2.33)	< 0.001		
Diabetes (present)	2.71 (1.93–3.81)	< 0.001		
CKD (present)	2.10 (0.72–6.10)	0.172		
NYHA III-IV (present)	2.02 (1.56–2.61)	< 0.001		
Preoperative SCr (per 1 µmol/L increase)	1.01 (1.01–1.02)	< 0.001	1.05 (1.03–1.08)	0.000
LVEF (per 1 unit decrease)	1.02 (1.01–1.04)	0.010	1.10 (1.04–1.17)	0.001
Off-pump surgery (present)	5.35 (4.03–7.09)	< 0.001		
CPB (per 1 min increase)	1.00 (1.00–1.01)	0.061		
Aortic cross-clamp (per 1 min increase)	1.01 (1.00–1.01)	0.014		
Oliguria (present)	1.87 (0.66–5.29)	0.239		
Cumulative fluid balance (per 1 L increase)	1.00 (1.00–1.01)	< 0.001	1.007 (1.005–1.009)	0.000

## Discussion

In this retrospective cohort study, we found that when after adjusting serum creatinine for the cumulative fluid balance, more patients met KDIGO criteria for CSA-AKI. Patients with underestimation of serum

creatinine had worse outcomes in terms of length of ICU stay, total length of hospital stay and mechanical ventilation dependent days, but not in the incidence of CRRT and in-hospital mortality rate.

Since minimal increase of serum creatinine was associated with adverse outcomes in patients within the ICU setting, precise recognition and accurate assessment of AKI may contribute to prevention and early intervention of reversible risk factors [19–22]. Serum creatinine may normally be influenced by several factors, including renal creatinine clearance or creatinine formation or both [23]. Importantly, serum creatinine level also can be affected by dilution effect of fluid resuscitation, which is frequently occurring in critically ill patients [22, 24]. Our results indicate that cumulative fluid balance in patients with cardiac surgery somewhat underestimate the diagnosis and staging of AKI, which is in accordance with previous studies [11, 15].

Post hoc analysis of Fluids and Catheters Treatment study illustrated that incidence of AKI with acute respiratory distress syndrome was greater in those managed with liberal fluid protocol than conservative fluid protocol after adjustment for fluid balance [11]. Moreover, mortality rate of these patients was similar to patients diagnosed with AKI before and after adjustment for fluid balance. Macedo et al. conducted analysis in patients underwent nephrology consultation for AKI in the ICU, which showed dilution effect of fluid overload on serum creatinine may delay diagnosis time for AKI [15]. Study focusing on cardiac surgery patients also demonstrated that patients with AKI only after adjustment for fluid balance had intermediate outcomes between non-AKI and classical AKI patients [12]. Our study demonstrated that patients with underestimation of serum creatinine had prolonged mechanical ventilation dependent days, longer length of ICU stay as well as longer length of hospital stay.

Multivariate analysis of our study also found that after adjustment for relevant risk factors, patients with older age, lower left ventricular ejection fraction, higher baseline serum creatinine and severe extent of cumulative fluid balance after cardiac surgery were independently associated with underestimation of serum creatinine. Thus, to reduce underestimation of serum creatinine along with subsequent prediction of poor prognosis, risk factors including age, baseline cardiac function as well as baseline kidney function should be taken into account during fluid administration in patients undergoing cardiac surgery. There are several limitations in our study. First, as a single center study, regardless of a large cohort of cardiac surgery patients, inherent bias of study design still remains to concern. Second, types of fluid given to patients during perioperative period were not included in the analysis owing to lacking of detailed record for fluid profile. Third, cause of fluid administration was not easily distinguished from our database. Excess fluid administration may be in an effort to manage low cardiac output, and fluid accumulation may be secondary to inflammatory response. Both low cardiac output and inflammation might be responsible for the development of AKI. Last, insensitive fluid loss during study period was not taken into account, which may influence accurate measurement of fluid balance, especially when patients were incubated during ICU period.

Regardless of these limitations, our study highlights dilution effect of cumulative fluid balance on serum creatinine and further illustrates associated outcomes in cardiac surgery patients, which may benefit

physicians to recognize mild AKI via adjustment for cumulative fluid balance. Strikingly, our study identified for the first time the risk factors including age, baseline cardiac function, and preoperative kidney function were independently associated with the underestimation of serum creatinine, which could contribute to screening patients at high risk for misinterpretation of postoperative serum creatinine. However, much more advanced studies should be designed to clear the underlying association between concealed mild AKI and actual change of renal function, such as using combination of serum creatinine and kidney injury biomarkers to timely detect deterioration of kidney function after cardiac surgery [25, 26].

## Conclusions

Cumulative fluid overload in patients after cardiac surgery is very common and leads to underestimation of postoperative AKI. Patients with underestimation of postoperative serum creatinine were associated with a substantial detrimental prognosis.

## Abbreviations

ACS

Acute coronary syndrome; AKI:Acute kidney injury; ANOVA:Analysis of variance; APACHE II:Acute Physiology and Chronic Health Evaluation II; BMI:Body mass index; CI:Confidence interval; CKD:Chronic kidney disease; CPB:Cardiopulmonary bypass; CRRT:Continuous renal replacement therapy; CSA-AKI:Cardiac surgery-associated acute kidney injury; DM:Diabetes mellitus; GFR:Glomerular filtration rate; ICU:Intensive care unit; KDIGO:Kidney Disease Improving Global Outcomes; LOS:Length of stay; LVEF:Left ventricular ejection fraction; NYHA:New York Heart Association; SCr:Serum creatinine; TBW:Total body water; VDD:Ventilation dependent days.

## Declarations

### *Ethics approval and consent to participate*

The study was approved by the Institutional Ethics Committee of Zhongshan Hospital, Fudan University, Shanghai, China (B2018-175). The need for informed consents were waived by the Institutional Ethics Committee because this was a retrospective analysis of data collected prospectively with no breach of privacy or anonymity.

### *Consent for publication*

Not applicable.

### *Availability of data and materials*

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

### ***Competing interest***

The authors declare that they have no competing interests.

### ***Funding***

This study was supported by the research grants from the Science and Technology Commission of Shanghai Municipality (14DZ2260200), Shanghai Clinical Medical Center for Kidney Disease Project support by Shanghai Municipal Health Commission (2017ZZ01015), and Shanghai Municipal Hospital Frontier Technology Project supported by Shanghai ShenKang Hospital Development Center (SHDC12018127). No financial support was provided from industrial companies.

### ***Authors' contributions***

JJ, JX, SX, and JH were responsible for the study design, data acquisition and analysis. JJ, JX, JH, and WJ drafted the first manuscript. BS, CW, JT, and XD conducted a critical revision of manuscript. All authors read and approved the final manuscript.

### ***Acknowledgements***

Not applicable.

## **References**

1. Chertow GM, Levy EM, Hammermeister KE, Grover F, Daley J. Independent association between acute renal failure and mortality following cardiac surgery. *Am J Med.* 1998;104:343–8.
2. Hashemzadeh K, Hashemzadeh S, Dehdilani M. Risk factors and outcomes of acute renal failure after open cardiac surgery. *Asian Cardiovasc Thorac Ann.* 2012;20:275–80.
3. Druml W. Acute renal failure is not a “cute” renal failure! *Intensive Care Med.* 2004;30:1886–90.
4. Thakar CV, Arrigain S, Worley S, Yared JP, Paganini EP. A clinical score to predict acute renal failure after cardiac surgery. *J Am Soc Nephrol.* 2005;16:162–8.
5. Perko MJ, Jarnvig IL, Højgaard-Rasmussen N, Eliassen K, Arendrup H. Electric impedance for evaluation of body fluid balance in cardiac surgical patients. *J Cardiothorac Vasc Anesth.* 2001;15:44–8.
6. Bouchard J, Weidemann C, Mehta RL. Renal replacement therapy in acute kidney injury: intermittent versus continuous? How much is enough? *Adv Chronic Kidney Dis.* 2008;15:235–47.
7. Grams ME, Estrella MM, Coresh J, Brower RG, Liu KD. Fluid balance, diuretic use, and mortality in acute kidney injury. *Clin J Am Soc Nephrol.* 2011;6:966–73.
8. Goldstein SL, Somers MJ, Baum MA, Symons JW, Brophy PD, Blowery D, et al. Pediatric patients with multi-organ dysfunction syndrome receiving continuous renal replacement therapy. *Kidney Int.* 2005;67:653–8.

9. Foland JA, Fortenberry JD, Warshaw BL, Pettignano R, Merritt RK, Heard ML, et al. Fluid overload before continuous hemofiltration and survival in critically ill children: a retrospective analysis. *Crit Care Med*. 2004;32:1771–6.
10. National Heart, Lung, and Blood Institute Acute Respiratory Distress Syndrome (ARDS) Clinical Trials Network. Comparison of two fluid-management strategies in acute lung injury. *N Engl J Med*. 2006;354:2564–75.
11. Liu KD, Thompson BT, Ancukiewicz M, Steingrub JS, Douglas IS, Matthay MA, et al. Acute kidney injury in patients with acute lung injury: impact of fluid accumulation on classification of acute kidney injury and associated outcomes. *Crit Care Med*. 2011;39:2665–71.
12. Moore E, Tobin A, Reid D, Santamaria J, Paul E, Bellomo R. The impact of fluid balance on the detection, classification and outcome of acute kidney injury after cardiac surgery. *J Cardiothorac Vasc Anesth*. 2015;29:1229–35.
13. Basu RK, Andrews A, Krawczeski C, Manning P, Wheeler DS, Goldstein SL. Acute kidney injury based on corrected serum creatinine is associated with increased morbidity in children following the arterial switch operation. *Pediatr Crit Care Med*. 2013;14(5):e218-24.
14. Jin J, Chang SC, Xu S, Xu J, Jiang W, Shen B, et al. Early postoperative serum creatinine adjusted for fluid balance precisely predicts subsequent acute kidney injury after cardiac surgery. *J Cardiothorac Vasc Anesth*. 2019;33(10):2695–702.
15. Macedo E, Bouchard J, Soroko SH, Chertow GM, Himmelfarb J, Ikizler TA, et al. Fluid accumulation, recognition and staging of acute kidney injury in critically-ill patients. *Crit Care*. 2010;14:R82.
16. Knaus WA, Wagner DP, Draper EA, Zimmerman JE, Bergner M, Bastos PG, et al. The APACHE III prognostic system: risk prediction of hospital mortality for critically ill hospitalized adults. *Chest*. 1991;100:1619–36.
17. Khwaja A. KDIGO clinical practice guidelines for acute kidney injury. *Nephron Clin Pract*. 2012;120:c179–84.
18. Moran SM, Myers BD. Course of acute renal failure studied by a model of creatinine kinetics. *Kidney Int*. 1985;27:928–37.
19. Kellum JA, Levin N, Bouman C, Lameire N. Developing a consensus classification system for acute renal failure. *Curr Opin Crit Care*. 2002;8:509–14.
20. Chertow GM, Burdick E, Honour M, Bonventre JV, Bates DW. Acute kidney injury, mortality, length of stay, and costs in hospitalized patients. *J Am Soc Nephrol*. 2005;16:3365–70.
21. Vaara ST, Pettilä V, Kaukonen KM, Bendel S, Korhonen AM, Bellomo R, et al. The attributable mortality of acute kidney injury: a sequentially matched analysis. *Crit Car Med*. 2014;42:878–85.
22. Lassnigg A, Schmidlin D, Mouhieddine M, Bachmann LM, Druml W, Bauer P, et al. Minimal changes of serum creatinine predict prognosis in patients after cardiothoracic surgery: a prospective cohort study. *J Am Soc Nephrol*. 2004;15:1597–605.
23. Singri N, Ahya SN, Levin ML. Acute renal failure. *JAMA*. 2003;289:747–51.

24. Lassnigg A, Schmid ER, Hiesmayr M, Falk C, Druml W, Bauer P, et al. Impact of minimal increases in serum creatinine on outcome in patients after cardiothoracic surgery: do we have to revise current definitions of acute renal failure? *Crit Care Med.* 2008;36:1129–37.
25. Devarajan P. Neutrophil gelatinase-associated lipocalin (NGAL): A new marker of kidney disease. *Scand J Clin Lab Invest Suppl.* 2008;241:89–94.
26. Han WK, Bailly V, Abichandani R, Thadhani R, Bonventre JV. Kidney Injury Molecule-1 (KIM-1): a novel biomarker for human renal proximal tubule injury. *Kidney Int.* 2002;62:237–44.

## Figures




Figure 1

The normal distribution of  $\Delta\text{Crea}$  Mean of  $\Delta\text{Crea} = 0.11$ ; Standard deviation of  $\Delta\text{Crea} = 4.81$ ;  $N = 1334$