

# Psychological distress and internet-related behaviors between schoolchildren with and without overweight during the COVID-19 outbreak

**Chao-Ying Chen**

Hong Kong Polytechnic University <https://orcid.org/0000-0002-9632-6659>

**I-Hua Chen**

Minnan Normal University

**Kerry O'Brien**

Monash University <https://orcid.org/0000-0003-3145-6038>

**Janet D Latner**

University of Hawaii at Manoa

**Chung-Ying Lin** (✉ [cylin36933@gmail.com](mailto:cylin36933@gmail.com))

Hong Kong Polytechnic University <https://orcid.org/0000-0002-2129-4242>

---

## Research Article

**Keywords:** stigma, overweight, obesity, COVID-19, anxiety, depression, internet, stress

**Posted Date:** May 7th, 2020

**DOI:** <https://doi.org/10.21203/rs.3.rs-27322/v1>

**License:**   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

**Version of Record:** A version of this preprint was published on January 25th, 2021. See the published version at <https://doi.org/10.1038/s41366-021-00741-5>.

# Abstract

**Objective:** The novel coronavirus disease 2019 (COVID-19) pandemic, and its resulting social policy changes, may result in psychological distress among schoolchildren with overweight. This study thus aimed to (1) compare psychological distress (including fear of COVID-19 infection, stress, anxiety, and depression), perceived weight stigma, and problematic internet-related behaviors between schoolchildren with and without overweight; (2) assess whether perceived weight stigma and problematic internet-related behaviors explained psychological distress.

**Methods:** Schoolchildren (n=1 357; mean age=10.7 years) with overweight (n=236) and without overweight (n=1 121) completed an online survey assessing their fear of COVID-19 infection, stress, anxiety, depression, perceived weight stigma, problematic smartphone application use, problematic social media use, and problematic gaming.

**Results:** Schoolchildren with overweight had significantly higher levels of COVID-19 infection fear, stress, depression, perceived weight stigma, and problematic social media use than those without overweight. Regression models showed that perceived weight stigma and problematic internet-related behaviors were significant predictors of psychological distress among schoolchildren with overweight.

**Conclusion:** Strategies to manage perceived weight stigma and problematic internet-related behaviors may have a positive influence on mental health among schoolchildren with overweight under health-threatening circumstances, such as the current COVID-19 pandemic.

## Introduction

The rapid growth of novel coronavirus disease 2019 (COVID-19) infection worldwide (>3.4 million COVID-19 infections) has caused more than 230 000 deaths as of May 1, 2020.<sup>1</sup> The absence of an effective treatment means that healthcare providers have paid high level attention to people who are particularly vulnerable to the COVID-19. This includes those with chronic disease (e.g., diabetes, coronary heart disease), compromised immune systems, and those with overweight (including obesity). Patients with overweight may disproportionately develop severe comorbidities after COVID-19 infection.<sup>2</sup> This suggestion is supported by previous work investigating the relationship between influenza and overweight. Specifically, the altered immune response and sedentary lifestyle that influences the physiological health contribute to poorer clinical outcomes for patients with overweight following influenza.<sup>3</sup> Recent evidence from France found that the severity of COVID-19 infection was positively related to patients' body mass index (BMI). That is, patients with a higher BMI, especially those with a BMI higher than 35kg/m<sup>2</sup>, are more likely to require support from invasive mechanical ventilators.<sup>4</sup>

Because of the health consequences mentioned above, people with overweight may additionally suffer from fears about their health and associated psychological distress. Specifically, people with overweight who feel they are more vulnerable to COVID-19 related morbidity and mortality may also display greater

psychological distress. Moreover, lifestyle and living patterns have by necessity changed during the COVID-19 outbreak. For example, many governments have implemented social distancing and lockdowns laws to prevent expansion of COVID-19 infection.<sup>5,6</sup> Mainland China has implemented temporary school closures and required primary schoolchildren to study online at home for several months during the COVID-19 outbreak. Such changes in lifestyles may have negative consequence on individuals' mental health. Prior research has shown that major life events, such as the loss of a parent or friend can be associated with psychological distress, and people with overweight can be more vulnerable to such life stressors than their counterparts with normal-weight.<sup>7</sup> Accordingly, children with overweight may encounter quite different levels of psychological distress and associated behavioral outcomes due to COVID-19 than children without overweight. Even during pre-pandemic times, children with overweight experience greater psychological distress and impaired psychosocial outcomes due to weight stigma (weight bias).<sup>8</sup> Research shows that people with overweight have a higher chance of being bullied and teased than those without overweight.<sup>8</sup> These stigmatizing experiences and the ongoing fear and perception of weight stigma can lead to significant psychological distress.<sup>9</sup>

One major lifestyle and behavioral change that has occurred during school closures is increased internet usage by schoolchildren. Schoolchildren are very likely to engage in more internet-related activities for educational, entertainment, and social-engagement purposes during periods of physical distancing and isolation. However, there can be unintended negative consequences if internet behaviors are not monitored or constrained. For example, a study across eight Asian and European countries found more problematic internet use to be associated with higher levels of depressive and anxiety symptoms.<sup>10</sup> Moreover, increased severity of gaming disorder and social media addiction may lead to psychological and sleep problems.<sup>11</sup> During the unprecedented period of COVID-19 self-isolation and distancing an increase home-based online activities is likely, and it seems important to examine the relationship between internet-related activities and psychological distress in schoolchildren with overweight during the pandemic.

There is little research examining whether weight status and weight stigma may influence perceptions of vulnerability to infectious diseases, and associated psychological distress. And whether problematic internet related behaviors during the COVID-19 lockdown are associated with greater fear of COVID-19 infection, and psychological distress. The present study recruited a large sample of schoolchildren with and without overweight from mainland China to examine the relationships between weight status, weight stigma, problematic internet and social media use, and psychological distress (including fear of COVID-19) during the COVID-19 lockdown.

## Methods

### *Participants, procedure, and ethical concerns*

The study was part of an ongoing longitudinal project assessing problematic internet-related behaviors among primary school children. This research was approved by the ethics committee of the Hong Kong

Polytechnic University (IRB ref: HSEARS20190718001). The longitudinal project routinely assesses primary school children's characteristics, problematic internet-related behaviors (smartphone applications, social media, and gaming), perceived weight stigma, and psychological distress across a one-year period. During the period of the longitudinal project, the COVID-19 outbreak occurred in mainland China and the primary schoolchildren were instructed to stay at home for online learning. Consequently, an item on fear of COVID-19 infection was incorporated into the longitudinal project (see *Instruments* section for details) to assess the extent of children's fear related to COVID-19. The present study was cross-sectional, with questionnaire data from the same wave that included the COVID-19 question analyzed.

The survey was conducted online with the distribution of the survey facilitated by teachers from three primary schools in Sichuan, China. The online survey was generated by the research team and checked by the teachers to ensure that the survey hyperlink worked. Schoolteachers then sent the hyperlink to their students for participation. The first page of the online survey stated the objectives and participants' rights, and sought consent from the students and one parent. Following the provision of participation consent from students and one of their parents, access to the survey proper was opened. Eligibility of the primary schoolchildren was defined the following inclusion criteria (i) they could read and understand the survey, which was written in Chinese and (ii) they owned and used at least one smartphone with internet access. The participants were excluded if they misperceived their weight status; that is, their perceived weight status did not fit with their body mass index (BMI) defined weight status. The reason for the exclusion criterion is because prior research shows that misperception of weight status is a strong predictor of children's psychological distress, and we did not wish for this association to confound the present study.<sup>12-14</sup>

### *Measures*

**Fear of COVID-19 infection.** Using a visual analogue scale (from 0 [not at all afraid] to 10 [completely afraid]), a question "Are you afraid of being infected by COVID-19?" was used to understand to what extent participants fear the COVID-19 infection.

**Depression, Anxiety, Stress Scale-21 (DASS-21).** Using 21 items rated on a 4-point Likert scale (ranging from 0 [did not apply to me at all] to 3 [applied to me very much or most of the time]), the DASS-21 assesses three types of psychological distress: stress (7 items), anxiety (7 items), and depression (7 items). Example items for the DASS-21 are "I found it hard to wind down" (for stress), "I was aware of dryness of my mouth" (for anxiety), and "I could not seem to experience any positive feeling at all" (for depression). The item scores are summated in each subscale to indicate the level of psychological distress, where higher scores indicate a higher level of depression, anxiety, or stress. The psychometric properties of the DASS-21 and the Chinese DASS-21 have been found to be good.<sup>15,16</sup> The present study found that the Chinese DASS-21 had good internal consistency ( $\alpha = 0.82$  for stress subscale; 0.79 for anxiety subscale; and 0.84 for depression subscale).

**Perceived weight stigma.** A 10-item measure was used to assess perceived weight stigma. Participants responded to items such as “people act as if you are inferior because of your weight” using a dichotomous scale (0 = No and 1 = Yes). Item scores were summated to indicate the level of perceived weight stigma (scores ranging from 0-10), with higher scores indicating a higher level of perceived weight stigma. The linguistic validity and internal consistency of the Chinese perceived weight stigma items have been found to be satisfactory.<sup>17</sup> The present study also showed that the Chinese perceived weight stigma items had good internal consistency ( $\alpha = 0.83$ ).

**Smartphone Application-Based Addiction Scale (SABAS).** The six-item SABAS was used to assess level of problematic smartphone application use. The SABAS was designed based on the addiction component model criteria (i.e., salience, mood modification, tolerance, withdrawal conflict and relapse) proposed by Griffiths.<sup>18,19</sup> Participants responded to items such as “During the past week, my smartphone is the most important thing in my life” using a Likert scale ranging from 1 (strongly disagree) to 6 (strongly agree). The item scores are summed to indicate the level of problematic smartphone application use, where a higher score indicates a higher level of problematic use. The psychometric properties of the SABAS and the Chinese SABAS have been found to be satisfactory.<sup>20-24</sup> The present study also showed that the Chinese SABAS had good internal consistency ( $\alpha = 0.88$ ).

**Bergen Social Media Addiction Scale (BSMAS).** The BSMAS assesses the level of problematic social media use with six items rated on a 5-point Likert scale ranging from 1 (very rarely) to 5 (very often). The scale was developed based on the addiction component model criteria proposed by Griffiths.<sup>18,19</sup> The following is an example item from the BSMAS “How often during the last week have you spent a lot of time thinking about social media or planned use of social media?”. Item scores are summated to indicate the level of problematic social media use, where a higher score indicate a higher level of problematic use. The psychometric properties of the BSMAS and Chinese BSMAS have been found to be satisfactory.<sup>25-28</sup> The present study also showed that the Chinese BSMAS had good internal consistency ( $\alpha = 0.88$ ).

**Internet Gaming Disorder Scale-Short Form (IGDS9-SF).** The nine-item IGDS9-SF assesses the level of disordered internet gaming behavior. The scale items were designed based on the nine criteria indicating internet gaming disorder [IGD] in the *Diagnostic and statistical manual of mental disorders, fifth edition* (DSM-5).<sup>29</sup> Items such as “During last week, do you feel more irritability, anxiety or even sadness when you try to either reduce or stop your gaming activity?” were presented to participants who responded using a 5-point Likert scale ranging from 1 (never) to 5 (very often). The item scores are summed to indicate the level of problematic gaming, where higher scores indicating higher levels of problematic gaming. The psychometric properties of the IGDS9-SF have been found to be good.<sup>30-36</sup> The present study also showed that the Chinese IGDS9-SF had good internal consistency ( $\alpha = 0.926$ ).

**Time spent on internet-related behaviors.** Time spent on internet-related behaviors was assessed with the question: “How much time did you spend on each of the following internet-related behaviors daily?”. Participants were asked to indicate time spent on each of the following internet-related behaviors in the

past week: smartphone, social media, and gaming. The time spent on each internet-related behavior was then converted into hours per day in use.

**Weight status.** The schoolchildren self-reported their height in cm and weight in kg. BMI was then calculated to determine whether a schoolchild belongs to a group with overweight or a group without overweight.<sup>37</sup> In addition to reporting height and weight, each schoolchild was asked to report their self-perceived weight status using the question: “What do you think your weight status is?” Answer choices included “Very thin”, “Thin”, “Normal weight”, “Overweight”, or “Obese”. Then, the self-perceived weight status was reclassified into two categories of “overweight” or “nonoverweight”.

### *Data analysis*

Independent t-tests were used to detect the significant differences between the group with overweight and that without overweight in their psychological distress, perceived weight stigma, problematic internet-related behaviors, and time spent on internet-related behaviors. In addition, multivariate linear regression models were used to examine the associations between psychological distress (i.e., fear of COVID-19 infection, stress, anxiety, and depression), perceived weight stigma, and problematic internet-related behaviors (including problematic smartphone application use, problematic social media use, and problematic gaming). We divided the participants into two groups (participants with overweight and without overweight) to conduct the regression models for each group. Age and gender were adjusted in all the regression models. Moreover, a regression model using all participants (i.e., including those with and without overweight) was used to examine the associations between weight status, psychological distress, perceived weight stigma, and problematic internet-related behaviors.

## **Results**

Table 1 presents participants’ characteristics among schoolchildren with overweight (n = 236; 132 boys), and without overweight (n = 1 121; 557 boys), and noneligible participants (n = 657; 318 boys). Table 2 shows the means and SDs for psychological distress, perceived weight stigma, problematic internet-related behaviors, and time spent on internet-related behaviors between the two groups. As can be seen in Table 2, schoolchildren with overweight had significantly higher levels of COVID-19 infection fear, stress, and depression than did those without overweight. Schoolchildren with overweight also had significantly higher levels of perceived weight stigma than did those without overweight. With regards to problematic internet-related behaviors, schoolchildren with overweight had significantly higher levels of problematic social media use than did those without overweight. There were no significant differences between overweight and non-overweight in time spent on internet-related behaviors.

Regression models for schoolchildren without overweight (Table 3) indicated that perceived weight stigma was a significant predictor in models, with higher levels of weight stigma associated with greater fear of COVID-19 infection, and higher levels of stress, anxiety, and depression. Problematic smartphone,

social media, and gaming use were also all significant positive predictor in stress, anxiety and depression, but not COVID-19 fear.

Regression models for schoolchildren with overweight (Table 3) indicated that perceived weight stigma was a significant predictor in fear of COVID-19 infection at the  $p < 0.10$  level, but not at the conventional 0.05 level. Weight stigma was a significant predictor of stress, anxiety, and depression. With higher levels of weight stigma associated with higher levels of stress, anxiety, and depression. Similar to non-overweight, problematic smartphone and social media application use were also significant positive predictor in stress, anxiety and depression, but not fear of COVID-19 infection. Problematic gaming was not a significant predictor of COVID-19 infection fear, but was a significant predictor of stress, and only reached the 0.10 level of significance for anxiety and depression.

Regression models for schoolchildren regardless of weight status (Table 3) indicated that overweight was a significant predictor in fear of COVID-19 infection. Weight stigma was a significant predictor of all types of psychological distress. With higher levels of weight stigma associated with higher levels of fear of COVID-19 infection, stress, anxiety, and depression. Additionally, problematic smartphone, social media, and gaming use were also all significant positive predictor in stress, anxiety and depression, but not COVID-19 fear.

## Discussion

Consistent with previous findings,<sup>8,12-14</sup> the present study found that schoolchildren with overweight had higher levels of psychological distress and perceived weight stigma than their counterparts with normal-weight. Moreover, schoolchildren with overweight had greater problematic social media use than did those without overweight. Regression models indicated that greater perceived weight stigma was associated with greater psychological distress in schoolchildren regardless of their weight status. Moreover, problematic smartphone application use and problematic social media use were associated schoolchildren's stress, anxiety, and depression, regardless of their weight status. Among schoolchildren without overweight, problematic gaming was associated with higher levels of stress, anxiety, and depression, which is in line with previous findings.<sup>10,11</sup> However, among those with overweight, problematic gaming was associated with lower levels of stress, anxiety, and depression.

There are several explanations for higher psychological distress in schoolchildren with overweight than their peers without overweight. One of the major reasons might be because of the perceived weight stigma in this population. This speculation is supported by the recent meta-analysis conclusion that perceived weight stigma is significantly associated with depression and anxiety in people with overweight.<sup>9</sup> Apart from perceived weight stigma, schoolchildren with overweight might be afraid of the serious health outcomes resulting from COVID-19 infection.<sup>4</sup> In the present study, schoolchildren with overweight demonstrated more fear of being affected by COVID-19 than did those without overweight. However, the present study did not investigate specific reasons why they fear COVID-19 infection.

Therefore, the speculation of fearing serious health outcomes resulted from COVID-19 infection needs corroboration from future studies. Moreover, the present study revealed that problematic social media use was more severe in schoolchildren with overweight than those without overweight. Thus, the higher level of psychological distress may come from a greater level of problematic social media use, as prior evidence showing the positive association between psychological distress and problematic social media use.<sup>11</sup>

Problematic social media use in children with overweight requires additional attention during prolonged school suspension after COVID-19 outbreak. Addiction or prolonged use of social media has been found to be a great risk of overweight.<sup>38,39</sup> People with overweight and resultant stigmatizing experiences may be more likely to use social media for virtual interaction instead of real interpersonal activities, and they might be more avoidant of live social interactions, possibly due to anxiety or fear of stigma. They may also heavily rely on social media to search information relating to overweight, which may increase the risk of problematic internet use.<sup>40,41</sup> Therefore, schoolchildren with overweight might heavily depend on social media because they want to search COVID-19-related information and seek emotional support during the current pandemic period. Unfortunately, such dependence on social media, even when not increasing significantly the time spent on social media, may trigger the development of psychological distress, especially for schoolchildren with overweight in the present study.

In our results, perceived weight stigma was an important factor contributing to psychological distress regardless of weight status. Children with high perceived weight stigma might fear that they will not receive high quality health care should they get COVID-19. However, because weight stigma is closely associated with anxiety and depression, the association of weight stigma with fear of COVID infection may simply reflect higher anxiety and poorer mood in those who are stigmatized. Interestingly, the association between perceived weight stigma and the fear of COVID-19 infection was significant in schoolchildren without overweight but only significant at the 0.10 level in those with overweight. The relationship between perceived weight stigma and fear of COVID-19 infection among schoolchildren without overweight is in line with prior evidence showing associations between weight stigma and psychological distress among individuals without overweight.<sup>43</sup>

Surprisingly, problematic gaming seemed to be a protective factor against psychological distress in schoolchildren with overweight. Specifically, increased problematic use of gaming was significantly associated with less stress, anxiety, and depression. It is possible that the feature of hiding one's identity (i.e., without revealing real information) during gaming may ease psychological distress for schoolchildren with overweight. Contrary to gaming, social media usually needs to reveal some personal information and may expose weight information. Thus, for schoolchildren with overweight who suffered from weight stigma, anonymous gaming could be an anxiety-alleviating activity, which may possibly override the more typical negative influence of problematic online behaviors on psychological well-being.<sup>11</sup> Moreover, gaming does not offer a platform for sharing COVID-19-related information while social media does. Considering the potential impact of COVID-19 on psychological distress, gaming

engagement may protect schoolchildren with overweight. Future research might explore the potential benefits of certain gaming activities in specific populations.

There are some limitations in the present study. First, all the data including demographics (e.g., body weight and height), psychological constructs, perceived weight stigma, and internet-related behaviors are self-reported. Hence, some information may be susceptible to memory-related bias or social desirability bias. Second, the societal perception of discrimination toward people with overweight varies across countries. Therefore, the perceived weight stigma and resulting consequences during this health crisis may also be different. That is to say, the results should be generalized to other populations with caution. Lastly, some important confounding variables could not be controlled in the present study. Specifically, schoolchildren's screen behaviors are highly associated with their parents' screen behaviors.<sup>44,45</sup> However, we did not collect information regarding parents' screen use.

## Conclusion

In conclusion, schoolchildren with overweight suffered from stress, depression, perceived weight stigma, and fear of COVID-19 infection during the COVID-19 outbreak period. They also demonstrated problematic social media use. Perceived weight stigma was positively associated with fear of COVID-19 infection, while problematic use of social media and smart phones were also correlated with increased psychological distress, regardless of weight status. However, problematic gaming appeared to be associated with protective effects in schoolchildren with overweight. Therefore, strategies to manage perceived weight stigma may have a positive influence on mental health under health-threatening circumstances, such as the current COVID-19 pandemic. Future studies are warranted on the causes, correlates, and amelioration of fears related to public health crises including COVID-19. Finally, it would be important to develop and test the efficacy of psychological support and monitoring programs for internet-related activities among schoolchildren with overweight.

## References

1. World Health Organization. Coronavirus disease (COVID-2019): Situation report-85. [https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200504-covid-19-sitrep-105.pdf?sfvrsn=4cdda8af\\_2](https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200504-covid-19-sitrep-105.pdf?sfvrsn=4cdda8af_2). Accessed May 5, 2020.
2. Ryan DH, Ravussin E, Heymsfield COVID 19 and the patient with obesity – The editors speak out. *Obesity*. 2020. doi: 10.1002/oby.22808. [Epub ahead of print]
3. Luzi L, Radaelli MG. Influenza and obesity: its odd relationship and the lessons for COVID-19 *Acta Diabetol*. 2020. doi: 10.1007/s00592-020-01522-8. [Epub ahead of print]
4. Simonnet A, Chetboun M, Poissy High prevalence of obesity in severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) requiring invasive mechanical ventilation. *Obesity*. 2020. doi: 10.1002/oby.22831. [Epub ahead of print]

5. Alvarez E. COVID-19 concerns lead to unusual bans in Dallas County; 2020.  
[https://www.wfaa.com/article/news/health/coronavirus/covid-19-concerns-lead-to-unusu al-bans-in-dallas-county/287-a8a50a42-7e86-4094-82a4-6ca5aeb80e5e](https://www.wfaa.com/article/news/health/coronavirus/covid-19-concerns-lead-to-unusual-bans-in-dallas-county/287-a8a50a42-7e86-4094-82a4-6ca5aeb80e5e). Accessed April 16,
6. Safer at home order for control of COVID-19. California Health and Safety Code §120295; Los Angeles County Code §11.02.080.; March 19, 2020.  
[http://publichealth.lacounty.gov/media/Coronavirus/COVID-19\\_SaferAtHome\\_HealthOfficerOrder\\_20200319\\_Signed.pdf](http://publichealth.lacounty.gov/media/Coronavirus/COVID-19_SaferAtHome_HealthOfficerOrder_20200319_Signed.pdf). Accessed April 16, 2020.
7. Salleh Life event, stress and illness. *Malays J Med Sci.* 2008;15(4):9-18.
8. Puhl RM, Heuer The stigma of obesity: A review and update. *Obesity.* 2009;17:941-964.
9. Alimoradi Z, Golboni F, Griffiths MD, Broström A, Lin CY, Pakpour Weight-related stigma and psychological distress: A systematic review and meta-analysis. *Clin Nutr.* 2019 doi: 10.1016/j.clnu.2019.10.016. [Epub ahead of print]
10. Balhara PS, Doric A, Stevanovic D, et Correlates of problematic internet use among college and university students in eight countries: An international cross-sectional study. *Asian J Psychiatr.* 2019;45:113-120.
11. Wong HY, Mo HY, Potenza MN, et Relationships between severity of internet gaming disorder, severity of problematic social media use, sleep quality and psychological distress. *Int J Environ Res Public Health.* 2020;17(6):1879.
12. Isomaa R, Isomaa AL, Marttunen M, Kaltiala-Heino R, Björkqvist K. Longitudinal concomitants of incorrect weight perception in female and male *Body Image* 2011;8:58-63.
13. Lin Y-C, Latner JD, Fung XCC, Lin C-Y. Poor health and experiences of being bullied in adolescents: Self-perceived overweight and frustration with appearance *Obesity.* 2018;26(2):397-404.
14. Ojala K, Tynjala J, Valimaa R, Villberg J, Kannas Overweight adolescents' self-perceived weight and weight control behaviour: HBSC Study in Finland 1994-2010. *J Obes* 2012;2012:180176.
15. Shaw T, Campbell MA, Runions KC, Zubrick Properties of the DASS-21 in an Australian community adolescent population. *J Clin Psychol.* 2017;73(7):879-892.
16. Wang K, Shi H-S, Geng F-L, et Cross-cultural validation of the Depression Anxiety Stress Scale–21 in China. *Psychol Assess.* 2016;28(5):e88-e100.
17. Cheng MY, Wang SM, Lam YY, Luk HT, Man YC, Lin The relationships between weight bias, perceived weight stigma, eating behavior and psychological distress among undergraduate students in Hong Kong. *J Nerv Ment Dis.* 2018;206(9):705-710.
18. Griffiths Internet addiction-time to be taken seriously? *Addict Res.* 2000;8(5):413-418.
19. Griffiths A 'components' model of addiction within a biopsychosocial framework. *J Subst Use.* 2005;10(4):191-197.

20. Yam C-W, Pakpour AH, Griffiths MD, et Psychometric testing of three Chinese online-related addictive behavior instruments among Hong Kong university students. *Psychiatr Q.* 2019;90(1):117-128.
21. Csibi S, Griffiths MD, Cook B, Demetrovics Z, Szabo The psychometric properties of the Smartphone Application-Based Addiction Scale (SABAS). *Int J Ment Health Addict.* 2018;16(2):393-403.
22. Gökler ME, Bulut Validity and reliability of the Turkish version of the Smart Phone Application Based Addiction Scale. *Journal of Cognitive-Behavioral Psychotherapy and Research.* 2019;8(2):100-106.
23. Lin C-Y, Imani V, Broström A, et Smartphone Application-Based Addiction among Iranian adolescents: A psychometric study. *Int J Ment Health Addict.* 2019;17(4):765-780.
24. Soraci P, Ferrari A, Urso A, Griffiths Psychometric properties of the Italian version of the Smartphone Application-Based Addiction Scale (SABAS). *Int J Ment Health Addict.* doi: 10.1007/s11469-020-00222-2 [Epub ahead of print]
25. Bányai F, Zsila Á, Király O, et Problematic social media use: Results from a large-scale nationally representative adolescent sample. *Plos One.* 2017;12(1):e0169839.
26. Chen IH, Strong C, Lin Y-C, et Time invariance of three ultra-brief internet-related instruments: Smartphone Application-Based Addiction Scale (SABAS), Bergen Social Media Addiction Scale (BSMAS), and the nine-item Internet Gaming Disorder Scale- Short Form (IGDS-SF9) (Study Part B). *Addict Behav.* 2020;101:105960.
27. Leung H, Pakpour AH, Strong C, et Measurement invariance across young adults from Hong Kong and Taiwan among three internet-related addiction scales: Bergen Social Media Addiction Scale (BSMAS), Smartphone Application-Based Addiction Scale (SABAS), and Internet Gaming Disorder Scale-Short Form (IGDS-SF9) (Study Part A). *Addict Behav.* 2020;101:105969.
28. Lin C-Y, Broström A, Nilsen P, et al. Psychometric validation of the Persian Bergen Social Media Addiction Scale using classic test theory and Rasch *J Behav Addict.* 2017;6(4):620-629.
29. American Psychiatric *Diagnostic and statistical manual of mental disorders: DSM-5* (5th ed.). Arlington, VA: American Psychiatric Association; 2013.
30. Beranuy M, Machimbarrena JM, Vega MA, et al. Spanish Validation of the Internet Gaming Disorder Scale-Short Form (IGDS9-SF): Prevalence and relationship with online gambling and quality of *Int J Environ Res Public Health.* 2020;17:1562.
31. Evren C, Dalbudak E, Topcu M, et Psychometric validation of the Turkish nine-item Internet Gaming Disorder Scale- Short Form (IGDS9-SF). *Psychiatry Res.* 2018;265:349-354.
32. Monacis L, de Palo V, Griffiths MD, Sinatra M. Validation of the Internet Gaming Disorder Scale-Short Form (IGDS9-SF) in an Italian-speaking *J Behav Addict.* 2016;5(4):683-690.
33. Pontes HM, Griffiths Measuring DSM-5 internet gaming disorder: Development and validation of a short psychometric scale. *Comput Human Behav.* 2015;45:137-143.
34. Pontes HM, Stavropoulos V, Griffiths MD. Measurement invariance of the Internet Gaming Disorder Scale-Short-Form (IGDS9-SF) between the United States of America, India and the United Kingdom.

*Psychiatry Res.* 2017;257:472-478.

35. T'ng ST, Pau Validation of a translated Internet Gaming Disorder Scale (short form) and measurement invariance across sex groups in Malaysian samples. *Current Psychol.* doi: 10.1007/s12144-020-00668-6 [Epub ahead of print]
36. Wu T-Y, Lin C-Y, Årestedt K, et al. Psychometric validation of the Persian nine-item Internet Gaming Disorder Scale-Short Form: Does gender and hours spent online gaming affect the interpretations of item descriptions? *J Behav* 2017;6(2):256-263.
37. So HK, Nelson EA, Li AM, et Secular changes in height, weight and body mass index in Hong Kong Children. *BMC Public Health.* 2008;8:320.
38. Robinson TN, Banda JA, Hale L, et Screen media exposure and obesity in children and adolescents. *Pediatrics.* 2017;140(Suppl 2):S97-S101.
39. Tsitsika AK, Andrie EK, Psaltopoulou T, et Association between problematic internet use, socio-demographic variables and obesity among European adolescents. *Eur J Public Health.* 2016;26(4):617-622.
40. Kaewpradub N, Kiatrungrit K, Hongsanguansri S, Pavasuthipaisit Association among internet usage, body image and eating behaviors of secondary school students. *Shanghai Arch Psychiatry.* 2017;29(4):208-217.
41. Park S, Lee Associations of body weight perception and weight control behaviors with problematic internet use among Korean adolescents. *Psychiatry Res.* 2017;251:275-280.
42. Tsenkova VK, Carr D, Schoeller DA, Ryff CD. Perceived weight discrimination amplifies the link between central adiposity and nondiabetic glycemic control (HbA1c). *Ann Behav Med.* 2011;41(2):243-251.
43. Lin C-Y, Strong C, Latner JD, Lin Y-C, Tsai M-C, Cheung Mediated effects of eating disturbances in the association of perceived weight stigma and emotional distress. *Eat Weight Disord.* 2020;25(2):509-518.
44. Lin Y-C, Tsai M-C, Strong C, Hsieh Y-P, Lin C-Y, Lee Exploring mediation roles of child screen-viewing between parental factors and child overweight in Taiwan. *Int J Environ Res Public Health.* 2020;17(6):1878.
45. Lin Y-C, Fung XCC, Tsai M-C, Strong C, Hsieh Y-P, Lin C-Y. Insufficient physical activity and overweight: Does caregiver screen-viewing matter? *J Child Fam Stud.* 2019;28:286-297.

## Tables

Table 1. Participants' characteristics

M (SD) or n (%)

F or  $\chi^2$  (p-value)

	Non-OW n = 1121	OW n = 236	Noneligible children <sup>a</sup> n = 657	
Age (year)	10.67 (1.11)	10.78 (0.93)	10.75 (1.06)	1.74 (0.18)
Gender (boy)	557 (49.7%)	132 (55.9%)	318 (48.4%)	4.04 (0.13)
Height (cm)	144.47 (9.00)	149.80 (8.81)	145.20 (9.64)	32.95 (<0.001)
Weight (kg)	35.94 (10.51)	58.88 (22.33)	51.66 (19.06)	342.52 (<0.001)
Body mass index (kg/m <sup>2</sup> )	17.09 (4.11)	26.02 (8.90)	24.35 (8.08)	374.95 (<0.001)
Ethnicity (Han)	1106 (98.7%)	235 (99.6%)	647 (98.5%)	1.69 (0.43)
Current sick (yes)	18 (1.6%)	6 (2.5%)	7 (1.1%)	2.57 (0.28)

a Noneligible children were those who misperceived their weight status; 123 perceived themselves overweight but were classified as nonoverweight according to body mass index; 534 perceived themselves as nonoverweight but were classified as overweight according to body mass index.

Non-OW = group with nonoverweight; OW = group with overweight.

Table 2. Comparisons of psychological distress and internet-related behaviors between groups

	M (SD)/n		t (p-value)
	Non-OW	OW	
Fear of COVID-19 infection	4.00 (3.64)/1042	5.08 (9.03)/210	2.87 (0.004)
Stress	9.00 (2.77)/1121	9.61 (3.04)/236	2.85 (0.005)
Anxiety	8.15 (2.12)/1121	8.39 (2.07)/236	1.58 (0.12)
Depression	8.26 (2.42)/1121	8.62 (2.61)/236	1.97 (0.049)
Perceived weight stigma	0.92 (1.78)/1121	1.30 (2.12)/236	2.59 (0.01)
Problematic smartphone application use	12.43 (6.17)/1121	12.77 (6.25)/236	0.78 (0.44)
Problematic social media use	8.90 (3.81)/1121	9.61 (4.12)/236	2.56 (0.01)
Problematic gaming	12.82 (5.41)/1121	13.10 (5.49)/236	0.72 (0.47)
Time spent on smartphone	2.29 (2.60)/1121	2.57 (2.78)/236	1.48 (0.14)
Time spent on social media	1.11 (2.03)/1121	1.20 (1.65)/236	0.65 (0.52)
Time spent gaming	0.92 (2.00)/1121	1.02 (1.95)/236	0.72 (0.47)

Non-OW = group with nonoverweight; OW = group with overweight.

Table 3. Regression models investigating relationships between internet-related behaviors and psychological distress among children with or without overweight

Coefficient (SE)/ Standardized coefficient

	Fear of COVID-infection	Stress	Anxiety	Depression
<b>Group without overweight</b>				
Age	-0.016 (0.101)/ -0.005	-0.033 (0.061)/ -0.013	0.017 (0.048)/ 0.009	-0.072 (0.055)/ -0.033
Gender (Ref: boys)	-0.372 (0.229)/ -0.051	-0.141 (0.139)/ -0.025	0.116 (0.109)/ 0.027	-0.089 (0.124)/ -0.018
Perceived weight stigma	0.279 (0.065)/ 0.138***	0.426 (0.040)/ 0.274***	0.382 (0.032)/ 0.320***	0.398 (0.036)/ 0.293***
PSAU	-0.010 (0.027)/ -0.017	0.101 (0.017)/ 0.226***	0.032 (0.013)/ 0.092*	0.044 (0.015)/ 0.113**
PSMU	0.050 (0.04)/ 0.052	0.068 (0.024)/ 0.094**	0.049 (0.019)/ 0.088**	0.085 (0.022)/ 0.134***
PG	0.018 (0.031)/ 0.027	0.086 (0.019)/ 0.168***	0.086 (0.015)/ 0.221***	0.087 (0.017)/ 0.194***
<b>R<sup>2</sup> (Adj. R<sup>2</sup>)</b>	<b>0.030 (0.024)</b>	<b>0.335 (0.331)</b>	<b>0.298 (0.294)</b>	<b>0.307 (0.303)</b>
<b>Group with overweight</b>				
Age	0.485 (0.668)/ 0.050	-0.370 (0.161)/ -0.113*	-0.143 (0.114)/ -0.064	-0.303 (0.144)/ -0.108*
Gender	-1.798 (1.291)/ -0.099	0.014 (0.311)/ 0.002	0.062 (0.219)/ 0.015	0.021 (0.277)/ 0.004
Perceived weight stigma	0.592 (0.312)/ 0.141 <sup>#</sup>	0.346 (0.077)/ 0.241***	0.345 (0.054)/ 0.353***	0.442 (0.068)/ 0.359***
PSAU	-0.093 (0.150)/ -0.618	0.178 (0.036)/ 0.362***	0.107 (0.026)/ 0.323***	0.129 (0.032)/ 0.309***
PSMU	-0.232 (0.213)/ -0.106	0.287 (0.053)/ 0.388***	0.130 (0.037)/ 0.259***	0.157 (0.047)/ 0.248***
PG	0.286 (0.182)/ 0.175	-0.115 (0.044)/ -0.208**	-0.055 (0.031)/ -0.147 <sup>#</sup>	-0.067 (0.039)/ -0.140 <sup>#</sup>
<b>R<sup>2</sup> (Adj. R<sup>2</sup>)</b>	<b>0.047 (0.019)</b>	<b>0.451 (0.436)</b>	<b>0.410 (0.394)</b>	<b>0.404 (0.388)</b>
<b>All participants</b>				
Weight (Ref: non overweight)	0.297 (0.125)/ 0.067*	0.108 (0.055)/ 0.043 <sup>#</sup>	0.008 (0.042)/ 0.004	0.035 (0.049)/ 0.016
Age	0.024 (0.128)/ 0.005	-0.068 (0.058)/ -0.026	0.003 (0.044)/ 0.002	-0.094 (0.051)/ -0.042
Gender (Ref: boys)	-0.615 (0.286)/ -0.062*	-0.097 (0.128)/ -0.017	0.121 (0.098)/ 0.029	-0.056 (0.113)/ -0.011
Perceived weight stigma	0.326 (0.079)/ 0.122***	0.420 (0.036)/ 0.275***	0.377 (0.027)/ 0.330***	0.412 (0.032)/ 0.310***
PSAU	-0.021 (0.034)/ -0.027	0.113 (0.015)/ 0.246***	0.044 (0.012)/ 0.129***	0.058 (0.013)/ 0.145***
PSMU	0.008 (0.049)/ 0.007	0.105 (0.022)/ 0.143***	0.061 (0.017)/ 0.112***	0.097 (0.020)/ 0.154***
PG	0.057 (0.039)/ 0.062	0.055 (0.018)/ 0.105**	0.064 (0.014)/ 0.165***	0.062 (0.016)/ 0.138***
<b>R<sup>2</sup> (Adj. R<sup>2</sup>)</b>	<b>0.032 (0.026)</b>	<b>0.348 (0.344)</b>	<b>0.309 (0.305)</b>	<b>0.320 (0.316)</b>

<sup>#</sup> p<0.1; \* p<0.05; \*\* p<0.01; \*\*\* p<0.001

PSAU = problematic smartphone application use; PSMU = problematic social media use; PG = problematic gaming.

## Declarations

**Disclosure:** The authors declared no conflict of interest.

**Correspondence:** C.-Y. Lin, PhD, Department of Rehabilitation Sciences, Faculty of Health and Social Sciences, The Hong Kong Polytechnic University, 11 Yuk Choi Rd, Hung Hom, Hong Kong. **E-mail:** cylin36933@gmail.com; **Tel:** 852-2766-6755; **Fax:** 852-2330-8656

**Conflict of interests:** All the authors declare that there is no conflict of interest.

**Funding resources:** None.