Communicable Disease Pandemic: A Simulation Model Based on Community Transmission and Social Distancing

Sourav Kumar Bhoi

BPUT: Biju Patnaik University of Technology

Kalyan Kumar Jena

BPUT: Biju Patnaik University of Technology

Debasis Mohapatra

BPUT: Biju Patnaik University of Technology

Munesh Singh

Indian Institute of Information Technology: International Institute of Information Technology

Raghvendra Kumar

GIET: Gandhi Institute of Engineering and Technology

Hoang Viet Long (✉ hoangvietlong@tdtu.edu.vn)

Ton Duc Thang University <https://orcid.org/0000-0001-9883-9506>

Research Article

Keywords: COVID-19, Community Transmission, Social Distancing, Simulation, GNU Octave, Number of Infected, Number of Health

Posted Date: July 6th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-274502/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Communicable Disease Pandemic: a simulation model based on community transmission and social distancing

Sourav Kumar Bhoi · Kalyan Kumar Jena · Debasis Mohapatra · Munesh Singh · Raghvendra Kumar · Hoang Viet Long

Received: date / Accepted: date

Abstract Communicable disease pandemic is a severe disease outbreak all over the countries and continents. Swine Flu, HIV/AIDS, COVID-19 (CORONA VIRUS DISEASE-19), etc. are some of the global pandemics in the world. The major cause of becoming pandemic is community transmission and lack of social distancing. Recently COVID-19 is such an largest outbreak all over the world. This disease is a communicable disease which is spreading fastly due to Community Transmission, where the affected people in the community affects the healthy people in the community. Government is taking precautions by imposing Social Distancing in the countries or state to control the impact of COVID-19. Social Distancing can reduce the community transmission of COVID-19 by reducing the number of infected persons in an area. This is performed by staying at home and maintaining social distance with people. It reduces the density of people in an area by which it is difficult for the virus to spread from one person to other. In this work, the community transmission

¹Sourav Kumar Bhoi, ¹Kalyan Kumar Jena, ¹Debasis Mohapatra, ²Munesh Singh,

³Raghvendra Kumar, ^{4,5,*}Hoang Viet Long

¹High Performance Computing Lab

Department of Computer Science and Engineering

Parala Maharaja Engineering College (Govt.), Berhampur 761003, BPUT University, India

E-mail: sourav.cse@pmec.ac.in, kalyan.cse@pmec.ac.in, debasis.cse@pmec.ac.in

²Department of Computer Science and Engineering

Indian Institute of Information Technology Design & Manufacturing, Kancheepuram, India

E-mail: muneshpal24@gmail.com

³Department of Computer Science and Engineering,

GIET University, India

E-mail: raghvendra@giuet.edu

⁴Division of Computational Mathematics and Engineering, Institute for Computational Science,

Ton Duc Thang University, Ho Chi Minh City, Vietnam;

⁵Faculty of Mathematics and Statistics,

Ton Duc Thang University, Ho Chi Minh City, Vietnam;

E-mail: hoangvietlong@tdtu.edu.vn

is presented using simulations. It shows how an infected person affects the healthy persons in an area. Simulations also show how social distancing can control the spread of COVID-19. The simulation is performed in GNU Octave programming platform by considering number of infected persons and number of healthy persons as parameters. Results show that using the social distancing the number of infected persons can be reduced and healthy persons can be increased. Therefore, from the analysis it is concluded that social distancing will be a better solution of prevention from community transmission.

Keywords COVID-19 · Community Transmission · Social Distancing · Simulation · GNU Octave · Number of Infected · Number of Healthy

1 Introduction

Communicable diseases are the fast growing disease outbreaks which happens when a person comes in contact of another diseased person and it becomes community transmission. Nowadays, COVID-19 is a dangerous disease spreading rapidly all over the world [1] by means of community transmission. Community transmission means when the disease is transferred from a local person to another person and spreads over the society. World Health Organization (WHO) has declared it as global pandemic as it is affecting peoples life and causing many deaths. Currently, according to the reports of WHO there are 106M confirmed cases, 2.31M deaths, and it affects nearly 208 countries. The developed countries like USA, Spain, Italy, Germany, France, etc. are mostly affected of this. It is rapidly affecting the health as well as economy of the country.

The people affected with COVID-19 experience respiratory illness and they recover themselves after a certain time without any special treatment [1]. The older people needs special treatment if they have other medical problems such as heart disease, diabetes, respiration problem, blood pressure, and cancer. The common symptoms of COVID-19 include fever, tiredness, dry cough, and other symptoms include shortness of breath, pains and aches, sore throat, and some people may report diarrhoea, runny nose, and nausea. The people with mild cases may contact medical helpline or can use self-isolation. The people with critical case should go immediately to hospitals for proper and sudden treatment.

COVID-19 can be prevented by following some common practices such as [1]:

- **Cleaning:** Washing hands and fingers are regularly with soap and water or cleaning them using alcohol based hand rubs.
- **Social Distancing:** Maintaining distance from others at least 1 meter from other (persons sneezing or coughing or any unknown persons)
- **Avoid Touching Face:** Avoiding touching eyes, nose, ears, and mouth.
- **Cover:** Covering the mouth with hygienic masks or any other material that can work as mask.

- **Refrain:** Refraining from smoking and other activities which weakens the respiratory system.
- **Practice:** Practicing the above points in a right manner to serve humanity.

From the above points, it is observed that social distancing will be a better solution and key to combat with this pandemic. From the reports of WHO, the countries which implemented this effectively for weeks have a great control over the disease. Social distancing means keeping ourselves away from a person or group of persons. This can be performed by staying at home for weeks by following government rules. That means density of the people in an area is reduced to control the number of infected persons. If there are no persons or very little density in an area then the chance of COVID-19 spread can be controlled in a massive way.

The major contributions in this work are stated as follows:

1. In this work, the community transmission is presented using simulations. It shows how an infected person affects the healthy persons in an area. This is performed by random walk model where an infected person is randomly moved in an area A by affecting the healthy persons in the neighborhood. The person moves from point A to point B , point B to point C , and then from point C to point D .
2. Simulations also show how social distancing can control the spread of COVID-19. This is performed by reducing the density of healthy persons in that area, and analyzing the number of healthy persons and infected persons in that area.
3. The simulation is performed in GNU Octave programming platform by considering number of infected persons and number of healthy persons as parameters.
4. Results show that using the social distancing the number of infected persons can be reduced and healthy persons can be increased.

The rest of the sections is presented as follows. Section 2 shows the related works. Section 3 presents the problem and methodology to solve the problem. Section 4 presents the simulations and results. Section 5 presents the conclusion and future scope.

2 Related Works

Many research works are currently going on to combat COVID-19 which rapidly which is rapidly increasing day by day all over the world. Many updated articles are published on COVID-19 regarding spreading, medicines, vaccinations, preventive measures, etc. Chen et al. [2] present the research on control of COVID-19 in China during the new year. Anderson et al. [3] presented the influence of country based mitigation methods over COVID-19 spread. Remuzzi et al. [4] presents the articles on social distancing in UK due to rise of deaths over the world. Wilder-Smith et al. [5] proposed the article by showing

how the old methods such as isolation, quarantine, social distancing, community containment reduces the effect of COVID-19. Wu et al. [6] presents the characteristics of lesson from COVID-19 outbreak at China. Dalton et al. [7] presents that the low cost social distancing can reduce the severity of cars in COVID-19. Roosa et al. [8] predicted the spread of COVID-19 in China from 5th February 2020 to 24th February 2020. Leung et al. [9] presented the mass masking for guidance in COVID-19 spread. Ferguson et al. [10] presented the impact of non-pharmaceutical interventions over the death rate in COVID-19. Thomson [11] presented a survey in social distancing that how it works in this situation. Roosa et al. [12] forecasted the spread of COVID-19 in Guangdong and Zhejiang from Feb 13-Feb 23 2020. YAP et al. [13] presented the social distancing and lockdowns report on COVID-19. Stein [14] presented the idea about rationally layered social distancing. Jiang et al [15] presented a review of clinical characteristics on COVID-19. Many such works are presented in regarding social distancing and COVID-19 [16–26, 1, 27, 15, 28–31]. From the study, according to our knowledge, no such work is performed by the researchers based on simulations of random walk by the infected persons or use of social distancing to reduce the infected persons in an area.

3 Methodology

This section describes clearly about the problem definition and the methodology or solution to solve it.

3.1 Problem Statement

Let, an infected person moves randomly in an area A of size $l \times b$ where l and b are the length and breadth respectively. The person moves from point A to point B , point B to point C , and then from point C to point D . In this move, the person affects another healthy person in that area based on neighborhood with a certain distance d . The main problem is to present this situation with an illustration and solving the problem using social distancing solution by reducing the number of infected persons and increasing the number of healthy persons in an area.

3.2 Methodology

In this section, the above problem is solved using an illustration and social distancing approach. To represent the problem with an illustration an area of A is considered. In that area, let there are h number of healthy persons and we consider each person in that area is healthy. We assumed this situation because no COVID-19 cases appears in that area till now. So, the density of that area is considered to be λ where it is represented as:

Fig. 1: A COVID-19 infected person infecting healthy persons in its neighborhood.

Fig. 2: Use of social distancing to reduce the number of infected persons.

$$\lambda = h + n; \quad (1)$$

where, n is the number of infected persons. From Fig. 1, it is observed that a COVID-19 infected person (orange) starts his journey from point A to point B . That means in this example, the coordinates of point A and point B are

randomly generated. The points are generated using pseudorandom number generation function, that generates the coordinates (x, y) randomly as follows:

$$x \text{ or } y = randi([a \ b], 1, c) \quad (2)$$

where, a and b are the numbers between which the random number is generated and c is the number of coordinates generated. The coordinates of all persons in that are is generated using Eq. (2).

The infected person (orange) affects those persons who are in the neighborhood. The *neighborhood* is defined as the persons nearer with maximum threshold distance d . The distance is calculated by using the Euclidean distance formula as follows:

$$d = \sqrt{(x1 - x2)^2 + (y1 - y2)^2} \quad (3)$$

From, Fig. 1 it is observed that the person affects $\{n_1, n_2, n_3, n_4, n_5, n_6, n_7\}$ in the journey because they are all in the neighborhood. 7 number of healthy persons are affected due to a single person. However, if the density of people in this area is less than the the number of infected case will be less than 7. This can be performed by social distancing where the density of persons in an area is reduced.

Social Distancing is the solution to reduce the number of infected persons in an area by reducing the density of people in an area. Let, say in an area A the density of people is λ where all persons are healthy initially. Then, after the move of that infected person n_k number of healthy persons get affected and the rest number of healthy persons are represented as:

$$h_1 = \lambda - n_k \quad (4)$$

where h_1 is the number of healthy persons after movement of that infected person from A to B . From Eq. (1) and Eq. (4), $h_1 < h$. By applying social distancing as shown in Fig. 2, the number of healthy persons can be increased and infected persons can be decreased. According to Fig. 2, the person only affects n_2 and n_5 because the density λ reduces. The total number of healthy persons after social distancing is represented as:

$$h_2 = \lambda - n_p \quad (5)$$

where n_p is the number of infected after social distancing. From Eq. (4) and Eq. (5) $h_2 > h_1$. Hence, by using social distancing in this example the COVID-19 spread can be controlled.

Algorithm 1 shows the implementation of community transmission and the impact of social distancing over this. Step-2 to Step-4 shows the deployment of λ (number of healthy persons) in area A by generating the coordinates randomly using `randi()`. Step-6 to Step-11 moves the person from point A to point B . Step-13 to Step-20 moves the person from point B to point C . Step-22 to Step-27 moves the person from point C to point D . Step 29 to step-40 shows the implementation of community transmission and social distancing

with distance d . The number of infected persons n and number of healthy persons h are calculated using step-41 and step-46 respectively. This method can be a better solution for reducing the number of infected case. This is analyzed by simulations which is presented in the next section.


```

Input:  $A, \lambda, d$ 
Output:  $n$ : number of infected persons,  $h$ : number of healthy persons
1  /* Deployment of number of healthy persons in area A */
2  xc=randi([a b],1, $\lambda$ )
3  yc=randi([a b],1, $\lambda$ )
4  plot(x,y,'bo')
5  /* random walk model starts with point A to point B */
6  c1=0;
7  for k=1:a1 do
8      c1=c1+1;
9      x(k)=c1;
10     y(k)=c1;
11 end
12                                     /* point B to point C */
13 d=a1;
14 e=a1;
15 for k=a1+1:a2 do
16     d=d+1;
17     e=e-1;
18     x(k)=d;
19     y(k)=e;
20 end
21                                     /* point C to point D */
22 g=0;
23 for k=a2+1:a3 do
24     g=g+1;
25     x(k)=a2;
26     y(k)=g;
27 end
28     /* Community Transmission and Social Distancing with
        distance d */
29 count=0;
30 for i=1:position do
31     for j=1: $\lambda$  do
32         d=sqrt((x(i)-xc(j))^2 + (y(i)-yc(j))^2 );
33         if d<10 && d!=0 then
34             circle(xc(j),yc(j),5);
35             count=count+1;
36             buffer(count,1)=xc(j);
37             buffer(count,2)=yc(j);
38         end
39     end
40 end
41 C=unique(buffer);
42 [m,n]=size(C);
43                                     /* Number of infected persons n */
44 n=round(m/2);
45                                     /* Number of healthy persons h */
46 h= $\lambda$ -n;

```

Algorithm 1: Community Transmission and Social Distancing

4 Results and Discussion

The simulation is performed in GNU Octave programming platform. It is based on MATLAB programming. This platform is installed in a machine of Intel core i3 processor of speed 2.4 GHz, RAM 8 Gb, and 64 bit Windows operating system. In this simulation, the community transmission is implemented by moving an infected person in an area A where the area has a density of healthy persons. The person moves from point A to point B , point B to point C , and then from point C to point D . The points are generated randomly to implement the random walk. The infected person affects those persons in the move who are in the neighborhood. This can be reduced by the implementation of social distancing where the density (number of healthy people) of an area is reduced. In this section, a study has been made by varying the density of people in an area. The main parameters are considered to be number of healthy persons and number of infected persons. To setup the simulation, major parameters are set and represented in Table 1 as follows.

Table 1: Parameter and Value

Sl. no.	Parameter	Value
1	A	$1000 \times 1000 \text{ m}^2$
2	λ	100-1000
3	d	10 m
4	Mobility pattern of infected person	Random Walk

In this simulation, an area of $1000 \times 1000 \text{ m}^2$ is considered for implementation of community transmission and social distancing. The community transmission is implemented by deploying λ density of healthy people in the area using `randi()` to generate the coordinates. The position of the healthy persons generated are assumed to be the current position. An infected person moves randomly by starting at a random position and moves to other random positions straightly. It moves in the route $A - B - C - D$. The person transfers the disease to that person if that person's position is at a distance lower than 10 m. After that, social distancing is implemented by reducing the density from $\lambda = 1000$ to $\lambda = 200$ and observing the impact of this on number of infected persons and number of healthy persons. The simulation results are average of 10 simulation runs.

From Fig. 3, it is observed that when the infected person moves from point A to point B by varying the density from 1000 to 200 then the number of infected people and number of healthy people are observed to be $\{14, 11, 7, 4, 2\}$ and $\{986, 989, 993, 996, 998\}$ respectively. From Fig. 4, it is observed that when the infected person moves from point A to point C by varying the density from 1000 to 200 then the number of infected people and number of healthy people

Fig. 3: Simulation results of impact of social distancing on community transmission when the infected moves from point A to point B. Green line: Move of infected person, Red circle: infected people, and Blue: Healthy people.

are observed to be $\{18, 18, 20, 6, 2\}$ and $\{982, 982, 980, 994, 998\}$ respectively. From Fig. 5, it is observed that when the infected person moves from point A to point D by varying the density from 1000 to 200 then the number of infected people and number of healthy people are observed to be $\{24, 20, 11, 11, 5\}$ and $\{976, 980, 989, 989, 995\}$ respectively. From the above results represented in Fig.

Fig. 4: Simulation results of impact of social distancing on community transmission when the infected moves from point A to point B then point C. Green line: Move of infected person, Red circle: infected people, and Blue: Healthy people.

Fig. 5: Simulation results of impact of social distancing on community transmission when the infected moves from point A to point B, point B to point C, then point D. Green line: Move of infected person, Red circle: infected people, and Blue: Healthy people.

Fig. 6: Impact of social distancing on community transmission when the infected moves from point A to point B.

Fig. 7: Impact of social distancing on community transmission when the infected moves from point A to point B then point C.

Fig. 8: Impact of social distancing on community transmission when the infected moves from point A to point B, point B to point C, then point D.

Table 2: Impact of social distancing on community transmission.

Random Walk	Parameters	$\lambda = 1000$	$\lambda = 800$	$\lambda = 600$	$\lambda = 400$	$\lambda = 200$
point A-point B	infected	14	11	7	4	2
	healthy	986	989	993	996	998
point A-point B-point C	infected	18	18	20	6	2
	healthy	982	982	980	994	998
point A-point B-point C-point D	infected	24	20	11	11	5
	healthy	986	980	989	989	995

6, Fig. 7, and Fig. 8 it is concluded that when the density of people in an area decreases the number of infected people reduces and number of healthy persons increases. It is also concluded that when that persons moves more then it is affecting more people. Therefore, social distancing should be implemented well by the government to reduce the number of infected cases. Table 2 shows the impact of social distancing on community transmission, when density of people decreases number of infected people decreases and number of healthy persons increases.

5 Conclusion

In this work, the community transmission is presented using simulations. It shows how an infected person affects the healthy persons in an area. This is performed by random walk model where an infected person is randomly moved in an area A by affecting the healthy persons in the neighborhood. The

person moves from point A to point B , point B to point C , and then from point C to point D . Simulations also show how social distancing can control the spread of COVID-19. This is performed by reducing the density of healthy persons in that area, and analyzing the number of healthy persons and infected persons in that area. Results show that using the social distancing the number of infected persons can be reduced and healthy persons can be increased. It is also concluded that when the person moves more in that area he affects more people. Therefore, social distancing, isolation, and home quarantine will be better solutions to be implemented well by the government to reduce the number of infected cases.

Ethics declarations

Conflict of interest

On behalf of all authors, the corresponding author states that there is no conflict of interest.

Authorship contributions

All authors contributed equally to this work.

Data availability

Data can be shared if needed.

References

1. World Health Organization et al. Coronavirus disease (covid-19) outbreak. URL <https://www.who.int/emergencies/diseases/novel-coronavirus-2019>, 2020.
2. Simiao Chen, Juntao Yang, Weizhong Yang, Chen Wang, and Till Barnighausen. Covid-19 control in china during mass population movements at new year. *The Lancet*, 2020.
3. Roy M Anderson, Hans Heesterbeek, Don Klinkenberg, and T Déirdre Hollingsworth. How will country-based mitigation measures influence the course of the covid-19 epidemic? *The Lancet*, 395(10228):931–934, 2020.
4. Andrea Remuzzi and Giuseppe Remuzzi. Covid-19 and italy: what next? *The Lancet*, 2020.
5. A Wilder-Smith and DO Freedman. Isolation, quarantine, social distancing and community containment: pivotal role for old-style public health measures in the novel coronavirus (2019-ncov) outbreak. *Journal of travel medicine*, 27(2):taaa020, 2020.
6. Zunyou Wu and Jennifer M McGoogan. Characteristics of and important lessons from the coronavirus disease 2019 (covid-19) outbreak in china: summary of a report of 72 314 cases from the chinese center for disease control and prevention. *Jama*, 2020.
7. Craig Dalton, Stephen Corbett, and Anthea Katelaris. Pre-emptive low cost social distancing and enhanced hygiene implemented before local covid-19 transmission could decrease the number and severity of cases. *SSRN*, 3549276, 2020.

8. K Roosa, Y Lee, R Luo, A Kirpich, R Rothenberg, JM Hyman, P Yan, and G Chowell. Real-time forecasts of the covid-19 epidemic in china from february 5th to february 24th, 2020. *Infectious Disease Modelling*, 5:256–263, 2020.
9. Chi Chiu Leung, Tai Hing Lam, and Kar Keung Cheng. Mass masking in the covid-19 epidemic: people need guidance. *The Lancet*, 2020.
10. Neil M Ferguson, Daniel Laydon, Gemma Nedjati-Gilani, Natsuko Imai, Kylie Ainslie, Marc Baguelin, Sangeeta Bhatia, Adhiratha Boonyasiri, Zulma Cucunubá, Gina Cuomo-Dannenburg, et al. Impact of non-pharmaceutical interventions (npis) to reduce covid-19 mortality and healthcare demand. *London: Imperial College COVID-19 Response Team*, March, 16, 2020.
11. Oxford Analytica. Covid-19 social distancing will be in place for months. *Emerald Expert Briefings*, (oxan-es).
12. Kimberlyn Roosa, Yiseul Lee, Ruiyan Luo, Alexander Kirpich, Richard Rothenberg, James M Hyman, Ping Yan, and Gerardo Chowell. Short-term forecasts of the covid-19 epidemic in guangdong and zhejiang, china: February 13–23, 2020. *Journal of Clinical Medicine*, 9(2):596, 2020.
13. Jason Chin-Huat YAP, Ian Yi Han ANG, Sharon Hui Xuan TAN, I Jacinta, CHEN Pei, Ruth Frances LEWIS, YANG Qian, Rowena Kah Sin YAP, Bob Xian Yi NG, and Hao Yi TAN. Covid-19 science report: Social distancing & lockdowns. 2020.
14. Richard Stein. Covid-19 and rationally layered social distancing. *International Journal of Clinical Practice*, page e13501, 2020.
15. Fang Jiang, Liehua Deng, Liangqing Zhang, Yin Cai, Chi Wai Cheung, and Zhengyuan Xia. Review of the clinical characteristics of coronavirus disease 2019 (covid-19). *Journal of General Internal Medicine*, pages 1–5, 2020.
16. Elisabeth Mahase. Covid-19: Uk starts social distancing after new model points to 260 000 potential deaths, 2020.
17. Anthony S Fauci, H Clifford Lane, and Robert R Redfield. Covid-19—navigating the uncharted, 2020.
18. Shahul H Ebrahim and Ziad A Memish. Covid-19: preparing for superspreader potential among umrah pilgrims to saudi arabia. *The Lancet*, 2020.
19. Marc Lipsitch, David L Swerdlow, and Lyn Finelli. Defining the epidemiology of covid-19—studies needed. *New England Journal of Medicine*, 2020.
20. Kenji Mizumoto and Gerardo Chowell. Transmission potential of the novel coronavirus (covid-19) onboard the diamond princess cruises ship, 2020. *Infectious Disease Modelling*, 2020.
21. Annelies Wilder-Smith, Calvin J Chiew, and Vernon J Lee. Can we contain the covid-19 outbreak with the same measures as for sars? *The Lancet Infectious Diseases*, 2020.
22. George Thomson. Covid-19: social distancing, ace 2 receptors, protease inhibitors and beyond? *International journal of clinical practice*, page e13503, 2020.
23. Angel N Desai and Payal Patel. Stopping the spread of covid-19. *JAMA*, 2020.
24. John N Nkengasong and Wessam Mankoula. Looming threat of covid-19 infection in africa: act collectively, and fast. *The Lancet*, 395(10227):841–842, 2020.
25. Rose Baker. Reactive social distancing in a sir model of epidemics such as covid-19. *arXiv preprint arXiv:2003.08285*, 2020.
26. Yuzhen Zhang, Bin Jiang, Jiamin Yuan, and Yanyun Tao. The impact of social distancing and epicenter lockdown on the covid-19 epidemic in mainland china: A data-driven seiqr model study. *medRxiv*, 2020.
27. World Health Organization et al. Coronavirus disease 2019 (? covid-19)? : situation report, 51. 2020.
28. Qiurong Ruan, Kun Yang, Wenxia Wang, Lingyu Jiang, and Jianxin Song. Clinical predictors of mortality due to covid-19 based on an analysis of data of 150 patients from wuhan, china. *Intensive Care Medicine*, pages 1–3, 2020.
29. Yan Bai, Lingsheng Yao, Tao Wei, Fei Tian, Dong-Yan Jin, Lijuan Chen, and Meiyun Wang. Presumed asymptomatic carrier transmission of covid-19. *Jama*, 2020.
30. Coronavirus Pneumonia Emergency Response Epidemiology Novel et al. The epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (covid-19) in china. *Zhonghua liu xing bing xue za zhi= Zhonghua liuxingbingxue zazhi*, 41(2):145, 2020.

-
31. Jianfeng Xie, Zhaohui Tong, Xiangdong Guan, Bin Du, Haibo Qiu, and Arthur S Slutsky. Critical care crisis and some recommendations during the covid-19 epidemic in china. *Intensive Care Medicine*, pages 1–4, 2020.