

Aedes Aegypti: Morphology the Egg and Embryonic Development

Ana Paula Miranda Mundim-Pombo

Universidade de Sao Paulo Faculdade de Medicina Veterinaria e Zootecnia

Marisol León

USP FMVZ: Universidade de Sao Paulo Faculdade de Medicina Veterinaria e Zootecnia

Hianka Jasmyne Costa de Carvalho

USP FMVZ: Universidade de Sao Paulo Faculdade de Medicina Veterinaria e Zootecnia

Durvanei Augusto Maria

Butantan Institute: Instituto Butantan

Maria Angelica Miglino (✉ miglino@usp.br)

Universidade de Sao Paulo Faculdade de Medicina Veterinaria e Zootecnia <https://orcid.org/0000-0003-4979-115X>

Research

Keywords: *Aedes aegypti*, Egg morphology, Embryonic development, Ultrastructural analysis.

Posted Date: March 3rd, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-274534/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: *Aedes aegypti* (Diptera: *Culicidae*) has great relevance in public health worldwide due to its performance as a vector of arboviruses, which makes the knowledge about its cycle fundamental. It is known that the acquaintance of embryonic kinetics has great potential to assist in the development of new vector control technologies and, consequently, in the control of arboviruses.

Methods: This research proposes to analyze the egg morphology and embryonic development of *Aedes aegypti*. Eggs from the insectary of the Institute of Biomedical Sciences of the University of São Paulo (n = 46) were used. The methodological procedure involved morphological and ultrastructural analyzes using equipment for optical microscopy (light and confocal) and electronic (scanning). Eggs and embryos were observed in initial, intermediate and final thirds of development, kept at a temperature of 28°C, 1°C until collection for processing.

Results: The embryos had different morphological characteristics according to the stage of gestation, in the initial and intermediate stages of development. In the initial third, the presence of primordial epithelium and characteristics suggestive of intense cellular activity were found.

Conclusion: According to the results obtained in the eggs, it is concluded that the great resistance of the chorion in the embryo is a factor that generates difficulties for microscopic analysis of the embryo (mainly in fixation), with a strong protective barrier.

Background

The *Aedes aegypti* has great importance in public health worldwide, due mainly to its involvement in the transmission of arboviruses (Arthropod-borne-viruses) such as Dengue (WHO 2018), *Chikungunya* (Charrel et al. 2014; WHO 2017) and *Zika* (WHO 2018). The simultaneous circulation of its etiological agents in various locations characterizes the epidemiological scenario of these diseases, which increases the risk of spread, the possibility of coinfection and transmission of more than one virus at a time in humans, complicating diagnosis and recovery and the relevance of this vector (Gloria-Soria et al. 2016; Chaves et al. 2018; Carrillo-Hernández et al. 2018).

The lack of an effective vaccine for vast majority of arboviruses (Ferguson et al. 2016), makes clear that there is a need for high levels of vector-directed control (Brazil 2015; Lima-Camara 2016), based on epidemiological and entomological control (WHO 2018; WHO 2017; WHO 2018). Nevertheless, keeping mosquito populations under control is still inefficient. In these cases, eggs are of special importance, as they are easily located, and they are very resistant and can remain viable for more than a year (Brazil 2015), which justifies the fact that the new technologies are destined to destroy eggs or to fail the hatch (Caragata et al. 2019). Therefore, new in-depth knowledge of the characteristics of eggs and the embryonic development would contribute to creating the basis for new control and eradication strategies.

In general, it is known that eggs have an external coating structure called a chorion, with a protective function, allowing gas exchange and minimizing water loss. The chorion is made of distinct layers: the endochorion and exochorion (Clements 1992), these structures are known to have different characteristics between the species and the genus *Aedes*, originating from Gwalior, India (Suman et al. 2011). Inside, embryogenesis occurs after fusion of the female pro-nucleus with the male pro-nucleus, after the oocyte passes through the ovarian tube (Fonseca, Gomes and Araújo 2012).

Previous studies reveal that the external structure of eggs (exochorion) from insects in general, differentiates in several orders such as culicids of medical interest (Obara et al. 2007; Famesi *et al.* 2017; Faull and Williams 2016; Mello et al. 2018; Alencar et al. 2003). Insects, especially those considered to be of medical importance, show phenotypic variation when subjected to environmental changes (Reinhold, Lazzari and Lahondère 2018). Currently, the evolutionary significance of genetic and epigenetic assimilation is widely discussed (Mathews *et al.* 2018). *Drosophila melanogaster* is considered the model insect for the study of embryonic development of insects, (Fonseca, Gomes and Araújo 2012), for which the precarious description about the embryonic morphology of *Aedes aegypti* has been attributed to the problems related to the permeability of the egg, plus the embryonic development has not yet been properly reported. This study was in charge of describing characteristics morphological of *Aedes aegypti* eggs and describing your embryonic development.

Material And Method

Obtaining biological material

Aedes aegypti eggs were made available by the Parasitology Department of the Institute of Biomedical Sciences of the University of São Paulo, USP, collected during the period from November / 2014 to February / 2015 (n=46). For the analyses of the embryonic development, there were synchronization of egg-laying, three groups of eggs of this insect were collected at different times of embryogenesis, without ending/initial, intermediate and final stages of its development. Embryogenesis is considered be completed in approximately 61.6 hours when the temperature is 28 ° C, according to a study by FARNESI *et al.* (2017). All samples were kept at a temperature of 28°C, and this reason does remain in the original insectary from the laying of eggs until the moment of collection for processing. This pattern was maintained for three different collection times. Each group of samples collected, had its development interrupted when submitted to the tissue fixation procedure. Morphological analysis was performed using resources such as optical microscopy (light and laser scanning confocal) and electronic microscopy (transmission and scanning).

Optical Microscopy: Light

Samples were fixed in a 10% formaldehyde solution. After complete fixation, they were dehydrated in a series of ethanol in increasing concentrations (70 to 100%) and diaphanized in xylol, with subsequent inclusion in histological paraffin. 3µm thick cuts were made in the microtome (Leica, German) and

stained with Hematoxylin-Eosin. The images were obtained through the Nikon Eclipse E-800 light microscope at the Advanced Diagnostic Imaging Center / FMVZ-USP.

Optical Microscopy: Laser Scanning Confocal

The eggs were initially processed with water washing, followed by immersion in 3% sodium hypochlorite until clarification (approximately 30 minutes), with subsequent washing in PBS, 0.02% triton for 5 minutes and again washing in PBS, the material then it was fixed in 3.7% formaldehyde for 20 minutes, after this procedure a new washing sequence was performed with PBS, permeabilization using 1% triton at room temperature and washing with PBS twice. The incubation was carried out in a dark room for 60 minutes and phalloidin (FITC) was used. RNase was added in the final 30 minutes of incubation. After this treatment, the embryos were again washed with PBS and the nuclei were stained with Propidium Iodide. Fluorescent images were obtained by laser scanning confocal microscopy (Zeiss LSM 510) at the Cellular and Molecular Biology laboratory (BioCeM) of the Institute of Biomedical Sciences-USP.

Transmission Electron Microscopy

The samples were fixed in 2% glutaraldehyde, post-fixed in a 1% osmium tetroxide solution at 4 ° C and a 5% aqueous solution of uranyl acetate at room temperature. Then, the samples were dehydrated in ethanol, increasing concentration, immersed in propylene oxide and soaked in Spurr resin. For light microscopy, semi-thin sections were cut in a Cut Ultra Reichert® ultramicrotome and stained with a 1% toluidine blue solution. The thin 90 nm sections were cut and collected in 200 mesh (Sigma ®) and contrasted with a 4% uranyl acetate solution and a 0.4% aqueous solution of lead citrate. The grids were examined using a transmission electron microscope at the Advanced Center for Diagnostic Imaging, Faculty of Veterinary Medicine and Animal Science (FMVZ) -USP. Specifically, for the analysis of the semi-fine cut, the eggs were subjected to previous processing with the washing in water, followed by immersion in sodium hypochlorite 3% until its clarification (approximately 30 minutes), with subsequent washing in PBS, 0.02 triton % for 5 minutes and again washed in PBS, only then were they fixed in glutaraldehyde giving continuity to the processing.

Scanning Electron Microscopy

The biological material was fixed in a modified Karnovsky solution (5% glutaraldehyde and 4% paraformaldehyde in 0.1 M cacodylate buffer, pH 7.2), followed by washing in sodium cacodylate buffer, pH 7.2 and posterior fixation in tetroxide osmium (OsO₄) at 1% in 0.2M sodium cacodylate buffer. After carrying out a new series of washing and cleaning controls, the eggs were dehydrated in an increasing series of ethanol to absolute ethanol (50%, 75%, 90% and 100%). After passing at a critical point, the material was mounted on Stubs using double-sided carbon adhesive tape followed by gold-plated coverage, using the sputtering system, and the FEI scanning electron microscope was used to analyze the material. Quanta 250 at the Cell Biology Laboratory of the Butantan Institute. - SP, Capital where they were photomicrographed for further analysis.

Morphometric analysis

Morphometric analysis of the eggs was performed from the measurements of the images obtained in Scanning Electron Microscopy. Linear dimensions of the egg were considered for this analysis:

- Length - distance between the micropyle and the opposite end, Width - largest distance perpendicular to the length,
- Egg index - length to width ratio,
- Diameter of the microcapillary disc - diameter of the anterior structure in the egg

After obtaining measurements were perform analysis of measures of central tendency (mean) and measures of dispersion (standard deviation, maximum and minimum values). Statistical analyzes were performed using the Graph Prism software, 95% Confidence Interval was considered.

Results

Eggs

Macroscopically, eggs of *Aedes aegypti* are whitish in color at the time of oviposition, which quickly becomes black. Some characteristics were identified, such as shiny appearance, tapered extremities, bilateral symmetry, in addition to a flattened surface opposite to a convex surface (Fig. 1A).

Morphometrically we observed that the linear dimension of the population of eggs referring to the length was $581.45 \pm 39.73 \mu\text{m}$ and the width of 175.36 ± 11.59 . The egg index (length / width ratio) was 3.32 ± 0.26 . Concerning the measurement of the micropyle (DM) disk diameter of the evaluated eggs, it remained at $18.75 \pm 1.92 \mu\text{m}$ (Table 1). The results were also compared with those from the studies by Suman et al., (13), Linley (22) and Faull (Fig. 1B).

Table 1

Morphometric parameters related to the linear size of the population of *Aedes aegypti* eggs coming from the municipality of São Paulo-SP in relation to measures of central tendency and dispersion. [This Table 1 should be attached between the 236 and 237 lines]

Atributes Morphometric		Central tendency and dispersion measures			
		Mean	IC95%	Maximum Value	Minimum Value
Eggs collected in Sao Paulo	Length	581.45 ± 39.73	569.65-593.25	655.20	521.40
	Width	175.36 ± 11.59	171.92-178.80	199.60	156.50
	Length/width	3.32 ± 0.26	3.24-3.40	4.13	2.78
	Diameter	18.75 ± 1.92	18.18-19.32	22.18	14.27

Table 2. Comparative analysis of the morphometric findings of *Aedes aegypti* eggs identified in this study and the results of Suman et al, (2011), Linley (1989), Faull and Williams (2016).

	<i>A. aegypti</i> ^{a *}	<i>A. aegypti</i> ^{b **}	<i>A. aegypti</i> ^{c ***}	<i>A. aegypti</i> ^{d ****}	<i>A. aegypti</i> ^{e *****}
Length	581.45±39.73	625.65±19.91	670.2±7.2	554.41±36.56	562.62 ± 30.85
Width	175.36±11.59	183.30±11.04	186.3±2.2	167.65±7.05	160.15 ± 9.73
Length/Width	3.32±0.26	————	3.61± 0.05	3.31±0.18	3.52 ± 0.27
Diameter of micropilar disc	18.75±1.92	————	————	33.49 ± 3.9	34.19 ± 5.4

a.Mundim-Pombo et al b.Suman et al, 2011 c. Linley,1989 d. Faull and Wiliams.,2016 e. Faull and Williams, 2016

* Population of São Paulo eggs, **Population of India eggs ***Population of Florida USA eggs **** Population of Cairns-Australia eggs *****Population of Charters Towers eggs - Australia

Ultrastructural SEM showed that the extremities are characterized by poles, the anterior pole, where the entire micropylar apparatus (disc/crown of the micropyle, sectors of the disc of the micropyle and micropyle) is located and is slightly prominent when compared to the opposite pole. The microcapillary device maintains a prominent and continuous circle shape (Figs. 1C and 1D), while the posterior pole is more tapered in relation to the opposite side (Figs. 1C and 1E). In the external coating of the eggs, regularity was identified about the aspect of distribution and shape of their cells, most of them maintaining a hexagonal shape (Figs. 1C and 1E).

In the central region of the chorionic cells, larger diameter tubers are identified. The tubers are symmetrically arranged, there are small peripheral tubers arranged in an organized manner around larger central tubers (Fig. 1F, 1G and 1H). Both appear prominently in the exochorion, thus giving the appearance of high relief or texture.

Transmission microscopy was used in a complementary manner to analyze the chorion (Fig. 1I-L), where it was possible to identify the ornamentations of the exochorion and confirm that the impermeable characteristic of this structure prevents the entry of fixers and resin inside the cell in the sample processing, causing the compromise of these cells. Another finding in relation to the exochorion is that it was more electron dense when compared to its adjacent layer, the endochorion.

The impermeability characteristic of chorion drew attention, which made the analyzes related to it, as being extremely difficult and generating several repetitions in the different stages of this study.

Embryo

The embryonic development of *Aedes aegypti* occurs inside eggs of up to 1 mm in length (Fig. 1A), dark in color and capable of adhering firmly to various substrates. In the period that comprises the initial third of the development of the *Aedes aegypti* embryo, it was possible to verify the scarce presence of bristles in the cephalic region and relatively homogeneous structures in terms of their general morphological aspects. Rupture of the chorion (Figs. 2A and 2B), which in turn will open the area outside the egg and release the larva. The identification of this region as being the cephalic region is possible because this region coincides with the micropillar apparatus.

In the histological analysis, elements characteristic of the ornamentation of the exochorion were identified, since surrounding this region are the central (larger) and peripheral tubercles (Fig. 2I and 2J). In the interior of the egg, embryonic cells are identified, and specifically in the initial third of development (5 hours), it was possible to observe in a semi-thin section, the presence of round cells with a peripheral nucleus and several cells presenting two nuclei, a process suggestive of cell division (Fig. 2K, 2M and 2N). Still in the initial stage of embryonic development, after 18 hours from the analysis of the semi-fine cut, folding of the embryo was found, around the egg internally in about 60% of it, a situation called the extension of the germ band and occurs at the moment of gastrulation when the ventral blastoderm undergoes extension (Fig. 2C, 2D and 2E), it is also suggested that the process of cellular activity is maintaining itself intensely after 18 hours of the onset of cell development, as a large number of cells were found in the process of division (Fig. 2G and 2H).

The intermediate phase (Fig. 3A-F) of development brings characteristics such as the increase in the presence of bristles, including palatal brushes, which in the future will allow the larva to perform movements promoting water flow to bring food to be consumed. There is also a separation of the cephalic and thoracic region and loss of homogeneity of the general morphological aspect. At the developing embryo stage, is evident the presence of a membrane surrounding the embryo and, therefore, located between the embryo and the chorion, this extraembryonic membrane is the serous layer (Fig. 3G).

Still at this stage, embryos were analyzed after 25 hours (Fig. 19C) and 30 hours of development (Fig. 3H to 3M). During this period, the process of formation of the abdominal and thoracic segments and structures such as the respiratory siphon and the last abdominal segment in the form of a caudal appendix (Fig. 3L), a spike that will promote the cleft to release the larva at the time of hatching, was observed of the egg, in addition to the presence of bristles in the cephalic region (Fig. 3M).

In the final third of development (Fig. 4A-D) the body of the future larva already has the divisions in the head, thorax and abdominal segments, but the indicator for completion of growth is mainly the presence of the chorion-breaking spike (Fig. 4C-H). This structure persists until the first larval stage (Figs. 4I and

4J), a factor that differentiates the first and second larval stages. In the period of completion of embryogenesis, prominence in the chorion was identified (Fig. 4E), specifically in the anterior region, which indicates that the larva is ready to gain the aquatic environment. The prominence identification site was the one that gave rise to the rupture of the transverse fissure in this region by the chorion-breaking spike (Fig. 4F-H), in the egg after completion of the embryonic development. This region will be the one where the larva will leave the egg. Histologically, the final stage of development (Fig. 4K, 4L, 4Q) shows that segmentation is better defined, as well as the primitive intestine (Fig. 4M), the discreet presence of bristles in the cephalic region is replaced by the abundant presence of these structures (Fig. 4P).

It is known that an important element of the cytoskeleton is actin, and this, in turn, was marked by the reaction with phalloidin, which then made it possible to understand its distribution by cells, while the nucleus was marked by propidium iodide. The images obtained by confocal microscopy made it possible to follow the sequence in embryonic development after 18 hours of development (initial phase) until the moment when the larva is ready to gain the environment (Fig. 5A to 5F). In this analysis it was possible to identify the embryo in a situation of extension of the germ band (Fig. 5B), after 30 hours of development the serous cell layer surrounding the embryo (Fig. 5B) was clear, and at the end of the development it was verified in 40 hours, the process of segmentation and dorsal closure (Fig. 5C) and within 50 hours an embryo was observed whose thoracic and abdominal segmentation was well defined (Fig. 5D), it is possible to observe the moment of the larvae hatching (Fig. 5E) and for end the embryo after 7 days diapause, with its development finished, ready to gain the external environment. In this case, the presence of visible segmentation was found and between the fifth and sixth abdominal segment, it was possible to show the intestine (Fig. 5F).

Discussion

The characteristics of eggs represented by *Aedes aegypti*, evaluated in this study were black or brown in color and measured less than 1.0 mm in length (Service 2003; Bova, Paulson and Paulson 2016). Despite the fact at the time of egg-laying, they had a white color, darkening afterward (Sallum, Barata and Santos 2007). Other characteristics inherent to the egg were: the presence of oval or elliptical outline, with bilateral symmetry (Suman et al. 2011; Bova, Paulson and Paulson 2016; Sallum, Barata and Santos 2007; de Morais et al. 2019), a characteristic identified in this research.

Fresh eggs are susceptible to water loss and this condition can impair its viability (Vargas et al. 2014). Such a statement suggests that there would be greater permeability in eggs with shorter embryo development time, so there would be greater ease in fixing embryonic tissues and obtaining results consistent with the proposed objectives. However, in this research, it was not possible to identify any easier processing of eggs in any of the evaluated phases, all of them were equally laborious since the resistance of the chorion was a constant complicating factor.

The exochorion generally maintain ornamentations that make it possible to identify the species. It is an excellent parameter for comparing species, as it has ability to reveal significant differences (Suman et al.

2011; Service 2003; Bova, Paulson and Paulson 2016; de Morais et al. 2019). In the figures presented (Figs. 1C, 1E, 1F and 1G) the ornamentations on the exochorion maintained the same pattern of organization in the species studied (Isoe et al. 2019). In addition to *Aedes aegypti*, other culicids of the same genus present ornamentation with polygonal chorionic cells in the exochorion of the egg, where a large central tubercle and other smaller and peripheral ones are observed, except for the anterior surface containing the micropylar apparatus (Suman et al. 2011; Faull and Williams 2016). The central tubercles on the external surface of the egg joined the peripheral tubercles employing thick projections forming a line; however, these differ in terms of their dimensions according to the location in the exochorion (Linley 1989; Isoe et al. 2019). In this study, this projection can be seen in Fig. 1.

When rejecting the hypothesis of equality between the two egg populations, it is suggested that a similarity identified in two distinct populations occurred due to chance, and given the above, the hypothesis arises that there may be microevolution in these populations considering these attributes (Wilk-Da-Silva et al. 2018; Louise, Vidal and Suesdek 2015). However, the use of other taxonomic and phylogenetic tools is necessary to deepen the discussion on this evolutionary process. When comparing morphologically, using scanning electron microscopy, eggs of *Aedes aegypti* in relation to eggs of *Aedes albopictus* (Suman et al. 2011; de Morais et al. 2019), it's affirmed that the former are greater concerning the micropyle disc and pore attribute in relation to the latter and the chorion polygonal cells have little capacity to assist in distinguishing between these species alone. Agreeing with Faull and Williams (2016) who expanded the analysis in different Australian cities regarding *Aedes notoscriptus* (Table 2).

Table 2

Comparative analysis of the morphometric findings of *Aedes aegypti* eggs identified in this study and the results of Suman et al, (2011), Linley (1989), Faull and Williams (2016). [This Table 2 should be attached between the 278 and 279 lines]

	<i>A. aegypti</i> ^{a *}	<i>A. aegypti</i> ^{b **}	<i>A. aegypti</i> ^{c ***}	<i>A. aegypti</i> ^{d ****}	<i>A. aegypti</i> ^{e *****}
Length	581.45 ± 39.73	625.65 ± 19.91	670.2 ± 7.2	554.41 ± 36.56	562.62 ± 30.85
Width	175.36 ± 11.59	183.30 ± 11.04	186.3 ± 2.2	167.65 ± 7.05	160.15 ± 9.73
Length/Width	3.32 ± 0.26	—	3.61 ± 0.05	3.31 ± 0.18	3.52 ± 0.27
Diameter of micropilar disc	18.75 ± 1.92	—	—	33.49 ± 3.9	34.19 ± 5.4
a. Mundim-Pombo et al b. Suman et al, 2011 c. Linley, 1989 d. Faull and Williams, 2016 e. Faull and Williams, 2016					
* Population of São Paulo eggs, ** Population of India eggs *** Population of Florida USA eggs **** Population of Cairns-Australia eggs ***** Population of Charters Towers eggs - Australia					

Table 2
Comparative
analysis
between the
morphometric
findings of
Aedes aegypti
eggs
identified in
this research
and the
results of
Suman et al,
(2011), Linley
(1989) and
Faull and
Williams
(2016).

The length of the eggs of *Aedes Aegypti*, identified dimensions of smaller than as for samples collected in India the Summan, et al. (2011) study. In another analysis, *Aedes aegypti* from Florida USA, presented a length of less than 700 μm (Linley 1989). The length identified in the egg population of São Paulo also reached lower levels, with a maximum value of 612.39 μm and 621.18 μm respectively. Regarding the chorionic coating, the findings were similar (Table 2), although there are mathematical differences in their measurements. Regarding the chorionic coating, the findings were similar (Table 2), although there are mathematical differences in their measurements.

It is known that *Aedes aegypti* eggs from different regions maintain the possibility of undergoing a process of adaptation to the environment (level of pollution, quality of pollutants in the air, water quality, morphological variation in terms of climate and the city), more studies of molecular biology must be carried out to confirm this (Dujardim, 2008).

The identified resistance of the chorion is consistent (Fonseca, Gomes and Araújo 2012) in which he attributes the precarious description of the embryonic morphology of *Aedes aegypti* to problems related to the permeability of the egg. External extraembryonic membrane, the serous one, can be easily seen (Panfilio 2008). This can be seen not only in the histological analysis (Fig. 3G) in which the serous layer was between the chorion and embryonic tissues, but also in the analysis using confocal microscopy (Fig. 5B) in which it was characterized as a distinct group of cells when compared to the others. However, in both images, the serosa was better evident in the intermediate phase of embryonic development.

After the end of the embryonic development, under favorable environmental conditions (Clean water, Temperature of 27 ± 2 ° C and neutral pH) the eggs of *Aedes aegypti* hatch, with 50% of larval hatching (Falcão et al. 2016; Famesi *et al.* 2009). In this study, some eggs with finished embryonic development were analyzed (Fig. 5E and 5F), in which case approximately 40% of the eggs hatched even after immersion in 3% sodium hypochlorite, that is, even when subjected to unfavorable environmental conditions, approximately 1 hour after the beginning of the sample processing, some embryos were still alive and the eggs hatched.

At the time of hatching, the chorion is ruptured due to the larval muscle activity, which increases in volume and consequently increases the pressure exerted from the spike, a specialized structure, then the rupture of the chorion occurs, from a crack in the part corresponding to the cited coating (Sallum, Barata and Santos 2007; Guindo-Coulibaly et al. 2018). This description was verified in this study (Fig. 4E-H) and the spike mentioned was evidenced histologically in Fig. 3M.

Conclusions

Given the above, it is concluded that there is great resistance of the chorion in the embryo is still a factor that generates difficulties for the microscopic analysis of the embryo (mainly in fixation process), with a strong protective barrier, which makes it difficult to use its embryonic cells study. No weaknesses were identified in the egg phase of the biological cycle, which would be extremely important for further research in order to identify new ways of effectively combating this important vector disease.

Declarations

Ethics approval and consent to participate

This research was approved by the Ethics Committee on the Use of Animals of the Faculty of Veterinary Medicine and Animal Science from São Paulo University, under protocol nº 2889/2013.

Consent for publication

The authors are able to share all dates information if necessary.

Availability of data and materials

The data will be available when requested.

Competing interest statement

The authors declare no conflict of interests.

Funding

This manuscript was funding by FAPEAM, Brazil.

Authors contributions

Ana Paula Miranda Mundim-Pombo (paulamedvet@gmail.com): concept and design; collect samples; samples preparation; data interpretation; immunofluorescence analysis; scanning electron microscopy analysis and compliance analysis.

Marisol León: helped data interpretation and compliance analysis.

Hianka Jasmyne Costa de Carvalho (hiankacarvalho@usp.br): essay redaction, translation and revision, reference research and discussion development;

Durvanei Augusto Maria (durvanei@usp.br): helped scanning electron microscopy and immunofluorescence analysis.

Maria Angelica Miglino (miglino@usp.br): macroscopic and microscopic description, study supervision, critical revision of the article and approval of article.

Acknowledgments

We would also like to thank the Advanced Diagnostic Imaging Center (CADI) of the School of Veterinary Medicine and Animal Science of University of São Paulo and Rose Eli Graci, PhD, the technician that helped the acquisition of the images used to analyze the results of the research.

Author's information

References

1. WHO. World Health Organization. Dengue and severe dengue. Fact Sheets. 2018. Available from: <http://www.who.int/news-room/fact-sheets/detail/dengue-and-severe-dengue>. Accessed 29 Jan 2021.
2. Charrel RN, Leparc-Goffart I, Gallian P, de Lamballerie X. Globalization of Chikungunya: 10 years to invade the world. *Clin Microbiol Infect*. 2014;20(7):662–3. DOI:10.1111/1469-0691.12694.
3. WHO. World Health Organization. Chikungunya. 2017. Available from: <https://www.who.int/news-room/fact-sheets/detail/chikungunya>. Accessed 8 Jan 2021.
4. WHO. World Health Organization. Zika virus. Fact Sheets. 2018. Available from: <http://www.who.int/news-room/fact-sheets/detail/zika-virus>. Accessed 8 Jan 2021.
5. Gloria-Soria A, Ayala D, Bheecarry A, Calderon-Arguedas O, Chadee DD, Chiappero M, et al. Global genetic diversity of *Aedes aegypti*. *Mol Ecol*. 2016;25(21):5377–95. DOI:10.1111/mec.13866.
6. Chaves BA, Orfano AS, Nogueira PM, et al. Coinfection With Zika Virus (ZIKV) and Dengue Virus Results in Preferential ZIKV Transmission by Vector Bite to Vertebrate Host. *J Infect Dis*. 2018;218(4):563–71. DOI:10.1093/infdis/jiy196.
7. Carrillo-Hernández MY, Ruiz-Saenz J, Villamizar LJ, Gómez-Rangel SY, Martínez-Gutierrez M. Co-circulation and simultaneous co-infection of dengue, chikungunya, and zika viruses in patients with febrile syndrome at the Colombian-Venezuelan border. *BMC Infect Dis*. 2018;18(1):1–12. DOI:10.1186/s12879-018-2976-1.
8. Ferguson NM, Rodríguez-Barraquer I, Dorigatti I, Mier-Y-Teran-Romero L, Laydon DJ, Cummings DAT. Benefits and risks of the sanofi-pasteur dengue vaccine: Modeling optimal deployment. *Science*. 2016;353(6303):1033–6. DOI:10.1126/science.aaf9590.

9. Brasil. MINISTÉRIO DA SAÚDE. SECRETARIA DE VIGILÂNCIA EM. SAÚDE. Monitoramento dos casos de dengue até a semana epidemiológica 41e febre de Chikungunya até a semana epidemiológica 15 de 2015. Bol Epidem. 2015;46(15).
10. Lima-Camara TN. Emerging arboviruses and public health challenges in Brazil. Rev Saúde Publica. 2016;50:1–7. DOI:10.1590/S1518-8787.2016050006791.
11. Caragata EP, Rocha MN, Pereira TN, Mansur SB, Dutra HLC, Moreira LA. Pathogen blocking in wolbachia-infected *Aedes aegypti* is not affected by zika and dengue virus co-infection. PLoS Negl Trop Dis. 2019;13(5):1–26. DOI:10.1371/journal.pntd.0007443.
12. CLEMENTS A. The biology of mosquitoes: development, nutrition and reproduction. London. 1992.
13. Suman DS, Shrivastava AR, Pant SC, Parashar BD. Differentiation of *Aedes aegypti* and *Aedes albopictus* (Diptera: Culicidae) with egg surface morphology and morphometrics using scanning electron microscopy. Arthropod Struct Dev. 2011;40(5):479–83. DOI:10.1016/j.asd.2011.04.003.
14. Fonseca RN, Gomes H, Araújo H. Aspectos morfofuncionais da embriologia dos artrópodes. In: INSTITUTO NACIONAL DE CIÊNCIA E TECNOLOGIA EM ENTOMOLOGIA MOLECULAR, editor. Tópicos avançados em entomologia molecular. Rio de Janeiro. 2012.
15. Obara MT, Da Rosa JA, Da Silva NN, et al. Estudo morfológico e histológico dos ovos de seis espécies do gênero *Triatoma* (Hemiptera: Reduviidae). Neotrop Entomol. 2007;36(5):798–806. DOI:10.1590/S1519-566X2007000500023.
16. Farnesi LC, Vargas HCM, Valle D, Rezende GL. Darker eggs of mosquitoes resist more to dry conditions: Melanin enhances serosal cuticle contribution in egg resistance to desiccation in *Aedes*, *Anopheles* and *Culex* vectors. PLoS Negl Trop Dis. 2017;11(10):1–20. DOI:10.1371/journal.pntd.0006063.
17. Faull KJ, Williams CR. Differentiation of *Aedes aegypti* and *Aedes notoscriptus* (Diptera: Culicidae) eggs using scanning electron microscopy. Arthropod Struct Dev. 2016;45(3):273–80. DOI:10.1016/j.asd.2016.01.009.
18. Mello CF, Santos-Mallet JR, Tátilla-Ferreira A, Alencar J. Comparing the egg ultrastructure of three *psorophora ferox* (Diptera: Culicidae) populations. Brazilian J Biol. 2018;78(3):505–8. DOI:10.1590/1519-6984.171829.
19. Alencar J, Guimarães AE, Mello RP, Lopes CM, Dégallier N, Santos-Mallet JR. [Scanning electron microscopy of eggs of *Haemagogus leucocelaenus* (Diptera: Culicidae)]. Rev Saude Publica. 2003;37(5):657–61. DOI:10.1590/s0034-89102003000500017.
20. Reinhold JM, Lazzari CR, Lahondère C. Effects of the environmental temperature on *Aedes aegypti* and *Aedes albopictus* mosquitoes: A review. Insects. 2018;9(4). DOI:10.3390/insects9040158.
21. Matthews BJ, Dudchenko O, Kingan SB, et al. Improved reference genome of *Aedes aegypti* informs arbovirus vector control. Nature. 2018;563(7732):501–7. DOI:10.1038/s41586-018-0692-z.
22. Linley JR. Comparative fine structure of the eggs of *Aedes albopictus*, *Ae. aegypti*, and *Ae. bahamensis* (Diptera: Culicidae). J Med Entomol. 1989;26(6):510–21. DOI:10.1093/jmedent/26.6.510.

23. Service Mike. 978-1-107-66818-8 - Medical Entomology for Students: Fifth Edition Mike Service Front Matter More information Medical Entomology for Students © in this web service Cambridge University Press Cambridge University Press 978-1-107-66818-8\$4- Medical Entomology. 2003.
24. Bova J, Paulson S, Paulson G. Morphological differentiation of the eggs of north American container-inhabiting *Aedes* mosquitoes. *J Am Mosq Control Assoc.* 2016;32(3):244–6. DOI:10.2987/15-6535.1.
25. Sallum MAM, Barata JMS, Santos RLC dos. Oswaldo Paulo Forattini: epidemiologista, entomologista e humanista. *Rev Saúde Pública.* 2007;41(6):885–913.
26. Li JS, Li J. Major chorion proteins and their cross-linking during chorion hardening in *Aedes aegypti* mosquitoes. *Insect Biochem Mol Biol.* 2006;36(12):954–64. DOI:10.1016/j.ibmb.2006.09.006.
27. de Moraes LMO, Jussiani EI, Zequi JAC, dos Reis PJ, Andrello AC. Morphological study of *Aedes aegypti* and *Aedes albopictus* (Diptera: Culicidae) eggs by X-ray computed microtomography. *Micron.* 2019. DOI:10.1016/j.micron.2019.102734.
28. Vargas HCM, Farnesi LC, Martins AJ, Valle D, Rezende GL. Serosal cuticle formation and distinct degrees of desiccation resistance in embryos of the mosquito vectors *Aedes aegypti*, *Anopheles aquasalis* and *Culex quinquefasciatus*. *J Insect Physiol.* 2014;62(1):54–60. DOI:10.1016/j.jinsphys.2014.02.001.
29. Isoe J, Koch LE, Isoe YE, et al. Identification and characterization of a mosquito-specific eggshell organizing factor in *Aedes aegypti* mosquitoes. *PLoS Biol.* 2019;17(1):1–23. DOI:10.1371/journal.pbio.3000068.
30. Wilk-Da-Silva R, De Souza Leal Diniz MMC, Marrelli MT, Wilke ABB. Wing morphometric variability in *Aedes aegypti* (Diptera: Culicidae) from different urban built environments. *Parasites Vectors.* 2018;11(1):1–9. DOI:10.1186/s13071-018-3154-4.
31. Louise C, Vidal PO, Suesdek L. Microevolution of *Aedes aegypti*. *PLoS One.* 2015;10(9):1–16. DOI:10.1371/journal.pone.0137851.
32. Panfilio KA. Extraembryonic development in insects and the acrobatics of blastokinesis. *Dev Biol.* 2008;313(2):471–91. DOI:10.1016/j.ydbio.2007.11.004.
33. Falcão A, Anjolette F, De Lourdes Da M, Macoris G. Técnicas para manutenção de *Aedes aegypti* em laboratório Techniques for the *Aedes aegypti* maintenance in the laboratory. *Bepa.* 2016;13(156):19–29.
34. Farnesi LC, Martins AJ, Valle D, Rezende GL. Embryonic development of *Aedes aegypti* (Diptera: Culicidae): Influence of different constant temperatures. *Mem Inst Oswaldo Cruz.* 2009;104(1):124–6.
35. Guindo-Coulibaly N, Diakite NR, Adja AM, et al. Biology of two larval morphological phenotypes of *Aedes aegypti* in Abidjan, Côte d'Ivoire. *Bull Entomol Res.* 2018;108(4):540–6. DOI:10.1017/S0007485317001109.

Figures




Figure 1

Aedes aegypti egg. (A) Macroscopic appearance; (B-G) SEM (H-K) Transmission microscopy, Chorion: (B) *Aedes aegypti* egg in anteroposterior position, with emphasis on the anterior position. Here you can see the microcapillary disc; (C) Microcapillary disc (DM) and the micropyle in the center (trilobulated pore in the center of the micropyle disc), an orifice through which the sperm penetrates to fertilize the oocyte; (D) The posterior end of the egg shows its ornamentation, called the exochorionic network. They are

prominent on the surface, giving the appearance of high relief or texture.; (E) Central tubercle (CT) has a larger dimension and is surrounded by smaller peripheral tubercles (TP); In the exochorionic network (F), the disposition of the Peripheral Tuber (TP) in relation to the Central Tuber (CT) is observed. (G) Central tubercle (TC); (H) Exochorion (top banner) more electrodense when compared to the endochorion (lower band); (I) the difference between the internal environment of the egg (but light) and the external one (darker with the presence of central, larger, and peripheral, smaller tubers). Shows the impermeability of the chorion, since the resin did not reach the interior of the egg (J) Chorion thickness of the egg (K) Central tubercle (larger) peripheral tubercle (smaller).




Figure 2

Embryo in the early stage of development of *Aedes aegypti* (A-B) SEM; (C-H) Embryo after 18 hours of development, semi-fine cut (I-J) Histological aspects of an embryo at an early stage (5 and 18 hours) (KN) Embryonic cells of *Aedes aegypti* at an early stage of development (5 hours) observed under microscopy of light from semi-fine cut. (A) Sectioned chorion exposes the embryonic structure at an early stage of development (B) Anterior region of the embryo, corresponding to the cephalic structure. Chorion-

breaking spike forming in the center and few bristles. (C) Discreet embryonic folding around the egg internally in about 60% of it (extension of the germ band). (D) Embryo cranial region (E) embryo caudal region (F) in the center of the highlighted area, sequential steps are presented in the embryonic development of distinct embryos. (G) Embryonic cells bypassing the corium internally (H) presence of calf surrounded by embryonic cells in cell activity; (I) Exochorion surrounded by the central tubercle and the presence of a calf inside the egg (J) *Aedes aegypti* egg after 18 hours of embryonic development. Embryo in a situation of extension of the germ band, on the surface of the chorion, tubers are present; (K, M and N) embryo cells at the beginning of embryogenesis, (L) cells of the egg exochorion (central tubercle, peripheral tubercle, ornamental structures that surround the egg).




Figure 3

Embryo in the middle stage of development (middle third of embryogenesis); (A-F) SEM; (G) Histological aspects, intermediate phase (25 hours); (H-M) Embryo after 30 hours of development from the analysis of the semi-fine cut; (A) Sectioned chorion, cephalic region of the embryo and part of the thoracic region. (B) cephalic region, presence of bristles and tapering separator from the cephalic and thoracic region. (C) anterior end of the cephalic region (D) sectioned chorion and approximation of the separating tapering of

the head and chest. (E) palatal brushes (F) cells of the cephalic region. (G) Embryo in the middle stage of development, in evidence is the serous. (H) In a smaller increase, there is an overview of the embryos after 30 hours of development (I and J) forming abdominal segments. (K) Adjacent to the forming abdominal segments, there are cells with nutritional reserves (L) respiratory siphon and caudal appendix in formation (M) cephalic region with the presence of bristles and formation of the chorion-breaking spike in the center.




Figure 4

Final stage of development. (A-J) Ultrastructure (SEM) of the embryo (A-D) and the first stage larva (after hatching) (I-J); (A-B) Embryo ventral view; (C-D) Embryo side view (E-H) process of hatching of the larva. (K) Histological aspects of an embryo at the final stage (45 hours) of development (L-Q) Embryo after 50 hours of development (semi-fine cut); (A) embryo with formed abdominal segments, body divided into head, chest and abdomen. (B) Head and part of the chest. Presence of bristles mainly in the cephalic region; (C) Spike in the center of the dorsal surface for hatching of the egg (D) Antenna, spike in the center and bristles in the cephalic region; (E) prominent point (arrow) where the chorion will break. (F and G) Chorion ruptured, making evident the structure responsible for its rupture, the disruptive spike (H) close image of the chorion-breaking spike; (I) 1-instar larva in dorsal view (J) head of the larva (dorsal face) Spike that breaks the chorion in the center. (L) Overview of the embryo after 50 hours of development. (M) Segments defined in the abdominal region, in the central region elongated structures composed of differentiated cells are part of the intestine of the insect. (N) Different groups of cells and better definition of the abdominal segments (O) cephalic region and thoracic embryo (P) cephalic region and presence of bilateral setae (Q) Defined abdominal segments.



Figure 5

Initial, intermediate and final stages of embryonic development of *Aedes aegypti*, hatching of the egg and first instar larvae analyzed by confocal microscopy. (A) Extension of the germ band in the embryo after 18 hours of embryonic development. (B) Embryo after 30 hours of development, dorsal and superficially, cells of the serous layer. (C) Embryo of 40 hours of development, dorsal closure and abdominal segmentation process (D) embryo after 50 hours of development, thoracic and abdominal segmentation

completed at the time of the larvae hatching (F) embryo after 7 days diapause, development completed. Presence of visible segmentation and between the fifth and sixth abdominal segment it is possible to show the intestine of the insect.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Addicionalfigure1.pdf](#)
- [Addicionalfigure2.pdf](#)
- [Addicionalfigure3.pdf](#)
- [Addicionalfigure4.pdf](#)
- [Addicionalfigure5.pdf](#)