

# Knockdown by *MSH2* and *EPCAM* siRNA suppress Wnt/ $\beta$ -Catenin Pathway in HCT116 Cell Line

Wan Khairunnisa Wan Juhari (✉ [khairunnisa.juhari@gmail.com](mailto:khairunnisa.juhari@gmail.com))

Universiti Sains Malaysia - Kampus Kesihatan

Khairul Bariah Ahmad Amin Noordin

Universiti Sains Malaysia - Kampus Kesihatan <https://orcid.org/0000-0002-0787-8532>

Wan Faiziah Wan Abdul Rahman

Universiti Sains Malaysia - Kampus Kesihatan

Andee Dzulkarnaen Zakaria

Universiti Sains Malaysia - Kampus Kesihatan

Wan Muhammad Mokhzani Wan Muhammad Mokhter

Universiti Sains Malaysia - Kampus Kesihatan

Bin Alwi Zilfalil

Universiti Sains Malaysia - Kampus Kesihatan

---

## Research Article

**Keywords:** Small interfering RNA (siRNA), colorectal cancer, MSH2, EPCAM,  $\beta$ -catenin

**Posted Date:** March 10th, 2021

**DOI:** <https://doi.org/10.21203/rs.3.rs-274862/v1>

**License:**   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# Abstract

**Purpose:** Small interfering RNA (siRNA) has the potential as a therapeutic approach against selective pathways in colorectal cancer. *EPCAM*, a transmembrane glycoprotein mediating cell adhesion, was known to be involved in suppressing Wnt/ $\beta$ -*catenin* pathway, an important pathway for tumour progression in colon cancer cells. *EPCAM* deletions caused a transcriptional read-through that may silence its neighbouring gene, *MSH2*. This study aimed to investigate the effect of co-siRNA targeted genes, *MSH2* and *EPCAM*, in colon cancer cell line, HCT116, and their effect in regulating the Wnt/ $\beta$ -*catenin* pathway.

**Methods:** Pre-designed siRNA of *MSH2* and *EPCAM* were transfected into HCT116 cells. The cells were divided into six groups of treatments: untreated cell group, cells treated with negative control siRNA, *MSH2*-siRNA treated cells, *EPCAM*-siRNA treated cells, cells treated with both *EPCAM* and *MSH2*-siRNA, and cells treated with transfection reagent (mock control). The mRNA and protein expression following the individual and combined siRNA treatments were assessed by quantitative polymerase chain reaction and Western blot.

**Results:** The mRNA and protein expression levels of *MSH2*, *EPCAM* and  $\beta$ -*catenin* were reduced in the individual *MSH2* and *EPCAM*-siRNA treated samples as compared to the untreated sample. Further reduction of mRNA and protein expressions for *MSH2*, *EPCAM* and  $\beta$ -*catenin* were identified in combined siRNA treatments.

**Conclusion:** Reduction of  $\beta$ -*catenin* expression by simultaneous silencing of *MSH2* and *EPCAM* suggested that these genes may play a role in suppressing the Wnt/ $\beta$ -*catenin* pathway in cancer cells.

## Introduction

Colorectal cancer (CRC) is the third most common cancer worldwide and the fourth most common cause of death [1, 2]. The most common hereditary colon cancer is Lynch syndrome (LS) which is caused by germline mutations in any of the four mismatch repair (MMR) genes namely *MLH1*, *MSH2*, *MSH6* and *PMS2* [3] and to date, *EPCAM* gene was also reported to be associated with LS [4]. *EPCAM* functions in intracellular signalling, differentiation and proliferation of normal and cancer tissues [4]. This was suggested by the *EPCAM* signalling that is activated by intra-membrane proteolysis in which the extracellular domain is cleaved off and the intracellular domain (EpicD) is then released into the cytoplasm. In the cytoplasm, *EPCAM* formed a nuclear complex with transcriptional regulators  $\beta$ -*catenin* and Lef, both are parts of Wnt signalling pathway [5]. Germline deletions in the last exon of the non-MMR gene, *EPCAM*, may silence its neighbouring gene, *MSH2*, through promoter hypermethylation [6]. Previous study has shown that in *EPCAM*-silenced colon cancer cells, the expression level of  $\beta$ -*catenin* was decreased, and the result showed that silencing of *EPCAM* could inactivate the Wnt/ $\beta$ -*catenin* pathway in tumour cells [7]. Hence, *EPCAM* was identified as a principal target for the treatment of tumours, through the mechanism of suppressing the Wnt/ $\beta$ -*catenin* pathway. *EPCAM* deletions cause a transcriptional

read-through and silenced the *MSH2* gene [8]. The mechanism of promoter hypermethylation happens when a number of methyl groups bound to the *MSH2* promoter region, and consequently decrease the expression of *MSH2* protein products in the epithelial cells. Considering the major role of *MSH2* protein in DNA repair mechanism, the loss of this protein may subsequently affect the process of DNA repair, and mistakes accumulated upon cell division [4].

Wnt signalling pathway has become one of the major pathway in CRC and LS predisposition with about 65% overactivation in colorectal cancers associated to LS and activating  $\beta$ -catenin mutations were identified in approximately 40% of these tumours [9]. Although CRC was commonly triggered by the alterations in the signalling components of the Wnt signalling pathway [10], the relationship between *MSH2* and *EPCAM* genes towards the effect of associated pathway is yet to be discovered. Therefore, this study aimed to investigate the effect of *MSH2* and *EPCAM* *siRNA* mediated gene knockdown in regulating the Wnt/ $\beta$ -catenin pathway.

## Materials And Methods

### Culture of HCT116 cell line

HCT116 cell line was purchased from RIKEN BioResource Centre, Japan, and was established according to manufacturer's protocol. The cell suspension was centrifuged in complete culture medium (DMEM supplemented with 1.0 g/l of glucose, L-glutamine, sodium pyruvate and 10% FBS) for centrifugation at 1000 rpm for 3 minutes at room temperature. Followed by centrifugation and resuspension, the cell suspension was seeded into two 25 cm<sup>2</sup> culture flasks and incubated at 37°C in humidified incubators with 5% CO<sub>2</sub>. The growth of cells was maintained daily and fresh culture medium was changed for every two days or when necessary. Sub-culturing was carried out when the cells reached 80% confluence. When the cells reached 80% confluence or 80% of the flask surface area was covered by the cells, sub-culturing was carried to ensure continuous growth of cells during their exponential growth. The first few cultures were sub-cultured with split ratio of 1:2 to 1:3 depending on the growth of cells. Before sub-culturing, the culture medium was discarded and the adherent cells were washed with 1 ml of PBS. After two to three washing steps, the cells were digested with 0.25% trypsin-EDTA for 5 minutes at 37°C in humidified incubators with 5% CO<sub>2</sub>. Upon confirmation of cell detachment, 1 ml of complete culture medium was added and gently dispersed by pipetting over the surface of the cell layer to deactivate the trypsin activity. The trypsin and cell suspension were mixed thoroughly and transferred into 15 ml centrifuge tube prior to centrifugation at 1200 rpm for 5 minutes at room temperature. Supernatant was then discarded and the cell pellet was re-suspended in 1 ml of complete culture medium. The cell suspension was sub-divided according to appropriate split ratio into the new culture flasks containing fresh complete culture medium. The cells were then maintained at 37°C in humidified incubators with 5% CO<sub>2</sub>.

### SiRNA Preparation And Transfection

Hiperfect® Transfection Reagent (Qiagen, Denmark) was used for siRNA transfection of MSH2 and EPCAM genes into the HCT116 cells. SiRNA was prepared in an RNase-free environment. The target sequence of four pre-designed siRNAs (FlexiTube GeneSolutions, Qiagen, Denmark) for each *MSH2*, *EPCAM* and negative control siRNA were listed in Table 1. RNase-free water in a total volume of 100 µl was added to 1 nmol of lyophilized siRNA for MSH2 and EPCAM genes to obtain a final stock concentration of 10 µM. For the negative control siRNA, a total volume of 250 µl of RNase-free water was added for a final stock concentration of 20 µM. For the preparation of 1 µM working solution, 2 µl of siRNA was added to the stock solution for a final volume of 20 µl. All siRNAs were stored at -20°C until further use.

Table 1  
List of siRNA sequences for MSH2 and EPCAM gene  
(FlexiTube GeneSolution siRNA, Qiagen).

<b><i>MSH2</i>_siRNA ID</b>	<b>Target sequence</b>
Hs_ <i>MSH2</i> _6	TCCAGGCATGCTTGTGTTGAA
Hs_ <i>MSH2</i> _5	CCCATGGGCTATCAACTTAAT
Hs_ <i>MSH2</i> _10	AAGAAGATGCAGTCAACATTA
Hs_ <i>MSH2</i> _12	TTGGATATTACTTTCGTGTAA
<b><i>EPCAM</i>_siRNA_ID</b>	<b>Target sequence</b>
Hs_ <i>EPCAM</i> _6	AACTATATAATTTGAAGATTA
Hs_ <i>EPCAM</i> _5	CTGGATCATCATTGAACTAAA
Hs_ <i>EPCAM</i> _10	AGGGA ACTCAATGCATAACTA
Hs_ <i>EPCAM</i> _9	TTTGTGAATAATAATCGTCAA

Before transfection, the cells were harvested in 1 ml complete culture medium. The viability of cells were assessed by dye exclusion test using Trypan Blue solution (GIBCO®, Invitrogen™, Canada). Approximately 60 000 cells were seeded in each well of 24-well plate containing 500 µl of complete culture medium. The cells were then incubated overnight under normal growth condition at 37°C with 5% CO<sub>2</sub>. The samples were divided into six groups with different treatments at a total siRNA concentration of 40 nmol per group of samples. The six groups were untreated cell group (Group 1), cells treated with negative control siRNA (Group 2), *MSH2*-siRNA treated cells (Group 3), *EPCAM*-siRNA treated cells (Group 4), cells treated with both *EPCAM* and *MSH2* siRNA (Group 5) and cells treated with HiPerfect Transfect Reagent which served as a mock control (Group 6). Each group of samples were run in triplicates. A set of four different siRNAs for each gene were used. A total of 37.5 ng (5 nmol) of each siRNA were diluted in 100 µl DMEM culture medium without serum and 3 µl of HiPerfect Transfect Reagent was added to the diluted siRNA. After gently mixed, the mixture was incubated for 10 minutes at room temperature for the formation of transfection complexes. The complexes were dispensed dropwise into each well and the plate was gently swirled to ensure uniform distribution of complexes throughout the cells.

# Rna Extraction And Cdna Synthesis

RNA extraction was performed after 48 hours of transfection by using a commercial RNA extraction kit, RNeasy Mini Kit (QIAGEN, Germany). The protocols of RNA extraction were carried out according to the manufacturer's protocols. The quality of the RNA was assessed by gel electrophoresis. Synthesis of cDNA was performed by using a commercial kit, SensiFAST™ cDNA Synthesis Kit (Bioline, USA) according to manufacturer's protocols. A total of 1 µg RNA served as an initial concentration for cDNA synthesis followed by the following conditions; 25°C for 10 minutes (primer annealing) followed by reverse transcription at 42°C for 15 minutes and the inactivation step at 85°C for 5 minutes with a final hold at 4°C. The final volume of 20 µl cDNA in RNase-free was stored in -20°C until further use.

## Gene Expression Using Quantitative Real-time Pcr

The relative quantification of gene expression for *β-actin* and the targeted genes, *MSH2*, *EPCAM* and *β-catenin* was performed using Stratagene Mx3000P qPCR System (Agilent Technologies, USA). cDNA was diluted in 10-fold dilution of 5 different concentrations with the initial concentration of 100 ng/µl to further determine the PCR efficiency of each targeted gene. Real-time PCR (qPCR) amplification was carried out using commercial kit, Quantinova™ SYBR® PCR Kit (QIAGEN, Germany). The kit was used in combination of real-time predesigned primers of Quantitect Primer Assay (QIAGEN, Germany). The reaction mastermix was prepared in 96 well plate according to the proposed reaction setup according to the manufacturer's protocol. *β-actin* was used as a reference gene for the relative quantification of *MSH2*, *EPCAM* and *β-catenin* genes. The reaction mixture for *β-actin*, *MSH2*, *EPCAM* and *β-catenin* was prepared in a total volume of 20 µl containing 10 ng of cDNA, 10 µl of 1X SYBR Green PCR buffer, 2 µl of 1X QN ROX reference dye, 1X primer assay and RNase free water. The samples were run in triplicates for each set of the primers. The conditions for the amplification were 95°C for 2 min for the initial heat inactivation, denaturation at 95°C for 5 sec and 45 cycles of combined annealing and extension at 60°C for 10 sec followed by dissociation stage at 95°C for 1 min, 55°C for 30 sec and 95°C for 30 sec.

## Relative Quantification Analysis And Statistical Analysis

Relative quantification analysis was performed to determine the expression of the targeted genes (*MSH2*, *EPCAM* and *β-catenin*) in all samples by using comparative double  $\Delta\text{Ct}$  method.

Statistical analysis was performed by SPSS® statistical package, version 24.0 (SPSS Inc., Chicago, IL, USA). All data were presented as mean  $\pm$  SD. The statistical comparison of more than two groups in this experiment was tested using one-way ANOVA.  $P < 0.05$  was considered significant.

## Western Blot Analysis

Protein extraction was performed by using RIPA Buffer (Nacalai Tesque, Japan) according to the manufacturer's protocols. The cells were washed twice with cold PBS and 1X RIPA buffer was added to the culture depending on the number of cells in the culture flask. The protein concentration in a sample was measured using Bradford protein assay. Each sample replicates (n = 3) for each treatment were pooled separately with concentration of 20 µg. The protein samples and the standard marker subsequently subjected to 10% SDS-PAGE electrophoresis at 100 V for 80 minutes. The samples were transferred to PVDF membrane and was run at 25 V for 2 hours prior to incubation in 5% blocking buffer for 1 hour at room temperature. The membrane was then incubated with 5% blocking buffer mixed with an appropriate dilution of each primary antibody; *MSH2* (1:2000), *EPCAM* (1:2000) and *β-catenin* (1:1800) and *β-actin* (1:7500), for overnight at 4°C with gentle agitation. After an overnight incubation, the primary antibody was discarded. The membrane was washed twice at room temperature with TBS-T buffer followed by one time washing with TBS buffer. The membrane was then incubated with Goat Anti-Mouse IgG secondary antibody conjugated to horseradish peroxidase (HRP) in 5% blocking buffer at 1:20 000 dilutions for 1 hour with gentle agitation. The samples were visualized using chemiluminescent detection kit (Nacalai Tesque, Japan).

## Densitometry Analysis

The intensity of the protein bands was semi-quantitatively analysed by Image J tool (NIH, Bethesda, MD, USA) (<http://rsb.info.nih.gov/ij/index.html>) to compare the intensity of each protein band across different samples. A profile plot for each band that represented the density of the band was then created by using the Image J tool. The relative protein level of each sample was then calculated based on the following equation [11].

$$\text{Relative protein level} = \frac{\text{Intensity of GOI (treated sample) / Intensity of GOI (non-treated sample)}}{\text{Intensity of } \beta\text{-actin (treated sample) / Intensity of } \beta\text{-actin (non-treated sample)}} \times 100\%$$

## Results

### Effect of siRNA transfection

The effect of *MSH2* and *EPCAM* on the Wnt/*β-catenin* was investigated by transfecting the HCT116 cell lines with siRNA against the *MSH2* and *EPCAM* gene. The comparison of cell growth and morphology in each group of treated samples were observed after 48 hours of treatment (Fig. 1). The number of attached cells in the *MSH2* and *EPCAM* treated cells or *MSH2* and *EPCAM* knockdown cells as well as cells which were simultaneously treated by both genes were found to decrease notably. From the microscopic observation, the cells in Group 1, Group 2 and Group 6 was appeared as epithelial in shape as compared to the group of cells treated with the targeted genes. The morphology of cells in Group 3, Group 4 and Group 5 were observed to change into rounded bodies with a few smaller cells in epithelial

shape. In the treated samples, the detached cells were observed to be aggregated with a few cells remained attached to the surface of the flask. As compared to Group 3 and Group 4 samples, the cells in Group 5 displayed the highest cell aggregation with a few numbers of cells attached on the surface of the flask.

### Gene expression analysis of MSH2 and EPCAM knockdown

The gene expression level of *MSH2*, *EPCAM* and  $\beta$ -*catenin* in six different groups were determined by comparing the  $2^{-\Delta\Delta Ct}$  values of the target genes normalized to the housekeeping genes,  $\beta$ -*actin* with respective to the control group. The expression level of individual gene knockdown, *MSH2*-siRNA treated group and *EPCAM*-siRNA treated group and simultaneous double gene knockdown, *MSH2*+*EPCAM*-siRNA treated group were shown to be statistically significant with  $p$ -value < 0.05 as compared to the control (Fig. 2). *MSH2*-siRNA significantly inhibited the *MSH2* mRNA expression at the level of  $0.17 \pm 0.04$  ( $p$ -value = 0.005) as compared to the untreated one (Fig. 2a). Meanwhile, a slight decrease which was not much altered were observed for mRNA expression level in negative control siRNA with no significant difference compared to the untreated control. A notable knockdown of *MSH2* expression was also observed by *EPCAM*-siRNA at the level of  $0.25 \pm 0.05$  ( $p$ -value = 0.012) (Fig. 2a). Similarly, *EPCAM* expression level was inhibited by both *MSH2*-siRNA and *EPCAM*-siRNA with the level of  $0.25 \pm 0.03$  ( $p$ -value = 0.007) and  $0.11 \pm 0.04$  ( $p$ -value = 0.010), respectively (Fig. 2b). A significant knockdown was also observed in  $\beta$ -*catenin* in which both *MSH2*-siRNA and *EPCAM*-siRNA has reduced  $\beta$ -*catenin* expression at the level of  $0.34 \pm 0.07$  ( $p$ -value = 0.044) and  $0.32 \pm 0.04$  ( $p$ -value = 0.016), respectively (Fig. 2c). The effect of simultaneous gene expression was determined in these targeted genes. The mRNA expression level in these three targeted genes were observed to have significantly reduced by *MSH2*+*EPCAM*-siRNA (Fig. 2).

### Protein expression analysis of MSH2 and EPCAM knockdown

Protein expression was observed after 48 hours of transfection, including the untreated control. Reduced band intensity was observed for *MSH2*-siRNA and *EPCAM*-siRNA treated samples that indicated less protein was expressed for these three targeted genes as compared to untreated control. Reduced band intensity of *MSH2*+*EPCAM* siRNA treated samples were further observed that may indicate the possible effect of simultaneous gene knockdown resulted in low expression of the two targeted proteins (Fig. 3). The percentage of *MSH2* protein level transfected by the individual *MSH2*-siRNA and *EPCAM*-siRNA were reduced at 33% and 32% respectively (Fig. 4a). Increased reduction of protein level was observed in the simultaneous knockdown of *MSH2* and *EPCAM* siRNA with the percentage protein level of 14% (Fig. 4a). For *EPCAM* gene, the protein level was also identified to decrease in both targeted siRNA by 44% in *MSH2*-siRNA transfected sample and 43% of protein level in the *EPCAM*-siRNA transfected sample (Fig. 4.2b). Reduced protein level was identified in both genes transfected sample with the remaining protein level of 10% (Fig. 4.2b). In addition, for  $\beta$ -*catenin* gene, the *MSH2* and *EPCAM*-siRNA transfected sample was decreased to 38% and 33% respectively (Fig. 4c). Notable reduction of  $\beta$ -*catenin* protein level was

observed in the simultaneous gene knockdown with the percentage remaining protein level of 5% (Fig. 4c).

## Discussions

Germline deletion that cause the *MSH2* inactivation was considered a novel findings in the predisposition of LS [4, 5] including a novel large duplication of *MSH2-EPCAM* previously reported in a LS patient [12]. However, the association of these two genes in the pathway associated to CRC and LS have not yet being elucidated. Previous study showed that *EPCAM* was suggested as an important target gene that can disrupt the mechanism of Wnt/ $\beta$ -catenin pathway in HCT116 cells [7] but tumour cells commonly involved multiple genetic and epigenetic alterations and single inhibition of one tumour associated gene as a therapeutic strategy may inadequate to inhibit the development of tumour [13].

The present study was carried out in colon cancer cell line, HCT116 to evaluate the role of *MSH2* and *EPCAM* gene in modulating the Wnt/ $\beta$ -catenin pathway using siRNA target gene. The resulted gene silencing after siRNA transfection was evaluated by analysis of gene expression and protein expression by Western blot analysis which further verified the efficient gene knockdown in the mRNA and protein level. The expression of individual gene knockdown has reduced the mRNA and protein expression of *MSH2*, *EPCAM* and  $\beta$ -catenin in *EPCAM*-siRNA and *MSH2*-siRNA transfected cells. This finding was concordant with previous study that reported a significant reduction of proliferative activity and reduced mRNA and protein expression in *EPCAM*-siRNA transfected in HCT116 cells [7]. The expression level of  $\beta$ -catenin was also reduced in the *EPCAM*-siRNA transfected cells [7]. In addition, *MSH2* knockdown was previously carried out in other colon cancer cell lines, SW480 using shRNA-mediated gene silencing which resulted in the reduction of cell proliferative activity and decreased mRNA expression [14].

The simultaneous knockdown of *MSH2* and *EPCAM* may also be associated with the epigenetic modifications that usually occurs at the promoter region or enhancers of tumour-suppressor genes that often cause tumorigenesis [15]. *MSH2* methylation was reported as disease specific due to the absence of *MSH2* methylation in normal tissues as well as in sporadic CRC cases [16]. In addition, promoter region of *MSH2* was suggested as a target of aberrant methylation in LS due to the presence of high level of promoter methylation [16]. Previous study demonstrated the role of 3'-end *EPCAM* deletion that may cause *MSH2* methylation in patients with no LS germline mutations in *MSH2* gene [8]. Due to the downstream position of *MSH2* gene to *EPCAM* gene, gene silencing by transcriptional read-through of a neighbouring gene could represent a general mutational mechanism and also caused by a second somatic hit that inactivates *MSH2* in tumours with *EPCAM* deletion [8].

A significant reduction of  $\beta$ -catenin expression was observed in the simultaneous gene knockdown of *MSH2* and *EPCAM* which suggested that the silencing of these two genes may interrupt the activation of Wnt/ $\beta$ -catenin pathway in cancer cells. It has been well known that the degradation of Tcf/ $\beta$ -catenin complex formation decreased *EPCAM* gene expression in normal human hepatocytes culture and HCC cell lines [17] and human colon cancer line [7]. The key transcription regulator of Wnt signalling pathway

is *CTNNB1* gene which was identified to encode for  $\beta$ -catenin [18]. The mechanism took place when the phosphorylation of  $\beta$ -catenin by CK1 $\alpha$  followed by GSK3 $\beta$  mediated phosphorylation of the destruction complex and targeting  $\beta$ -catenin for ubiquitination and subsequent proteolysis [19]. However, the subsequent degradation of  $\beta$ -catenin will be avoided by the point mutations at these amino acids by hindering the  $\beta$ -catenin from being phosphorylated [18]. Mutations in *CTNNB1* and *AXIN2* in CRC were mostly arise in tumours with MMR genes; *MLH1*, *MSH2* or *PMS2* inactivation [20]. Although previous *in-vivo* study has showed that the deficiency of *MSH2* led to an enhance of  $\beta$ -catenin activity and cellular hyperproliferation in colon epithelial cells, however, this activity was dependent on gut microbes [21].

In conclusion, Wnt pathway played an important role as molecular signalling in various cancers and due to frequent abnormality of Wnt activation in colorectal cancer, it has been proposed as one of the key pathway in CRC predisposition [22, 23]. The activation of Wnt signalling commonly occurs by the presence of genetic alterations in *APC*,  *$\beta$ -catenin gene*, *AXIN1* and *AXIN2* that further caused  $\beta$ -catenin to be accumulated in the cytoplasm [20]. Based on the findings in the *in-vitro* study, the profound effect of the simultaneous gene knockdown of *MSH2* and *EPCAM* gene to the reduction of  $\beta$ -catenin expression may indicate the ability of these two genes to become co-target genes in regulating the Wnt/ $\beta$ -catenin pathway. This combined siRNA approach has also suggested to be a new therapeutic approach in the treatment of CRC as well as LS through suppression of Wnt/ $\beta$ -catenin pathway.

## Declarations

### Funding acknowledgement

This study was supported by Toray Science and Technology Research Grant by Malaysia Toray Science Foundation (MTSF) (304/PPSG/ 6150153/M126), Universiti Sains Malaysia (USM) Research University Grant (1001/PPSP/812112) and USM Short term grant (304/PPSP/61313202).

**Authors' contributions-** WK Wan Juhari performed the experiment, drafted the manuscript, performed the analysis. KB Ahmad Amin Noordin have been involved in the conception of idea, drafting the manuscript, performed the analysis and revising it critically for important intellectual content. WF Wan Abdul Rahman, AZ Zakaria, WMM Wan Muhamad Mokhter, BA Zilfalil involved in the conception of idea and drafting of the manuscript. All authors approved the final manuscript.

**Conflict of Interest** - The authors declare no conflict of interest

**Ethical Approval** – Not applicable

**Consent to Participate** – Not applicable

**Consent to Publish** – Not applicable

**Availability of data and materials** – Not applicable

**HCT116 culture information**

HCT116 culture information – Human cell line derived from colon cancer (RIKEN BioResource Centre, Japan)	
Year of deposit	2009
Animal	Human< Mammals
Gender	Male
Tissue	Colon
Classification	Cancer
Lifespan	Infinite
Morphology	Epithelial-like
STR (Human)	OK
Animal PCR	OK

## References

1. Hagggar FA, Boushey RP (2009) Colorectal cancer epidemiology: incidence, mortality, survival, and risk factors. *Clin Colon Rectal Surg* 22(4):191
2. Rawla P, Sunkara T, Barsouk A (2019) Epidemiology of colorectal cancer: Incidence, mortality, survival, and risk factors. *Przegląd Gastroenterologiczny* 14(2):89
3. Peltomäki P (1997) and A. De la Chapelle, Mutations predisposing to hereditary nonpolyposis colorectal cancer. *Adv Cancer Res* 71:94–119
4. Tutlewska K, Lubinski J, Kurzawski G (2013) Germline deletions in the EPCAM gene as a cause of Lynch syndrome–literature review. *Hereditary cancer in clinical practice* 11(1):9
5. Ligtenberg MJ et al (2013) EPCAM deletion carriers constitute a unique subgroup of Lynch syndrome patients. *Familial cancer* 12(2):169–174
6. Guarinos C et al (2010) EPCAM germ line deletions as causes of Lynch syndrome in Spanish patients. *J Mol Diagn* 12(6):765–770
7. Zhou F et al (2015) Expression of EpCAM and Wnt/beta-catenin in human colon cancer. *Genet Mol Res* 14(2):4485–4494
8. Ligtenberg MJ et al., Heritable somatic methylation and inactivation of MSH2 in families with Lynch syndrome due to deletion of the 3' exons of TACSTD1. *Nature genetics*, (2009). 41(1): 112
9. Miyaki M et al (1999) Frequent mutation of  $\beta$ -catenin and APC genes in primary colorectal tumors from patients with hereditary nonpolyposis colorectal cancer. *Cancer research* 59(18):4506–4509
10. Polakis P, Wnt signaling in cancer. *Cold Spring Harbor perspectives in biology*, (2012): a008052
11. Liang W, Mason AJ, Lam JK, *Western blot evaluation of siRNA delivery by pH-responsive peptides*, in *Target Identification and Validation in Drug Discovery*. 2013, Springer. p. 73–87

12. Pirini F et al (2019) Identification of a novel large EPCAM-MSH2 duplication, concurrently with LOHs in chromosome 20 and X, in a family with Lynch syndrome. *Int J Colorectal Dis* 34(11):1999–2002
13. Doan CC et al., Simultaneous silencing of VEGF and KSP by siRNA cocktail inhibits proliferation and induces apoptosis of hepatocellular carcinoma Hep3B cells. *Biological research*, (2014). 47(1): 70
14. Shen K et al (2012) MSH2 is required for cell proliferation, cell cycle control and cell invasiveness in colorectal cancer cells. *Chin Sci Bull* 57(20):2580–2585
15. Bultman SJ, Interplay between diet, gut microbiota, epigenetic events, and colorectal cancer. *Molecular nutrition & food research* (2017) **61**(1): 1500902
16. Nagasaka T et al (2010) Somatic hypermethylation of MSH2 is a frequent event in Lynch Syndrome colorectal cancers. *Cancer research* 70(8):3098–3108
17. Yamashita T et al (2007) Activation of hepatic stem cell marker EpCAM by Wnt- $\beta$ -catenin signaling in hepatocellular carcinoma. *Cancer research* 67(22):10831–10839
18. Novellasdemunt L, Antas P, Li VS (2015) Targeting Wnt signaling in colorectal cancer. A review in the theme: cell signaling: proteins, pathways and mechanisms. *American Journal of Physiology-Cell Physiology* 309(8):C511
19. Liu C et al (2002) Control of  $\beta$ -catenin phosphorylation/degradation by a dual-kinase mechanism. *Cell* 108(6):837–847
20. Castiglia D et al (2008) Concomitant activation of Wnt pathway and loss of mismatch repair function in human melanoma. *Genes Chromosom Cancer* 47(7):614–624
21. Belcheva A et al (2014) Gut microbial metabolism drives transformation of MSH2-deficient colon epithelial cells. *Cell* 158(2):288–299
22. Ji J, Yamashita T, Wang XW (2011) Wnt/beta-catenin signaling activates microRNA-181 expression in hepatocellular carcinoma. *Cell bioscience* 1(1):4
23. Network CGA (2012) Comprehensive molecular characterization of human colon and rectal cancer. *Nature* 487(7407):330