Prevention of UV-Induced Skin Cancer in Mice by Gamma Oryzanol-Loaded Nanoethosomes

Mahdi Zeinali

Biotechnology Research Center, Student Research Committee, and Faculty of Pharmacy, Tabriz University of Medical Sciences, Tabriz.

Soheil Abbaspour-Ravasjani

Drug Applied Research Center, Tabriz University of Medical Sciences, Tabriz.

Tannaz Soltanfam

Biotechnology Research Center, Student Research Committee, and Faculty of Pharmacy, Tabriz University of Medical Sciences, Tabriz.

Ana Cláudia Paiva-Santos

Department of Pharmaceutical Technology, Faculty of Pharmacy, University of Coimbra, Coimbra.

Hossein Babaei

Drug Applied Research Center, Tabriz University of Medical Sciences, Tabriz.

Francisco Veiga

Department of Pharmaceutical Technology, Faculty of Pharmacy, University of Coimbra, Coimbra.

Hamed Hamishehkar (✉ Hamishehkarh@tbzmed.ac.ir)

Drug Applied Research Center, Tabriz University of Medical Sciences, Tabriz.

Research Article

Keywords: Skin cancer, ultraviolet radiation (UV), cosmetics , γ -Oryzanol (GO), GO-loaded nanoethosomes (GO-NEs)

Posted Date: March 5th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-275658/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Prevention of UV-Induced Skin Cancer in Mice by Gamma Oryzanol-Loaded Nanoethosomes

Mahdi Zeinali ^{a,b}, Soheil Abbaspour-Ravasjani ^b, Tannaz Soltanfam ^a, Ana Cláudia Paiva-Santos ^{c,d}, Hossein Babaei ^b, Francisco Veiga ^{c,d}, Hamed Hamishehkar ^{b,*}

^a Biotechnology Research Center, Student Research Committee, and Faculty of Pharmacy, Tabriz University of Medical Sciences, Tabriz, Iran.

^b Drug Applied Research Center, Tabriz University of Medical Sciences, Tabriz, Iran.

^c Department of Pharmaceutical Technology, Faculty of Pharmacy of the University of Coimbra, University of Coimbra, Coimbra, Portugal

^d REQUIMTE/LAQV, Group of Pharmaceutical Technology, Faculty of Pharmacy of the University of Coimbra, University of Coimbra, Coimbra, Portugal

* Corresponding author:

Prof. Hamed Hamishehkar, Pharm D, Ph.D., Professor of Pharmaceutics, Drug Applied Research Center, Tabriz University of Medical Sciences, Tabriz, Iran.

Tele/Fax: +98 (41) 33363231, E-mails: Hamishehkarh@tbzmed.ac.ir, Hamishehkar.hamed@gmail.com

Abstract

Skin cancer is the most widespread cancer worldwide, mainly caused by exposure to ultraviolet radiation (UV) in sunlight. Utilizing topical preventive agents in routinely daily used cosmetics may prevent UV-related skin damages and skin cancers. γ -Oryzanol (GO) is a natural component derived from rice bran oil, with potential antioxidant and skin anti-aging properties. We biologically thorough studied the antioxidant and anticancer effects of GO *in vitro* to found the effective signaling pathways, then evaluated the sun protection factor of prepared formulation, and finally investigated the long-term preventive effects of GO-loaded nanoethosomes (GO-NEs) against UVB-induced skin cancer in mice. Our results suggest that GO-NEs could effectively prevent UVB-induced skin cancer and could be utilized as an innovative ingredient in cosmetics.

Introduction

Cancer is one of the serious human health problems globally, and its treatment is always associated with a variety of challenges ¹. Skin is the important and largest organ in the human body ² which, undergoes the most exposure to the environment, including ultraviolet (UV) radiation from sunlight. Consequently, skin cancer is the most common cancer diagnosed in Caucasians in the United States and Australia ³. Exposure to solar UVB radiation is a prominent cause of several skin disorders, particularly skin cancers caused by DNA damage and the generation of free radicals, i.e., reactive oxygen species (ROS) ⁴. Cosmetic products are increasingly being used throughout the world and are now being developed using nanotechnology ⁵. Nanotechnology-based cosmetics ⁶, and new ingredients with anticancer ⁷, antioxidant ⁸, and anti-inflammatory properties ⁹ have been introduced into cosmetic products to prevent and/or treat skin diseases. Recently, a few studies have demonstrated the effect of specific compounds e.g., naproxen ¹⁰, silver nanoparticles ¹¹, nicotinamide ¹², etc., on the prevention of UVB-induced skin cancers. The possible toxicity of chronic application of these compounds on the skin limits their routine administration as daily applied cosmeceuticals such as sunscreens. Natural products may overcome the problems mentioned above. In this regard, the anticancer activities of herbal extracts ¹³ like pterostilbene ¹⁴, honokiol ¹⁵, luteolin ¹⁶, tea polyphenols ¹⁷, sesame oil ¹⁸, quercetin ¹⁹, etc., have been shown against UVB-induced skin cancers. The main limitations of these natural products as far as the applications in the cosmeceutical industry is concerned, are dermal delivery obstacle due to considerable molecular weight, being expensive, having limited availability, and being obtained from difficult natural sources. Gamma-Oryzanol (GO) is a natural product derived from rice bran oil and consists of a mixture of at least ten phytosterol ferulates ²⁰. GO has been revealed to has good antioxidant

activity²¹, anti-inflammatory²², skin anti-aging²³, anticancer²⁴ properties, and its positive effect on increasing innate immune response in humans²⁵ has been reported. Despite the advantages of GO as mentioned above, its use as a novel agent for preventing solar UV-related skin cancer is hindered by the fact that its high molecular weight (>500 Da) and its hydrophobicity restrict dermal delivery across the stratum corneum (SC)²⁶. It has been shown that nanotechnology can be efficiently used for the delivery of such materials across the SC down into the epidermis and the dermis. Ethosomes are a modified type of liposomes with high ethanol added (20-45%) (Fig. 1) that can function as a novel dermal drug delivery system²⁷. The presence of a high amount of ethanol makes ethosomes more elastic and deformable than liposomes²⁸, and consequently, it makes ethosomes an appropriate carrier for enhanced dermal drug delivery.

In this study, we prepared nanoethosomes (NEs) containing GO (GO-NEs) and carried out completely cellular studies on murine melanoma cell line (B16F10 cells) for the first time and found its apoptosis mediated signaling pathways. Besides, we evaluated possible sun protection factor (SPF) of GO-NEs. Moreover, long-termly (22 weeks) investigated the potentiality of GO-NEs to prevent UVB-induced skin cancer *in vivo* to introduce a novel cosmeceutical formulation. The findings of the present study may open a new horizon on the formulation of novel sunscreens and other cosmeceuticals that could be used in populations faced with a high risk of skin cancers, such as Australians.

Results

Characterization of nanoethosomes. As illustrated in Fig. 2A, the characterization analysis reported the mean particle size (Fig. 2A-a) and zeta potential value (Fig. 2A-b) of NEs 80 ± 5 nm and -5 ± 0.9 mV, respectively. The SEM analysis displayed spherical morphology and confirmed particle size reports (Fig. 2A-c) and showed the narrow size distribution of particles reported by the PDI size study. TEM also revealed that NEs had ethosomal morphology (Fig. 2A-d). EE% and DLC% indexes of GO-NEs evaluated $96.36 \pm 2.14\%$ and $13.56 \pm 1.46\%$.

Evaluation of sun protection factor. The SPF of gel-mediated GO-NEs and free gel-medium were recorded as 3.68 ± 0.3 and 0.75 ± 0.3 , respectively.

Antioxidant stability assay and cellular ROS detection. The antioxidant activity for freshly prepared and stored for one year of GO, NEs, and GO-NEs, showed the concentration-dependent activity and the lack of significant scavenging activity of plain NEs (Fig. 2B-a). The IC_{50} for GO and GO-NEs according to DPPH assay on the first day of preparation were 90.8 and 103, respectively, and for GO-NEs was almost four times lower than free GO after one year (110 and 419.9 μ M, respectively.).

Moreover, the ROS induction capacity of GO and GO-NEs was studied on B16F10 cells with LC and HC of GO and GO-NEs. According to Fig. 2B-b, the ROS levels significantly increased in H_2O_2 (100 μ M) treated cells as compared to untreated cells ($p < 0.01$). Outcomes showed that the Vit. C, GO-NEs, and GO could reduce the oxidative stress in LC treatment around 29, 36, and 15 times compared with control positive, respectively. The statistical results between Vit C, GO-NEs, and GO showed a significant difference between GO and GO groups ($p > 0.01$), demonstrate the high ROS scavenging ability of the GO-NEs as compared to GO. Also, GO, and GO-NEs in HC treatments substantially increase the ROS and approve the double-edged sword hypothesis²⁹.

GO, and GO-NEs could increase significantly substantially increase the ROS substantially in B16F10 cells, and these amounts were 14 and 34 times higher than control, respectively.

Cellular cytotoxicity studies. Evaluation of cell viability with MTT assay showed a reduction in the cell viability by free GO and GO-NEs after 24 and 48 hours of treatment. As illustrated in Fig. 2C, it revealed that 1280 μ M of free GO and GO-NEs were able to decrease the cell viability to 35% and 30% after 48 h of treatment, respectively. The IC_{50} values results showed a significant difference ($P < 0.01$) between free GO and GO-NEs with 838 μ M and 255 μ M after 48 h, respectively. GO-NEs could improve the IC_{50} value 3.27 times more than free GO after 48 h of incubation. NEs revealed no cytotoxic effects.

The consequences from cell cycle investigation showed a significant cell cycle arrest in the G2/M phase in GO ($p < .01$) and GO-NEs ($p < .001$) after 48 h of treatment. Encapsulation of the GO significantly decreased G0/G1 phase cells from 67.4% to 47.3%. On the other hand, the GO's encapsulation increased the number of Sub G1 and G2/M cells from 4.47 % to 7.81 % and 16.4 to 29.7%, respectively (Fig. 3a).

B16F10 cells were stained by MDC, and cellular morphology changes were observed to investigate the effect of GO, GO-NEs, and blank NEs on autophagy induction in the cells. As shown in Fig. 3b, weak green fluorescence points are observed in blank NEs, GO, and control groups. While, cells treated with GO-NEs are entirely covered by fluorescence points, which indicated that GO-NEs affected a noticeable upsurge in the number of MDC-positive cells and a hallmark of autophagy.

Apoptosis and necrosis effects of GO and GO-NEs are shown in Fig. 4. Necrosis and apoptosis rate for GO were 15.7% and 14.23%, respectively, which were approximately equal, while for

120 GO-NEs reported 2.18% and 29.23%, respectively. B16F10 cells treated with blank NEs showed
121 15% apoptosis and less than 1% necrosis.

122 From the results of DAPI staining, the blank NEs showed a low rate of DNA alteration. Also,
123 GO-treated cells showed a low rate of DNA fragmentation, while GO-NEs increased the DNA
124 fragmentation rate as compared to GO (Fig. 4b).

125 A quantitative cellular uptake study (Fig. 5a) showed that the NEs had considerable cell uptake,
126 and its internalizations showed a dose-dependent response (Fig. 5b). These results were
127 confirmed by fluorescent imaging of the cells as qualitative results (Fig. 5c).

128 The inhibitory effect of GO and GO-NEs on cell migration was measured using a scratch healing
129 assay method. According to Fig. 6a and 6b, during 36 h of monitoring, both GO and GO-NEs
130 could display an inhibitory effect on cell migration. The migration rates during 12, 24, and 36 h
131 for GO were found to be 12.9, 26.4, and 36.8 %, respectively, additionally for GO-NEs were
132 found to be 5.2, 6.5, and 14.31 %, respectively. According to Fig. 6c, GO-NEs could inhibit
133 migration almost 1.4 times more than GO.

134 Moreover, utilizing formation assay, the effect of plain NEs, GO, and GO-NEs were evaluated.
135 As shown in Fig. 6d, the number of colonies decreased when treated by plain NEs and
136 significantly reduced by GO. The number of colonies drastically diminished after treatment with
137 GO-NEs.

138 **Molecular cytotoxicity studies.** Furthermore, we investigated the mRNA expression profile of
139 apoptosis-related pathway genes to prove and reach support that GO-NEs have perceived the
140 levels of apoptosis in B16F10 cells. Additionally, we completed the WB assay to quantify the
141 targeted proteins to verify the consequences of the RT-PCR assay. As shown in Fig. 7, we

142 remarked that both mRNA and protein expression levels of Bcl-2 and survivin were declined,
143 and the levels of P53, BAX, caspase-3, caspase-8, and caspase-9 were increased under treatment
144 B16F10 cells with GO and GO-NEs, while GO-NEs showed the largely effective in comparison
145 GO. Cells under treatment with free NEs showed no significant change in selected mRNA and
146 protein expression.

147 **Weight changes monitoring.** Weekly body weight measurements for 22 weeks are reported in
148 Fig. 8. After week 15, all groups showed weight loss. In week 15, the mice exposed with UV
149 showed significant weight decreases compared with the Ctrl group. By continuation of the
150 experiment, significant changes were observed between the groups. In the 18th week of the
151 investigation, all groups except the group treated with GO-NEs showed substantial weight loss
152 compared to the control group (Fig. 8 A). Finally, in the last week of the experiment, our results
153 showed that mice treated with GO-NEs did not lose significant weight compared to the control
154 group (Fig. 8 a).

155 **A qualitative study of GO-NEs protective effect against UV radiation.** Possible observational
156 preventative effects of formulations are presented in Fig. 9A. By these results, the skin cancer
157 prevention effect is seen in a group treated by GO-NEs, so that only a tiny lesion is visible (Fig.
158 9A-a). Skin lesions are observed in the groups treated by free GO (Fig. 9A-b) and NEs (Fig. 9A-
159 c), but a bit lower than those in the non-treated group (negative control group). Skin injury with a
160 large area and intensity was generated in the negative control group (Fig. 9A-d).

161 **Histopathological studies.** Histopathological results for skin samples from each group are
162 presented in Fig. 9B. By these results, in the group treated by GO-NEs, the epithelial and
163 connective tissues are entirely healthy. The hair follicle is reasonable, and there is no evidence

for any hyperplasia (Fig. 9B-a). In the group treated by free GO, although the epidermal layer was completely healthy, there were still adnexal tumors indicated by red narrow in Fig. 9B-b. In a group where only free NEs were applied, there are a large number of adnexal tumors. But in this sample, the epidermis layer was less destroyed, and irregular connective tissue arrangement is shown (Fig. 9B-c). As shown in Fig. 9B-d, which belongs to the positive control group, the epidermal and connective tissues are destroyed, and a high number of adnexal tumors have been formed. Many tumors were generated in the lower layers of the dermis, indicating the lesion's depth in the positive control group.

Discussion

The increased incidence of skin cancer proposes the usage of sunscreens lonely is not adequate in preventing skin cancer. It needs additional strategies like adding novel ingredients to daily cosmetics with skin cancer-preventing effects. GO is a natural product derived from rice bran oil. Previous studies verified that GO had well skin anti-aging effects²³. Therapeutic, cosmetic agents must penetrate through SC to have beneficial effects. Studies have demonstrated that particles in the size range of sub 100 nm present the best skin penetration³⁰. In this study, the dermal penetration problem was overwhelmed by nanotechnology to gain sub 100 nm and near-zero charged vesicular structure. The negatively close zero surface charge of particles is also promising for enhancing transdermal delivery³¹. Zero surface charge is necessary for colloidal stability; then the formulations were introduced in a gel-medium immediately after preparation to minimize dynamic changes in the particles and stabilize the NEs. DLC% of liposomes are usually not more than 10%³². In this case, more than 13% of DLC% was reported because of

186 higher hydrophobicity of GO, and addition hydrogen/oxygen bonds of GO structure, which could
187 form hydrogen bonds with the polar head group of lecithin ³³ in the outer layer of the ethosomal
188 system. These consequences could be the reason for the low negative zeta potential value and the
189 higher ethosomal system DLC%. Free radical scavenging assessment displayed a bit reduced
190 activity of the GO-NEs on the first day of preparation, which was possibly due to a temperature
191 effect during the preparation procedure. However, after one year of storage showed a strong
192 antioxidant protective effect. These results highlighted the carrier's effect ³⁴ and displayed an
193 advantage for the chemical stability of GO encapsulation into the ethosomal system. Studies
194 demonstrated that the nuclear factor erythroid 2–related factor 2 (Nrf2) plays a critical role in
195 acting as an emergent controller of cellular oxidants resistance to anti-inflammatory responses ³⁵.
196 Rungratanawanich et al. investigated the cellular antioxidant activity of GO and its related
197 signaling pathway. This study revealed that GO could successfully increase the upregulation of
198 Nrf2, additionally enhanced glutathione peroxide and superoxide dismutase, which are
199 antioxidant enzymes ³⁶. Antioxidant activity of our investigations at the cellular level showed
200 that the GO-NEs and free GO could reduce the oxidative stress and the supremacy of oxidative
201 stress reduction of GO-NEs in LC treatment, which highlighted the effect of the nanoethosomal
202 system. Moreover, ROS have been identified as signaling molecules in various pathways
203 regulating both cell survival and cell death ³⁷. Our results in HC treatment results shown that the
204 GO and GO-NEs can significantly increase the ROS levels; this phenomenon can be justified by
205 dubbing sword theory ³⁸.Recent research demonstrated that there is an effective relationship
206 between increased oxidative stress and induction of autophagy ³⁹. Following further
207 investigation, we found that GO and GO-NEs could activate the autophagy pathway in B16F10
208 cells. Our results can confirm that by increasing the ROS level, the autophagy pathway is

209 activated in B16F10 cells. As mentioned, antioxidants can have a double effect on inhibiting or
210 causing oxidative stress. To that end, due to GO effects on Nrf2 modulation and its role in anti-
211 inflammatory possesses, it is suggested to investigate the anti-inflammatory behavior of GO-NEs
212 as well. Our results in most of the tests showed that the formulated GO performance was better
213 than its free form. These results could be attributed to better cellular internalization of GO-NEs
214 compared to GO, highlighting the impact of nanoparticles (especially sub 100 nm size) on
215 cellular internalization ⁴⁰. Most of the antioxidants induce apoptosis in cancer cells ⁴¹. Our
216 apoptotic assessment results indicated that both GO and GO-NEs were able to induce apoptosis.
217 From the results, the blank NEs showed a low rate of DNA alteration and apoptosis rate due to
218 ethanol in the nanoethosomal formulation. These outcomes indicated the effect of sub 100 nm
219 GO-NEs that successfully increased B16F10 cells DNA alteration, apoptosis rate, and
220 consequently effectively decreased necrosis rate. The progression of apoptosis is linked with
221 higher endocytosis of the GO. Cell cycle plays a critical role in cell migration ⁴²; cell migration
222 occurred with a high velocity and directionality during the S–G2-phase of the cell cycle ⁴³. Our
223 results showed that the GO increases the cell population in the G2 phase, and it can lead to the
224 reduction of the cell migration rate. Scratch and colony formation assay tests were performed to
225 confirming this claim. The following results showed that the GO could inhibit the colony-
226 forming and cell migration. The obtained results proved its anticancer status and supremacy of
227 GO-NEs. The most important issue in the introduction of an anticancer agent is the identification
228 of the apoptosis pathways by alteration of the levels of apoptotic and anti-apoptotic genes. In our
229 study, the protein level of caspase-8, as apoptosis promoter, effectively increased, and
230 simultaneously, the levels of apoptotic inducers, including P53, BAX, and caspase-3,
231 dramatically increased. Increasing the levels of cleaved caspase-3/caspase-8 and concomitantly

describing the levels of pro-caspase-3/caspase-8 confirmed the activation of caspase-3 and caspase-8, respectively. Following, significantly describing in the protein level of survivin and Bcl-2 as anti-apoptotic inducers proved the anticancer behavior of GO and the superiority of GO-NEs by reducing anti-apoptotic proteins and increasing the levels of apoptotic proteins. *In vitro*, SFP measurements showed a bit of boost protection effects. It is important to note that the ingredient, which wants to be a novel cosmeceutical candidate, has sun protection effects, and these results highlighted the advantages of GO-NEs. Attained results proved *in vitro* cytotoxicity effects and showed a bit SPF benefit of GO-NEs that could be a justification to investigate the potential effect of GO-NEs on skin cancer prevention in animals. The safety of cosmetics that are taken routinely daily is more highlighted. Chaudhary et al. demonstrated the positive effect of naproxen on UVB-induced skin cancer ¹⁰. This study did not investigate the safety of naproxen for daily usage. Although adding chemical products to daily cosmetics are controversial due to toxicity issues. Besides, rice derived components generally are well-thought-out to be safe and non-allergenic, as well as investigations had reported no irritation when rice components tested on the skin ⁴⁴. Nowadays, the safety of natural products is a fascinating topic in these issues ⁴⁵. Our study, mean animal weight measurement during 22 weeks of the study, displayed no significant changes in GO-NEs treated group. Outcomes demonstrated a) the therapeutical effect of GO-NEs on preventing skin cancer and reducing its related stress, b) the safety of GO-NEs on long-term daily usage. Pathology results confirmed our hypothesis about the prevention of UV-Induced Skin cancer in mice by GO-NEs. Healthy epithelial tissue in the group treated by GO verified the potentiality therapeutical effect of GO. The generation of adnexal tumors in this group is due to GO could not penetrate through SC. Less destroyed epithelial tissue in the group

treated only NEs could be related to the nature of lipid used in ethosomal structure, which could cover the wound and speed up recovery.

Conclusion

In conclusion, GO-NE could effectively prevent UVB-induced skin cancer and suggest a new nanoethosomal formulation of GO. Our investigation opened up a new horizon, which may have novel clinical relevance as a pharmacologically active ingredient of cosmetics and sunscreens for the first time.

Experimental

Study design. This study has been approved by institutional review board of Faculty of Pharmacy and Drug Applied Research center, Tabriz University of Medical Sciences, Tabriz, Iran. All methods were carried out in accordance with relevant international regulations and guidelines.

Materials. The Soy lecithin was purchased from Lipoid GmbH, Ludwigshafen, Germany. GO was a generously gifted gift from Tsuno Rice Fine Chemicals Co., Ltd, Japan. The xanthan gum, 4',6-diamidino-2-phenylindole (DAPI), monodansylcadaverine (MDC), Rhodamine B (RhB), propidium iodide (PI), 2',7'-dichlorofluorescein diacetate (DCFH-DA), 2,2-diphenyl-1-picrylhydrazyl (DPPH), 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT), and ethanol 96% were purchased from Sigma-Aldrich co., St Louis, MO. RNase A was obtained from CinnaGen co., Iran. RPMI 1640, fetal bovine serum (FBS), Trizol, trypsin EDTA (0.25%),

and Penicillin-Streptomycin were purchased from Thermo Fisher Scientific, Waltham, Massachusetts. All of the WB reagents were purchased from Santa Cruz Biotechnology, Inc. Dallas, Texas.

Preparation of nanoethosomes. NEs were prepared via a modified ethanol injection method⁴⁶ using soy lecithin as a surfactant. Briefly, 350 mg of soy lecithin, and 50 mg of GO, were dissolved entirely in 20 ml of ethanol as lipid/GO solution (Fig. 1a) and slowly injected with 1ml/min flow rate into the 50 ml of heated distilled water with 65 °C temperature with mixing by a homogenizer (Silent Crusher M, Heidolph, Germany) (20,000 rpm) (Fig. 1b). The homogenization step was continued for 15 min after finishing the lipid/GO solution to achieve GO-NEs in sub-100 nm size. The empty nanoethosomal formulation (NEs without GO) was also prepared using the same method.

Preparation of gel-mediated nanoethosomal formulation. Prepared hydroethanolic suspension of GO-NEs was immediately transferred into a 250 ml beaker and 450 mg of xanthan gum was added slowly under stirring with a mixer (Ultraturrax, IKA, Germany) to prepare a gel-medium on a laboratory scale (Fig. 1d). For preparing GO gel-medium, GO (50 mg) was wholly dissolved at 20 ml of ethanol and added to 50 ml of distilled water. Next, xanthan gum was then added (1g of gel contains 0.085 % w/w GO). Gel-medium for plain NEs was also prepared in the same manner without the addition of GO.

Characterization of prepared ethosomes. Particle size and zeta potential of nanoethosomal formulations were assessed by laser diffraction technique (Nano ZS, Malvern, UK). Scanning electron microscope (SEM) (MIRA3 TESCAN, UK, LEO 84 1430VP, UK & Germany)

operating at 15 kV ⁴⁷, transmission electron microscopy (TEM) (TEM; LEO 906; Zeiss, Oberkochen, Germany) were used to study the morphology of prepared NEs.

Encapsulation efficacy. The amount of loaded GO was estimated to evaluate the encapsulation efficacy (EE%) and drug loading capacity (DLC%) indexes. For this, due to the low density of GO (0.32 g/cm³) ⁴⁸, the supernatant of nanoethosomal dispersion was precisely collected after gentle centrifugation at 500 rpm for 10 min. Then, 1 ml from the middle of the container was collected and mixed with 5 ml of chloroform, vortexed (LS100, Labtron, Iran) for 10 min for the destruction of the nanoethosomal structure and subsequently dissolving GO in chloroform. By measuring the absorbance of loaded GO spectrophotometrically (Ultrospec 2000, Pharmacia Biotech, UK) at 319 nm, the amount of entrapped GO was determined ⁴⁹. Following the EE% and DLC% were calculated using the following equations (1).

Radical scavenging activity. The antioxidant activity and stability during one year storage time of the GO and GO-NEs were determined using DPPH radical scavenging activity test ⁵⁰. Different concentrations of nanoethosomal formulation (10, 30, 50, 70, 90, 110, 130, 150, 170 µM) were prepared and mixed with DPPH solution (0.4 mM) by 1:1 ratio and incubated for 30 min in darkness and room temperature. Afterward, the absorbance at 520 nm was measured via a microplate reader system (Cytation 5.5, Biotek, USA).

Determination of sun protection factor. The SPF of GO-NEs was determined using the classical *in vitro* spectrophotometric method ⁵¹. To this purpose, 2 mg/cm² of gel-mediated GO-NEs was applied by a Transpore™ membrane, which mimics natural skin, and UV absorption of the sample was recorded from 290 to 400 nm.

Cell culture and seeding. Skin melanoma (B16F10) cells (Pasteur Institute, Iran) were maintained in 10% FBS contain RPMI-1640 and cultured at 37°C with 5% CO₂. The cells were sub-cultured 24–48 h later with an initial concentration of 4×10^4 cells/ml and used in the logarithmic phase in all experiments. The cells seeded at the density of 1.2×10^4 , 1×10^5 , and 2×10^5 cell/well for 96, 12, and 6-well plates, respectively. The treatment was done when confluency reached 70% for MTT assay, flow cytometry, and fluorescent imaging experiments. For the scratch migration assay, the required confluency was 100%, and the seeding concentration for colony formation assay was 1×10^3 cell/well.

Cell viability study. The MTT assay technique was used to evaluate the cytotoxicity effect of GO and GO-NEs on cell viability⁵². After seeding the cells in a 96-well plate and incubation for 24 h, the cells were treated with 100 µl different concentrations of GO and GO-NEs (5-1280 µM). After 24 and 48 h of treatment, the media was discarded, and cells were washed with PBS twice. Next, 150 µl of fresh media containing 50 µl of MTT solution (2 mg/ml in PBS) was added to each well of the plate. After four h, the incubated medium was replaced with 180 µl of DMSO and 20 µl of Sorensen's phosphate buffer. Then, the plates' absorption was read using a wavelength of 570 nm and a reference wavelength of 630 nm by the Elisa Reader instrument (TCAN, Austria).

Cellular ROS detection. DCFH-DA was employed as a fluorescent dye to evaluate intracellular ROS accumulation through flow cytometry analysis⁵³. Briefly, cells were cultured on a 6-well plate and treated for 48 h with Blank NEs, GO in low concentration (LC, based on DPPH assay IC₅₀ concentration) and high concentration (HC, Based on MTT assay IC₅₀ concentration), LC and HC of GO-NEs, Vit C as antioxidant control and H₂O₂ as a positive control. Concentrations were selected based on DPPH IC₅₀ for LC treatments and MTT assay IC₅₀ for HC treatments.

Then, cells were exposed to 500 μ l of 10 μ M of DCFH-DA. Next, de-attached after two h incubation and washed twice with PBS. Cells were resuspended in 500 μ l PBS and subjected to the flow cytometry (Becton Dickinson FACSCalibur, Franklin Lakes, New Jersey, United States) analysis using an FL2-H band-pass filter (FITC).

Cell uptake study. RhB as a fluorescent agent was used for labeling NEs to measure the cellular uptake intensity of the prepared formulations by cells ²⁸. Briefly, RhB was added to the NEs (0.05% w/w RhB ratio to lipid) during the same procedure carried out to prepare of GO-NEs, and unload RhB was separated with ultra-filtration method (Amicon® tube, 30 kDa Mw; Millipore, Germany). The cells were seeded in a 6-well plate, incubated overnight, and treated with 10, 100, 200, 400, 800, 1600, 3200, 6400, and 12800 μ g/ml (based on the lipid weight) concentrations of RhB labeled NEs for four h. The quantitative cellular uptake of the prepared formulations was measured by a flow cytometry instrument (MACS Quant 10, Miltenyi Biotech GmbH). For qualitative evaluation of the cell uptake, the B16F10 cells were seeded into lamel coated six-well palate and incubated with the same concentrations of the quantitative cell uptake measurements method and incubated for four h; after the incubation, the upper media was discarded and replaced with 1 ml f formaldehyde 4% solution and fixed for four h. after fixing the cells were photographed with fluorescence imaging microscopy system to identify cell uptake intensity (BX50, Olympus, Japan).

DAPI staining. The DAPI staining method was carried out for evaluating the DNA fragmentation in B16F10 cells ²⁸. For this purpose, cells were seeded onto 12 mm coverslips at the seeding density of 2×10^5 cells/well. After 48 h incubation, the cultured cells were treated with the IC₅₀ concentration of blank NEs, GO, and GO-NEs. After 48 h of treatment, the cells were washed with PBS twice and fixed with formalin 4% solution for four hours. Next, the cells

were washed with PBS once again and incubated for 15 min with Triton X100 solution (0.01% V/V) and DAPI solution (1 µg/ml) for 15 min. Finally, the stained cells were monitored by a fluorescence imaging microscopy system to identify apoptotic nuclei.

Detection of autophagic vacuoles by monodansylcadaverine. MDC is a fluorescent dye as the marker for visualization and identification of the autophagic process that was used for detection of the autophagy pathway activation level in the B16F10 cells. For this purpose, the cells were seeded onto 12 mm coverslips at the seeding density of 2×10^5 cells/well. After 48 h of incubation, the cultured cells were treated with the IC₅₀ concentration of blank NEs, GO, and GO-NEs. After 48 h of treatment, the cells were washed with PBS twice and stained with 0.05 mmol/L of MDC at 37°C for 10 min. Subsequently, the cells were washed three times to remove MDC in PBS and observed under a fluorescence imaging microscopy system⁵⁴.

Cell cycle analysis. B16F10 cells were cultivated at a density of 2×10^5 cells/well into six-well plates in four groups, including 1) negative control cells, 2) blank NEs, 3) GO, and 4) GO-NEs. The cells were harvested 48 h after the treatment, and following fixed with 75% of ice-cold ethanol overnight, treated with 500 µl of 20 µg/ml RNase A, stained with 500µl of 0.1 mg/ml PI contain 0.01% of Triton X-100 solution. Then the cells were washed with PBS and finally analyzed by flow cytometry for DNA synthesis and cell cycle status⁵⁵.

Apoptosis assay. The flow cytometry method was used to evaluate the cell death pathway treated with GO and GO-NEs. After seeding the cells into six-well plates and incubating overnight. The upper media of the plate wells were changed with 2 ml of fresh media containing the 48 h IC₅₀ concentration of each treatment group. Then cells were incubated for 48 h. Following, cells were detached and stained via an annexin V/PI apoptosis detection kit (Exbio,

385 Czech Republic) according to the manufacturer's protocol ⁵⁶. The rate of apoptosis was analyzed
386 by flow cytometry (MACS Quant 10, Miltenyi Biotech GmbH).

387 **In vitro scratch healing assay.** The scratch healing assay was employed to evaluate the *in vitro*
388 cell migration rate ⁵⁷. Briefly, cells were seeded into a 12-well plate and cultured until cells
389 formed a confluent monolayer. Afterward, a linear wound was created in the monolayer with a 3
390 mm thickness scratch pen. Next, cells were treated with 48 h IC₅₀ concentration of the free GO
391 and GO-NEs and incubated. The migration percent of cells was measured according to the
392 wound distance, which was recorded via an inverted microscope (LABOMED, TCM 400, USA)
393 at 4X magnification during various times (0, 6, 12, 24, and 36 h) after incubation, and compared
394 to the wound distance at the initial time.

395 **Colony formation assay.** For evaluating the ability of the GO and GO-NEs in colony formation
396 inhibition, the colony formation assay was performed. To perform this test, 2×10^3 cells per well
397 of B16F10 cells were seeded into a 6-well plate. They were then incubated and followed for 15
398 days. The colonies were fixed and stained with crystal violet ten times diluted solution with PBS,
399 and then photographed by iPhone 8 (Apple Inc. Cupertino, California, United States) and
400 counted via ImageJSoftware. ⁵⁸.

401 **Real-Time PCR.** Expression of subjected genes at the cells was assessed by real-time PCR (RT-
402 PCR) ⁵⁵. TRIzol was used to extracting RNA from the cells. At the first step, 5×10^4 cells per
403 well of B16F10 cells were seeded into six-well plates and incubated overnight; then the cells
404 were treated with 48 h IC₅₀ concentration of GO and GO-NEs. The cells were harvested after
405 48h and incubated with 1 ml of TRIzol solution (Thermo Fisher Scientific, Waltham,
406 Massachusetts, U.S) for 20 min; at the next step, the cells 200 μ l of chloroform was added to the

TRIZol cocktail and mixed gently, then the cocktail incubated in -20 refrigerator for 20 min. At the next step, the TRIzol/chloroform cocktail centrifuged at 12,000 RPM speed at 4-8°C for 15 min and the colorless upper aqueous phase was harvested and mixed with 500 µl of isopropyl alcohol and blended gently and incubated for 10 min then the mRNA cocktail for 10 minutes centrifuged at 12,000 RPM and the upper media was discarded and washed with isopropyl alcohol once again. At the last step, the mRNA sediment dried at room temperature and was solved with 20 µl Rnase free water. The concentration of the isolated mRNA was measured by a Nano-Drop spectrophotometer (2000, Thermo Fisher Scientific Life Sciences, Waltham, Massachusetts). The complementary DNA was generated by the QuantiTect Reverse Transcription kit (Qiagen, Hilden, Germany). RT-PCR was done by the Power SYBR GREEN mix (Applied Biosystems, Foster City, California). β -Actin was amplified as a reference gene. mRNA expression was measured with the $2^{-\Delta\Delta CT}$ method. Table 1 addressed the primer sequences used in RT-PCR reactions.

Western blot analysis. Total protein levels were extracted and quantified from the cells by RIPA protein extraction kit, and protein concentration was assessed by Nano-Drop spectroscopy. Before loading proteins into the wells, they mixed with sample buffer and boiled at 100°C for 5 min. Following, 50µg micrograms protein/lane whole cell lysates were electrophoresed in SDS-PAGE and transferred to a nitrocellulose membrane (Millipore, Billerica, Massachusetts, U.S). 2% non-fat milk was used to block non-specific antigens, and then specific primary antibodies (1:1000, SCB) were added, then incubated 18 h at 4 °C. The membrane was washed three times with buffer TBS-T for 15 min, incubated with anti-rabbit secondary antibody (1:1000, SCB), shake for 75 min at room temperature, chemiluminescence reaction was performed ⁵⁹.

Animal care and treatment. A bank of two TL 40W/12 RS UVB lamps (Philips, Holland) with a peak emission of 313 nm was used to mimic sunlight UV and expose mice to UV irradiation for inducing skin damage. Thirty male BALB/C mice (6 weeks old with a mean weight around 36 g) were randomly divided into five groups and housed in a standard room (temperature 25-27 ° C, 12h daylight cycle with air conditioning) one week before the start of animal studies. Animals in groups A, B, and C, were applied once daily GO-NEs gel, plain GO gel (1g of gel containing 0.085 % w/w GO), and blank nanoethosomal gel (gel-medium without GO), respectively. Group D and E did not receive any formulation as positive and negative control groups, respectively. Dorsal areas of animals were shaved weekly to diminish any UV covering and/or protective effects of animal hair and ease of applying the gels. For producing UVB induced skin lesions, groups A, B, C, and D were placed 25 cm under the UVB lamps and irradiated 200 J/m²/day, one h after topical application of formulations ¹⁵. The 25 cm was the best distance to gain a calculated UV dose of 200 J/m². While irradiating, to immobilize the animals and protect other areas from irradiation, mice were fixed in the mice restrainer was covered with aluminum foil and black tape. The limited upper part of the restrainer (the region in contact with the dorsal area of the animals) was only exposed area to UV light. Animal studies continued until skin lesions were observed, which were completed at 22 weeks. Weekly animal body weights were recorded during 22 weeks of study.

Histopathological study. After 22 weeks of study, the animals were sacrificed, and dorsal skin samples were removed and fixed in 10% formalin solution for hematoxylin and eosin (H&E) staining and subsequent examination under a light microscope in cooperation with a pathologist.

Statistical and data analysis. All experiments were repeated three times in succession. Statistical analysis was done using the independent t-test and Two-way analysis of variance

452 (Two-way ANOVA) with multiple assessments between confession data using a Tukey honest
453 noteworthy difference test (Graphpad Prism, version 8, San Diego, CA). A P-value of <0.05 was
454 considered significant. Data obtained from flow cytometry were analyzed using the FlowJo
455 software package (V10.5.3, Treestar, Inc., San Carlos, CA).

456 **References**

- 457 1 Zeinali, M. *et al.* Nanovehicles for co-delivery of anticancer agents. *Drug Discov. Today* **25**, 1416-
458 1430, doi:10.1016/j.drudis.2020.06.027 (2020).
- 459 2 Paiva-Santos, A. C. *et al.* Plant-mediated green synthesis of metal-based nanoparticles for
460 dermatopharmaceutical and cosmetic applications. *International Journal of Pharmaceutics* **597**,
461 120311, doi:<https://doi.org/10.1016/j.ijpharm.2021.120311> (2021).
- 462 3 Cronin, K. A. *et al.* Annual Report to the Nation on the Status of Cancer, part I: National cancer
463 statistics. *Cancer* **124**, 2785-2800, doi:10.1002/cncr.31551 (2018).
- 464 4 Lee, J. W., Ratnakumar, K., Hung, K. F., Rokunohe, D. & Kawasumi, M. Deciphering UV-induced
465 DNA Damage Responses to Prevent and Treat Skin Cancer. *Photochemistry and Photobiology* **96**,
466 478-499, doi:10.1111/php.13245 (2020).
- 467 5 Rahimpour, Y., Javadzadeh, Y. & Hamishehkar, H. Solid lipid microparticles for enhanced dermal
468 delivery of tetracycline HCl. *Colloids and Surfaces B: Biointerfaces* **145**, 14-20,
469 doi:10.1016/j.colsurfb.2016.04.034 (2016).
- 470 6 Rahimpour, Y. & Hamishehkar, H. Liposomes in cosmeceutics. *Expert Opinion on Drug Delivery* **9**,
471 443-455, doi:10.1517/17425247.2012.666968 (2012).
- 472 7 Su, S. *et al.* The pharmacological properties of chrysophanol, the recent advances. *Biomed.*
473 *Pharmacother.* **125**, doi:10.1016/j.biopha.2020.110002 (2020).
- 474 8 De Jesus E Silva De Almendra Freitas, B. *et al.* High 15-F2t-isoprostane levels in patients with a
475 previous history of nonmelanoma skin cancer: The effects of supplementary antioxidant
476 therapy. *BioMed Research International* **2015**, doi:10.1155/2015/963569 (2015).
- 477 9 Reinau, D., Surber, C., Jick, S. S. & Meier, C. R. Nonsteroidal anti-inflammatory drugs and the risk
478 of nonmelanoma skin cancer. *International Journal of Cancer* **137**, 144-153,
479 doi:10.1002/ijc.29357 (2015).
- 480 10 Chaudhary, S. C. *et al.* Naproxen Inhibits UVB-induced Basal Cell and Squamous Cell Carcinoma
481 Development in Ptch1+/-/SKH-1 Hairless Mice. *Photochemistry and Photobiology* **93**, 1016-
482 1024, doi:10.1111/php.12758 (2017).
- 483 11 Arora, S. *et al.* Silver nanoparticles protect human keratinocytes against UVB radiation-induced
484 DNA damage and apoptosis: potential for prevention of skin carcinogenesis. *Nanomed.*
485 *Nanotechnol. Biol. Med.* **11**, 1265-1275, doi:10.1016/j.nano.2015.02.024 (2015).
- 486 12 Chhabra, G., Garvey, D. R., Singh, C. K., Mintie, C. A. & Ahmad, N. Effects and Mechanism of
487 Nicotinamide Against UVA- and/or UVB-mediated DNA Damages in Normal Melanocytes.
488 *Photochemistry and Photobiology*, doi:10.1111/php.12994 (2018).
- 489 13 Babazadeh, A., Zeinali, M. & Hamishehkar, H. Nano-phytosome: A developing platform for
490 herbal anti-cancer agents in cancer therapy. *Current Drug Targets* **19**, 170-180,
491 doi:10.2174/1389450118666170508095250 (2018).
- 492 14 Sirerol, J. A. *et al.* Topical treatment with pterostilbene, a natural phytoalexin, effectively
493 protects hairless mice against UVB radiation-induced skin damage and carcinogenesis. *Free*
494 *Radical Biology and Medicine* **85**, 1-11, doi:10.1016/j.freeradbiomed.2015.03.027 (2015).
- 495 15 Chilampalli, S. *et al.* Chemopreventive effects of honokiol on UVB-induced skin cancer
496 development. *Anticancer Res.* **30**, 777-784 (2010).
- 497 16 Byun, S. *et al.* Luteolin inhibits protein kinase C ϵ and c-Src activities and UVB-induced skin
498 cancer. *Cancer Research* **70**, 2415-2423, doi:10.1158/0008-5472.CAN-09-4093 (2010).
- 499 17 Sharma, P. *et al.* Tea polyphenols for the prevention of UVB-induced skin cancer.
500 *Photodermatology Photoimmunology and Photomedicine* **34**, 50-59, doi:10.1111/phpp.12356
501 (2018).

502 18 Ghorbanzadeh, M., Farhadian, N., Golmohammadzadeh, S., Karimi, M. & Ebrahimi, M.
503 Formulation, clinical and histopathological assessment of microemulsion based hydrogel for UV
504 protection of skin. *Colloids and Surfaces B: Biointerfaces* **179**, 393-404,
505 doi:10.1016/j.colsurfb.2019.04.015 (2019).

506 19 Bagde, A. *et al.* Combination of UVB Absorbing Titanium Dioxide and Quercetin Nanogel for Skin
507 Cancer Chemoprevention. *AAPS PharmSciTech* **20**, doi:10.1208/s12249-019-1424-x (2019).

508 20 Xu, Z. & Godber, J. S. Purification and identification of components of γ -oryzanol in rice bran oil.
509 *Journal of Agricultural and Food Chemistry* **47**, 2724-2728, doi:10.1021/jf981175j (1999).

510 21 Toorani, M. R., Golmakani, M. T. & Hashemi Gahrue, H. Antioxidant activity and inhibitory
511 mechanism of γ -oryzanol as influenced by the unsaturation degree of lipid systems. *LWT* **133**,
512 doi:10.1016/j.lwt.2020.109930 (2020).

513 22 Zolali, E. *et al.* Effects of gamma oryzanol on factors of oxidative stress and sepsis-induced lung
514 injury in experimental animal model. *Iranian Journal of Basic Medical Sciences* **18**, 1257-1263
515 (2015).

516 23 Heydari, S. *et al.* Nanoethosomal formulation of gammaoryzanol for skin-aging protection and
517 wrinkle improvement: a histopathological study. *Drug Dev. Ind. Pharm.* **43**, 1154-1162,
518 doi:10.1080/03639045.2017.1300169 (2017).

519 24 Hirsch, G. E. *et al.* γ -oryzanol reduces caveolin-1 and PCGEM1 expression, markers of
520 aggressiveness in prostate cancer cell lines. *Prostate* **75**, 783-797, doi:10.1002/pros.22960
521 (2015).

522 25 Shin, S. Y. *et al.* γ -Oryzanol-Rich Black Rice Bran Extract Enhances the Innate Immune Response.
523 *Journal of Medicinal Food* **20**, 855-863, doi:10.1089/jmf.2017.3966 (2017).

524 26 Manosroi, A., Chutoprapat, R., Abe, M., Manosroi, W. & Manosroi, J. Transdermal absorption
525 enhancement of rice bran bioactive compounds entrapped in niosomes. *AAPS PharmSciTech* **13**,
526 323-335, doi:10.1208/s12249-012-9751-1 (2012).

527 27 Das, S. K. *et al.* Ethosomes as novel vesicular carrier: An overview of the principle, preparation
528 and its applications. *Current Drug Delivery* **15**, 795-817,
529 doi:10.2174/1567201815666180116091604 (2018).

530 28 Gaballu, F. A. *et al.* Comparative of in-vitro evaluation between erlotinib loaded nanostructured
531 lipid carriers and liposomes against A549 lung cancer cell line. *Iran. J. Pharm. Res.* **18**, 1168-
532 1179, doi:10.22037/ijpr.2019.1100775 (2019).

533 29 Asadi-Samani, M., Farkhad, N. K., Mahmoudian-Sani, M. R. & Shirzad, H. in *Antioxidants*
534 (IntechOpen, 2019).

535 30 Adib, Z. M., Ghanbarzadeh, S., Kouhsoltani, M., Khosroshahi, A. Y. & Hamishehkar, H. The effect
536 of particle size on the deposition of solid lipid nanoparticles in different skin layers: A
537 histological study. *Advanced Pharmaceutical Bulletin* **6**, 31-36, doi:10.15171/apb.2016.006
538 (2016).

539 31 Gillet, A. *et al.* Liposome surface charge influence on skin penetration behaviour. *International*
540 *Journal of Pharmaceutics* **411**, 223-231, doi:10.1016/j.ijpharm.2011.03.049 (2011).

541 32 Cheng, C. *et al.* Improvement on stability, loading capacity and sustained release of rhamnolipids
542 modified curcumin liposomes. *Colloids and Surfaces B: Biointerfaces* **183**,
543 doi:10.1016/j.colsurfb.2019.110460 (2019).

544 33 Tai, K. *et al.* The effect of sterol derivatives on properties of soybean and egg yolk lecithin
545 liposomes: Stability, structure and membrane characteristics. *Food Research International* **109**,
546 24-34, doi:10.1016/j.foodres.2018.04.014 (2018).

547 34 Viriyaroj, A. *et al.* Physicochemical properties and antioxidant activity of gamma-oryzanol-
548 loaded liposome formulations for topical use. *Pharmaceutical Development and Technology* **14**,
549 665-671, doi:10.3109/10837450902911937 (2009).

550 35 Kahroba, H. & Davatgaran-Taghipour, Y. Exosomal Nrf2: From anti-oxidant and anti-
551 inflammation response to wound healing and tissue regeneration in aged-related diseases.
552 *Biochimie* **171-172**, 103-109, doi:10.1016/j.biochi.2020.02.011 (2020).

553 36 Rungratanawanich, W. *et al.* Characterization of the antioxidant effects of γ -oryzanol:
554 Involvement of the Nrf2 Pathway. *Oxidative Med. Cell. Longevity* **2018**,
555 doi:10.1155/2018/2987249 (2018).

556 37 Dianat-Moghadam, H. *et al.* Cancer stem cells-emanated therapy resistance: Implications for
557 liposomal drug delivery systems. *J. Control. Release* **288**, 62-83,
558 doi:<https://doi.org/10.1016/j.jconrel.2018.08.043> (2018).

559 38 Bouayed, J. & Bohn, T. Exogenous antioxidants - Double-edged swords in cellular redox state:
560 Health beneficial effects at physiologic doses versus deleterious effects at high doses. *Oxidative*
561 *Med. Cell. Longevity* **3**, 228-237, doi:10.4161/oxim.3.4.12858 (2010).

562 39 Meghan B, A., Chen., Y. & Spencer B, G. Regulation of Autophagy by Reactive Oxygen Species
563 (ROS): Implications for Cancer Progression and Treatment. *Antioxidants & Redox Signaling* **11**,
564 777-790, doi:10.1089/ars.2008.2270 (2009).

565 40 Pattni, B. S., Chupin, V. V. & Torchilin, V. P. New Developments in Liposomal Drug Delivery.
566 *Chem. Rev.* **115**, 10938-10966, doi:10.1021/acs.chemrev.5b00046 (2015).

567 41 Ilghami, R. *et al.* The conundrum of dietary antioxidants in cancer chemotherapy. *Nutr. Rev.* **78**,
568 65-76, doi:10.1093/nutrit/nuz027 (2020).

569 42 Boehm, M. & Nabel Elizabeth, G. Cell Cycle and Cell Migration. *Circulation* **103**, 2879-2881,
570 doi:10.1161/01.CIR.103.24.2879 (2001).

571 43 Chu, T. L. H. *et al.* Cell cycle-dependent tumor engraftment and migration are enabled by
572 aurora-A. *Mol. Cancer Res.* **16**, 16-31, doi:10.1158/1541-7786.MCR-17-0417 (2018).

573 44 Amended final report on the safety assessment of Oryza sativa (Rice) Bran Oil, Oryza Sativa
574 (Rice) Germ Oil, Rice Bran Acid, Oryza Sativa (Rice) Bran Wax, Hydrogenated Rice Bran Wax,
575 Oryza Sativa (Rice) Bran Extract, Oryza Sativa (Rice) Extract, Oryza Sativa (Rice) Germ Powder,
576 Oryza Sativa (Rice) Starch, Oryza Sativa (Rice) Bran, Hydrolyzed Rice Bran Extract Hydrolyzed
577 Rice Bran Protein. *International Journal of Toxicology* **25**, 91-120,
578 doi:10.1080/10915810600964626 (2006).

579 45 Saeedi, M., Eslamifar, M. & Khezri, K. Kojic acid applications in cosmetic and pharmaceutical
580 preparations. *Biomed. Pharmacother.* **110**, 582-593, doi:10.1016/j.biopha.2018.12.006 (2019).

581 46 Babaei, S. *et al.* Enhanced skin penetration of lidocaine through encapsulation into
582 nanoethosomes and nanostructured lipid carriers: A comparative study. *Pharmazie* **71**, 247-251,
583 doi:10.1691/ph.2016.5158 (2016).

584 47 Nazari, M., Ghanbarzadeh, B., Samadi Kafil, H., Zeinali, M. & Hamishehkar, H. Garlic essential oil
585 nanophytosomes as a natural food preservative: Its application in yogurt as food model. *Colloids*
586 *Interface Sci. Commun.* **30**, doi:10.1016/j.colcom.2019.100176 (2019).

587 48 Ishaka, A., Imam, M. U., Ismail, M., Mahmud, R. & Abu Bakar, Z. Z. Nanoemulsified gamma-
588 oryzanol rich fraction blend regulates hepatic cholesterol metabolism and cardiovascular
589 disease risk in hypercholesterolaemic rats. *Journal of Functional Foods* **26**, 338-349,
590 doi:10.1016/j.jff.2016.08.015 (2016).

591 49 Nazari, M. *et al.* Cinnamon nanophytosomes embedded electrospun nanofiber: Its effects on
592 microbial quality and shelf-life of shrimp as a novel packaging. *Food Packaging Shelf Life* **21**,
593 doi:10.1016/j.fpsl.2019.100349 (2019).

594 50 Akbarbaglu, Z. *et al.* Influence of spray drying encapsulation on the retention of antioxidant
595 properties and microstructure of flaxseed protein hydrolysates. *Colloids and Surfaces B:*
596 *Biointerfaces* **178**, 421-429, doi:10.1016/j.colsurfb.2019.03.038 (2019).

- 51 Diffey, B. L. & Robson, J. The influence of pigmentation illumination on the perception of erythema. *Photodermatology Photoimmunology and Photomedicine* **9**, 45-47 (1992).
- 52 Sabzichi, M. *et al.* A comparative cytotoxicity study of doxorubicin loaded PLGA, P (NIPAAm-MAA-DMAEMA) and solid lipid nanoparticles as anticancer drug delivery systems. *Lat. Am. J. Pharm.* **36**, 750-757 (2017).
- 53 Ghorbani, M. *et al.* Nanozyme antioxidants as emerging alternatives for natural antioxidants: Achievements and challenges in perspective. *Nano Today* **29**, doi:10.1016/j.nantod.2019.100775 (2019).
- 54 Chen, X., Zhang, L., Ding, S., Lei, Q. & Fang, W. Cisplatin combination drugs induce autophagy in HeLa cells and interact with HSA via electrostatic binding affinity. *RSC Adv.* **7**, 22270-22279, doi:10.1039/c7ra00056a (2017).
- 55 Mansoori, B. *et al.* Hyaluronic acid-decorated liposomal nanoparticles for targeted delivery of 5-fluorouracil into HT-29 colorectal cancer cells. *J. Cell. Physiol.*, doi:10.1002/jcp.29576 (2020).
- 56 Khaki-Khatibi, F. *et al.* Harnessing WYE-132 as an inhibitor of the mTOR signaling enriches the cytotoxicity effect of vinblastine in B16F10 melanoma cancer cells. *Process Biochem.* **99**, 123-130, doi:10.1016/j.procbio.2020.08.026 (2020).
- 57 Farzaneh Behelgard, M., Zahri, S., Mashayekhi, F., Mansouri, K. & Asghari, S. M. A peptide mimicking the binding sites of VEGF-A and VEGF-B inhibits VEGFR-1/-2 driven angiogenesis, tumor growth and metastasis. *Sci. Rep.* **8**, doi:10.1038/s41598-018-36394-0 (2018).
- 58 Masjedi, A. *et al.* Silencing of IL-6 and STAT3 by siRNA loaded hyaluronate-N,N,N-trimethyl chitosan nanoparticles potently reduces cancer cell progression. *International Journal of Biological Macromolecules* **149**, 487-500, doi:10.1016/j.ijbiomac.2020.01.273 (2020).
- 59 Joseph, M. M. *et al.* Anticancer activity of galactoxyloglucan polysaccharide-conjugated doxorubicin nanoparticles: Mechanistic insights and interactome analysis. *European Journal of Pharmaceutics and Biopharmaceutics* **93**, 183-195, doi:10.1016/j.ejpb.2015.04.001 (2015).

Acknowledgment. We are grateful to Behzad Mansoori, Mehdi Talebi, Seyed Kazem Forghani, Maryam Kuhsoltani, Shaho Ghahremani-Dehbokri and all of other colleagues assisted us in this project. This study was a part of Mahdi Zeinali's Pharm D thesis and financially supported by Faculty of Pharmacy and Drug Applied Research Center, Tabriz University of Medical Sciences, Tabriz, Iran (Grant number: 58549).

Author contributions. M.Z. NEs preparation and characterization, *in vitro* and *in vivo* experiments, investigation, methodology and writing the original draft. S.A.R. Cellular experiments, cellular and molecular section methodology, the original draft writing, figures preparation, data analysis. T.S. Characterization and toxicology advice. A.C.P.S Scientific editing the original draft. H.B Supervision. F.V Scientific editing the original draft. H.H. Supervision, project administration, and funding acquisition.

Competing interests. The authors declare no competing interests.

Ethics declarations. Animal studies were carried out in compliance with the ARRIVE guidelines and in agreement with the protocols of the care and use of laboratory animals of Tabriz University of Medical Sciences, Tabriz, Iran (National Institutes of Health Publication No. 85-23, revised 1985) and under license of the ethical committee of the university (issued certificate No. IR.TBZMED.VCR.REC.1397.152).

Data availability statement. The data that support the findings of this study are available from the corresponding authors upon reasonable request.

Figure legends

Figure 1. A brief illustration of the preparation of gamma oryzanol-loaded nanoethosomes (GO-NEs), *in vitro/in vivo* studies, and schematically comparison of ethosome and liposome.

Figure 2. (A): Size (a), zeta potential (b), scanning electron microscopy image (c), and transmission electron microscopy (d) of optimized gamma oryzanol-loaded nanoethosomes (GO-NEs). (B): Antioxidant activity (diphenyl-2-trinitrobenzene hydrazine (DPPH) inhibition) of GO, NEs, and GO-NEs at the first day of preparation (D1) and after one year (Y1) of storage (a). flow cytometry analysis of ROS induction in B16F10 (murine melanoma) cells and inhibitory effect of GO and GO-NEs (b). (C): Cell viability study of GO, NEs, and GO-NEs on B16F10 cells during 24 and 48 h. The data represent mean $\pm$ SD (n = 3); **p < .01, ***p < .001, and ****p < .0001, versus multiple comparisons.

Figure 3. (a): Cell cycle progression and the percent of the cell population of B16F10 (murine melanoma) cells treated with gamma oryzanol (GO), nanoethosomes (NEs), GO loaded NEs (GO-NEs), and untreated cells as the control group (Ctrl). (b): Autophagy study at B16F10 treated with GO, NEs, GO-NEs and Ctrl group after 48 h of treatment, which cells were stained with monodansylcadaverine (MDC) staining and observed under fluorescence imaging microscopy system (scale bars, 20 μ m). The data represent mean $\pm$ SD (n = 3); **p < .01, ***p < .001, and ****p < .0001, versus multiple comparisons.

Figure 4. (a): Apoptotic rates of the control group as untreated (Ctrl) and treated B16F10 (murine melanoma) cells with gamma oryzanol (GO), nanoethosomes (NEs), and GO-loaded NEs (GO-NEs) over 48 h by using Annexin V-FITC/PI staining. (b): DAPI stained DNA morphology images of Ctrl, DAPI staining B16F10 cells and treated with GO, blank NEs, and GO-NEs. Data represent mean $\pm$ SD (n=3).

Figure 5. (A): Flow cytometry analysis of Rhodamine B (RhB) labeled nanoethosomes (NEs) after 24 h of incubation. Histogram of B16F10 (murine melanoma) cells treated with different concentration of NEs (from 10 to 12800 $\mu\text{g/ml}$ based on lipid weight) (a); mean fluorescence intensity of the B16F10 cells (b); fluorescent images of RhB-labeled NE cellular uptake in different concentration (from 10 to 12800 $\mu\text{g/ml}$ based on lipid weight) (c). Data represent mean $\pm$ SD (n=3).

Figure 6. (a) *In vitro* wound-healing/scratch assay on B16F10 (murine melanoma) cells until 36 h and (b) the last morphology picture (36 h) of cells. (c) Cell migration percent values calculated from scratch test results. (d) Colony formation assay on untreated cells (Ctrl), blank nanoethosomes (NEs), free gamma oryzanol (GO) and GO-loaded NEs (GO-NEs). The data represent mean $\pm$ SD (n = 3); **p < .01, ***p < .001, and ****p < .0001, versus multiple comparisons.

Figure 7. mRNA and protein levels of β -Actin, Bcl-2, BAX, P53, survivin, caspase-3, caspase-8 and caspase-9 on B16F10 (murine melanoma) cells treated by free nanoethosome (NEs), gamma oryzanol (GO) and GO-loaded NEs (GO-NEs). The data represent mean $\pm$ SD (n = 3); **p < .01, ***p < .001, and ****p < .0001, versus multiple comparisons.

Figure 8. Mean mice body weight measurement of each group that applying (a) gamma oryzanol loaded nanoethosomes (GO-NEs), (b) GO, (c) blank NEs, (d) untreated UV-exposed mice, and (e) negative control group (received no UV and no treatment) during 22 weeks of study. The data represent mean $\pm$ SD (n = 6); **p < .01, ***p < .001, and ****p < .0001, versus multiple comparisons.

Figure 9. (A): The severity of skin cancer induction by ultraviolet B (UVB) lamps (200 J/m²/day) on treated BALB/c mice with (a) gamma oryzanol loaded nanoethosomes (GO-NEs), (b) GO, (c) blank NE, and (d) non-treated UV exposed mice after 22 weeks. The size of skin lesion in the free GO gel, and nano-formula groups is much smaller than the blank NEs and non-treated UV exposed group. (B): H and E (20×) of animal skin tissue samples; (a) GO-NEs, (b) GO, (c) blank NE, (d) positive control group (no treatment) and (e) negative control group (received no UV and no treatment). The black arrow shows normal epithelial tissue, red arrows indicate adnexal tumors, and the yellow arrow shows normal hair follicles. UV radiation destroyed the epidermal and connective tissues and induced a high number of adnexal tumors. Also, generate many tumors in the lower layers of the dermis, indicating the lesion's depth. Blank NEs gel does not show any remarkable protective effect against damage caused by ultraviolet radiation. Applying the Free GO gel on the skins of the mice protect the epidermal layer against UV radiation, but adnexal tumors remain. Using the GO-NEs gel on the skin of the mice protect epithelial and connective tissues against adnexal tumors establishment caused by UV radiation.

702 **Table 1.** Primer sequences.

Gene	Forward	Reverse
β-actin	5'-GGTCATCACTATTGGCAACG-3'	5'-ACGGATGTCAACGTCACACT-3'
Bax	TCCCTGGAGAAGAGCTACG	GTAGTTTCGTGGATGCCACA
Bcl2	5'- GGCTGGGGATGACTTCTCTC-3'	5'- ACAATCCTCCCCCAGTTCAC-3'
Caspase 3	TGGTGATGAAGGGGTCATTTATG	TTCGGCTTTCCAGTCAGACTC
Caspase 8	TGGTTCATCCAGTCGCTTTG	AATTCTGTTGCCACCTTTTCG
Caspase 9	CTGTCTACGGCACAGATGGAT	GGGACTCGTCTTCAGGGGAA
P ₅₃	5'-TCAGATCCTAGCGTCGAGCCC-3'	5'-GGGTGTGGAATCAACCCACAG-3'
Survivin	GACCACCGCATCTCTACATTC	TGCTTTTTTATGTTCCCTCTATGGG

703

704

705 **Equation 1:**

706 $EE\% = \frac{W_{EE}}{W_{Initial}} \times 100$

707 $DLC\% = \frac{W_{EE}}{W_{Lipid}} \times 100$

708 Where W_{EE} is the total amount of loaded GO, and $W_{Initial}$ is the amount initial GO used for
709 loading in the nanoethosomal system, W_{Lipid} is the amount of the lipid complex used in preparing
710 the.

711

712 **Fig. 1**

713

714

715 **Fig. 2**

716

717

Fig. 4

725

726

729 **Fig. 7**

730

731

732 **Fig. 8**

a

b

A

B

Figures

Fig. 1

Figure 1

A brief illustration of the preparation of gamma oryzanol-loaded nanoethosomes (GO-NEs), in vitro/in vivo studies, and schematically comparison of ethosome and liposome.

Fig. 2

Figure 2

(A): Size (a), zeta potential (b), scanning electron microscopy image (c), and transmission electron microscopy (d) of optimized gamma oryzanol-loaded nanoethosomes (GO-NEs). (B): Antioxidant activity (diphenyl-2-trinitrobenzene hydrazine (DPPH) inhibition) of GO, NEs, and GO-NEs at the first day of preparation (D1) and after one year (Y1) of storage (a). flow cytometry analysis of ROS induction in B16F10 (murine melanoma) cells and inhibitory effect of GO and GO-NEs (b). (C): Cell viability study of

GO, NEs, and GO-NEs on B16F10 cells during 24 and 48 h. The data represent mean $\pm$ SD (n = 3); **p < .01, ***p < .001, and ****p < .0001, versus multiple comparisons.

Fig. 3

Ctrl

Blank NEs

GO

GO-NEs

Figure 3

(a): Cell cycle progression and the percent of the cell population of B16F10 (murine melanoma) cells treated with gamma oryzanol (GO), nanoethosomes (NEs), GO loaded NEs (GO-NEs), and untreated cells as the control group (Ctrl). (b): Autophagy study at B16F10 treated with GO, NEs, GO-NEs and Ctrl group after 48 h of treatment, which cells were stained with monodansylcadaverine (MDC) staining and observed under fluorescence imaging microscopy system (scale bars, 20 μ m). The data represent mean $\pm$ SD (n = 3); **p < .01, ***p < .001, and ****p < .0001, versus multiple comparisons.

Fig. 4

Figure 4

(a): Apoptotic rates of the control group as untreated (Ctrl) and treated B16F10 (murine melanoma) cells with gamma oryzanol (GO), nanoethosomes (NEs), and GO-loaded NEs (GO-NEs) over 48 h by using Annexin V-FITC/PI staining. (b): DAPI stained DNA morphology images of Ctrl, DAPI staining B16F10 cells and treated with GO, blank NEs, and GO-NEs. Data represent mean $\pm$ SD (n=3).

Fig. 5

Figure 5

(A): Flow cytometry analysis of Rhodamine B (RhB) labeled nanoethosomes (NEs) after 24 h of incubation. Histogram of B16F10 (murine melanoma) cells treated with different concentration of NEs (from 10 to 12800 $\mu\text{g/ml}$ based on lipid weight) (a); mean fluorescence intensity of the B16F10 cells (b); fluorescent images of RhB-labeled NE cellular uptake in different concentration (from 10 to 12800 $\mu\text{g/ml}$ based on lipid weight) (c). Data represent mean $\pm$ SD (n=3).

Fig. 6

Figure 6

(a) In vitro wound-healing/scratch assay on B16F10 (murine melanoma) cells until 36 h and (b) the last morphology picture (36 h) of cells. (c) Cell migration percent values calculated from scratch test results. (d) Colony formation assay on untreated cells (Ctrl), blank nanoethosomes (NEs), free gamma oryzanol (GO) and GO-loaded NEs (GO-NEs). The data represent mean $\pm$ SD ($n = 3$); ** $p < .01$, *** $p < .001$, and **** $p < .0001$, versus multiple comparisons.

Fig. 7

Figure 7

mRNA and protein levels of β -Actin, Bcl-2, BAX, P53, survivin, caspase-3, caspase-8 and caspase-9 on B16F10 (murine melanoma) cells treated by free nanoethosome (NEs), gamma oryzanol (GO) and GO-loaded NEs (GO-NEs). The data represent mean $\pm$ SD ($n = 3$); ** $p < .01$, *** $p < .001$, and **** $p < .0001$, versus multiple comparisons.

Fig. 8

a

b

Figure 8

Mean mice body weight measurement of each group that applying (a) gamma oryzanol loaded nanoethosomes (GO-NEs), (b) GO, (c) blank NEs, (d) untreated UV-exposed mice, and (e) negative control group (received no UV and no treatment) during 22 weeks of study. The data represent mean $\pm$ SD (n = 6); **p < .01, ***p < .001, and ****p < .0001, versus multiple comparisons.

Fig. 9

Figure 9

(A): The severity of skin cancer induction by ultraviolet B (UVB) lamps (200 J/m²/day) on treated BALB/c mice with (a) gamma oryzanol loaded nanoethosomes (GO-NEs), (b) GO, (c) blank NE, and (d) non-treated UV exposed mice after 22 weeks. The size of skin lesion in the free GO gel, and nano-formula groups is much smaller than the blank NEs and non-treated UV exposed group. (B): H and E (20×) of animal skin tissue samples; (a) GO-NEs, (b) GO, (c) blank NE, (d) positive control group (no

treatment) and (e) negative control group (received no UV and no treatment). The black arrow shows normal epithelial tissue, red arrows indicate adnexal tumors, and the yellow arrow shows normal hair follicles. UV radiation destroyed the epidermal and connective tissues and induced a high number of adnexal tumors. Also, generate many tumors in the lower layers of the dermis, indicating the lesion's depth. Blank NEs gel does not show any remarkable protective effect against damage caused by ultraviolet radiation. Applying the Free GO gel on the skins of the mice protect the epidermal layer against UV radiation, but adnexal tumors remain. Using the GO-NEs gel on the skin of the mice protect epithelial and connective tissues against adnexal tumors establishment caused by UV radiation.