

Phytoremediation effect of *Medicago sativa* colonized by *Piriformospora indica* in the phenanthrene and cadmium co-contaminated soil

Liang Li (✉ liangli@hebut.edu.cn)

Pengyue Zhu

Tianjin University

Xiaoyang Wang

Hebei University of Technology

Zhenhua Zhang

Hebei University of Technology

Research article

Keywords: *Piriformospora indica*; *Medicago sativa*; Phytoremediation; Soil-contamination; Bioavailability; Rhizospheric soil

Posted Date: July 26th, 2019

DOI: <https://doi.org/10.21203/rs.2.11980/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Biotechnology on April 28th, 2020. See the published version at <https://doi.org/10.1186/s12896-020-00613-2>.

Abstract

Background: Coexistence of polycyclic aromatic hydrocarbons (PAHs) and heavy metals deleteriously threatens the quality of environmental health. Few reports uncover the mechanism of inoculation plants with *Piriformospora indica* for remediating PAH-metal co-contaminated soil by analyzing the chemical speciations of contaminants. This study investigated the influence of inoculation *Medicago sativa* with *Piriformospora indica* to remediate phenanthrene (kind of PAHs), and cadmium (one of heavy metals) co-contaminated soil by analyzing the plant growth, physiological parameters and chemical speciation in rhizospheric and non-rhizospheric soil. Results: The presence of *P. indica* significantly increased plants tolerance, Chlorophyll a, Chlorophyll b, maximum quantum efficiency of PSII photochemistry and electron transport rate values in phenanthrene and/or cadmium contaminated soil. *P. indica* inoculation in *M. sativa* root increased fluorescein diacetate activities in phenanthrene, cadmium and both of that co-contaminated soil, especially in non-rhizospheric soil. The presence of phenanthrene hindered the inoculated plant from accumulating cadmium to some extent; Whereas the presence of cadmium did not hinder the degradation of phenanthrene in both rhizospheric and non-rhizospheric soil after *P. indica* colonization. Although the poor bioavailability of cadmium in rhizospheric soil restricted the transportation into stem, *P. indica* colonization in plant efficiently increased cadmium accumulation in root in cadmium and phenanthrene co-contaminated soil. Conclusions: In conclusion, the work provides the theoretical basis that *Piriformospora indica* combined with *Medicago sativa* contributed to the remediation of PAH-Metal co-contaminated soil.

Background

Coexistence of heavy metals and polycyclic aromatic hydrocarbons (PAHs) in soil has caused worldwide environmental problems (Mansour et al. 2010). Heavy metals and PAHs are risky to human health because of their cytotoxicity, mutagenicity and teratogenicity (Rezek et al. 2008). Co-contamination of heavy metals and PAHs had higher toxicity in the environment, which increased the difficulty in remediating the polluted soil. For example, PAHs mineralization was inhibited by high content of heavy metals in co-contaminated soil (Obuekwe et al. 2013). Phytoremediation has been regarded as a cost-effective and environmentally-friendly technology and widely used for soil remediation (Ehsan et al. 2014). Plant associated microbiology significantly enhance the removal efficiency of PAHs and heavy metals from soil (Adediran et al. 2015; Palaniyandi et al. 2013). The meaning of removal contains two aspects: one aspect is fixation and isolation of PAHs and heavy metals by plant root; the other aspect is degradation of PAHs and heavy metals by microbial in rhizosphere. The presence of microbial could protect plants against damage from plant pathogens and promote plant growth; in turn, the presence of plants changes the microbial community structure and results in more conducive to remove the contamination in soil.

Arbuscular mycorrhizal fungi (AMF) have been introduced into phytoremediation of soil contaminated with diesel (Yang et al. 2009) or heavy metal (Tamayo et al. 2014) to enhance plant resistance and capacity to accumulate heavy metal. However, AMF are obligate endosymbionts which is unable to be

cultivated *in vitro*. *Piriformospora indica* which was first discovered in the rhizospheres of woody shrubs in the sandy desert soils of the Thar region in India, was classified as the *Sebacinales* in basidiomycete (Verma et al. 1998). *Piriformospora indica* is different from AMF that can be cultivated *in vitro*. With AMF-like characteristics, *P. indica* has the added advantage that could grow in axenic cultures. *P. indica* is able to colonize a broad spectrum of plant and conveys various beneficial effects to colonized host plants. *P. indica* acts as a bioprotector against pathogens (Das et al. 2012), alters plant secondary metabolites (Li et al. 2016), increases nutrient uptake and promotes plant growth (Vahabi et al. 2013; Varma et al. 2012), confers drought tolerance to Arabidopsis and barley (Ghabooli et al. 2013; Sherameti et al. 2008), and alleviates salt stress in barley and rice by increasing the activity of detoxifying enzymes and photosynthetic pigment content in colonized plants (Jogawat et al. 2013; Waller et al. 2005). Considering the various beneficial effects caused by *P. indica*, the fungus is regarded to have significant agronomical and high ecological application value (Schäfer and Kogel. 2009). However, there is few reports about the utilization of this endophyte for the remediation of the soil contaminated by heavy metal and PAHs.

No matter the remediation of heavy metal or PAHs, bioavailability determined the remediation efficiency (Chen et al. 2016; Hou et al. 2015). It was proved that the organic acids produced by root could efficiently increase the bioavailability of heavy metals and PAHs (Juwarkar et al. 2007; Braud et al. 2006). *P. indica* directly increase plant root biomass by producing indole-3 acetic (IAA) (Sirrenberg et al. 2007), which definitely contribute to more organic acids produced by root. Parallel, biosurfactants produced by plant growth promoting microbial (PGPM) could increase the bioavailability of heavy metals and PAHs (Juwarkar et al. 2007; Braud et al. 2006). As far as the remediation of heavy metal, the presence of heavy metals stimulated PGPM to produce siderophores, which in turn increased the bioavailability of heavy metals (Gaonkar et al. 2013; Tribedi et al. 2016; Chen et al. 2017). In case of remediating PAHs, plants provide a good platform to recruit more efficient microbes to digest or degrade the PAHs (Mansour et al. 2010). Therefore, PGPM combined with plant roots provide a major way for remediating the heavy metal and PAHs contaminated soil.

Some researchers have reported that using microbial to help promote the remediation of single contamination caused by heavy metal or PAH (Johnson. 2005), however, few study focuses on the influence of inoculation plants with *P. indica* on the chemical speciations of heavy metal and PAH in the metal and PAH co-contaminated soil. Notably, the selected *P. indica* was widely studied on its interaction with plant. However, the special views of this study were taken from other aspects: (1) to address the stress tolerance of *M. sativa* against phenanthrene and cadmium after *P. indica* colonization; (2) to speculate the rhizospheric effect of *P. indica* inoculation by identifying microorganism activity and enzyme activity in rhizospheric and non-rhizospheric soil; (3) to evaluate the effect of *P. indica* inoculation on phytoremediation efficiency of phenanthrene-cadmium co-contaminated soil.

Results

P. indica inoculation increased the biomass of *M. sativa* in contaminated soil

The biomass of *M. sativa* was recorded to reflect the tolerance to pollutants in soil. *P. indica* inoculation significantly promoted plant biomass both on root and stem in contrast to non-inoculated plant (control) (Fig. 1A). Besides, the leaf area was obviously increased compared to the control (Fig. 1B). However, obviously biomass (fresh weight) reduced was observed in Cd, Phe and Phe+Cd treatments without *P. indica* inoculation; whereas the addition of *P. indica* spores significantly enhanced the biomass in those respective treatment (Fig. 1C-D). One hundred plants for each different treatments were statistically recorded in Fig.2. Phe+Cd treatment led to the most serious biomass reduced. Contrastingly, plant biomass in these treatments including Piri+Cd, Piri+Phe and Piri+Phe+Cd was higher than those respective treatments without *P. indica* inoculation. Notably, the growth inhibition on Cd treatment was more serious than that of Phe treatment.

Chlorophyll (Chl) a, b contents and fluorescence parameters

The differences on Chl *a* and Chl *b* contents were significant ($p < 0.01$) between *P. indica* inoculation or not when plant exposure on Cd, Phe contamination (Table 2). Without Cd and Phe contamination, *P. indica* inoculated plant increased Chl *a* and Chl *b* contents in *M. sativa* leaves. Either Cd or Phe contamination reduced Chl *a* and Chl *b* contents significantly. Phe effects on Chl *a* and Chl *b* contents was serious than Cd. However, *P. indica* inoculation significantly increased Chl *a* and Chl *b* contents under Cd, Phe, Cd+Phe treatments in compare to non-inoculated plants. Presence of *P. indica* significantly reduced F_0 so that F_0 values of inoculation plants were 8%, 12.9%, 14% lower than non-inoculated plants under Cd, Phe, Cd+Phe treatments respectively. Additionally, all three fluorescence parameters including F_m (maximal fluorescence level in the dark-adapted state), F_v/F_m (maximum quantum efficiency of PSII photochemistry), and ETR (the relative PSII electron transport rate) were significantly decreased under Cd, Phe, Cd+Phe treatments. However, a noticeably increased on F_m , F_v/F_m , ETR was observed in *P. indica*-inoculated *M. sativa* in comparison to non-inoculated plants.

Auxin production of P. indica affected by phenanthrene and cadmium

The effects of IAA on the growth of plants showed promotion in low concentration and inhibition in high concentration. Therefore, we were interested to reveal whether IAA production in *P. indica* will be affected by phenanthrene and cadmium. Qualitative analysis by Salkowski-method improved that small amount of IAA was produced under Phe, Cd and both of Phe and Cd treatment (Fig. 3A). IAA content produced by *P. indica* was identified by HPLC. The result indicated that IAA content was affected by Phe and Cd in a certain extent (Fig. 3B). Especially, cadmium and phenanthrene co-treatment resulted in less IAA production ($0.69 \mu\text{M}$). However, reduction of IAA content still showed root growth promotion. Root was usually more sensitive to auxin than stem, therefore, appropriate auxin concentration for root would restrict stem growth. This result was consistent with the biomass compensation of root in *P. indica* colonization plant (Fig. 1).

P. indica inoculation increased FDA activities in non-rhizospheric soil

Fluorescein diacetate assay (FDA) was an easy and convincing method to determine soil microorganisms activities. Fig. 4. showed that FDA activity was higher in *P. indica* inoculated plant than non-inoculated plant. The FDA activity under *P. indica* treatment in non-rhizospheric the was the highest, whereas Cd+Phe treatment had the lowest FDA activity. An interesting phenomenon was observed that *P. indica* colonization in plant roots significantly increased microorganisms activity in non- rhizospheric soil than rhizospheric soil. The Cd and Phe contamination significantly reduced FDA activity both in rhizospheric and non- rhizospheric soil. By comparing Cd and Piri+Cd treatments, no obvious differences of FDA activity were observed in rhizospheric soil, adding of *P. indica* increased FDA activity in non-rhizospheric soil. Similar results were obtained between Phe and Piri+Phe treatments.

***P. indica* inoculation increased the polyphenol oxidase in rhizosphere**

Polyphenol oxidase in rhizospheric and non-rhizospheric soil was identified in Fig. 5. The activity of polyphenol oxidase in rhizospheric soil was higher than that in non-rhizospheric soil. *P. indica* colonization significantly increase polyphenol oxidase activity either in rhizospheric or non-rhizospheric. The order of polyphenol oxidase activity form low to high was: Phe+Cd < Phe < Cd < Piri+Phe < Piri+Phe+Cd < Piri+Cd < Con < Piri. According to these data, several aspects were revealed: 1) polyphenol oxidase activity in soil was easily affected by Phe than Cd contamination; 2) the addition of *P. indica* significantly increased polyphenol oxidase activity in rhizospheric soil; 3) for Phe and Cd co-contamination soil, the polyphenol oxidase activity was affected seriously, whereas after *P. indica* colonization in root of *M. sativa*, the polyphenol oxidase was obviously enhanced.

***P. indica* inoculation reduced phenanthrene content in rhizosphere**

Phenanthrene degradation rate was identified around the rhizospheric soil and non-rhizospheric soil planted with *M. sativa* (Fig. 6A). Generally, phenanthrene content in rhizospheric soil was less than the non-rhizospheric soil. For rhizospheric area, among four different treatments, phenanthrene concentration in soil from high to low was Phe+Cd, Phe, Piri+Phe+Cd, Piri+Phe. The concentration gradient in non-rhizospheric area was similar to those in rhizospheric area. Those data implies that *P. indica* colonization could effectively reduce phenanthrene content in soil.

In order to prove this evidence, phenanthrene concentration in plant roots and stem was measured additionally. Phenanthrene concentrations in roots were 135 ± 6 mg kg⁻¹, 165 ± 4 mg kg⁻¹, 108 ± 10 mg kg⁻¹, 156 ± 7 mg kg⁻¹ among the Phe, Piri+Phe, Phe+Cd and Piri+Phe+Cd treatment respectively (Fig. 6B). Obviously, phenanthrene concentrations in roots were higher than those in stem. In stem, the Phe+Cd treatment had the lowest phenanthrene content than the other treatment. There is no obvious difference of phenanthrene content in the stem among Phe, Piri+Phe and Piri+Phe+Cd treatments.

***P. indica* inoculation reduced cadmium content in rhizospheric soil**

The cadmium content in rhizospheric soil and non-rhizospheric soil of *M. sativa* was identified as shown in Fig.7A. Similarly, cadmium content in rhizospheric soil was less than non-rhizospheric soil. For

rhizospheric area, among four different treatments, cadmium concentration in soil from high to low was Phe+Cd, Piri+Phe+Cd, Cd, Piri+Cd. Adding of *P. indica* spores significantly reduced cadmium content in compare to the treatment without *P. indica* inoculation. For instance: either in rhizospheric soil or non-rhizospheric soil, Piri+Cd treatment had lower cadmium content than Cd treatment; and Piri+Phe+Cd treatment had lower cadmium content than Phe+Cd treatment. Remarkably, Cd content in rhizospheric soil of *P. indica* treatment was obviously reduced (less than 2 mg kg⁻¹) than the original Cd content in soil (10 mg kg⁻¹). The cadmium concentration gradient in non-rhizospheric area was similar to those in rhizospheric area. Those data implies that adding of *P. indica* colonization could effectively reduce the cadmium content in soil.

Additionally, cadmium concentration in plant roots and stem was explored in parallel (Fig.7B). Cadmium concentrations in roots were 26±4mg kg⁻¹, 34±6 mg kg⁻¹, 23±8 mg kg⁻¹, 29±5 mg kg⁻¹ in the Cd, Piri+Cd, Phe+Cd and Piri+Phe+Cd treatment respectively. Likewise, the cadmium concentrations in roots were notably higher than those in stem. In stem, Cd content was different from phenanthrene content: Piri+Cd treatment had lower Cd content than Cd treatment; and Cd content in Piri+Phe+Cd treatment was lower than Phe+Cd treatment. It suggested that adding of *P. indica* inhibit the transport of Cd into stem.

Discussion

Heavy metal and organic pollutants in soil do serious harm to plant growth and development. Cd treatment leads to the most serious affection to plant biomass accumulation. Shahabivand et al. (2017) reported that *P. indica* successfully colonized plants grown in 120 mg/kg of Cd contaminated soil, which suggest that this fungus is able to colonize roots under high Cd stress in soil. However, we found that the spores germination and hyphal growth of *P. indica* were seriously inhibited under 40 mg kg⁻¹ of Cd (data not shown). Therefore, 10 mg kg⁻¹ of Cd was selected as the test concentration.

Our data indicated that additional colonization of *P. indica* in Cd (10 mg kg⁻¹), Phe (40 mg kg⁻¹) and Phe+Cd treatments remits stress from heavy metal and PAHs. Though Cd concentration in roots of Piri+Cd treatment was higher than that in Cd treatment, the biomass was not significantly affected. The results implied that high concentration of Cd could be fixed or isolated in some degree by fungus and plant co-living system and therefore reduced the harmful affection to plant. Cd accumulation in stem was reduced after phenanthrene contamination in soil which indicated that the organic pollutants accumulated in roots may hinder Cd transport from root to above-ground parts. Piri+Phe treatment accumulated high concentration of phenanthrene in root which decreased the biomass of roots and stem compared to Piri treatment. However, even high concentration of phenanthrene in root did not hinder the growth promotion effect of *P. indica* to host plant compared to Phe treatment. This result was different from the previous report that the plant growth promotion bacterial (PGPB) accumulation of pyrene decreased the biomass of stem (Chen et al. 2017).

M. sativa plants subjected to Cd, Phe stress showed reduced growth in terms of shoot and root lengths, and shoot and root fresh weight. Cd toxic effects have widely been reported in different plant species and

Cd is known to reduce or inhibit plant growth because of Cd harmful impact on the processes of photosynthesis, respiration and essential elements uptake (Guo et al., 2016; Shahabivand., 2017). Chaoui and Ferjani (2005) reported that activity of indole-3-acetic acid (IAA) oxidase (as a growth-limiting enzyme) was increased under Cd toxicity, which result in the reduced plant growth due to diminution of endogenous content of plant growth-promoting hormone auxin. In our work, the elevation in photosynthesis efficiency of host via enhanced Chl *a* and *b* contents and elevated Fv/Fm and ETR values (Table 2) contributed to the *P. indica*-induced growth promotion and stress tolerance.

Soil microorganism activity could effectively reflect soil fertility. Soil microorganism activity was determined to speculate the phytoremediation efficiency. *P. indica* has been reported to inhibit the phytopathogens such as *Fusarium graminearum* infection to barley plant (Deshmukh et al. 2007; Zhang et al. 2012). Thereby it is reasonable to explain that because of the presence of *P. indica*, the microorganism activity in rhizosphere was lower than that in non-rhizosphere. Anyhow, the FDA activity in *P. indica* treatment was higher than non-*P. indica* treatment either in rhizospheric or non-rhizospheric. Phe treatment had lower FDA activity than Cd treatment which was in accordance with the previous report that petroleum hydrocarbons inhibit the microorganism activity in soil (Zhang et al. 2014). Though similar FDA activities in Cd and Piri+Cd or Phe and Piri+Phe were obtained in rhizosphere, the mechanism might be different. The reduction of FDA activities in Cd or Phe treatments was probably due to heavy metal or PAHs stress, whereas the reduction of FDA activities in Piri+Cd or Piri+Phe treatments probably depends on the phyto-pathogens growth inhibition. Enzymes, as part of soil composition, the degree of its active sensitively reflects the direction and strength of a biochemical reaction in soil (Liu et al. 2009). The changes of enzyme activity effectively reflect the degradation ability of organic pollutants by microbial and plant roots in soil. In fact, in our experiment, we identified several enzymes including urease, invertase and so on, and the results indicated those enzymes activity were better in the rhizosphere of *P. indica* colonization root (data not shown). Polyphenol oxidase is an important oxidoreductase enzyme in soil which participates in the process of decomposition and transformation of aromatic compounds. Enhanced polyphenol oxidase in this study implied that *P. indica* had a strong repairing ability for the contaminated soil.

Co-contamination of heavy metals and PAHs result in deleterious influence on microorganism activity in soil compared to single heavy metals or PAHs contamination (Maliszewska et al. 2003), which explained that Phe+Cd treatment had the lowest FDA activity in all treatment. The Piri+Cd and Piri+Phe+Cd treatments had higher FDA activities than Cd and Phe+Cd treatments respectively, which implied that the endophyte and plant co-living system contribute to a well-established micro ecosystem for microorganism in soil.

As for Phe, Piri+Phe, Cd+Phe and Piri+Phe+Cd four different treatments, Phe accumulation in plants were 135 ± 6 mg kg⁻¹, 165 ± 4 mg kg⁻¹, 108 ± 10 mg kg⁻¹, 156 ± 7 mg kg⁻¹ respectively. The high concentration of Phe accumulation in plant means the less Phe residual in soil. Besides plant, microorganism in soil played key roles in the removal of the PAHs. For Phe+Cd treatment, Phe concentration left in rhizospheric soil or non-rhizospheric soil was higher than Phe treatment, which was reasonable due to higher

microorganism activity in Phe treatment. Though no significant difference of FDA activities was obtained between Phe and Phe +Piri treatment, *P. indica* colonization obviously reduced Phe concentration in rhizospheric and non-rhizospheric part. Similar data was collected from Phe+Cd and Phe +Cd +Piri treatments. Totally, Phe concentration in rhizospheric soil was less compared to non-rhizospheric part, which explained that the rhizospheric effect was responsible for the less accumulation of PAHs in the rhizospheric soil than non-rhizospheric soil (Wang et al. 2016).

The order of Cd concentrations in root from high to low was as follows: Piri+Cd, Cd, Piri+Phe+Cd, Phe+Cd. The data indicated that *P. indica* colonization increased Cd accumulation in roots. However, by comparing Piri+Cd and Piri+Phe+Cd treatments, it was obvious that the addition of Phe reduced *P. indica* accumulation Cd in a certain degree. Bioavailability of heavy metal determines the phytoremediation efficiency (Megharaj et al. 2011). Microorganism in soil like fungi and bacterial is capable to produce biosurfactants, for instance, rhamnolipid which could increase the mobility of heavy metals, to enhance its bioavailability (Mulligan 2005). Therefore, it was reasonable to understand that *P. indica* colonization significantly reduced Cd concentration in rhizospheric than non-rhizospheric soil. In the other hand, plant roots exudates also contribute to the acidification of heavy metals, which increase the efficiency of phytoremediation. *P. indica* colonization could significantly promote root growth and development, the roots in turn secret exudates to increase bioavailability of heavy metals and enhance phytoremediation efficiency in rhizospheric soil, which a virtuous cycle was formed.

Conclusions

In conclusion, *P. indica* colonization posed positive effects on the processes of photosynthesis and increased the tolerance of *M. sativa* in phenanthrene and cadmium co-contaminated soil, especially in single cadmium contaminated soil. Phenanthrene and cadmium co-contaminated soil affected soil microorganism seriously than single phenanthrene or cadmium contamination. *P. indica* inoculation in *M. sativa* root increased the FDA activity in phenanthrene, cadmium and both of that co-contaminated soil, especially in non-rhizospheric soil. The addition of *P. indica* stimulated root of *M. sativa* to accumulate higher concentration of phenanthrene and cadmium, and thereby enhanced phytoremediation efficiency. One vividly conclusion picture (Fig. 8) was drawn about phytoremediation effect of *Medicago sativa* colonized by *Piriformospora indica* in phenanthrene and cadmium co-contaminated soil. This study suggested the application of *Piriformospora indica* combined with *M. sativa* contributed to the remediation of PAH-metal co-contaminated soil for sustainable agriculture.

Methods

Chemicals

Phenanthrene with a purity of 98% was purchased from Sigma (USA). The rest chemicals were purchased from Dingguo (Tian jin, China).

Soil used in the study

The experiment soil, never contaminated by PAHs or heavy metals was collected from the topsoil (0-20 cm) of Hebei University of Technology, China. The soil was a kind of phaeozems (alfisol) in Hebei Province (Soil Survey Staff, 2010). The test soil was sieved by 2 mm sieve after dried by air. Soil samples were measured using standard methods (Lu, 1999) before phytoremediation. The composition of the sample soil was physico-chemically characterized: $55.6 \pm 2.1\%$ silt, $30.8 \pm 1.8\%$ sand and $13.6 \pm 1.5\%$ clay, 0.06% total N, 8 mg/kg available P, 40 mg/kg available K, 1.1% organic matter. The soil pH was 7.36 ± 0.06 .

Preparation of phenanthrene contaminated soil

Phenanthrene was dissolved in acetone and added into a small part soil, 1 day later, after acetone volatilized, the small part soil was added into the whole sample soil and incorporated thoroughly, the final concentration of phenanthrene in soil was measured as $40 \pm 3 \text{ mg kg}^{-1}$.

Preparation of cadmium contaminated soil

The aqueous solution of cadmium nitrate was added to the prepared soil, the final concentration of cadmium in soil was measured as $10 \pm 2 \text{ mg kg}^{-1}$.

Preparation of phenanthrene and cadmium co-contaminated soil

The acetone stock solution containing phenanthrene was firstly added to the test soil. After acetone evaporated, cadmium nitrate aqueous solution was added to the previous soil polluted by phenanthrene. The final concentration of cadmium and phenanthrene in soil were measured as $10 \pm 2 \text{ mg kg}^{-1}$ and $40 \pm 3 \text{ mg kg}^{-1}$ respectively. The well prepared soil and a control soil without any pollution were shifted to boxes and aged in the dark for 15 days.

Fungal culture

P. indica was supplied by Karl-Heinz-Kogel (Institute of Plant Pathology and Applied Zoology, Giessen, Germany). *P. indica* was maintained at 23°C on CM medium (Pham et al. 2004). For solid medium, 14 g L⁻¹ agar was added; For liquid cultures, 100 mL medium was inoculated in a 300-mL Erlenmeyer flask. To test whether Cd and Phe will affect IAA production contents, 5 mg kg⁻¹ and 20 mg kg⁻¹ of cadmium and phenanthrene were added into the liquid cultures respectively. CM medium was inoculated with 20 mycelium plugs from the margin of a growing colony of *P. indica* on CM solid medium, Liquid cultures were incubated at 23°C at 150 rpm on a rotary shaker.

Quantification of IAA in fungal growth media by HPLC

IAA production ability of *P. indica* under Phe, Cd and Phe+ Cd treatment was performed using Salkowski method according to the literature (Glickmann and Dessaux. 1995). Quantification of IAA in fungal

growth media was performed by HPLC. Fungal culture filtrates were harvested, acidified and extracted twice with ethyl acetate as described in Sirrenberg et al. 2007. The parameters used for HPLC are as follows: 50% methanol: 45% water: 5% acetonitrile (v/v) was used as the mobile phase. The 0.2 mL min⁻¹ of flow rate was applied. The injection volume was 10 µl. Column temperature was kept at 40°C. The content of IAA was quantified by Agilent HPLC which equipped with a HC^R C18 (5 µm, 4.6 x 250 mm, Agilent, USA) reversed phase chromatographic column. IAA concentrations were always determined in parallel in medium in which no fungus had been cultured but which had been incubated under the same conditions.

***M. sativa* treatment and *P. indica* inoculation**

P. indica spores were collected according to the literature (Li et al. 2016). The seeds of *M. sativa* were surface sterilized in 70% alcohol for 1 min and then 3% NaClO for 15 min. After sterilized, seeds were washed by sterile deionized water repeatedly and planted into the MS medium for germination. Seven days when the roots were presence, *P. indica* spores were added into the root surface. Then the seedlings contained *P. indica* spores were transferred into different types of soil in pot (5 kg). Three replicates were applied for each treatment. The experiment design was listed in Table 1.

The plants were rejuvenated for one week in shade. After then, pots were transferred to green house for three month. The pots were irrigated 3-4 times a week to keep the soil moisty during this period. Three month later, plants were harvested and soil from rhizosphere and non- rhizosphere were collected respectively from the roots of *M. sativa* .

Measurement of chlorophyll contents and chlorophyll fluorescence

Chlorophyll content in the youngest fully expanded leaves (0.1 g) was extracted by 80% acetone, centrifuged at 4000 rpm for 20 min, and then the optical density of the supernatant was read at 663 and 645 nm wavelengths for Chl *a* and Chl *b*, respectively (Arnon, 1967). The following parameters of chlorophyll fluorescence were measured by analyzing the first fully grown leaves of *M. sativa* using a portable fluorometer (Hansatech, Instruments LTD, UK): F0 (minimal fluorescence level in the dark-adapted state), Fm (maximal fluorescence

level in the dark-adapted state), Fv/Fm (maximum quantum efficiency of PSII photochemistry) and ETR (the relative PSII electron transport rate). *M. sativa* plants were dark-adapted for 30 min to measure the influence of factors on photosystem II (PSII) efficiency.

Measure of soil microorganism activity

Soil microorganism activity was measured referenced to the literature Tribedi et al. (2013). Firstly, the soil was freeze-dried by Freeze drying machine (Alpha 1-2 L D plus, Germany). The 5 g freeze-dried soil was dissolved in 15 mL phosphate buffer solution (NaCl–8.5 g, Na₂HPO₄–2.2 g, NaH₂PO₄–0.1 g, pH 7.6) at room temperature. The turbid liquid was shaken at 180 rpm for 30 min and then 0.5 mL of fluorescein

diacetate (FDA) (2 g L^{-1} , in acetone) solution was added into the mixture. The absorbance value was recorded at $\text{OD}_{490\text{nm}}$.

Analysis of phenanthrene in rhizospheric and non-rhizospheric soil

The ultrasonic extraction and high performance liquid chromatograph (HPLC) ultraviolet detector detection method was used. Ten gram freeze-dried soil samples was dissolved in 50 mL of acetone-hexane (1:1, v/v) mixed extraction solvent. The progress of extraction was performed for 1 h in the ultrasonic cleaners. Extracted liquid was filled with filter funnel containing 10 g anhydrous Na_2SO_4 . The extracted liquid was concentrated to 5 mL by rotary evaporation instrument in 60°C water bath. Then 5 mL concentrated extracted liquid was transferred to silica gel-Alumina column chromatography and washed with methylene chloride-hexane (1:1, v/v) mixture elution. The condensed elution nearly drying was diluted to a final volume of 1 mL and used for HPLC determination. The parameters used for HPLC are as follows: methanol and water (87:13, v/v) was used as the mobile phase. The 1 mL min^{-1} of flow rate and 254 nm of detection wave length was applied. The content of phenanthrene was quantified by Agilent HPLC which equipped with a $\text{HC}^{\text{R}} \text{C18}$ ($5 \mu\text{m}$, $4.6 \times 250 \text{ mm}$, Agilent, USA) reversed phase chromatographic column.

Analysis of cadmium in rhizospheric and non-rhizospheric soil

Cadmium was determined according to the references (Tessier et al. 1979; Wang et al. 2016). Soil about the rhizosphere and non-rhizosphere of *M. sativa* was extracted by mixing 0.5 g soil with 10 mL HCl solution and then heating for 3 h (45°C). After cool down, Cadmium was extracted by mixing the soil with HNO_3 and HClO_4 (v/v, 4:1) for digestion (220°C , 1 h) and then adding HF and HClO_4 (v/v, 5:1) for further digestion (220°C , 2 h). The same method was used for cadmium extraction from plants. At the end of extraction step, the supernatant was harvested by centrifuging at 6000 rpm, 20 min. Then the supernatant was filtered through $0.45 \mu\text{m}$ microfiltration membrane and quantified by Inductively Coupled Plasma Optical Emission Spectrometry (ICPOES).

Analysis of enzyme activity in rhizospheric and non-rhizospheric soil

Enzyme activity of polyphenol oxidase was determined according to the protocol book of soil enzymes and analytical methods (Guan 1986). Ten gram freeze-dried soil was dissolved in 10 mL pyrogallol (1%) and shaken with 150 r. min^{-1} for 1 min. Then the sample was kept in dark at 30°C for 2 h. Four milliliter citric acid - phosphate buffer (disodium hydrogen phosphate- 35.61 g/L , citric acid- 21.01 g/L , pH4.5) was added into the sample. Finally, 35 mL ether was added and shaken for 2 min. The absorbance value at 430 nm was reordered after 30 min extraction.

Statistical analysis

In this study, all data are expressed as means \pm SE and represent at least three independent biological experiments. The significant differences were analyzed by using a one-way analysis of variance

(ANOVA). Duncan's multiple range test was used to compare the means at 0.05 and 0.01 probability level.

Abbreviations

PAHs: polycyclic aromatic hydrocarbons; *P. indica*: *Piriformospora indica*; AMF: Arbuscular mycorrhizal fungi; IAA: indole-3 acetic; PGPM: plant growth promoting microbial; Chl a/ Chl b: Chlorophyll (Chl) a, b; F0: minimal fluorescence level in the dark-adapted state; Fm: maximal fluorescence level in the dark-adapted state; Fv/Fm: maximum quantum efficiency of PSII photochemistry; ETR: the relative PSII electron transport rate; FDA: Fluorescein diacetate assay.

Declarations

Authors's contributions

Liang Li designed and guided the whole experiment, also performed the work of analyzing the data and writing the Ms. Pengyue Zhu was responsible for the *Medicago sativa* and *Piriformospora indica* cultivation. Xiaoyang Wang was responsible for the phenanthrene (Phe) and cadmium (Cd) co-contaminated soil preparation and Fluorescein diacetate (FDA) activities analysis. Zhenhua Zhang performed the Cd and Phe concentration analysis in soil and other relative work. All authors read and approved the final manuscript.

Funding

We thank Scientific Research Fund of Hebei Provincial Education Department (QN2018014) and National Natural Science Foundation of China (Grant No.31801948) for financial support.

Availability of data and materials

All data generated or analysed during this study are included in this article.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Acknowledgement

We would like to express deep thanks to Professor Karl-Heinz Kogel for providing the fungus of *Piriformospora indica* and guidance for this research work.

References

- Adediran, G. A., Ngwenya, B. T., Mosselmans, J. F., Heal, K. V., Harvie, B. A. 2015. Mechanisms behind bacteria induced plant growth promotion and Zn accumulation in brassica juncea. *J. Hazard. Mater.* 283, 490-499.
- Arnon, A., 1967. Method of extraction of chlorophyll in the plants. *Agron. J.* 23, 112–121.
- Braud, A., Jézéquel, K., Vieille, E., Tritter, A., Lebeau, T. 2006. Changes in extractability of Cr and Pb in a polycontaminated soil after bioaugmentation with microbial producers of biosurfactants, organic acids and siderophores. *Water Air Soil Pollut.* 6(3-4), 261-279.
- Chaoui, A., Ferjani, E. E., 2005. Effects of cadmium and copper on antioxidant capacities, lignification and auxin degradation in leaves of pea (*Pisum sativum* L.) seedlings. *Plant Biol. Pathol.* 328, 23-31.
- Chen, T., Liu, X., Zhang, X., Chen, X., Tao, K., Hu, X. 2016. Effect of alkyl polyglucoside and nitrilotriacetic acid combined application on lead/pyrene bioavailability and dehydrogenase activity in co-contaminated soils. *Chemosphere* 154, 515-520.
- Chen, X., Liu, X., Zhang, X., Cao, L., Hu, X. 2017. Phytoremediation effect of *Scirpus triqueter* inoculated plant-growth-promoting bacteria (PGPB) on different fractions of pyrene and Ni in co-contaminated soils. *J. Hazard. Mater.* 325, 319-326.
- Das, A., Kamal, S., Shakil, N. A., Sherameti, I., Oelmüller, R., Dua, M. 2012. The root endophyte fungus *Piriformospora indica* leads to early flowering, higher biomass and altered secondary metabolites of the medicinal plant, *coleus forskohlii*. *Plant Signal Behav.* 7(1), 103-112.
- Deshmukh, S. D., Kogel, K. H. 2007. *Piriformospora indica* protects barley from root rot caused by *Fusarium*. *J. Plant Dis. Prot.* 114 (6), 263-268.
- Ehsan, S., Ali, S., Noureen, S., Mahmood, K., Farid, M., Ishaque, W. 2014. Citric acid assisted phytoremediation of cadmium by *Brassica Napus* L. *Ecotoxicol. Environ. Saf.* 106, 164-172.
- Gaonkar, T., Bhosle, S. 2013. Effect of metals on a siderophore producing bacterial isolate and its implications on microbial assisted bioremediation of metal contaminated soils. *Chemosphere* 93 (9), 1835-1843.
- Ghabooli, M., Khatabi, B., Ahmadi, F. S., Sepehri, M., Mirzaei, M., Amirkhani, A. 2013. Proteomics study reveals the molecular mechanisms underlying water stress tolerance induced by *Piriformospora indica* in barley. *J. Proteomics.* 94 (20), 289-301.

- Glickmann, E., Dessaux, Y. 1995. A critical examination of the specificity of the salkowski reagent for indole compounds produced by phytopathogenic bacteria. *Applied. Environ. Microbiol.* 61, 793-796.
- Guan, S. Y. 1986. *Soil enzymes and analytical methods*. Beijing: Agric. press.
- Guo, H.P., Hong, C.T., Chen, X.M., Xu, Y.X., Liu, Y., Jiang, D.A., Zheng, B.S., 2016. Different growth and physiological responses to cadmium of the three *Miscanthus* species. *PLoS One* 11, e0153475.
- Hou, Y., Liu, X., Zhang, X., Chen, X., Tao, K., Chen, X. 2015. Identification of *Scirpus triqueter* root exudates and the effects of organic acids on desorption and bioavailability of pyrene and lead in co-contaminated wetland soils. *Environ. Sci. Pollut. Res. Int.* 22 (22), 17780-17788.
- Jogawat, A., Saha, S., Bakshi, M., Dayaman, V., Kumar, M., Dua, M. 2013. *Piriformospora indica* rescues growth diminution of rice seedlings during high salt stress. *Plant Signal Behav.* 8, 26891.
- Johnson, D. L., Mcgrath, S. P. 2005. Soil microbial response during the phytoremediation of PAH contaminated soil. *Soil Biol. Biochem.* 37 (12), 2334-2336.
- Juwarkar, A. A., Nair, A., Dubey, K. V., Singh, S. K., Devotta, S. 2007. Biosurfactant technology for remediation of cadmium and lead contaminated soils. *Chemosphere* 68 (10), 1996-2002.
- Karene, G., Huang, X. D., Bernardr, G., Brucem, G. 2009. Phytoremediation and rhizoremediation of organic soil contaminants: potential and challenges. *Plant Sci.* 176 (1), 20-30.
- Li, L., Chen, X., Ma, C., Wu, H., Qi, S. 2016. *Piriformospora indica* requires kaurene synthase activity for successful plant colonization. *Plant Physiol. Biochem.* 102, 151-160.
- Liu, S.L., Luo, Y. M., Wu, L. H. 2009. Degradation of phenanthrene in soil planted with ryegrass and the effect of phenanthrene on soil enzymes. *Acta. Pedologica Sinica.* 46(3), 419-425.
- Lu, R. K. 1999. *Analytical methods of soil agriculture chemistry*. China Agricultural Science and Technology Press, Beijing (in Chinese).
- Maliszewska-Kordybach, B., Smreczak, B. 2003. Habitat function of agricultural soils as affected by heavy metals and polycyclic aromatic hydrocarbons contamination. *Environ. Int.* 28(8), 719-728.
- Mansour, S. A., Gad, M. F. 2010. Risk assessment of pesticides and heavy metals contaminants in vegetables: a novel bioassay method using daphnia magna Straus. *Food Chem. Toxicol.* 48 (1), 377-389.
- Megharaj, M., Ramakrishnan, B., Venkateswarlu, K., Sethunathan, N., Naidu, R. 2011. Bioremediation approaches for organic pollutants: a critical perspective. *Environ. Int.* 37 (8), 1362-1375.
- Mulligan, C. N. 2005. Environmental applications for biosurfactants. *Environ. Pollut.* 133 (2), 183-198.

- Obuekwe, I. S., Semple, K. T. 2013. Impact of zinc-copper mixtures on the development of phenanthrene catabolism in soil. *Int. Biodeterior. Biodegrad.* 85(11), 228-236.
- Palaniyandi, S. A., Yang, S. H., Zhang, L., Suh, J. W. 2013. Effects of actinobacteria on plant disease suppression and growth promotion. *Appl. Microbiol. Biotechnol.* 97 (22), 9621-9636.
- Rezek J., Wiesche C. I. D., Mackova M., Zadrazil, F., Macek, T. 2008. The effect of ryegrass (*Lolium perenne*) on decrease of PAH content in long term contaminated soil. *Chemosphere.* 70(9), 1063-1608.
- Sahay, N. S., Varma, A. 1999. *Piriformospora indica*: a new biological hardening tool for micropropagated plants. *FEMS Microbiol. Lett.* 181, 297–302.
- Soil Survey Staff, 2010. Keys to Soil Taxonomy, 11th ed. USDA-Natural Resources Conservation Service, Washington, DC.
- Shahabivand, S., Parvaneh, A., Aliloo, A. A. 2017. Root endophytic fungus *Piriformospora indica* affected growth, cadmium partitioning and chlorophyll fluorescence of sunflower under cadmium toxicity. *Ecotox. Environ. Safe.* 145, 496-502.
- Schäfer, P., Kogel, K. H. 2009. The Sebacinoid Fungus *Piriformospora indica*: An orchid mycorrhiza which may increase host plant reproduction and fitness. *Plant Relationship.* 5, 99-112.
- Shahabivand, S., Parvaneh, A., Aliloo, A. A. 2017. Root endophytic fungus *Piriformospora indica* affected growth, cadmium partitioning and chlorophyll fluorescence of sunflower under cadmium toxicity. *Ecotoxicol. Environ. Saf.* 145, 496-502.
- Sherameti, I., Tripathi, S., Varma, A., Oelmüller, R. 2008. The root-colonizing endophyte *Piriformospora indica* confers drought tolerance in *Arabidopsis* by stimulating the expression of drought stress-related genes in leaves. *Mol. Plant-Microbe. Interact.* 21 (6), 799-807.
- Tamayo, E., Gómez-Gallego, T., Azcón-Aguilar, C., Ferrol, N. 2014. Genome-wide analysis of copper, iron and zinc transporters in the arbuscular mycorrhizal fungus *Rhizophagus irregularis*. *Front Plant Sci.* 5, 547
- Tessier, A., Campbell, P. G. C., Bisson, M. 1979. Sequential extraction procedure for the speciation of particulate trace metals. *Anal. Chem.* 51(7), 844-851.
- Tribedi, P., Sil, A. K. 2013. Bioaugmentation of polyethylene succinate-contaminated soil with *Pseudomonas*, sp. AKS2 results in increased microbial activity and better polymer degradation. *Environ. Sci. Pollut. Res.* 20, 1318–1326.
- Vahabi, K., Camehl, I., Sherameti, I. 2013. Growth of *Arabidopsis* seedlings on high fungal doses of *Piriformospora indica* has little effect on plant performance, stress, and defense gene expression in spite

of elevated jasmonic acid and jasmonic acid-isoleucine levels in the roots. *Plant Signal Behav.* 8, 26301-26311.

Verma, S., Varma, A., Rexer, K. H., Hassel, A., Kost, G., Sarbhoy, A. 1998. *Piriformospora indica*, gen. et sp. nov., a new root-colonizing fungus. *Mycologia*, 90(5), 896-903.

Waller, F., Achatz, B., Baltruschat, H., Fodor, J., Becker, K., Fischer, M. 2005. The endophytic fungus *Piriformospora indica* reprograms barley to salt-stress tolerance, disease resistance, and higher yield. *Proc. Natl. Acad. Sci.* 102(38), 13386-13391.

Wang, Q., Liu, X., Zhang, X., Hou, Y., Hu, X., Liang, X. 2016. Influence of tea saponin on enhancing accessibility of pyrene and cadmium phytoremediated with *Lolium multiflorum* in co-contaminated soils. *Environ. Sci. Pollut. Res.* 23(6), 5705-5711.

Wijayawardena, M. A. A., Naidu, R., Megharaj, M., Lamb, D., Thavamani, P., Kuchel, T. 2015. Using soil properties to predict in vivo, bioavailability of lead in soils. *Chemosphere* 138, 422-428.

Yang, T., Lin, X. G., Hu, J. L., Zhang, J., Lu, J. L., Wang, J. H. 2009. Effects of arbuscular mycorrhizal fungi on phytoremediation of PAHs-contaminated soil by *Medicago sativa* and *Lolium multiflorum*. *Ecol. Rural. Environ.* 25(4), 72-76.

Zhang, X. Y., Liu, X. Y., Liu, S. S., Liu, F. H., Chen, L. S., Xu, G. 2012. Response characteristics of *Scirpus triquetra* and its rhizosphere to pyrene contaminated soils at different growth stages. *Int. J. Phytorem.* 14(7), 691-702.

Zhang, X., Chen, L., Liu, X., Wang, C., Chen, X., Xu, G. 2014. Synergic degradation of diesel by *Scirpus triquetra* and its endophytic bacteria. *Environ. Sci. Pollut. Res.* 21(13), 8198-8205.

Figures




Fig.1

Figure 1

Biological effect on root and stem in Phe, Cd, and Phe-Cd co-contaminated soil with or without *P. indica* inoculation. A: Control with or without *P. indica* inoculation. B: Leaves area were compared between *P. indica* inoculation and non- *P. indica* inoculation without any contamination. C: Above-ground parts were compared between *P. indica* inoculation and non- *P. indica* inoculation in Cd contaminated soil. D: Above-ground parts were compared between *P. indica* inoculation and non- *P. indica* inoculation in Phe, and Phe-Cd contaminated soil.




Fig.2

Figure 2

The fresh weight of stems and roots of *M. sativa* was statistically analyzed in control, Phe/Cd and Phe-Cd contaminated soil with or without *P. indica* colonization. Control with (Piri) or without (Con) *P. indica* inoculation; Phe contaminated soil with (Piri+Phe) or without (phe) *P. indica* inoculation; Cd contaminated soil with (Piri+Cd) or without (Cd) *P. indica* inoculation; Phe-Cd co-contaminated soil with (Piri+Cd+Phe) or without (Cd+Phe) *P. indica* inoculation. *M. sativa* was planted in all treatments. Each value of fresh weight is the mean of three replicates. Error bars show standard error. The asterisk in the columns indicates significant difference in fresh weight between treatments according to Duncan's Multiple Range Test ($P < 0.01$).




Fig.3

Figure 3

IAA production by *P.indica* during growth in CM medium containing Cd or Phe contamination. IAA was determined in culture supernatants by HPLC-MS after 6 weeks containing Cd, Phe or Cd and Phe contamination medium respectively. IAA concentrations in CM medium were $1.34 \pm 0.03 \mu\text{M}$ (n=3), but in Cd containing medium were $0.71 \pm 0.05 \mu\text{M}$ (n=3); in Phe containing medium were $1.12 \pm 0.07 \mu\text{M}$ (n=3), in Cd and Phe containing medium were $0.69 \pm 0.04 \mu\text{M}$ (n=3), respectively. Error bars show standard error. The asterisk in the columns indicates significant difference in fresh weight between treatments according to Duncan's Multiple Range Test ($P < 0.01$).




Fig.4

Figure 4

FDA activities were identified in control, Phe/Cd and Phe-Cd contaminated soil with or without *P. indica* colonization. Control with (Piri) or without (Con) *P. indica* inoculation; Phe contaminated soil with (Piri+phe) or without (phe) *P. indica* inoculation; Cd contaminated soil with (Piri+Cd) or without (Cd) *P. indica* inoculation; Phe-Cd co-contaminated soil with (Piri+Cd+Phe) or without (Cd+Phe) *P. indica* inoculation. *M. sativa* was planted in all the treatments. Each value of FDA activities is the mean of three replicates. Error bars show standard error. The asterisk in the columns indicates significant difference in FDA activities between treatments according to Duncan's Multiple Range Test ($P < 0.01$).




Fig.5

Figure 5

Phenanthrene content was identified in soil and plant. A: Phenanthrene content was identified different treatments in rhizospheric soil and non-rhizospheric soil. B: Phenanthrene content was identified in different treatments in root and stem. Phe contaminated soil with (Piri+phe) or without (phe) *P. indica* inoculation; Phe-Cd co-contaminated soil with (Piri+Cd+Phe) or without (Cd+Phe) *P. indica* inoculation. *M. sativa* was planted in all treatments. Each value of phenanthrene content is the mean of three replicates. Error bars show standard error. The asterisk in the columns indicates significant difference in phenanthrene content between treatments according to Duncan's Multiple Range Test ($P < 0.01$).




Fig.6

Figure 6

Cadmium content was identified in soil and plant. A: Cadmium content was identified in different treatments in rhizospheric soil and non-rhizospheric soil. B: Cadmium content was identified in different treatments in root and stem. Cd contaminated soil with (Piri+Cd) or without (Cd) *P. indica* inoculation; Phe-Cd co-contaminated soil with (Piri+Cd+Phe) or without (Cd+Phe) *P. indica* inoculation. *M. sativa* was planted in all treatments. Each value of cadmium content is the mean of three replicates. Error bars show standard error. The asterisk in the columns indicates significant difference in cadmium content between treatments according to Duncan's Multiple Range Test ($P < 0.01$).




Fig.7

Figure 7

Polyphenol oxidase was identified in different treatments in rhizospheric soil and non-rhizospheric soil. Control with (Piri) or without (Control) *P. indica* inoculation; Phe contaminated soil with (Piri+Phe) or without (Phe) *P. indica* inoculation; Cd contaminated soil with (Piri+Cd) or without (Cd) *P. indica* inoculation; Phe-Cd co-contaminated soil with (Piri+Cd+Phe) or without (Cd+Phe) *P. indica* inoculation. *M. sativa* was planted in all treatments. Each value of fresh weight is the mean of three replicates. Error bars show standard error. The asterisk in the columns indicates significant difference in polyphenol oxidase between treatments according to Duncan's Multiple Range Test ($P < 0.01$).




Fig.8

Figure 8

The phytoremediation effect of *Medicago sativa* colonized by *Piriformospora indica* in phenanthrene and cadmium co-contaminated soil. Four benefits aspects: including increased biomass, well- developed root, increased FDA activities in non-/rhizospheric soil and increased enzyme activities in rhizospheric soil were obtained after *P. indica* colonization in *Medicago sativa* root under phenanthrene and cadmium co-contaminated soil.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [supplement1.pdf](#)
- [supplement2.pdf](#)