Topological Phase Transition Between Normal Insulator and Topological Metal State in a Quasi-one-dimensional System

Milad Jangjan

University of Zanjan

Mir Vahid Hosseini (✉ mv.hosseini@znu.ac.ir)

University of Zanjan

Research Article

Keywords: Topological phase, topological metal state, Fermi level

Posted Date: March 8th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-279154/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Topological phase transition between normal insulator and topological metal state in a quasi-one-dimensional system

Milad Jangjan¹ and Mir Vahid Hosseini^{1,*}

¹Department of Physics, Faculty of Science, University of Zanjan, Zanjan 45371-38791, Iran

*mv.hosseini@znu.ac.ir

ABSTRACT

We theoretically report the finding of a new kind of topological phase transition between a normal insulator and topological metal state where the closing-reopening of bandgap is accompanied by passing the Fermi level through an additional band. The resulting nontrivial topological metal phase is characterized by stable zero-energy localized edge states that exist within the full gapless bulk states. Such states living on a quasi-one-dimensional system with three sublattices per unit cell are protected by hidden inversion symmetry. While other required symmetries such as chiral, particle-hole, or full inversion symmetry are absent in the system.

In recent years, the exploring of topological phases has been at the center of attention, in particular, in condensed matter systems.¹ The key feature is the emergence of symmetry protected gapless boundary modes due to topological bulk states. Topological phases mainly have been categorized into topological insulators²⁻⁵ and topological superconductors⁶ that are studied theoretically and experimentally. In contrast to these topological phases where edge states reside within the gap of bulk states, there are other types of unconventional topological phases known as topological semi-metals⁷ and topological metals (TMs).^{8,9} In the former, there exist band touching nodes at Fermi energy occurring in three dimensional noncentrosymmetric or magnetic materials. While the latter case, which can also take place in low-dimensional systems, has a finite Fermi surface.

Depending on the properties of edge states, topological semimetals and metals can be regarded as two types. First, while topological edge states remain isolated in the bandgap, Fermi level crosses bulk and edge states at different momenta (quantum numbers). This situation, for instance, has been reported in TMs,⁸ semimetals,¹⁰ and even in a narrow energy window of topological insulators.¹¹ Second, gapless edge states coexist with gapless bulk states such that some edge and bulk states would have not only the same energy but also the same momentum (quantum number). This phase has been investigated in a quasi-one-dimensional (1D) system¹⁸ where the finite-energy edge states can penetrate into bulk states hybridizing with them¹² and in a 1D system where the coexistence of edge and bulk states occurs at Fermi energy in a single point of parameter space.⁹ However, this kind of TM phase deserves to be investigated further with a more stable feature.

Often, on the other hand, topological superconducting or insulating phase can be settled down, respectively, on the trivial superconductor or insulator through occurring topological phase transition when band inversion takes place. Furthermore, in previous works, it has been shown that the TM phase can be established on the trivial metallic ground state through closing-reopening of subband gap⁹ or main bandgap.⁸ So, it is indeed intriguing to have a situation in which the underlying state of matter in nontrivial and trivial topological phase of a system belonging to different states can be related to each other, possibly, with a new kind of topological phase transition.

In this paper, within the tight-binding approach in a quasi-1D model with three sublattices per unit cell (Fig. 1), the occurrence of a new kind of topological phase transition between normal insulator (NI) and TM is investigated. Interestingly, in such phase transition in addition to gap closing-reopening between two bands, another band passes the Fermi level (Fig. 2) resulting in the emergence of zero-energy edge states within bulk states (Fig. 3(a)). Although the system has no explicit symmetry, in a subspace of the Hilbert space, there is a hidden inversion symmetry protecting the TM phase.

Figure 1. (Color online) Schematic geometry of quasi-1D lattice comprising of three sublattices A, B, and C per unit cell. Intra and horizontal (oblique) inter unit cell hoppings are indicated by dark magenta and dark cyan (green) colors, respectively.

Model and Theory.

The model we consider is a three-component quasi-1D lattice as represented in Fig. 1. Using the tight-binding theory, the Hamiltonian of system can be defined as

$$\begin{aligned}
 H = & \sum_{n=1}^N t_1 (A_n^\dagger B_n + B_n^\dagger C_n + A_n^\dagger C_n) + \sum_{n=1}^{N-1} t_2' (A_n^\dagger A_{n+1} \\
 & + B_n^\dagger B_{n+1}) + \sum_{n=1}^{N-1} t_2 (C_n^\dagger A_{n+1} + C_n^\dagger B_{n+1}) + h.c.,
 \end{aligned} \quad (1)$$

where $X_n^\dagger$ (X_n) is the creation (annihilation) operator on the sublattice $X (= A, B, C)$ of the n th unit cell and N is the number of unit cell. Also, $t_1 = t(1 + \delta_1)$ is intra unit cell hopping and $t_2^{(\prime)} = t(1 - (+)\delta_2^{(\prime)})$ is oblique (horizontal) inter unit cell hopping to the nearest neighbors. δ_1 and $\delta_2^{(\prime)}$ are some parameters modulating hopping energies. We choose t as the unit of energy.

With periodic boundary conditions, H is invariant under translations by a unit cell. So, after Fourier transformation, the Hamiltonian can be written in the basis of $\psi_k^\dagger = (A_k, B_k, C_k)^\dagger$ yielding compact form $H = \sum_k \psi_k^\dagger \mathcal{H}(k) \psi_k$ with

$$\mathcal{H}(k) = \begin{pmatrix} 2t_2' \cos(k) & t_1 & t_1 + t_2 e^{ik} \\ t_1 & 2t_2' \cos(k) & t_1 + t_2 e^{ik} \\ t_1 + t_2 e^{-ik} & t_1 + t_2 e^{-ik} & 0 \end{pmatrix}. \quad (2)$$

Diagonalizing Eq. (2) gets the eigenvalues,

$$\begin{aligned}
 E^0 &= -2t_1 + \eta, \\
 E^\pm &= \frac{1}{2}(\eta \pm \sqrt{\eta^2 + 2(t_1^2 + 4t_2'^2 + 8t_1 t_2 \cos(k))}),
 \end{aligned} \quad (3)$$

where $\eta = t_1 + 2t_2' \cos(k)$. The possibility of emerging topological phases though gap closing/reopening conditions at the $k_s = (0, \pi)$ would be provided if $E^+ = E^-$ leading to

$$t_1 = e^{ik_s} \frac{2}{9} (t_2' + 4t_2 \pm \frac{\sqrt{-(-2t_2' + t_2)^2}}{2}). \quad (4)$$

Note, Eq. (4) implicates that topological phase transition is possible if the term under the square root is zero, i.e., $t_2' = t_2/2$. Under such condition, Eq. (4) indicates phase boundaries distinguishing topologically nontrivial phase from topologically trivial one.

In Fig. 2, we depicted the band structure of system showing closing and reopening of the energy gap between the two bands E^+ and E^- near the topological phase transition point. In Fig. 2(a), the gap of system is open and there are no energy states at Fermi energy. So, the system is a NI. With increasing t_1 , from Fig. 2(b), one can see that the gap closes leading to the topological phase transition. At the same time, surprisingly, the band E^0 shifts towards the Fermi level and touches it. After topological phase transition, as shown in Fig. 2(c), the energy gap between E^+ and E^- is reopened and the band E^0 crosses the Fermi level which represents a metallic state. This signals that the whole system will be a conductor in a nontrivial topological phase giving rise a new type of topological phase transition from NI to TM phase.

As already mentioned above, if $t_2' \neq t_2/2$ the system does not support any topological phase. Because, in general, Hamiltonian (2) does not have chiral, particle-hole, and/or inversion symmetries. As such, the system provides a trivial phase

Figure 2. (Color online) Band structure of the system for (a) $t_1 = -2t$, (b) $t_1 = -1.2t$, and (c) $t_1 = -0.6t$ under periodic boundary conditions. Horizontal thin line indicates Fermi level. Here, $t'_2 = t_2/2 = -0.6t$.

without any topological edge state in the gap or bulk states under the open boundary conditions. However, if $t'_2 = t_2/2$ then inversion symmetry in a subspace of the Hilbert space of system would be revived (as will be discussed below) and Eq. (4) reduces to $t_1 = t_2 e^{ik_s}$ reminiscing the gap closure condition of SSH model.¹³ So, hereafter, we set $t'_2 = t_2/2$ otherwise specified. Note that to satisfy the requirement $t'_2 = t_2/2$, the hoppings can also be effectively tuned by externally applied periodic fields using Floquet-Bloch theory^{14,15} (for details see¹⁶).

Furthermore, Hamiltonian $\mathcal{H}(k)$ has exchange symmetry with the corresponding exchange operator Υ defined as,

$$\Upsilon \psi_k = \psi'_k = \begin{pmatrix} B_k \\ A_k \\ C_k \end{pmatrix}, \quad (5)$$

which exchanges the sublattices A and B . Owing to the presence of such symmetry, $[\mathcal{H}(k), \Upsilon] = 0$, and then the Hamiltonian can be block-diagonalized in the basis of Υ under the transformation $\mathcal{U}^{-1} \mathcal{H}(k) \mathcal{U} = \mathcal{H}^{BD}(k)$ yielding

$$\mathcal{H}^{BD}(k) = \begin{pmatrix} h_1 & 0 \\ 0 & h_2 \end{pmatrix}, \quad (6)$$

where

$$h_1 = -2t_1 + \eta, \quad (7)$$

$$h_2 = \begin{pmatrix} \eta & \sqrt{2}(t_1 + t_2 e^{ik}) \\ \sqrt{2}(t_1 + t_2 e^{-ik}) & 0 \end{pmatrix}, \quad (8)$$

and

$$\mathcal{U} = \frac{1}{\sqrt{2}} \begin{pmatrix} -1 & 1 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & \sqrt{2} \end{pmatrix}. \quad (9)$$

Note that the obtained bases of exchange operator (9) can be related to a unitary matrix in block-diagonalizing a class of quasi-1D and -2D systems comprised of odd chains.^{17,18} In fact, the block-diagonalization (6) splits the Hilbert space of the system into two subspaces so that one can examine their topological properties independently. Interestingly, in one of the subspaces, the subsystem h_2 can be regarded as generalized SSH model with next-nearest hopping, t'_2 , and on-site potential, t_1 .¹⁹ Then, the subspace associated with h_2 would host topological phases which in combination with the metallic subspace of h_1 would make the whole system TM.

To discuss about the symmetry of the subsystem h_2 we rewrite its Hamiltonian as

$$h_2 = \hbar_0 \sigma_0 + \sum_i \hbar_i \sigma_i, \quad (10)$$

where σ_0 is identity matrix, σ_i is $i(=x, y, z)$ component of Pauli matrices, and

$$\hbar_0 = \hbar_z = \eta/2, \quad (11)$$

$$\hbar_x = \sqrt{2}\eta, \quad (12)$$

$$\hbar_y = \sqrt{2}t_2 \sin(k). \quad (13)$$

It is well-studied that the simultaneous presence of all three components $\mathfrak{h}_x$, $\mathfrak{h}_y$, and $\mathfrak{h}_z$ in 1D two-band Hamiltonians, for instance, SSH model, breaks inversion symmetry and, subsequently destroys topological edge states.²⁰ But, for our model although $\mathfrak{h}_z$ is nonzero, due to the relation $\mathfrak{h}_z = \frac{1}{2\sqrt{2}}\mathfrak{h}_x$, one can find effective inversion symmetry, $\Pi h_2(k)\Pi = h_2(-k)$, in the subspace of h_2 with inversion operator

$$\Pi = \frac{1}{3} \begin{pmatrix} 1 & 2\sqrt{2} \\ 2\sqrt{2} & -1 \end{pmatrix}. \quad (14)$$

This implies that there exists a hidden inversion symmetry in the system protecting the topological phase. The emergence of inversion-symmetry-protected topological phase can be verified by quantized $\mathbb{Z}$ invariant.²¹

Moreover, the h_2 has time-reversal symmetry fulfilling $\mathcal{T}h_2(k)\mathcal{T}^{-1} = h_2(-k)$ where the time-reversal operator is $\mathcal{T} = \sigma_0 \mathcal{K}$ with $\mathcal{K}$ being complex conjugate operator. Also, the reason for the breaking of chiral symmetry can also be seen from the energy dispersions of the two-band subsystem h_2 which are

$$E^\pm = \mathfrak{h}_0 \pm \sqrt{\mathfrak{h}_x^2 + \mathfrak{h}_y^2 + \mathfrak{h}_z^2} = \frac{\mathfrak{h}_x}{3} \pm \sqrt{\frac{9\mathfrak{h}_x^2}{8} + \mathfrak{h}_y^2}. \quad (15)$$

Here, the term $\mathfrak{h}_0$ breaks chiral symmetry so, as usual, one may expect that the zero-energy edge states will be shifted by this term. Nevertheless, interestingly, the relation between $\mathfrak{h}_0$ and $\mathfrak{h}_x$ holds the energy of edge states at zero energy which is in contrast to the previous studies. Therefore, due to $\mathcal{T}^2 = 1$ and $\Pi^2 = 1$, according to the modified periodic table,^{4,22,23} the symmetry class of subsystem h_2 is AI with topological index $\mathbb{Z}$.

The existence of the hidden inversion symmetry guarantees that the eigenvectors of h_2 have a well-defined parity at the inversion symmetric momenta $k_s = (0, \pi)$. So, one can define an integer invariant as $\mathbb{Z} = |n_0 - n_\pi|$ where n_0 and n_π denote the number of negative parities at $k_s = 0$ and $k_s = \pi$, respectively.²¹ The analytical expression of topological invariant $\mathbb{Z}$ for the subsystem h_2 can be obtained as (for details see¹⁶)

$$\mathbb{Z} = \begin{cases} 0, & \text{if } \text{sgn}(\eta(0)) = \text{sgn}(\eta(\pi)) \\ 1, & \text{if } \text{sgn}(\eta(0)) \neq \text{sgn}(\eta(\pi)) \end{cases}, \quad (16)$$

where $\text{sgn}(x)$ is the Sign function. For $\mathbb{Z} = 1$ a nontrivial topological phase will be revealed in the subspace of h_2 . In such a situation, due to bulk-edge correspondence, the gapless edge states associated with the nontrivial topological character of bulk states will be appeared at the boundary of the system under open boundary conditions. For $\mathbb{Z} = 0$ the subsystem h_2 has a trivial phase.

Generally, combining the two-band subsystem h_2 with the single-band subsystem h_1 leads to another topological aspect for which gapped topological bands would coexist with trivial metallic band resulting in TM phase.^{8,9,12,17,18} This means that when the system has open boundary conditions, in the topologically nontrivial phase of h_2 , there exist zero-energy edge states near the system boundaries while the subsystem h_1 has topologically trivial phase. So, the band structure of the whole system shows an interesting phenomenon that topological edge states of h_2 are in the metallic bulk states of h_1 instead of being in the system gap as will be shown below.

Numerical results.

To complement analytical results with numerical ones, we diagonalize the Hamiltonian (1) numerically under open boundary conditions. In order to identify the localized edge states penetrated into extended bulk states under open boundary conditions, in the following, we evaluate the inverse participation ratio (IPR) of states.²⁴ The IPR for an eigenstate $\psi_E(j)$ in the corresponding eigen energy E is given as

$$I_E = \frac{Ln \sum_j |\psi_E(j)|^4}{Ln 3N}. \quad (17)$$

If $\psi_E(j)$ is localized, then $I_E = 0$, whereas if $\psi_E(j)$ is extended, then $I_E = -1$.

The band structure of system along with its corresponding topological invariant $\mathbb{Z}$ as a function of t_1 is plotted in Figs. 3(a) and 3(b) for the cases $t'_2 = t_2/2$ and $t'_2 \neq t_2/2$, respectively. In Fig. 3(a), with establishing the hidden inversion symmetry due to $t'_2 = t_2/2$, the gap of h_2 closes and then reopens at the topological phase transition points. Subsequently, the topological invariant takes 0 and 1 values for trivial and nontrivial regimes, respectively. In the nontrivial region, the system hosts the degenerate zero-energy edge states within the bulk states in a width range of t_1 indicating stable TM phase. Since, in the trivial region, the system is trivial insulator, as a result, a topological phase transition has occurred between NI and stable TM phase. Also, when $t'_2 \neq t_2/2$, owing to violating the hidden inversion symmetry, as shown in Fig. 3(b), the degeneracy of edge states is

Figure 3. (Color online) Energy spectra with their relevant topological invariant as a function of t_1 under open boundary conditions for (a) $t'_2 = t_2/2$ and (b) $t'_2 \neq t_2/2 = -0.3$. Here, $t'_2 = -0.6t$. (c) Topological phase diagram as functions of intra and inter unit cell hoppings t_1 and t_2 . The red region represents the nontrivial TM phase while the gray region indicates NI.

Figure 4. (Color online) Dependence of local density of states on E and on unit cell index x in (a) TM phase with $t_1 = -0.9t$ and (b) NI phase with $t_1 = -1.55t$. Here, $t'_2 = t_2/2 = -0.6t$.

destroyed, and depending on values of t_1 , the system is either NI or trivial metal. Also, the invariant $\mathbb{Z}$ takes continuous values. In Fig. 3(c), topological phase diagram is shown as functions of t_1 and t'_2 . The regions represented by red (gray) colors indicate the nontrivial TM (NI) regime. One can easily see that in addition to the condition $t'_2 = t_2/2$, when $|t_2| > |t_1|$ the TM phase will be appeared.

In Fig. 4, the local density of states (LDOS) as functions of energy E and unit cell index x is displayed. In the TM phase, at zero energy, the localized states are revealed at the ends of the system (see Fig. 4(a)). One can also see, as mentioned above, these localized edge states coexist with the bulk states. Note that the peak intensities of localized edge states at both ends are not the same. This is because of breaking chiral symmetry. In the NI phase, at Fermi energy, the LDOS vanishes (see Fig. 4(b)) and the localized edge states disappear.

Summary and discussion

We introduced a quasi-1D model that presents a new topological phase transition between the TM phase and NI. In such topological phase transition, in addition to bandgap closing-reopening, another band crosses the Fermi level. The Hamiltonian of system can be block-diagonalized into two subsystems in the presence of exchange symmetry. One of these subsystems can be regarded as the generalized SSH model hosting topological edge states when the band of other subsystem passes the Fermi level resulting in the TM phase. Finally, the edge states created in the absence of chiral or particle-hole symmetry and protected by the hidden inversion symmetry. By breaking the hidden inversion symmetry, depending on the parameter, the system has either NI or metallic state with nondegenerate trivial edge states.

Experimentally, our model can be realized by coupled acoustic resonators,¹⁸ topoelectrical circuits,²⁵ optical lattices,²⁶ photonic crystals,^{27,28} and mechanical systems.^{29–33} Using cold atoms, it is possible to simulate quasi-1D chains³⁴ and to reveal the topological features employing density and momentum-distribution measurements.³⁵ Also, using spatially resolved

radio-frequency spectroscopy,³⁶ the topological edge states can be probed by the LDOS.

Acknowledgement—The authors would like to thank T. Neupert and L. Li for helpful comments.

References

1. M. Z. Hasan and C. L. Kane, Rev. Mod. Phys. **82**, 3045 (2010).
2. X.-L. Qi and S.-C. Zhang, Rev. Mod. Phys. **83**, 1057 (2011).
3. C. L. Kane and E. J. Mele, Phys. Rev. Lett. **95**, 226801 (2005).
4. A. P. Schnyder, S. Ryu, A. Furusaki, and A. W. W. Ludwig, Phys. Rev. B **78**, 195125 (2008).
5. L. Fu, C. L. Kane, and E. J. Mele, Phys. Rev. Lett. **98**, 106803 (2007).
6. M. Sato and Y. Ando, Rep. Prog. Phys. **80**, 076501 (2017).
7. A. A. Burkov, Nat. Mater **15**, 1145 (2016).
8. X. Ying and A. Kamenev, Phys. Rev. Lett. **121**, 086810 (2018).
9. M. Bahari and M. V. Hosseini, Phys. Rev. B **99**, 155128 (2019).
10. S.-Y. Xu, C. Liu, S. K. Kushwaha, R. Sankar, J. W. Krizan, I. Belopolski, M. Neupane, G. Bian, N. Alidoust, T.-R. Chang, Science **347**, 294 (2015).
11. L. A. Wray, S.-Y. Xu, Y. Xia, Y. S. Hor, D. Qian, A. V. Fedorov, H. Lin, A. Bansil, R. J. Cava, and M. Z. Hasan, Nat. Phys. **6**, 855 (2010).
12. M. Jangjan and M. V. Hosseini, Sci. Rep. **10**, 14256 (2020).
13. W. P. Su, J. R. Schrieffer, and A. J. Heeger, Phys. Rev. Lett. **42**, 1698 (1979).
14. J. H. Shirley, Phys. Rev. **138**, B979 (1965).
15. H. Sambe, Phys. Rev. A **7**, 2203 (1973).
16. See Supplemental Material for further details.
17. Y.-X. Xiao, Z.-Q. Zhang, and C. Chan, Sci. Rep. **8**, 5160 (2018).
18. Y.-X. Xiao, G. Ma, Z.-Q. Zhang, and C. T. Chan, Phys. Rev. Lett. **118**, 166803 (2017).
19. L. H. Li, Z. H. Xu, and S. Chen, Phys. Rev. B **89**, 085111 (2014).
20. J. K. Asbóth, L. Oroszlány, and A. Pályi, A Short Course on Topological Insulators (Springer, 2016).
21. T. L. Hughes, E. Prodan, and B. A. Bernevig, Phys. Rev. B **83**, 245132 (2011).
22. Y. X. Zhao, A. P. Schnyder, and Z. D. Wang, Phys. Rev. Lett. **116**, 156402 (2016).
23. Y.-M. Lu and D.-H. Lee, arXiv:1403.5558.
24. B. Kramer and A. MacKinnon, Rep. Prog. Phys. **56**, 1469 (1993).
25. Ch. H. Lee, S. Imhof, Ch. Berger, F. Bayer, J. Brehm, L. W. Molenkamp, T. Kiessling, and R. Thomale, Commun. Phys. **1**, 39 (2018).
26. M. Atala, M. Aidelsburger, J. T. Barreiro, D. Abanin, T. Kitagawa, E. Demler, and I. Bloch, Nat. Phys. **9**, 795 (2013).
27. M. Hafezi, S. Mittal, J. Fan, A. Migdall, and J. Taylor, Nat. Photonics **7**, 1001 (2013).
28. J. M. Zeuner, M. C. Rechtsman, Y. Plotnik, Y. Lumer, S. Nolte, M. S. Rudner, M. Segev, and A. Szameit, Phys. Rev. Lett. **115**, 040402 (2015).
29. L. M. Nash, D. Kleckner, A. Read, V. Vitelli, A. M. Turner, and W. T. Irvine, Proc. Natl. Acad. Sci. U.S.A. **112**, 14495 (2015).
30. M. Xiao, G. Ma, Zh. Yang, P. Sheng, Z. Q. Zhang, and C. T. Chan, Nat. Phys. **11**, 240 (2015).
31. B. G. Chen, B. Liu, A. A. Evans, J. Paulose, I. Cohen, V. Vitelli, and C. D. Santangelo, Phys. Rev. Lett. **116**, 135501 (2016).
32. S. D. Huber, Nat. Phys. **12**, 621 (2016).
33. Zh. Yang, F. Gao, and B. Zhang, Sci. Rep. **6**, 29202 (2016).
34. X. P. Li, E. Zhao, and W. V. Liu, Nat. Commun. **4**, 1523 (2013).
35. L. Mazza, M. Aidelsburger, H.-H. Tu, N. Goldman, and M. Burrello, New J. Phys. **17**, 105001 (2015).
36. Y. Shin, C. H. Schunck, A. Schirotzek, and W. Ketterle, Phys. Rev. Lett. **99**, 090403 (2007).

Author contributions

M. V. Hosseini conceived the idea of the research and directed the project. All authors developed the research conceptions, analysed, discussed the obtained results, and wrote the paper. M. Jangjan performed the calculations.

Additional information

Competing financial interests: The authors declare no competing financial and non-financial interests.

Figures

Figure 1

(Color online) Schematic geometry of quasi-1D lattice comprising of three sublattices A, B, and C per unit cell. Intra and horizontal (oblique) inter unit cell hoppings are indicated by dark magenta and dark cyan (green) colors, respectively.

Figure 2

See the Manuscript Files section for the complete figure caption.

Figure 3

See the Manuscript Files section for the complete figure caption.

Figure 4

See the Manuscript Files section for the complete figure caption.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryMaterialforTopologicalphasetransitionbetweennormalinsulatorandtopologicalmetalstateinaquasione-dimensional-system.pdf](#)