Topological phases, topological phase transition, and bulk-edge correspondence of magnetized cold plasmas

Yichen Fu (✉ yfu@pppl.gov)

Princeton Plasma Physics Laboratory <https://orcid.org/0000-0002-5205-9827>

Hong Qin

Princeton Plasma Physics Laboratory

Article

Keywords: Topological Materials, Langmuir Wave-L-wave Resonance, Langmuir Wave-cyclotron Wave Resonance, Zero Magnetic Field, Edge Modes, Fermi Arcs

Posted Date: March 4th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-279306/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Nature Communications on June 24th, 2021. See the published version at <https://doi.org/10.1038/s41467-021-24189-3>.

Topological phases, topological phase transition, and bulk-edge correspondence of magnetized cold plasmas

Yichen Fu* and Hong Qin

*Princeton Plasma Physics Laboratory and Department of Astrophysical Sciences,
Princeton University, Princeton, NJ 08540*

Abstract

Plasmas have been recently studied as topological materials. We attempt at a comprehensive picture of the topological phases and the bulk-edge correspondence of magnetized cold plasmas. We find that there are 10 topological phases in the parameter space of density n , magnetic field B , and parallel wavenumber k_z , separated by the surfaces of Langmuir wave-L wave resonance, Langmuir wave-cyclotron wave resonance, and zero magnetic field. For fixed B and k_z , only the phase transition at the Langmuir wave-cyclotron wave resonance corresponds to edge modes. A sufficient and necessary condition for the existence of this new type of edge modes is given and verified by numerical solutions. The edge modes fall into four categories characterized by different behaviors of the Fermi arcs. We demonstrate that edge modes exist not only on a plasma-vacuum interface but also on more general plasma-plasma interfaces. This finding broadens the possible applications of these exotic excitations in space and laboratory plasmas.

Recently, the relation between the topological properties of the bulk modes and the chiral (unidirectional) edge modes has attracted growing interest in classical fluid [1–5] and plasma physics [6–10]. Originating in condensed matter physics, the bulk-edge correspondence [11–13] predicts that at the interfaces between two topologically different materials, there exist gapless edge modes across the common band gap. For condensed matter systems, the gap Chern number is a topological invariant for bulk modes to give a $\mathbb{Z}$ classification [12]. However, continuum media, including plasmas, do not have well defined Brillouin zones, and compactification techniques in the wavenumber space [4, 5, 10, 14] are required to generate integer Chern numbers.

For cold magnetized plasmas, topological phases and topological phase transitions have not been systematically mapped out because of the complicated parameter dependency. The bulk-edge correspondence has not been well established due to the non-compactness of the wavenumber space. For cases with wave vectors perpendicular to the magnetic field, the bulk topology and corresponding edge states of the X waves (transverse-magnetic waves) have been extensively studied [13–19] when the O waves (transverse-electric waves) are ignored. It is reported that under certain conditions, the bulk-edge correspondence between the gap of X waves can be physically violated [20]. Another type of edge mode has been derived using a simplified analytical model [6] and linked to the Weyl degeneracies [7]. This edge mode was also numerically demonstrated [10] for a plasma-vacuum interface with continuous density falloff. However, the corresponding bulk topological phases and phase transition have not yet been identified.

In the present study, we attempt at a comprehensive picture of the topological phases, topological phase transitions, and the bulk-edge correspondence of magnetized cold plasmas in the absence of a solid boundary. We find that in the parameter space of magnetic field $\mathbf{B} = B_0 \hat{z}$, density n , and wavenumber k_z , there are 10 topological phases, separated by the Langmuir wave-L wave (LL) resonance, the Langmuir wave-Cyclotron wave (LC) resonance, and the $B = 0$ surface. Their topological properties are classified by the integer Chern numbers of the spectrum. For fixed non-vanishing B and k_z , there are two possible topological phase transitions due to the two resonances, while only the transition at the LC resonance produces edge modes. There exists a critical density n_c such that plasmas below and above n_c are in different topological phases across the LC resonance. We find that edge modes exist at more general plasma-plasma interfaces when a necessary and sufficient

condition, Eq. (5), is satisfied, and the edge modes can be categorized by different behaviors of the Fermi arcs or Fermi-arcs-like curves. This finding broadens the possible applications of these exotic edge modes in space and laboratory plasmas.

Results

Bulk dispersion relation and eigenmodes. Following Ref. [10], we use the linearized fluid equations for a magnetized cold plasma with stationary ion and uniform density n_e to study the eigenmodes of the system. The constant background magnetic field is in the z -direction, i.e., $\mathbf{B}_0 = B_0 \hat{\mathbf{z}}$, and there is no equilibrium flow. After spacetime Fourier transform, $\partial_t \rightarrow -i\omega$, $\nabla \rightarrow i\mathbf{k}$, and the governing equations can be written as $H(\mathbf{k})|\psi\rangle = \omega|\psi\rangle$, where $H(\mathbf{k})$ is a 9×9 Hermitian matrix, and $|\psi\rangle = (\mathbf{v}, \mathbf{E}, \mathbf{B})^T$ is a nine-dimensional vector consisting of the perturbed velocity and electromagnetic fields.

The dispersion relation is given by the vanishing determinant of $H(\mathbf{k}) - \omega\mathbf{I}_9$, which can be simplified as

$$\det(\mathbf{N}\mathbf{N} - N^2\mathbf{I}_3 + \boldsymbol{\epsilon}) = 0, \quad (1)$$

where $\mathbf{N} = c\mathbf{k}/\omega$, and $\boldsymbol{\epsilon}$ is the 3×3 cold plasma dielectric tensor. In terms of the standard notations in Ref. [21],

$$\boldsymbol{\epsilon} = \begin{bmatrix} S & -iD & 0 \\ iD & S & 0 \\ 0 & 0 & P \end{bmatrix}, \quad P = 1 - \frac{\omega_p^2}{\omega^2},$$

$$S = 1 - \frac{\omega_p^2}{\omega^2 - \Omega^2}, \quad D = -\frac{\Omega}{\omega} \frac{\omega_p^2}{\omega^2 - \Omega^2},$$

where $\omega_p = \sqrt{n_e e^2 / m_e \epsilon_0}$ is the plasma frequency, and $\Omega = -eB_0 / m_e$ is the signed electron cyclotron frequency. The plasma is called underdense if $\omega_p < |\Omega|$, and overdense if $\omega_p > |\Omega|$. It can be shown that the dispersion relation surfaces in the $(\omega, \mathbf{k})$ space are symmetric with respect to all four coordinate hyperplanes. For each given k_z , the system has 9 eigenvalues ω_n and eigenvectors $|\psi_n\rangle$ as functions of $\mathbf{k}_\perp$, where $\omega_{-n} = -\omega_n$ and $n = -4, -3, \dots, 3, 4$ is the index for the eigenmodes. Note that $\omega_0 = 0$ is the zero frequency eigenmode of the system. The dispersion surfaces $\omega(k_z, k_\perp)$ of the four positive-frequency branches for both overdense and underdense plasmas are shown in Figs. 1(a) and 1(c). To better illustrate the

crossing between branches, the dispersion curves $\omega(k_z)$ at different values of $k_\perp$ are shown in Figs. 1(b) and 1(d). We can observe that branch crossing occurs only when $k_\perp = 0$, represented by the coldest blue lines. In this case, the horizontal lines $\omega = \omega_p$ are the Langmuir waves given by $P = 0$. The other three branches are the R wave and the L wave given by $N^2 = R$ and $N^2 = L$, respectively, where $R = S + D$ and $L = S - D$. For convenience, we define functions

$$k^\pm := \frac{\omega_p/c}{\sqrt{1 \pm \omega_p/\Omega}}. \quad (2)$$

When $k_z > 0$, the Langmuir wave resonates with the L wave at $k_z = k^+$. In an underdense plasma, the Langmuir wave also resonates with the lower branch of the R wave, a.k.a electron cyclotron wave, at $k_z = k^-$. Notice that $k^- \rightarrow \infty$ when $\omega_p \rightarrow |\Omega|$. These four resonant points at $k_z = \pm k^\pm$, previously recognized as the Weyl points [7], play important roles in determining the topological properties of magnetized plasmas.

86

Topological phase diagram. When fixing k_z as a parameter, we can calculate the Chern number for each band, i.e., branch of the dispersion relation, in the $\mathbf{k}_\perp = (k_x, k_y)$ space. If the $\mathbf{k}_\perp$ space can be properly compactified [10, 14], Chern numbers in the system should be integers, which are invariant under continuous transforms. It means that each band's Chern number is a topological invariant that can change only when different bands cross. As shown in Fig. 1, the band crossing in a cold plasma is only possible at the points $(k_x, k_y, k_z) = (0, 0, \pm k^\pm)$ when $B \neq 0$ and $k_z \neq 0$. Therefore, the locations of $k_z = \pm k^\pm$ defines the boundaries between topologically different regions if separated regions have different Chern numbers. In Fig. 2(a), the surfaces of $k_z = k^\pm$ and $\Omega = 0$ in the (ω_p, Ω, k_z) space are shown, which separate the parameter space into 10 different regions, each represents a different phase characterized by a set of Chern numbers. Although the first band touches the zero-frequency mode at $k_z = 0$, we will show later that this band crossing does not affect the topology. The cross sections of the 3D surfaces at $(\omega_p = 1, \Omega > 0, k_z > 0)$ and $(\omega_p > 0, \Omega = 1, k_z > 0)$ are shown in Figs. 2(b) and 2(c), each of which is separated into three phases. Notice that phase I in these cross sections only exists in underdense plasmas, i.e, when $\omega_p < |\Omega|$.

We adopt the same formalism of Berry curvature and regularization strategy used in Ref. [10] to calculate the Chern numbers numerically. The Chern number of the n -th band is denoted by C_n . As functions of (ω_p, Ω, k_z) , Chern numbers admit the following symmetries,

FIG. 1. Dispersion relation surfaces of (a) an overdense plasma and (c) an underdense plasma. Only the $k_{\perp} > 0$ and $k_z > 0$ part of positive-frequency branches are shown. When fixing $k_{\perp}$, the dispersion curves $\omega(k_z)$ for both cases are shown in (b) and (d). The colors in (b) and (d) represent different values of $k_{\perp}$. Only the crossing of the curves with the same color indicates the crossing of different branches. $\omega_{1,2}/|\Omega| = (\sqrt{4r^2 + 1} \pm 1)/2$, where $r = \omega_p/|\Omega|$. $k^{\pm}$ are given by Eq. (2) assuming $\Omega > 0$.

$C_n(\Omega, k_z) = C_n(\Omega, -k_z) = -C_n(-\Omega, k_z) = -C_{-n}(\Omega, k_z)$. See additional details in Methods. Therefore, it suffices to calculate the Chern numbers for the three phases in the domain of $k_z > 0$ and $\Omega > 0$, as shown in Figs. 2(b) and 2(c).

The resulting Chern numbers for each positive-frequency band in all three phases are shown in Figs. 3(a)-3(c). We find that the Chern numbers in phases II and III are $(C_1, C_2, C_3, C_4) = (-1, 1, 1, -1)$ and $(-1, 2, 0, -1)$, respectively, as reported in Ref. [10]. For the special case of $k_z = 0$, the Chern numbers for phase III are consistent with those calculated in Refs. [14, 17]. For phase I, we discover that its Chern numbers are $(C_1, C_2, C_3, C_4) = (0, 0, 1, -1)$. This is a new finding that was not reported previously. When the parameters cross the boundary of $k_z = k^-$ and change from phase I to phase II, bands 1 and 2 cross at $k_{\perp} = 0$ and change their Chern numbers from $(0, 0)$ to $(-1, 1)$.

FIG. 2. (a) 3D phase diagram of a cold plasma in the (ω_p, Ω, k_z) space. There are 10 topological phases. (b) 2D cross section of (a) at $\omega_p = 1$. (c) 2D cross section of (a) at $\Omega = 1$. Only the $k_z > 0$ and $\Omega > 0$ part is shown. Dashed lines indicates $\Omega = \omega_p$. Dotted lines indicates $k_z = k^*$. The band structures and Chern numbers at three colored dots in (b) are shown in Fig. 3.

FIG. 3. Chern numbers of each positive frequency bands of the three phases of the phase diagram. Since the dispersion relation is isotropic in (k_x, k_y) plane, only the $k_{\perp} > 0$ part is shown. The parameters used are $\Omega = 1$ and $(\omega_p, ck_z) = (0.4, 1.2)$, $(1.5, 1.2)$, and $(1.5, 0.5)$, which are shown as colored dots in Fig. 2(b).

116 This change agrees with the fact that the Weyl point at $k_z = k^-$ has Chern number 1 [7].
 117 Similar behavior is observed between bands 2 and 3 when parameters cross the boundary
 118 of $k_z = k^-$ and change from phase II to phase III.

Of particular interest in the present study is the transition between phases I and II. The boundary between them, i.e., the LC resonance surface, defines a critical density n_c , which, expressed in terms of the corresponding plasma frequency $\omega_{p,c} = \sqrt{n_c e^2 / m_e \epsilon_0}$, is

$$\omega_{p,c} = \frac{|\Omega|}{2} \left[\sqrt{\left(\frac{ck_z}{\Omega}\right)^4 + 4\left(\frac{ck_z}{\Omega}\right)^2} - \left(\frac{ck_z}{\Omega}\right)^2 \right]. \quad (3)$$

For fixed k_z and Ω , transition between phases I and II occurs at $n = n_c$. Notice that when $k_z \rightarrow \infty$, $\omega_{p,c} \rightarrow |\Omega|$, and n_c becomes $n_c^\infty \equiv \Omega^2 m_e \epsilon_0 / e^2$.

The surface of $\Omega = 0$ is also a boundary between different topological phases because $C_n(-\Omega) = -C_n(\Omega)$. For the special case of $k_z = 0$, the boundary between phases I and II and the boundary between phases II and III collapse to the lines defined by $\Omega = 0$ and $\omega_p = 0$. In this case, the only possible nontrivial phase transition happens at the $\Omega = 0$ surface.

Band gaps.— Bulk-edge correspondence suggests that edge modes exist in the common band gaps at the interface between two topological materials with different gap Chern numbers. We now identify possible band gaps in a magnetized cold plasma. At fixed k_z , when $k_\perp \rightarrow \infty$, $(\omega_1, \omega_2, \omega_3, \omega_4) \rightarrow (0, \omega_{\text{uh}}, ck_\perp, ck_\perp)$, where $\omega_{\text{uh}}^2 = \omega_p^2 + \Omega^2$ is the upper hybrid frequency. Thus, when $|k_z| \neq k^\pm$, it is possible to have gaps between bands 1 and 2 and between bands 2 and 3. However, the gap between bands 2 and 3 does not always exist. When $k_\perp = 0$, $\omega_3(k_z)$ is given by $N^2 = L$. The non-overlapping of bands 2 and 3 requires $\omega_3(k_z) > \omega_{\text{uh}}$, which leads to

$$|k_z| > k^* := \frac{|\Omega|}{c} \left(1 + \frac{\omega_p^2}{\Omega^2} \right)^{1/4}. \quad (4)$$

The locations of $|k_z| = k^*$ in the parameter space are shown in Fig. 2(b) and 2(c). It is clear that the gap between bands 2 and 3 does not exist in phase III because condition (4) is not satisfied there.

The topology of a band gap is characterized by its gap Chern number, which is defined as $C_{i,i+1} = \sum_{n=-4}^i C_n$ for the gap between the i -th and $(i+1)$ -th bands. In phase I, both $C_{1,2}$ and $C_{2,3}$ are trivially zero, as in the phase of vacuum [11] that phase I neighbors at the boundary of $\omega_p = 0$. Thus, as far as the gap topology is concerned, phase I plasmas are identical to the vacuum, which is interesting if not surprising. In phase II, the gap

Chern numbers $(C_{1,2}, C_{2,3})$ become $(-1, 0)$, indicating a topological phase transition at the boundary between phases I and II due to the crossing of the gap between bands 1 and 2. In phase III, the gap Chern numbers $(C_{1,2}, C_{2,3})$ are $(-1, 1)$. Although $C_{2,3}$ is different between phase II and III, there is no band gap between bands 2 and 3 in phase III as proved above. When $|k_z| \neq 0$, there is another band gap between bands 0 and 1. However, the gap Chern number $C_{0,1}$ for this gap is zero for all three phases, and it is a trivial band gap. Therefore, only the band gap between bands 1 and 2 shared by phases I and II is interesting in the context of bulk-edge correspondence.

It is worth mentioning that when $k_z = 0$, band 3 becomes the O wave, and bands 2 and 4 are the X wave. If one chooses to ignore the O wave [13–15, 17], then a band gap shows up between bands 2 and 4, as long as $\Omega \neq 0$. The physical properties of this gap has been extensively studied, including the violation of bulk-edge correspondence under certain conditions [20]. However, as an important eigenmode in magnetized cold plasmas, band 3 always exists. Especially when $|k_z| > k^+$, band 3 has a non-zero Chern number and should not be ignored. In the present study, we include all 9 bands in magnetized cold plasmas.

Bulk-edge correspondence. Having established the topological phase diagram and possible band gaps in a magnetized cold plasma, we now investigate the edge modes at the interface between two different magnetized cold plasmas. As discussed above, at a fixed B , the only possible nontrivial gap that admits two different gap Chern numbers is the gap between bands 1 and 2 shared by phases I and II. The bulk-edge correspondence then predicts that edge modes exist in the common band gap at the interface between a phase I plasma and a phase II plasma. We now solve for these edge modes numerically in an 1D inhomogeneous plasma. The background magnetic field is constant, i.e., $\mathbf{B}_0 = B_0 \hat{z}$, and the plasma density is nonuniform only in the x -direction, as shown in Fig 4(a). The density profile is given by $n(x) = \frac{1}{2}(n_1 - n_2)\{\tanh[-(x-l)/\delta] + \tanh[(x+l)/\delta]\} + n_2$, where n_1 and n_2 are the densities of the inner and outer plasmas, and l and δ are the location and width of the interface. For realistic plasmas, the width of the interface is finite, i.e., $\delta > 0$.

Because the system is uniform in the z -direction, the parallel wavenumber k_z enters as a parameter. The inner and outer plasmas can be represented by two points, $(\omega_{p,1}, \Omega, k_z)$ and $(\omega_{p,2}, \Omega, k_z)$, in the phase diagram shown in Fig. 2. Here, $\omega_{p,1}$ and $\omega_{p,2}$ are the inner and outer plasma frequencies. As discussed above, when there is a common band gap shared

by the inner and outer plasmas, chiral edge modes exist in the gap if and only if the inner plasma is in phase I and outer plasma is in phase II such that they have different gap Chern numbers. Furthermore, the number of chiral edge modes at the interface should be equal to the difference of the gap Chern numbers, which is 1 in the present case. Since n_c is the critical density at the boundary between phases I and II, for given k_z and Ω , the chiral edge mode exists if and only if

$$n_1 > n_c > n_2, \quad (5)$$

which, expressed in terms of plasma frequencies, is

$$\frac{\omega_{p,1}}{|\Omega|} + \frac{\omega_{p,1}^2}{c^2 k_z^2} > 1 > \frac{\omega_{p,2}}{|\Omega|} + \frac{\omega_{p,2}^2}{c^2 k_z^2}. \quad (6)$$

Here, we observe that underdense and overdense plasmas behave differently. When both the inner and outer plasmas are overdense, edge mode cannot exist regardless of k_z . If the inner plasma is overdense while the outer plasma is underdense, edge modes can be found when $k_z > k^-(\omega_{p,2})$. If both the inner and outer plasma are underdense, the edge mode can only be found when $k^-(\omega_{p,1}) > |k_z| > k^-(\omega_{p,2})$. Notice that when the outer side is a vacuum, the criteria given by Eq. (6) can be satisfied in two scenarios, either k_z^2 is small enough or the inner plasma is overdense. Incidentally, all the parameters chosen in Ref. [10] belong to the first scenario.

To numerically verify the criteria in Eqs. (5) or (6), we Fourier-transform in y , z and t , then spatially discretize the Hamiltonian H in the x -direction using a finite difference method. A periodic boundary condition at $x = \pm 2l$ is adopted, as in Ref. [1]. More details of the numerical methods are provided in the methods. The structures of the positive-frequency bands for different (n_1, n_2) at $ck_z/|\Omega| = 0.7$ are shown in Fig. 4(d)-4(f). For the cases of $n_c > n_1 > n_2 = 0$ and $n_1 > n_2 > n_c$, there is no edge mode in the gap between the first and second bands. In particular, Fig. 4(f) shows that the edge modes can be absent at a plasma-vacuum interface. In Fig. 4(e), $n_1 > n_c > n_2$ and there are two gapless edge modes in the band gap. Due to the symmetry of $\omega(k_y) = \omega(-k_y)$, the two edge modes cross at $k_y = 0$. The electric field structure of the two gapless modes localized at different edges is shown in Figs. 4(b) and 4(c). The number of edge modes at each edge is the same as the difference of the gap Chern number, consistent with the prediction of bulk-edge correspondence.

To further understand the edge modes, band structures at $k_y = 0$ with various densities are plotted in Fig. 5, which shows that edge modes can be classified by behaviors of the

FIG. 4. The band structure of nonuniform plasma for $ck_z/|\Omega| = 0.7$. The critical plasma frequency given by Eq. (3) is $\omega_{p,c}/|\Omega| \approx 0.5$. (a) Schematic diagram of the density profile in x direction, where $cl/|\Omega| = 40$ and $c\delta/|\Omega| = 4$. (b) and (c) The non-zero components of electric fields of edge modes in (e) at $ck_y/|\Omega| = 0.05$. (d), (e), and (f) The band structure $\omega(k_y)$ with various inner and outer densities. The plasma frequencies $(\omega_{p,1}, \omega_{p,2})/|\Omega|$ are $(0.8, 0.6)$, $(0.6, 0.4)$, and $(0.4, 0)$ in (d), (e), and (f), respectively.

207 Fermi arc or Fermi-arc-like curves. The inner plasma is underdense in Figs. 5(a) and 5(c)
 208 and overdense in Figs. 5(b) and 5(d). The outer is underdense in Figs. 5(c) and 5(d) and
 209 vacuum in Figs. 5(a) and 5(b). The bulk dispersions are shown by orange and blue lines for
 210 inner and outer plasmas in each case. The topological edge modes are shown in red lines
 211 representing the crossing points between the left and right edge modes at each given k_z . We
 212 can see that the range where edge modes exist coincides with Eq. (6) exactly. Noticeably, in
 213 Fig. 5(a), edge modes exist when $|k_z| < k^-(\omega_{p,1})$ between a underdense plasma and a vacuum.
 214 The dispersion surface $\omega = \omega(k_z, k_y)$ of the edge modes connects the two Weyl points of the
 215 inner plasma at $k_z = \pm k^-(\omega_{p,1})$. The intersection of this dispersion surface and $\omega = \omega_{p,1}$
 216 is known as the Fermi arc connecting two Weyl points [22, 23]. When the inner plasma is
 217 overdense in Fig. 5(b), the Weyl points $\pm k^-(\omega_{p,1})$ disappear, but the edge modes still exist
 218 and the dispersion surface connects to $k_z = \pm\infty$. When the inner and outer plasmas are

FIG. 5. Band structure at $k_y = 0$, showing four possible types of Fermi-arc-like structures of edge mode connecting Weyl points. The bulk modes of the inner and outer plasmas when $k_\perp = 0$ are highlighted by orange and blue curves, respectively. The topological edge modes are highlighted by the red curves. $k_1^- = k^-(\omega_{p,1})$ and $k_2^- = k^-(\omega_{p,2})$. The plasma frequencies $(\omega_{p,1}, \omega_{p,2})/|\Omega|$ in (a)-(d) are $(0.75, 0)$, $(1.1, 0)$, $(0.75, 0.3)$, and $(1.1, 0.3)$.

all underdense in Fig. 5(c), edge modes are prohibited if $-k^-(\omega_{p,2}) < k_z < k^-(\omega_{p,2})$, then the dispersion surface no longer connects the Weyl points of the inner plasma at positive and negative k_z . Instead, a Fermi-arc-like curve connects $k_z = k^-(\omega_{p,1})$ and $k_z = k^-(\omega_{p,2})$, the Weyl points of the inner and outer plasmas. In Fig. 5(d), the inner is overdense and the outer underdense, a Fermi-arc-like curve connects the Weyl point of the outer plasma to infinity. These numerical results also confirm the condition given by Eqs. (5) or (6) for the existence of the edge modes.

As a side note, the gap between bands 1 and 2 of the inner plasma may overlap with the gap between bands 2 and 3 of the outer plasma. It is reasonable to suggest that if

228 $C_{\text{in},1,2} \neq C_{\text{out},2,3}$, edge modes might exist within this gap. However, this common gap is
 229 filled with other eigenmodes in reality. The upper hybrid frequency ω_{uh} , which depends on
 230 plasma density, sets the upper range for band 2. Since the density profile is continuous, the
 231 local upper hybrid frequency $\omega_{\text{uh}}(x)$ will always fill in the gap between the second bands of
 232 the inner and outer plasmas. An example is illustrated in Fig. 6, where the inner and outer
 233 plasmas belong to phases II and I, respectively.

FIG. 6. Band structures when the gap between bands 1 and 2 of the inner plasma overlaps with gap
 between bands 2 and 3 of the outer plasma. Here, $ck_z/|\Omega| = 1.4$ and $(\omega_{p,1}, \omega_{p,2})/|\Omega| = (1.4, 0.1)$.
 Orange and blue lines indicate the bulk modes of the inner and outer plasma when $k_x = 0$.

234 Varying density is a convenient but not the only way to create interfaces between
 235 topologically different plasmas. Since plasma topology varies with the strength and direction
 236 of the magnetic field, one can create topologically nontrivial interfaces by assembling two
 237 plasmas with different background magnetic field. In particular, when $k_z = 0$, the only
 238 possible topological phase transition in a cold plasma occurs at the $\Omega = 0$ surface, as
 239 discussed above. In this case, the edge modes at the interface between two plasmas with
 240 opposite magnetic fields has been studied for the X wave [13, 18]. However, as a whole
 241 system, such a setup is inhomogeneous, and the O wave cannot be decoupled. More
 242 thorough analysis is need.

243

244 Discussion

In summary, we found that a magnetized cold plasma has 10 different topological phases in the (ω_p, Ω, k_z) space. The different phases are separated by the surfaces of Langmuir wave-L wave resonance, Langmuir wave-cyclotron wave resonance, and $\Omega = 0$. We found that at fixed Ω and k_z , only the band gap between bands 1 and 2 shared by phases I and II is interesting in the context of bulk-edge correspondence, and that the necessary and sufficient condition for the existence of edge modes is $n_1 > n_c > n_2$. These findings were verified by numerical studies of the corresponding chiral edge modes in 1D inhomogeneous plasmas. The edge modes exist not only on the plasma-vacuum interface, but also on more general plasma-plasma interfaces. No violation of bulk-edge correspondence is observed in this system.

Our study of edge modes focused on the general gaseous plasma-plasma interfaces. We didn't consider gaseous plasma-solid interfaces, which, unlike the interfaces between solid state materials [23], involve more complex physical processes, such as the plasma sheath and plasma-wall interactions [24]. It is not appropriate to model gaseous plasma-solid interfaces only as simple interfaces between two different topological materials.

The present investigation on the topology phases and edge modes is carried out for the model of magnetized cold plasmas. When other physical effects, such as finite temperature, plasma collisions, and kinetic interactions, are important, band structures of the system can change significantly [21]. The bulk topology and the validity of the general bulk-edge correspondence in these regimes need to be further investigated. In particular, linear dynamics in these plasmas is expected to be non-Hermitian [25, 26], permitting unstable and damped eigenmodes. Applying the methods of topological phases for non-Hermitian systems will bring new insights and discoveries in the study of plasma instabilities for laboratory and astrophysical plasmas.

Methods

Basic equations. Here we outline the band structure of cold plasma waves. Assume the background plasma is uniform and stationary, the ions are motionless and the magnetic field

274 is constant, i.e., $\mathbf{B}_0 = B_0 \hat{z}$. The linearized fluid equations are [21]

$$\begin{aligned}
\partial_t \mathbf{v} &= \frac{e}{m_e} (\mathbf{E} + \mathbf{v} \times \mathbf{B}_0), \\
\partial_t \mathbf{E} &= c^2 \nabla \times \mathbf{B} - \frac{en_e}{\epsilon_0} \mathbf{v}, \\
\partial_t \mathbf{B} &= -\nabla \times \mathbf{E},
\end{aligned} \tag{7}$$

275 where $\mathbf{v}, \mathbf{B}, \mathbf{E}$ are perturbed velocity, magnetic field and electric field, e is the electron
276 charge, m_e and n_e are electron mass and density, and c is light speed. Let plasma frequency
277 $\omega_p^2 = n_e e^2 / \epsilon_0 m_e$ and signed cyclotron frequency $\Omega = -e B_0 / m_e$. Define new velocity $\tilde{\mathbf{v}} = \omega_p \mathbf{v}$,
278 reference electric field $\bar{\mathbf{E}}$ and reference frequency $\bar{\omega}$. We normalize time to $\bar{\omega}^{-1}$, frequency
279 to $\bar{\omega}$, length to $c/\bar{\omega}$, new velocity to $e\bar{\mathbf{E}}/m_e$, electric field to $\bar{\mathbf{E}}$, and magnetic field to $\bar{\mathbf{E}}/c$.
280 The equation system becomes

$$\begin{aligned}
\partial_t \tilde{\mathbf{v}} &= \omega_p \mathbf{E} - \Omega \tilde{\mathbf{v}} \times \hat{z}, \\
\partial_t \mathbf{E} &= \nabla \times \mathbf{B} - \omega_p \tilde{\mathbf{v}}, \\
\partial_t \mathbf{B} &= -\nabla \times \mathbf{E}.
\end{aligned} \tag{8}$$

281 Notice that when the density is not uniform, it suffices to change ω_p to $\omega_p(\mathbf{r})$ in Eq. 8. From
282 now on, we omit the tilde for convenience. After space Fourier transform, $\partial_t \rightarrow -i\omega$, $\nabla \rightarrow i\mathbf{k}$,
283 and the governing equations can be written as $H|\psi\rangle = \omega|\psi\rangle$, where $|\psi\rangle = (\mathbf{v}, \mathbf{E}, \mathbf{B})^T$ and

$$H(\omega_p, \Omega, \mathbf{k}) = \begin{pmatrix} i\Omega \hat{z} \times & i\omega_p & 0 \\ -i\omega_p & 0 & -\mathbf{k} \times \\ 0 & \mathbf{k} \times & 0 \end{pmatrix}. \tag{9}$$

284 Here, H is a 9×9 Hermitian matrix. The dispersion relation is given by $\det(H - \omega \mathbf{I}_9) = 0$,
285 which simplifies to Eq. (1).

286

287 **Symmetries of the system.** As a real system, the equations of motion for plasmas admit
288 an unbreakable particle-hole symmetry [9, 27]. For our system, the symmetry states that
289 $H(-\mathbf{k})^* = -H(\mathbf{k})$, where $*$ denotes complex conjugate. It ensures that the dispersion
290 relation has the symmetry of $\omega(-\mathbf{k}) = -\omega(\mathbf{k})$. Notice that Eq. (1) remains invariant under
291 $\mathbf{k} \rightarrow -\mathbf{k}$, so the dispersion relation satisfy $\omega(\mathbf{k}) = \omega(-\mathbf{k})$ as well. In addition, since the
292 system is isotropic in the direction perpendicular to the background magnetic field, ω only
293 depends on $k_z \equiv k_z$ and $k_\perp \equiv \sqrt{k_x^2 + k_y^2}$. Therefore, the dispersion relation surfaces in

the (ω, k_x, k_y, k_z) space are symmetric with respect to the reflections of all four coordinate hyperplanes.

The symmetries of eigenvalues can also be obtained, which will be useful during the calculation of the Chern numbers. $H(\omega_p, \Omega, \mathbf{k})$ has 9 eigenvalues, one of which is identically zero. The eigenvalues and corresponding eigenvectors can be labeled as $\omega_n, |\psi_n\rangle$, where $n = -4, -3, \dots, 3, 4$, $\omega_i < \omega_j$ if $i < j$. Assume that at some $(\omega_p, \Omega, \mathbf{k})$, the n -th eigenvalue and eigenvector are $\omega_n = \omega$ and $|\psi_n\rangle = (\mathbf{v}, \mathbf{E}, \mathbf{B})^T$. We can verify the following symmetries of eigenvalues and eigenvectors:

1. For the reflection of band number $n \rightarrow -n$, $\omega_{-n} = -\omega$ and $|\psi_{-n}\rangle = (\mathbf{v}^*, \mathbf{E}^*, -\mathbf{B}^*)^T$.
2. For the reflection of magnetic field $\Omega \rightarrow -\Omega$, $\omega_n = \omega$ and $|\psi_n\rangle = (-\mathbf{v}^*, \mathbf{E}^*, \mathbf{B}^*)^T$.
3. For the reflection of wavenumber $\mathbf{k} \rightarrow -\mathbf{k}$, $\omega_n = \omega$ and $|\psi_n\rangle = (\mathbf{v}, \mathbf{E}, -\mathbf{B})^T$.

The symmetries of Chern numbers. In this section, we briefly describe the calculation and symmetries of Chern numbers. For any given parallel wavenumber k_z , the Chern number can be calculated in the $\mathbf{k}_\perp$ space for each band by [12] $C_n = (2\pi)^{-1} \int d\mathbf{S} \cdot \mathbf{F}_n(\mathbf{k})$, where $\mathbf{F}_n = \nabla_{\mathbf{k}} \times \mathbf{A}_n$ is the Berry curvature, and $\mathbf{A}_n = i\langle\psi_n|\nabla_{\mathbf{k}}\psi_n\rangle$ is the Berry connection. Let $|\psi_n(\omega_p, \Omega, \mathbf{k})\rangle = (\mathbf{v}, \mathbf{E}, \mathbf{B})^T$. The Berry connection is $\mathbf{A}_n(\omega_p, \Omega, \mathbf{k}) = i(\mathbf{v}^\dagger \nabla_{\mathbf{k}} \mathbf{v} + \mathbf{E}^\dagger \nabla_{\mathbf{k}} \mathbf{E} + \mathbf{B}^\dagger \nabla_{\mathbf{k}} \mathbf{B})$, where $\dagger$ denotes conjugate transpose. Based on the symmetries of eigenvectors, we obtain the following symmetries of Chern numbers:

1. For the reflection of band number $n \rightarrow -n$,

$$\begin{aligned} \mathbf{A}_{-n}(\omega_p, \Omega, \mathbf{k}) &= i\langle\psi_{-n}|\nabla_{\mathbf{k}}\psi_{-n}\rangle \\ &= i(\mathbf{v} \nabla_{\mathbf{k}} \mathbf{v}^\dagger + \mathbf{E} \nabla_{\mathbf{k}} \mathbf{E}^\dagger + \mathbf{B} \nabla_{\mathbf{k}} \mathbf{B}^\dagger) \\ &= -\mathbf{A}_n(\omega_p, \Omega, \mathbf{k}). \end{aligned} \tag{10}$$

Thus, $C_{-n}(\omega_p, \Omega, k_z) = -C_n(\omega_p, \Omega, k_z)$.

2. For the reflection of magnetic field $\Omega \rightarrow -\Omega$, $\mathbf{A}_n(\omega_p, -\Omega, \mathbf{k}) = -\mathbf{A}_n(\omega_p, \Omega, \mathbf{k})$. Thus, $C_n(\omega_p, -\Omega, k_z) = -C_n(\omega_p, \Omega, k_z)$.

3. For the reflection of parallel wavenumber $k_z \rightarrow -k_z$, since the system is isotropic in $\mathbf{k}_\perp$ -plane, it is equivalent to the reflection of wavenumber $\mathbf{k} \rightarrow -\mathbf{k}$. Due to the symmetries

of the eigenvectors, we have $\mathbf{A}_n(\omega_p, \Omega, -\mathbf{k}) = -\mathbf{A}_n(\omega_p, \Omega, \mathbf{k})$ and $\mathbf{F}_n(\omega_p, \Omega, -\mathbf{k}) = \mathbf{F}_n(\omega_p, \Omega, \mathbf{k})$. Therefore, $C_n(\omega_p, \Omega, -k_z) = C_n(\omega_p, \Omega, k_z)$.

For numerical evaluation of the Chern numbers, the following alternative formula of Berry curvature is used,

$$\mathbf{F}_n = \mathrm{i} \sum_{m \neq n} \frac{\langle \psi_n | \nabla_{\mathbf{k}} H | \psi_m \rangle \times \langle \psi_m | \nabla_{\mathbf{k}} H | \psi_n \rangle}{(\omega_m - \omega_n)^2}. \quad (11)$$

To enforce integer Chern numbers, we adopt the same regularization strategy used in Ref. [10]. At large $k_{\perp}$, we regularize the plasma frequency in Eq. (9) by replacing ω_p with $\omega_p/(1 + k_{\perp}^2/k_c^2)$, where k_c is a large-enough cutoff wavenumber.

Numerical scheme. Here we introduce the numerical methods for eigenmodes calculation. When density $n(x)$ is nonuniform in the x -direction, we Fourier-transform Eq. (8) in y, z, t and but not in x . The simulation region in the x -direction is $[-2l, 2l]$ and is discretized into N grids. The interval of grids is $\Delta x = 4l/N$ and the grid points are $x_i = i\Delta x - 2l$, $i = 0, \dots, N-1$. Similar to Ref. [1], a periodic boundary condition is applied at $x = \pm 2l$. Next, we adopt the strategy of structure-preserving geometric algorithms in plasma physics to discretize $\mathbf{v}$ and $\mathbf{E}$ on integer grid points x_i and $\mathbf{B}$ on half-integer grid points $x_{i+1/2} \equiv (x_i + x_{i+1})/2$, i.e., $\mathbf{v}_i \equiv \mathbf{v}(x_i)$, $\mathbf{E}_i \equiv \mathbf{E}(x_i)$, and $\mathbf{B}_{i+1/2} \equiv \mathbf{B}(x_{i+1/2})$. Such discretization ensures centered discretization of x -derivatives and preserves the geometric relations between different components of the field. Periodic boundary condition enforces that $\mathbf{v}_N = \mathbf{v}_0$, $\mathbf{E}_N = \mathbf{E}_0$, $\mathbf{B}_{N+1/2} = \mathbf{B}_{1/2}$. Define $\omega_{p,j} \equiv \omega_p(x_j) \sim \sqrt{n(x_j)}$. Then, Eq. (8) is discretized as

$$\begin{cases} \omega v_{x,i} &= -\mathrm{i}\Omega v_{y,i} + \mathrm{i}\omega_{p,i} E_{x,i}, \\ \omega v_{y,i} &= \mathrm{i}\Omega v_{x,i} + \mathrm{i}\omega_{p,i} E_{y,i}, \\ \omega v_{z,i} &= \mathrm{i}\omega_{p,i} E_{z,i}. \end{cases} \quad (12)$$

$$\begin{cases} \omega E_{x,i} &= -\mathrm{i}\omega_{p,i} v_{x,i} + k_z \frac{B_{y,i+1/2} + B_{y,i-1/2}}{2} - k_y \frac{B_{z,i+1/2} + B_{z,i-1/2}}{2}, \\ \omega E_{y,i} &= -\mathrm{i}\omega_{p,i} v_{y,i} - k_z \frac{B_{x,i+1/2} + B_{x,i-1/2}}{2} - \mathrm{i} \frac{B_{z,i+1/2} - B_{z,i-1/2}}{\Delta x}, \\ \omega E_{z,i} &= -\mathrm{i}\omega_{p,i} v_{z,i} + k_y \frac{B_{x,i+1/2} + B_{x,i-1/2}}{2} + \mathrm{i} \frac{B_{y,i+1/2} - B_{y,i-1/2}}{\Delta x} \end{cases} \quad (13)$$

$$\begin{cases} \omega B_{x,i+1/2} &= -k_z \frac{E_{y,i+1} + E_{y,i}}{2} + k_y \frac{E_{z,i+1} + E_{z,i}}{2}, \\ \omega B_{y,i+1/2} &= k_z \frac{E_{x,i+1} + E_{x,i}}{2} + i \frac{E_{z,i+1} - E_{z,i}}{\Delta x}, \\ \omega B_{z,i+1/2} &= -k_y \frac{E_{x,i+1} + E_{x,i}}{2} - i \frac{E_{y,i+1} - E_{y,i}}{\Delta x}. \end{cases} \quad (14)$$

Equations (12)-(14) are now a standard matrix eigenvalue problem, which can be solved by standard matrix eigenvalue algorithms. We note that the method of structure-preserving geometric algorithms enables the reformulation of the eigenvalue problem of ODEs as an eigenvalue problem for Hermitian matrices.

Acknowledgments This research was supported by the U.S. Department of Energy (DE-AC02-09CH11466). We thank Jeff Parker and Fang Xie for fruitful discussions.

* yichenf@princeton.edu

- [1] P. Delplace, J. Marston, and A. Venaille, Topological origin of equatorial waves, [*Science* **358**, 1075 \(2017\)](#).
- [2] A. Souslov, B. C. Van Zuiden, D. Bartolo, and V. Vitelli, Topological sound in active-liquid metamaterials, [*Nat. Phys.* **13**, 1091 \(2017\)](#).
- [3] M. Perrot, P. Delplace, and A. Venaille, Topological transition in stratified fluids, [*Nat. Phys.* **15**, 781 \(2019\)](#).
- [4] A. Souslov, K. Dasbiswas, M. Fruchart, S. Vaikuntanathan, and V. Vitelli, Topological waves in fluids with odd viscosity, [*Phys. Rev. Lett.* **122**, 128001 \(2019\)](#).
- [5] C. Tauber, P. Delplace, and A. Venaille, A bulk-interface correspondence for equatorial waves, *J. Fluid Mech.* **868**, [10.1017/jfm.2019.233](https://doi.org/10.1017/jfm.2019.233) (2019).
- [6] B. Yang, M. Lawrence, W. Gao, Q. Guo, and S. Zhang, One-way helical electromagnetic wave propagation supported by magnetized plasma, [*Sci. Rep.* **6**, 1 \(2016\)](#).
- [7] W. Gao, B. Yang, M. Lawrence, F. Fang, B. Béri, and S. Zhang, Photonic Weyl degeneracies in magnetized plasma, [*Nat. Commun.* **7**, 1 \(2016\)](#).
- [8] J. B. Parker, J. Burby, J. Marston, and S. M. Tobias, Nontrivial topology in the continuous spectrum of a magnetized plasma, [*Phys. Rev. Research* **2**, 033425 \(2020\)](#).

- [9] J. B. Parker, Topological phase in plasma physics, [arXiv preprint arXiv:2009.06026](#) (2020).
- [10] J. B. Parker, J. Marston, S. M. Tobias, and Z. Zhu, Topological gaseous plasmon polariton in realistic plasma, [Phys. Rev. Lett. **124**, 195001 \(2020\)](#).
- [11] M. Z. Hasan and C. L. Kane, Colloquium: topological insulators, [Rev. Mod. Phys. **82**, 3045 \(2010\)](#).
- [12] B. A. Bernevig and T. L. Hughes, *Topological insulators and topological superconductors* (Princeton university press, 2013).
- [13] M. G. Silveirinha, Proof of the bulk-edge correspondence through a link between topological photonics and fluctuation-electrodynamics, [Phys. Rev. X **9**, 011037 \(2019\)](#).
- [14] M. G. Silveirinha, Chern invariants for continuous media, [Phys. Rev. B **92**, 125153 \(2015\)](#).
- [15] M. G. Silveirinha, Bulk-edge correspondence for topological photonic continua, [Phys. Rev. B **94**, 205105 \(2016\)](#).
- [16] S. A. H. Gangaraj, A. Nemilentsau, and G. W. Hanson, The effects of three-dimensional defects on one-way surface plasmon propagation for photonic topological insulators comprised of continuum media, [Sci. Rep. **6**, 30055 \(2016\)](#).
- [17] S. A. H. Gangaraj, M. G. Silveirinha, and G. W. Hanson, Berry phase, Berry connection, and Chern number for a continuum bianisotropic material from a classical electromagnetics perspective, [IEEE JMMCT **2**, 3 \(2017\)](#).
- [18] S. A. H. Gangaraj and F. Monticone, Coupled topological surface modes in gyrotropic structures: Green's function analysis, [IEEE AWPL **17**, 1993 \(2018\)](#).
- [19] S. A. H. Gangaraj, G. W. Hanson, M. G. Silveirinha, K. Shastri, M. Antezza, and F. Monticone, Unidirectional and diffractionless surface plasmon polaritons on three-dimensional nonreciprocal plasmonic platforms, [Phys. Rev. B **99**, 245414 \(2019\)](#).
- [20] S. A. H. Gangaraj and F. Monticone, Physical violations of the bulk-edge correspondence in topological electromagnetics, [Phys. Rev. Lett. **124**, 153901 \(2020\)](#).
- [21] T. H. Stix, *Waves in plasmas* (Springer Science & Business Media, 1992).
- [22] N. Armitage, E. Mele, and A. Vishwanath, Weyl and Dirac semimetals in three-dimensional solids, [Rev. Mod. Phys. **90**, 015001 \(2018\)](#).
- [23] T. Ozawa, H. M. Price, A. Amo, N. Goldman, M. Hafezi, L. Lu, M. C. Rechtsman, D. Schuster, J. Simon, O. Zilberberg, *et al.*, Topological photonics, [Rev. Mod. Phys. **91**, 015006 \(2019\)](#).
- [24] J. Wesson and D. J. Campbell, *Tokamaks*, Vol. 149 (Oxford university press, 2011).

- 394 [25] H. Qin, R. Zhang, A. S. Glasser, and J. Xiao, Kelvin-Helmholtz instability is the result of
395 parity-time symmetry breaking, [Phys. Plasmas](#) **26**, 032102 (2019).
- 396 [26] Y. Fu and H. Qin, The physics of spontaneous parity-time symmetry breaking in the Kelvin-
397 Helmholtz instability, [New J. Phys.](#) **22**, 083040 (2020).
- 398 [27] D. Jin, L. Lu, Z. Wang, C. Fang, J. D. Joannopoulos, M. Soljačić, L. Fu, and N. X. Fang,
399 Topological magnetoplasmon, [Nat. Commun.](#) **7**, 13486 (2016).

Figures

Figure 1

Please see manuscript .pdf for full caption.

Figure 2

Please see manuscript .pdf for full caption.

Figure 3

Please see manuscript .pdf for full caption.

Figure 4

Please see manuscript .pdf for full caption.

Figure 5

Please see manuscript .pdf for full caption.

Figure 6

Please see manuscript .pdf for full caption.