

Effect of capsular tension ring implantation on capsular stability after phacoemulsification in patients with weak zonules: a randomized controlled trial. CTR implantation in cataract patients with weak zonules

Shangfei Yang

Sichuan University West China Hospital <https://orcid.org/0000-0002-9306-5609>

Hui Jiang

Sichuan University West China Hospital

Kailai Nie

Sichuan University West China Hospital

Liwen Feng

Sichuan University West China Hospital

Wei Fan (✉ fanwei55@yahoo.com)

West China Hospital of Sichuan University <https://orcid.org/0000-0002-9674-114X>

Research article

Keywords: capsular tension ring, weak zonules, pars plana vitrectomy, high myopia, phacoemulsification

Posted Date: May 27th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-27972/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on January 7th, 2021. See the published version at <https://doi.org/10.1186/s12886-020-01772-8>.

Abstract

Background

Patients who had previously undergone pars plana vitrectomy (PPV) or who had severe myopia may have weaker zonules, for which a capsular tension ring (CTR) implantation is still controversial. This study is to test the effect of CTR implantation on capsular stability after phacoemulsification in those patients.

Methods

A total of 42 patients who underwent phacoemulsification and received an intraocular lens (IOL) were randomized to undergo CTR implantation or not. The primary outcomes were uncorrected distant visual acuity (UDVA), best corrected distant visual acuity (BCDVA), refractive prediction error, the area of anterior capsulorhexis, and IOL inclination angle. Follow-up visits were conducted postoperatively at 1 day, 1 week, 1 month and 3 months. Outcomes were compared between all CTR and control patients, as well as between subgroups who had previously undergone PPV or who had severe myopia.

Results

Surgery significantly improved UDVA and BCDVA to similar extents in CTR and control patients, and refraction prediction error was similar between the two groups at all follow-up times. At 3 months after surgery, the area of anterior capsulorhexis was significantly larger in CTR patients than in controls ($p = 0.0199$). These differences were also significant between the subgroups of patients with strong myopia. Vertical IOL inclination was less within CTR groups at 3 months after surgery, especially in patients with severe myopia ($P = 0.0286$). At 1 week postoperatively, the proportion of individuals whose posterior lens capsule that had completely adhered to the posterior IOL surface was significantly higher among CTR patients ($p = 0.023$). No serious surgical complications were observed.

Conclusion

CTR implantation can benefit cataract patients with weak zonules by maintaining the shape of the capsular bag, reducing capsule shrinkage and stabilizing IOL inclination.

Trail Registration:

This study was registered at the Chinese Clinical Trial Registry (ChiCTR-INR-17011217).

Background

Zonular fibers connect the ciliary body to the equator of the lens and maintain the position of the lens as well as adjust its curvature. Zonules are considered weak if zonular fibers are looser than normal, making them more susceptible to damage or rupture, or if they have already been ruptured. Zonular weakness makes the lens unstable, which can complicate intraocular procedures such as cataract surgery and increase risk of intraocular lens (IOL) dislocation [1].

Many factors predispose individuals to weak zonules, such as vitrectomy [2–6], high myopia [4, 7–9], ageing [10], pseudoexfoliation syndrome [5, 11, 12], retinal pigment degeneration [13], Marfan syndrome [14], and eye injury [4, 5]. In China, vitrectomy and strong myopia are the common causes of a weak zonule, thus this study focused on patients who underwent previous pars plana vitrectomy (PPV), or who had severe high myopia. During pars plana vitrectomy (PPV), zonular fibers near pars plana are unavoidably damaged and become weaker [8, 9, 15, 16]. People with strong myopia also have longer zonular fibers because the axial length is longer, and the wall of the eyeball is thinner. Therefore, these patients are at increased risk of loosened capsular bag, unstable anterior chamber, and lens dislocation during cataract surgery. The chance of capsular shrinkage is also higher after surgery [6, 17, 18]. Capsular shrinkage increases tension on zonular fibers, further lengthening and weakening them [1, 4, 6].

Severe zonular fiber rupture requires extracapsular cataract extraction, intracapsular cataract extraction, or pars plana lensectomy, and the IOL can be implanted in the anterior chamber or sulcus posterior chamber, or it can be sutured to the iris or sclera. However, these surgical procedures are more difficult and time-consuming, and also involve high risk of post-operative complications [19–21].

For zonular fiber relaxation or mild rupture, there is no effective treatment. The most promising approach is the use of a capsular tension ring (CTR) together with small-incision phacoemulsification and in-the-bag posterior chamber IOL implantation. The CTR maintains the shape of the capsular bag, balances the tension in zonular fibers, and decreases the risk of capsular shrinkage and IOL decentration or dislocation [22–29]. While a CTR has been used for focal zonular rupture or diffuse zonulopathy either to stretch out a capsular bag or to facilitate scleral suture fixation [7, 22–34], other studies suggested that whether the capsular bag and IOL can remain stable after CTR implantation during cataract surgery in patients with strong myopia are controversial [7, 22, 34]. In addition to these uncertainties, the effects of CTR on cataract surgery in patients with an abnormal zonule after PPV are unclear. Therefore, the current study analyzed the effects of CTR on capsular bag changes and complications after cataract surgery, in patients with weak zonules, specifically with subgroup analyses focused on patients who underwent PPV or who had strong myopia.

Methods

Patient enrollment

Consecutive cataract patients referred to the Department of Ophthalmology at West China Hospital of Sichuan University between December 2017 and December 2018 were eligible for enrollment if (1)

phacoemulsification was proposed as a treatment; (2) patients had axial length > 28 mm, or more than 3 months had passed since PPV; (3) zonules were considered weak when the average zonular length > 0.7 mm [35, 36], as measured by ultrasound biomicroscopy; and (4) patients were eligible for phacoemulsification alone or with CTR implants. Exclusion criteria included: (1) zonular issues caused by pseudoexfoliation syndrome, uveitis, retinitis pigmentosa, trauma, other intraocular surgeries than PPV, or connective tissue disorders; (2) zonular rupture > 90°; and (3) unsuitability for CTR implants. Only data on the first operated eye were included in this study, regardless of whether patients underwent operations in both eyes. Patients were randomly assigned to phacoemulsification alone or with CTR implants based on permuted-block randomization by an optometrist who was not involved in patient selection or surgery. A computer program for randomization that provided random permuted blocks was used by the optometrist. The sample size was calculated with $\alpha = 0.05$ (two-sided), $\beta = 0.2$ and determined to be approximately 21 patients in each group. Assuming an expected dropout rate of 10%, 23 patients in each group were needed.

Patients were provided thorough information about the surgery and possible complications, and patients were not blinded to treatments.

Preoperative eye examination

Uncorrected distant visual acuity (UDVA) and best corrected distant visual acuity (BCDVA) of all patients were recorded using a Snellen chart. Visual acuity values were converted to logarithms of the minimum angle of resolution (logMAR). Refractive prediction error was determined by measuring the difference between the preoperative refractive target and actual refraction (D, diopters) after surgery. Eye pressure was determined by non-contact tonometry before pupillary dilation. Axial length of the eye was measured by optical biometry (IOLMaster 500, Zeiss, Oberkochen, Germany). Zonular weakness was considered if there was an abnormal zonular length which was measured using an ultrasound biomicroscope [35, 36]. Zonular length was reported as an average of 4 measurements at 3, 6, 9, 12 o'clock. Measurements were made between the mid-point of the ciliary process and the end of zonular fibers at the lens equator (Fig. 1). All measurements were made by the same investigator.

Surgical procedures

All surgical procedures were performed by an experienced surgeon (Wei Fan). All enrolled patients received phacoemulsification and IOL implantation (Akreos MI60, Bausch & Lomb, Rochester, NY, USA). A 2.0-mm corneal incision was made, a 5.0-5.5 mm CCC was performed and the IOL was implanted using a Stellaris System (Bausch & Lomb). Patients in the intervention group underwent CTR implantation (ACPi-11, Bausch & Lomb) before IOL implantation.

Postoperative eye examination

Postoperative eye examination was performed by the same ophthalmologist and optometrist in each patient at 1 day, 1 week, 1 month and 3 months after the surgery. Examination parameters were the same as in the preoperative eye examination. Postoperative complications, if any, were recorded. Micrographs

of anterior continuous curvilinear capsulorhexis and IOL optics were taken using a slit-lamp camera, and areas of anterior capsulorhexis were analyzed by Image J (National Institutes of Health, Bethesda, MD, USA). The position of IOL and the attachment ratio between posterior lens capsule and posterior surface of IOL were evaluated by anterior segment optical coherence tomography (Carl Zeiss Meditec, Jena, Germany). IOL inclination angle, defined as the angle between the posterior surface of the iris and the anterior surface of the IOL, was measured vertically and horizontally [37]. The angle was measured from micrographs using Adobe PDF Editor (San Jose, CA, USA).

Statistical analysis

Data were analyzed using SAS 9.4 (IBM, Armonk, NY, USA). Continuous results are presented as mean \pm standard deviation, and categorical results are presented as number (percentage). Inter-group and intragroup differences were assessed for significance using repeated measurement variance analysis in the case of repeated-measurement continuous data. Inter-group differences were assessed for significance using the chi-squared test in the case of categorical data. All tests were 2-sided. Differences were considered significant when $p < 0.05$.

Results

Patient characteristics

A total of 42 patients (21 men) were enrolled in the study and randomized into a CTR group [T(total)-CTR, $n = 22$, 12 men] and control group (T-CON, $n = 20$, 9 men). Subgroup analyses within these two groups were performed based on whether patients suffered zonular problems because of PPV (P-CTR vs. P-CON) or myopia (M-CTR vs. M-CON). Patients showed the following primary vitreo-retinal diseases: retinal detachment, vitreous hemorrhage, macular pucker, and macular hole. Average length of zonules was (1.08 ± 0.28) mm in the T-CTR group and (1.03 ± 0.19) mm in the T-CON group, and the two values were not statistically different (Table 1).

Table 1. Patient baseline characteristics.

Groups	N	Sex		Age(y)	AL(mm)	Length of zonules
		Male	Female			
T-CTR	22	12(54.5%)	10(45.5%)	59.35±12.15	28.32±3.65	1.08±0.28
T-CON	20	9(45.0%)	11(55.0%)	61.35±9.35	27.91±3.11	1.03±0.19
<i>t</i>				-0.526	0.457	0.628
<i>P</i>			0.379	0.605	0.653	0.534
P-CTR	11	6(54.5%)	5(45.5%)	59.36±10.99	25.36±1.93	1.01±0.21
P-CON	11	4(36.4%)	7(63.4%)	59.36±10.51	26.05±2.67	1.00±0.21
<i>t</i>				0.000	-0.694	0.152
<i>P</i>			0.335	1.000	0.503	0.881
M-CTR	11	6(54.5%)	5(45.5%)	60.27±11.16	30.76±2.36	1.13±0.34
M-CON	9	5(55.6%)	4(44.4%)	61.77±9.57	30.19±1.86	1.06±0.18
<i>t</i>				-0.916	0.560	0.413
<i>P</i>			0.658	0.387	0.591	0.684

AL, axial length; T, Total groups, P, post-PPV; M, myopia; CTR, capsular tension ring groups; CON, control groups.

Vision acuity

UDVA did not differ significantly between CTR and control patients overall (Fig. 2a) or in the subgroups with PPV (Fig. 2b) or strong myopia (Fig. 2c) at any of the time points examined. Similar results were obtained for BCDVA (Fig. 2d-f). Nevertheless, UDVA and BCDVA tended to be better in the CTR group than the control group at all time points, and visual acuity was more stable in the CTR group. In both CTR and control patients, UDVA and BCDVA were significantly better after the operation than before.

Refractive prediction error (RPE)

Refractive prediction error was not significantly different between CTR and control patients overall (Fig. 3a), or in the subgroup with PPV (Fig. 3b) or strong myopia (Fig. 3c) at any of the time points examined. Data from two patients were discarded because silicone oil was present in the vitreous body.

Area of continuous curvilinear capsulorhexis (A_{CCC})

A_{CCC} was significantly larger in total CTR patients than control patients at 3 months ($p = 0.0199$), but not at 1 week or 1 month after surgery (Table 2). Among CTR patients, A_{CCC} was lower at 3 months than at 1 week but not 1 month after surgery. Among control patients, A_{CCC} was lower at 3 months than at 1 week or 1 month after surgery. These findings suggest earlier stabilization of A_{CCC} in the CTR group.

Further detailed analysis was conducted in subgroups with PPV or myopia. Among CTR and control patients with PPV, A_{CCC} was similar between the two subgroups at 1 week, 1 month, and 3 months after

surgery. Among CTR patients with PPV, A_{CCC} was lower at 3 months than at 1 week and 1 month after surgery, while among control patients with PPV, A_{CCC} was lower at 3 months than at 1 week after surgery.

Among CTR and control patients with myopia, while A_{CCC} showed a trend of greater decrease in the control subgroup than in the CTR subgroup from 1 week ($P = 0.0709$) to 1 month ($P = 0.0529$) postoperatively, it was significantly smaller in the control subgroup than in the CTR subgroup at 3 months ($P = 0.0113$) after surgery. Within the CTR subgroup with myopia, A_{CCC} was smaller at 1 and 3 months than at 1 week after surgery, but from 1 month on, A_{CCC} was stable. However, within the control subgroup with myopia, A_{CCC} became smaller with time and was not stable until 3 months after surgery. Moreover, it is noteworthy that when comparing mean A_{CCC} at 1 day to 1 week after surgery in 3 random cases from the CTR subgroup with myopia (23.55 mm^2) and 3 random cases from the control subgroup with myopia (23.05 mm^2), A_{CCC} did not seem to start to shrink in the CTR subgroup (23.31 mm^2) until 1 week after surgery, whereas it showed contraction in the control subgroup (20.55 mm^2), suggesting an early capsular contraction within a week after surgery in patients with high myopia.

Table 2. Area of continuous curvilinear capsulorhexis (A_{CCC} , mm^2).

Groups	N	1d	1wk	1m	3m
T-CTR	22		22.56±3.61	21.96±3.38	21.53±3.47 ^b
T-CON	20		20.70±2.45	20.04 ±2.83	18.76±3.93 ^{bc}
<i>t</i>			1.94	1.99	2.43
<i>P</i>			0.0599	0.0532	0.0199
Groups(F/P)			F=4.97, P=0.0315		
Time(F/P)			F=16.05, P<0.0001		
Group*Time			F=1.85, P=0.1642		
P-CTR	11		21.81±3.58	21.43±3.27	20.58 ±3.09 ^{bc}
P-CON	11		20.82 ±2.49	20.64±2.42	20.22±1.98 ^b
<i>t</i>			0.75	0.65	0.32
<i>P</i>			0.4612	0.5241	0.7494
Groups(F/P)			F=0.36, P=0.5557		
Time(F/P)			F=8.78, P=0.0007		
Group*Time			F=1.03, P=0.3665		
M-CTR	11	23.55±0.22*	23.31±3.64	22.49±3.57 ^b	22.48±3.71 ^b
M-CON	9	23.05±0.43*	20.55±2.56	19.30±3.25 ^b	16.97±5.03 ^{bc}
<i>t</i>		1.325	1.92	2.07	2.82
<i>P</i>		0.316	0.0709	0.0529	0.0113
Groups(F/P)			F=5.86, P=0.0263		
Time(F/P)			F=12.44, P<0.0001		
Group*Time			F=5.56, P=0.0079		

*.N=3/group. ^bp<0.05 vs. 1 week after surgery within the same group; ^cp<0.05 vs. 1 month after surgery within the same group.

Ratio of A_{CCC} / area of IOL optics (A_{IOL})

A_{CCC} was further analyzed using the A_{CCC}/A_{IOL} ratio, because IOLs of two diameters were used (6.0 and 6.2 mm). When IOL diopter was $\leq +15.0$ D, CCC was relatively large (5.5-6.0 mm). The A_{CCC}/A_{IOL} ratio showed the same change as A_{CCC} : the ratio was higher in CTR patients than controls at 3 months after surgery, both in the total group of patients ($P = 0.0172$) as well as in the subgroup with myopia ($P = 0.0124$) (Table 3).

Table 3. Ratio of A_{CCC} / A_{IOL} .

Groups	N	1wk	1m	3m
T-CTR	22	0.77 ±0.13	0.75 ±0.12	0.74±0.12 ^b
T-CON	20	0.70±0.09	0.68 ±0.10	0.64 ±0.14 ^{bc}
<i>t</i>		1.99	2.01	2.49
<i>P</i>		0.0529	0.0524	0.0172
Groups(F/P)		F=5.22, P=0.0277		
Time(F/P)		F=16.32, P<0.0001		
Group*Time		F=1.78, P=0.1761		
P-CTR	11	0.77 ±0.13	0.75±0.12	0.72 ±0.11 ^{bc}
P-CON	11	0.72 ±0.10	0.71 ±0.10	0.70 ±0.08 ^b
<i>t</i>		1.03	0.94	0.68
<i>P</i>		0.3144	0.3582	0.5066
Groups(F/P)		F=0.83, P=0.3743		
Time(F/P)		F=8.6, P=0.0008		
Group*Time		F=1.06, P=0.3557		
M-CTR	11	0.78±0.13	0.75±0.12	0.75±0.13
M-CON	9	0.69±0.08	0.65±0.10	0.57±0.16 ^{bc}
<i>t</i>		1.79	2.03	2.78
<i>P</i>		0.0908	0.0577	0.0124
Groups(F/P)		F=5.51, P=0.0306		
Time(F/P)		F=12.69, P<0.0001		
Group*Time		F=5.68, P=0.0072		

^bp<0.05 vs. 1 week after surgery within the same group; ^cp<0.05 vs. 1 month after surgery within the same group.

Attachment ratio between posterior lens capsule and IOL surface

Attachment ratio between posterior lens capsule and IOL surface was higher in CTR patients than in control patients (Table 4). This was only true among patients with PPV. The ratio was higher in the P-CTR group than in the P-CON group at 1 week and 1 month after surgery. The ratio was not significantly different between the M-CTR and M-CON groups at 1 week, 1 month or 3 months after surgery.

Table 4. Attachment ratio between posterior lens capsule and IOL surface

Groups	N	1wk	1m	3m
T-CTR	22	54.5% ¹²	86.4% ¹⁹	90.9% ²⁰
T-CON	20	20.0% ⁴	60.0% ¹²	75.0% ¹⁵
<i>X</i> ²		5.301	3.767	1.909
<i>P</i>		0.023	0.055	0.167
P-CTR	11	72.7% ⁸	100.0% ¹¹	100.0% ¹¹
P-CON	11	27.3% ³	63.6% ⁷	81.8% ⁹
<i>X</i> ²		4.545	4.889	2.200
<i>P</i>		0.043	0.045	0.238
M-CTR	11	36.4% ⁴	72.7% ⁸	81.8% ⁹
M-CON	9	11.1% ¹	55.6% ⁵	66.7% ⁶
<i>X</i> ²		1.684	0.642	0.606
<i>P</i>		0.221	0.370	0.396

IOL inclination angle

Horizontal and vertical IOL inclination did not differ significantly between CTR and control patients in the total sample or in subgroups with PPV or myopia at 1 week, 1 month or 3 months after surgery. Within the total group and within each subgroup, horizontal IOL inclination angle did not vary significantly across the time points from 1 week to 3 months after surgery. Similarly, vertical IOL inclination did not differ significantly across CTR subgroups (total group and subgroup with PPV or myopia) at any of the postoperative time points. Within the total group of control patients, vertical IOL inclination was significantly higher at 3 months than at 1 week after surgery ($0.73 \pm 0.46^\circ$ vs. $1.84 \pm 2.15^\circ$; $P = 0.0224$; Fig. 4a). Similar results were observed in the subgroup of control patients with myopia ($0.74 \pm 0.59^\circ$ vs. $2.82 \pm 2.59^\circ$; $P = 0.0286$; Fig. 4b).

Depth of central anterior chamber

Depth of the central anterior chamber did not differ significantly between CTR and control patients in the total sample or in subgroups with PPV or myopia at 1 week, 1 month or 3 months after surgery.

Post-operative complications

All the patients in this study showed well-controlled postoperative inflammation. One CTR patient experienced capsular block syndrome 1 day after surgery. Two CTR patients and one control patient showed transient elevated intraocular pressure (>21 mmHg). By 3 months after surgery, one CTR patient (4.55%, in the PPV subgroup) and three control patients (15.00%, in the myopia subgroup) showed significant capsular contraction syndrome. No other severe complications were occurred.

Discussion

CTR is widely used in clinical treatment of cataracts complicated with lens dislocation. CTR can effectively balance the tension of zonular fibers, uniformly distribute the tension of the capsular bag, maintain the shape of the capsular bag, reduce loss of the vitreous body, and increase attachment between posterior capsule and IOL surface. These effects of CTR increase IOL stability after cataract extraction involving IOL implantation and reduce the occurrence of posterior capsule opacification and IOL dislocation [7, 22–34]. Whether and how CTR implantation can also benefit patients with weakened zonular fibers but no lens dislocation is unclear. The present randomized controlled trial suggests that CTR in such patients can reduce the incidence of capsular shrinkage, maintain capsular bag stability, make IOL inclination more manageable and stable, as well as increase the rate of complete attachment between the posterior capsule of the lens and posterior surface of the IOL.

Vanags et al [27] demonstrated that CTR implantation during cataract surgery with lens dislocation can lead to better capsular shape, IOL position and lower risk of capsular shrinkage at 1 week to 2 years after surgery than without CTR implantation. Studies in patients [37, 38] and animals [39] showed similar results even in cases without weak zonules. However, there are other studies showing no improvement in the incidence of IOL dislocation by CTR implantation at 3 months after cataract surgery [29, 30]. In the current study, CTR implantation showed the ability to prevent capsular shrinkage starting from 1 week postoperatively in patients with high myopia. At 3 months after surgery, this effect was even more significant, suggesting that CTR is effective in delaying shrinkage and maintaining capsular bag stability in the longer term, especially in patients with severe myopia.

In contrast to these results in patients with myopia, our analysis of patients with PPV showed no clear beneficial effects of CTR, especially on the anterior capsulorhexis shrinkage, likely due to the adverse effects of silicone oil or gas bubbles injected after vitrectomy. Thus, the benefits of CTR implantation may not compensate the adverse effects of silicone oil or gas bubbles, which reduce elasticity and alter capsule structure as well as its response to capsulorhexis.

Capsular contraction syndrome is a more serious complication following capsular shrinkage [27, 28], which is usually accompanied by posterior capsule opacification and even causes IOL dislocation. Timely intervention is needed in order to prevent further visual impairment. Incidence of significant capsular contraction syndrome was lower in among our CTR patients (4.55%, 1 case in PPV subgroup, 0 in myopia subgroup) than among control patients (15.00%, 0 in PPV subgroup, 3 cases in myopia subgroup) at 3 months after surgery, which made it mandatory for these patients to have Nd:YAG treatment. This finding suggests that CTR implantation can reduce the incidence of, but not completely eliminate, capsular shrinkage and contraction, especially in patients with severe myopia [39].

We also found that the rate of complete attachment between the posterior capsule of the lens and posterior surface of the IOL was higher in the CTR group than the control group after only 1 week and even higher after 3 months. This may be because the CTR can mechanically compress the capsular bag closer to the IOL surface [26, 34, 40, 41]. However, this effect of CTR was not obvious in patients with

strong myopia, perhaps because such eyes are axially longer, and the capsular bag is larger. We speculate that the CTR used in this study, with a diameter of 11 mm, may not be large enough to completely open the capsular bag in patients with strong myopia.

Previous animal and human studies demonstrated that IOL eccentricity and inclination were small and remained stable within 2 years after CTR implantation in cataract surgery [37, 39]. We found that the vertical inclination angle gradually decreased in CTR patients but increased in control patients. Horizontal IOL inclination remained stable within 3 months after surgery in CTR and control patients. These results support the idea that IOL inclination after CTR implantation is manageable and stable [22, 42].

Whether CTR implantation during cataract surgery changes postoperative refractive prediction error is unclear. A retrospective study on CTR implantation in 25 patients with abnormal zonules found that the position of the posterior chamber IOL exceeded the predicted value by + 0.5 to + 2.0 D [31]. A randomized controlled trial in 52 cataract patients without other complications showed hyperopia drift after CTR implantation [29]. Those authors recommended reducing preoperative refractive predictions by 0.5 D. The present study suggests, in contrast, that preoperative refractive predictions should not be adjusted for patients undergoing CTR implantation: we did not observe substantial differences in the values for CTR and control patients, consistent with other studies [33, 43–45]. It is noteworthy that both UDVA and BCDVA were better in CTR patients than in controls.

There are a few noteworthy limitations in this study. First, CTR with 11 mm diameter was used in all patients. Individualized optimization of CTR diameter is more desirable especially in case of high myopia and may benefit cataract patients more. Second, the IOL profile of some patients was not fully visible in anterior segment optical coherence tomography because of limited pupil dilation. Only IOL tilt was measured in our study, making IOL inclination angle measurements less conclusive. Third, patients were followed up for only 3 months, based on studies suggesting that many beneficial and adverse effects of CTR implantation occur within that time. Fourth, we did not collect complete data on the area of anterior continuous curvilinear capsulorhexis at 1 day after surgery, which prevented us from studying early changes in anterior continuous curvilinear capsulorhexis, such as within 1 week.

Despite these limitations, our study provides evidence that the benefits of CTR implantation outweigh its disadvantages in cataract patients with mild abnormal zonules such as those with high myopia or previous PPV.

Conclusion

CTR is effective in delaying shrinkage and maintaining capsular bag stability in patients with severe myopia in the longer term, but this effect is not obvious in patients after PPV. CTR implantation can promote the adhesion between the posterior capsule and the posterior surface of IOL in patients with weak zonules, especially those after PPV. In a word, CTR implantation can benefit cataract patients with weak zonules by maintaining the shape of the capsular bag, reducing capsule shrinkage and stabilizing IOL inclination.

Abbreviations

CTR: Capsular tension ring; IOL: Intraocular lens; PPV: pars plana vitrectomy; UDVA: uncorrected distant visual acuity; BCDVA: best corrected distant visual acuity; logMAR: logarithms of the minimum angle of resolution; RPE: Refractive prediction error; A_{CCC} : Area of continuous curvilinear capsulorhexis; A_{IOL} : Area of IOL optics.

Declarations

Availability of data and materials

All data generated or analyzed during this study are included in this published article. Any additional information can be requested from the corresponding author.

Authors' contributions

SFY: Conceptualization, Methodology, Investigation, Resources, Data curation, Writing Original draft preparation, Project administration;

WF: Conceptualization, Methodology, Writing-Reviewing and Editing, Supervision, Project administration;

HJ: Resources, Investigation, Data curation;

KN: Investigation, Data curation, Software;

WF: Investigation, Resources.

All authors read and approved the final manuscript.

Ethics approval and consent to participate

These procedures were approved by the Ethics Committee of West China Hospital of Sichuan University. Written informed consent was obtained from all participants before the surgery. This study was registered at the Chinese Clinical Trial Registry (ChiCTR-INR-17011217).

Consent for publication

Was taken in writing from all the patients before surgery. All the patients in the study consented for their individual data to be published.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by the National Natural Science Foundation of China (No. 81670869). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

1. Saori Y, Shigeo Y, Yukari YY, Tadahiko K, Hiroko BM, Al-Ghoul KJ. Objective classification of zonular weakness based on lens movement at the start of capsulorhexis[J]. *Plos One*. 2017;12(4):e0176169.
2. Koike T, Baba T, Nizawa T, Oshitari T, Yamamoto S. Characteristics of patients with spontaneous dislocation of in-the-bag intraocular lens after pars plana vitrectomy[J]. *Jpn J Ophthalmol*. 2017;61(3):1–4.
3. Hocaoglu M, Karacorlu M, Sayman MI, et al. Incidence and factors associated with complications of sutured and sutureless cataract surgery following pars plana vitrectomy at a tertiary referral centre in Turkey[J]. *Br J Ophthalmol*. 2016;100(9):1206–10.
4. Fernández-Buenaga R, Alio JL, Pérez-Ardoy AL, et al. Late in-the-bag intraocular lens dislocation requiring explantation: risk factors and outcomes[J]. *Eye (Lond)*. 2013;27(7):795–801.
5. Davis D, Brubaker J, Espandar L, et al. Late in-the-bag spontaneous intraocular lens dislocation: evaluation of 86 consecutive cases[J]. *Ophthalmology*. 2009;116(4):664–70.
6. Yasuda A, Ohkoshi K, Orihara Y, Kusano Y, Sakuma A, Yamaguchi T. Spontaneous luxation of encapsulated intraocular lens onto the retina after a triple procedure of vitrectomy, phacoemulsification, and intraocular lens implantation[J]. *Am J Ophthalmol*. 2000;130(6):836–7.
7. Zhang K, Zhu X, Chen M, et al. Elevated Transforming Growth Factor- β 2 in the Aqueous Humor: A Possible Explanation for High Rate of Capsular Contraction Syndrome in High Myopia[J]. *Journal of Ophthalmology*. 2016;2016(7):1–6.
8. Lam JK, Chan TC, Ng AL, et al. Outcomes of cataract operations in extreme high axial myopia[J]. *Graefes Arch Clin Exp Ophthalmol*. 2016;254(9):1–7.
9. Chong EW, Mehta JS. High myopia and cataract surgery[J]. *Curr Opin Ophthalmol*. 2016;27(1):45–50.
10. Bron AJ, Vrensen GF, Koretz J, Maraini G, Harding JJ. The Ageing Lens[J] *Ophthalmologica*. 2000;214(1):86–104.
11. Kristianslund O, Råen M, Østern AE, Drolsum L. Late In-the-Bag Intraocular Lens Dislocation: A Randomized Clinical Trial Comparing Lens Repositioning and Lens Exchange[J]. *Ophthalmology*. 2017;124(2):151–9.
12. Vazquez-Ferreiro P, Carrera-Hueso FJ, Fikri-Benbrahim N, Barreiro-Rodriguez L, Diaz-Rey M, Ramón Barrios MA. Intraocular lens dislocation in pseudoexfoliation: a systematic review and meta-analysis[J]. *Acta Ophthalmol*. 2017;95(3):e164–9.
13. Najjar DM, Igbre AO, Tsai FF. Late capsular bag contraction and intraocular lens subluxation in retinitis pigmentosa: a case report[J]. *J Med Case Rep*. 2011;5(1):65.

14. Ladewig MS, Robinson PN, Neumann LM, Holz FG, Foerster MH. Ocular manifestations and surgical results in patients with Marfan syndrome[J]. *Der Ophthalmologe*. 2006;103(9):777–82.
15. Matsumoto M, Yamada K, Uematsu M, et al. Spontaneous dislocation of in-the-bag intraocular lens primarily in cases with prior vitrectomy[J]. *Eur J Ophthalmol*. 2012;22(3):363–7.
16. Do DV, Gichuhi S, Vedula SS, Hawkins BS. Surgery for post-vitrectomy cataract [J]. *The Cochrane database of systematic reviews*, 2018;10(1), No.:CD006366.
17. Hsuan JD1, Brown NA, Bron AJ, Patel CK, Rosen PH. Posterior subcapsular and nuclear cataract after vitrectomy. *J Cataract Refract Surg*. 2001;27(3):437–44.
18. Grusha YO, Masket S, Miller KM. Phacoemulsification and lens implantation after pars plana vitrectomy. [J]. *Ophthalmology*. 1998;105(2):287–94.
19. Blecher MH, Kirk MR. Surgical strategies for the management of zonular compromise[J]. *Curr Opin Ophthalmol*. 2008;19(1):31–5.
20. Yoshida K, Kiryu J, Kita M, Ogura Y. Phacoemulsification of Dislocated Lens and Suture Fixation of Intraocular Lens Using a Perfluorocarbon Liquid[J]. *Jpn J Ophthalmol*. 1999;42(6):471–5.
21. Hayashi K, Hayashi H, Nakao F, Hayashi F. Intraocular lens tilt and decentration, anterior chamber depth, and refractive error after trans-scleral suture fixation surgery[J]. *Ophthalmology*. 1999;106(5):878–82.
22. Zhao Y, Li J, Yang K, Li X, Zhu S. Combined Special Capsular Tension Ring and Toric IOL Implantation for Management of Astigmatism and High Axial Myopia with Cataracts[J]. *Semin Ophthalmol*. 2018;33(3):389–94.
23. Wilkie DA, Colitz CMH. Surgery of the lens[J]. *Veterinary Ophthalmology*, 2013:1234–1286.
24. Trikha S, Agrawal S, Saffari SE, et al. Visual outcomes in patients with zonular dialysis following cataract surgery[J]. *Eye*. 2016;30(10):1331–5.
25. Rai G, Sahai A, Kumar PR. Outcome of Capsular Tension Ring (CTR) Implant in Complicated Cataracts[J]. *Journal of Clinical Diagnostic Research Jcdr*. 2015;9(12):NC05–07.
26. Menapace R, Findl O, Georgopoulos M, Rainer G, Vass C, Schmetterer K. The capsular tension ring: designs, applications, and techniques[J]. *J Cataract Refract Surg*. 2000;26(6):898–912.
27. Vanags J, Erts R, Laganovska G. Anterior capsulorhexis opening reduction after cataract surgery with subluxated lenses[J]. *Medicina*. 2017;53(5):310–5.
28. Morenomontañés J, Sáncheztocino H, Rodríguezconde R. Complete anterior capsule contraction after phacoemulsification with acrylic intraocular lens and endocapsular ring implantation[J]. *J Cataract Refract Surg*. 2002;28(4):717–9.
29. Jakobsson G, Zetterberg M, Lundström M, Stenevi U, Grenmark R, Sundelin K. Late dislocation of in-the-bag and out-of-the bag intraocular lenses: Ocular and surgical characteristics and time to lens repositioning[J]. *J Cataract Refract Surg*. 2010;36(10):1637–44.
30. Ramón Lorente, Rojas VD, Parga PVD, et al. Management of late spontaneous in-the-bag intraocular lens dislocation: Retrospective analysis of 45 cases[J]. *J Cataract Refract Surg*. 2010;36(8):1270–

82.

31. Baranwal VK, Kumar S, Mishra A, et al. A study to evaluate whether CTR increases refractive unpredictability between predicted and actual IOL position[J]. *Medical Journal Armed Forces India*. 2014;70(1):36–8.
32. Park HJ, Lee H, Kim do W, Kim EK, Seo KY, Kim TI. Effect of Co-Implantation of a Capsular Tension Ring on Clinical Outcomes after Cataract Surgery with Monofocal Intraocular Lens Implantation[J]. *Yonsei Med J*. 2016;57(5):1236–42.
33. Schild AM, Rosentreter A, Hellmich M, Lappas A, Dinslage S, Dietlein TS. Effect of a capsular tension ring on refractive outcomes in eyes with high myopia[J]. *J Cataract Refract Surg*. 2010;36(12):2087–93.
34. Halili I, Mutlu FM, Erdurman FC, et al. Influence of capsular tension ring on posterior capsule opacification in myopic eyes[J]. *Indian J Ophthalmol*. 2014;62(3):311–5.
35. Yang SF, Huang YZ, Chen H, Jiang H, Fan W. Influence of pars plana vitrectomy on the length of lens zonules and anterior chamber depth [J]. *Int Eye Sci*. 2018;18(07):87–91.
36. Miao N, Tang XL, Huang YZ, Li RL, Fan W. Study on preoperative observation of cataract patients' zonules by ultrasound biomicroscopy[J]. *West China Medical Journal*. 2016;31(12):41–5.
37. Lee DH, Shin SC, Joo CK. Effect of a capsular tension ring on intraocular lens decentration and tilting after cataract surgery[J]. *J Cataract Refract Surg*. 2002;28(5):843–6.
38. Nishi O, Nishi K, Menapace R, Akura J. Capsular bending ring to prevent posterior capsule opacification: 2 years follow-up[J]. *Cataract Refract Surg*. 2001;27(9):1359–65.
39. Lee DH, Lee HY, Lee KH, Chung KH, Joo CK. Effect of a capsular tension ring on the shape of the capsular bag and opening and the intraocular lens[J]. *J Cataract Refract Surg*. 2001;27(3):452–6.
40. Findl O, Drexler W, Menapace R, et al. Accurate determination of effective lens position and lens-capsule distance with 4 intraocular lenses[J]. *J Cataract Refract Surg*. 1998;24(8):1094–8.
41. Nishi O, Nishi K. Intraocular lens encapsulation by shrinkage of the capsulorhexis opening[J]. *J Cataract Refract Surg*. 1993;19(4):544–5.
42. Mastropasqua R, Toto L, Vecchiarino L, Falconio G, Nicola MD, Mastropasqua A. Multifocal IOL implant with or without capsular tension ring: study of wavefront error and visual performance[J]. *Eur J Ophthalmol*. 2013;23(4):510–7.
43. Boomer JA, Jackson DW. Effect of the morcher capsular tension ring on refractive outcome[J]. *J Cataract Refract Surg*. 2006;32(7):1180–3.
44. Sun R, Gimbel HV. In vitro evaluation of the efficacy of the capsular tension ring for managing zonular dialysis in cataract surgery[J]. *Ophthalmic Surgery Lasers*. 1998;29(6):502.
45. Alio J, Elkady BD, Bernabeu G. Microincision multifocal intraocular lens with and without a capsular tension ring: optical quality and clinical outcomes[J]. *J Cataract Refract Surg*. 2008;34(9):1468–75.

Figures



Figure 1

Micrographs showing zonular length by ultrasound biomicroscopy. Arrows indicate zonule.




Figure 2

Visual acuity comparison at baseline, 1 week, 1 month, and 3 months after surgery. Comparison of uncorrected distant visual acuity between (A) all CTR and control patients, (B) the subgroups of CTR and control patients with PPV, and (C) the subgroups of CTR and control patients with myopia. Comparison of best corrected distant visual acuity between (D) all CTR and control patients, (E) the subgroups of CTR and control patients with PPV, and (F) the subgroups of CTR and control patients with myopia. a- p<0.05 vs. before surgery within the same group, b- p<0.05 vs. 1 week after surgery within the same group.




Figure 3

Comparison of refractive prediction error (RPE) at 1 week, 1 month, and 3 months after surgery between (A) all CTR and control patients, (B) the subgroups of CTR and control patients with PPV, and (C) the subgroups of CTR and control patients with myopia.




Figure 4

Comparison of intraocular lens vertical inclination angle between (A) all CTR and control patients, and (B) the subgroups of CTR and control patients with myopia. b- $p < 0.05$ vs. 1 week after surgery within the same group.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [CONSORT2010Checklist.doc](#)
- [CONSORT2010FlowDiagram.doc](#)