

# Imported Mansonella Perstans Infection in Spain.

**Sabino Puente**

ISCI

**Mar Lago**

Instituto de Salud Carlos III

**Mercedes Subirats**

Instituto de Salud Carlos III

**Ismael Sanz-Esteban**

Universidad Europea de Madrid

**Marta Arsuaga**

Instituto de Salud Carlos III

**Belen Vicente**

Universidad de Salamanca

**Montserrat Alonso-Sardon**

Universidad de Salamanca

**Moncef Belhassen-Garcia** (✉ [belhassen@usal.es](mailto:belhassen@usal.es))

CAUSA <https://orcid.org/0000-0002-2230-1256>

**Antonio Muro**

Universidad de Salamanca

---

## Research Article

**Keywords:** Mansonellosis, Mansonella perstans, Clinical study, Immigrant, Imported diseases, Spain

**Posted Date:** May 14th, 2020

**DOI:** <https://doi.org/10.21203/rs.3.rs-28069/v1>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

---

**Version of Record:** A version of this preprint was published on July 23rd, 2020. See the published version at <https://doi.org/10.1186/s40249-020-00729-9>.

# Abstract

**Background:** *Mansonella perstans* infection can be considered one of the most neglected tropical infectious diseases. Very few studies have reported on the clinical picture caused by infection with this nematode.

**Methods:** The present study evaluated a large cohort of migrants who have been diagnosed, examined and treated for imported *M. perstans* infection at a Spanish reference center (Hospital Carlos III Tropical Medicine Unit, Madrid).

**Results:** Five hundred three cases of migrants from tropical and subtropical areas with *M. perstans* infection were identified. Two hundred sixty-four patients were female (52.5%). The mean age ( $\pm$ SD) was  $44.6 \pm 18.2$  years (range, 16–93 years). The mean time ( $\pm$ SD) between the arrival in Spain and the first consultation was  $8.6 \pm 18.0$  months. The origin of the patients was Equatorial Guinea (97.6%). Regarding the clinical picture, 257 patients were asymptomatic (54.7%) and 228 were symptomatic (45.3%); 190 patients had pruritus (37.8%), 50 (9.9%) had arthralgia, 18 patients had Calabar-like swelling (3.6%), and 15 (3%) had abdominal pain. Four hundred forty-two (87.9%) migrants had hyper-IgE, and 340 (67.6%) had eosinophilia. One hundred ninety-five patients had coinfections with other filarial nematodes (38.8%), and 308 migrants had only *M. perstans* infection (61.2%). Four hundred thirty-seven cases (86.9%) had been treated with anti-filarial drugs: 292 cases were treated with one anti-filarial drug, and 145 cases were treated with combined anti-filarial therapy. No significant differences were found in the healing outcomes ( $p > 0.05$ ). Additionally, 20 (4%) cases received steroids and 38 (7.6%) cases received antihistamines. The combined anthelmintics and steroid treatment presented significantly better results ( $p = 0.042$ ); however, no significant results were noted for the combined anthelmintic and antihistamine treatment ( $p = 0.191$ ). No statistical association was found between the percentage of side effects to treatment and cases healed ( $p = 0.094$ ).

**Conclusions:** A long series of *M. perstans* infections is presented in sub-Saharan immigrants whose data indicate that it should be included in the differential diagnosis in patients with pruritus or analytical alterations such as eosinophilia or hyper-IgE presentation, and they also have a high number of coinfections with other microorganisms whose treatment needs to be protocolized.

## Introduction

*Mansonella perstans* is a human filarial parasite endemic in 33 countries in western, eastern, and central Africa, as well as in regions of South and Central America[1]. Flies of the genus *Culicoides* transmit infective larvae to humans. The adult parasites are thought to live in serous body cavities, and the female parasites release microfilariae into the blood[2]. The diagnosis of *M. perstans* infection is usually by detection and identification of the microfilariae that circulate in the blood. Because the microfilariae are present in the peripheral blood in almost equal concentrations during the day and night [3], blood samples for the diagnosis can be obtained at any time. The prevalence and intensity of microfilaremia increase gradually with age. Conflicting reports exist concerning the presence of *Wolbachia* endosymbionts in the microfilariae of *M. perstans*[4-7]. Few and old studies have reported on the clinical picture caused by infection with *M. perstans* nematodes because the parasite is widespread in remote areas[1,8]. Usually, infected people have other parasitic infections that could contribute to clinical manifestations [1,8]. Clinical manifestations seem to be related to adult parasites than to the microfilariae, and the symptoms are probably related to the migration of the worms, including transient subcutaneous swellings (similar to the Calabar swellings caused by *Loa loa*), pruritus, rash, urticaria, arthralgia, abdominal pain, eosinophilia, fatigue, pericarditis, pleuritis and inflammatory granulomatous nodules surrounding dead adult worms[1]. Currently, no standard treatment exists for mansonellosis and its handling is still debatable. Therefore, many drugs have been used alone or combined, such as diethylcarbamazine, ivermectin, mebendazole, levamisole, albendazole, and thiabendazole[1,9].

In summary, the health-related impact on people living with these filariae remains unknown, and it can be considered one of the most neglected tropical infectious diseases [10]. Therefore, our study was aimed to describe the clinical patterns and treatment guidelines of imported *M. perstans* infection by migrants from Africa.

## Material And Methods

### Study

The La Paz-Carlos III Hospital in Madrid, Spain, is a tropical disease referral unit. Most patients voluntarily attend the emergency unit or are referred from primary care or general hospitals in Madrid. A very small percentage of patients come from other regions.

A retrospective study was conducted on the data regarding immigrants diagnosed with *M. perstans* infection. The diagnosis of *M. perstans* infections was established with confirmed microfilaremia or adult identification following extraction. The direct detection of circulating microfilaria was performed on fresh venous blood obtained around midday with a thick film and/or thin smear after Giemsa staining. Microfilaremia was occasionally estimated on thin smears. The exclusion criteria included diagnosis in travelers, unspecified diagnosis methods (i.e., clinical data only), and medical records with missing data. The data included demographics (age, gender, nationality, time of the first consultation) and clinical characteristics (symptoms and when the symptoms first appeared). The eye examination results and analytical data regarding serologic tests for syphilis, HIV, hepatitis B and C, eosinophil counts, IgE levels and stool test results regarding ova and parasites were reviewed. Other laboratory test results were also recorded. Systematic ophthalmology exploration was performed in patients with a clinical suspicion of onchocerciasis. Hyper-IgE was defined as an increase in peripheral blood IgE to more than 200 U/mL. Hyper-IgE was classified as being mild (>200–399 U/mL), moderate (>399–999 U/mL) and/or severe (>1000 U/mL). Relative eosinophilia was defined as an elevated percentage of eosinophils (>5%) in individuals with  $<450 \times 10^6$  eosinophils/L. Absolute eosinophilia was defined as an increase in the peripheral blood eosinophilic leukocytes to more than  $450 \times 10^6$  eosinophils/L of blood. Mild eosinophilia was defined as  $>450 \times 10^6$  eosinophils/L to  $999 \times 10^6$  eosinophils/L. Moderate eosinophilia was defined as  $>1,000 \times 10^6$  eosinophils/L to  $2,999 \times 10^6$  eosinophils/L, and severe eosinophilia was defined as  $>3,000 \times 10^6$  eosinophils/L.

## Statistical analysis

Categorical variable results were expressed as percentages and as the mean and standard deviation (SD) for continuous variables. Chi-square test was used to compare the association between the categorical variables (i.e., clinical and demographic variables). The measured outcomes were expressed as the odds ratio (OR) with a 95% CI. The continuous variables were compared by Student's t-test or the Mann–Whitney test for two groups depending on their normal or nonnormal distribution. The corresponding regression models were used for multivariate analysis considering  $p < 0.05$  for a statistically significant difference. The Statistical Package for the Social Sciences (SPSS 23.0®; IBM Corp., Armonk, New York, USA) was used to analyze all the data.

## Results

### Demographic & Epidemiological data

In total, 503 cases of *M. perstans* infection were identified at the Carlos III Hospital. The main epidemiological data are shown in **Table 1**. Two hundred sixty-four patients were female (52.5%), and 239 (47.5%) were male. The migrants' mean age ( $\pm$ SD) was  $44.6 \pm 18.2$  years (range, 16–93); the median (25<sup>th</sup>, 75<sup>th</sup> percentiles) age was 43 years (28, 60). Most of them (97.6%) came from Equatorial Guinea. All the patients were infected in their country of origin. The mean time ( $\pm$ SD) between their arrival to Spain and their first medical consultation was  $8.6 \pm 18.0$  months (range, 1–180); the median (25<sup>th</sup>, 75<sup>th</sup> percentiles) time was 2 months (1, 7). Half of the patients (50.3%) were evaluated during the first month of stay in Spain. The mean number of cases ( $\pm$ SD) was  $25.1 \pm 11.8$  (5–45), with an irregular chronological distribution.

	All patients	Microbiological association			Clinical association		
	N=503 (100%)	Only M. perstans n <sub>1</sub> =308 (61.2%)	Co-infections n <sub>2</sub> =195 (38.8%)	p-value*	Symptomatic n <sub>3</sub> =228 (45.3%)	Asymptomatic n <sub>4</sub> =275 (54.7%)	p-value*
Epidemiological data							
Male	239 (47.5)	149 (48.4)	90 (46.2)	0.627	94 (41.2)	145 (52.7)	0.010*
Female	264 (52.5)	159 (51.6)	105 (53.8)		134 (58.8)	130 (47.3)	
Age, mean±SD, years	44.6±18.2	45.4±17.8	43.3±18.7	0.228	44.8±18.6	44.4±17.8	0.771
0-24	93 (18.5)	50 (16.2)	43 (22.1)		44 (19.3)	49 (17.8)	
25-50	205 (40.8)	129 (41.9)	76 (39.0)		93 (40.8)	112 (40.7)	
>50	205 (40.8)	129 (41.9)	76 (39.0)		91 (39.9)	114 (41.5)	
Months in Spain, mean±SD	8.6±18.0	8.2±19.9	9.2±14.4	0.540	8.4±15.7	8.7±19.7	0.850
Clinical data							
Asymptomatic	275 (54.7)	194 (63.0)	81 (41.5)	<0.001*			
Symptomatic	228 (45.3)	114 (37.0)	114 (58.5)				
Pruritus	190 (37.8)	88 (28.6)	102 (52.3)				
Arthralgia	50 (9.9)	33 (10.7)	17 (8.7)				
Abdominal pain	15 (3.0)	12 (3.9)	3 (1.5)				
Subcutaneous step (calabar swelling)	18 (3.6)	3 (1.0)	15 (7.7)				
Laboratory data							
Eosinophilia	N=503	n <sub>1</sub> =308	n <sub>2</sub> =195		n <sub>3</sub> =228	n <sub>4</sub> =275	
Without eosinophilia (<450)	105 (20.9)	69 (22.4)	36 (18.5)	<0.001*	54 (23.7)	51 (18.5)	<0.001*
Relative eosinophilia (<450 + >5%)	58 (11.5)	54 (17.5)	4 (2.1)		22 (9.6)	36 (13.1)	
Mild eosinophilia (450-999)	159 (31.6)	104 (33.8)	55 (28.2)		53 (23.2)	106 (38.5)	
Moderate eosinophilia (1000-2999)	154 (30.6)	75 (24.4)	79 (40.5)		78 (34.2)	76 (27.6)	
Severe eosinophilia (≥3000)	27 (5.4)	6 (1.9)	21 (10.8)		21 (9.2)	6 (2.2)	
Mean±SD	1151.7±1296.5	819.8±863.0	1465.4±1707.2		1252.0±1387.7	919.2±1197.4	
Immunoglobulin E	N=491	n <sub>1</sub> =300	n <sub>2</sub> =191		n <sub>3</sub> =222	n <sub>4</sub> =269	
Normal (<200)	49 (10.0)	37 (12.3)	12 (6.3)	0.001*	20 (9.0)	29 (10.8)	0.749

Mild hyper-IgE (200-399)	57 (11.6)	46 (15.3)	11 (5.8)		23 (10.4)	34 (12.6)	
Moderate hyper-IgE (400-999)	105 (21.4)	61 (20.3)	44 (23.0)		48 (21.6)	57 (21.2)	
Severe hyper-IgE ( $\geq 1000$ )	280 (55.7)	156 (52.0)	124 (64.9)		131 (59.0)	149 (55.4)	
Mean $\pm$ SD	1417.3 $\pm$ 1132.9	1310.0 $\pm$ 1111.7	1585.8 $\pm$ 1148.2		1442.2 $\pm$ 1153.7	1396.8 $\pm$ 1117.2	
Evolution							
Healing	240 (47.7)	147 (47.7)	93 (47.7)	0.599	118 (51.8)	122 (44.4)	0.247
No	18 (3.6)	9 (2.9)	9 (4.6)		7 (3.1)	11 (4.0)	
No follow-up	245 (48.7)	152 (49.4)	93 (47.7)		103 (45.2)	142 (51.6)	
*Statistical significance level of 5% (p <0.05)							

Table 1

Epidemiological, clinical, laboratory and evolution data, according to microbiological (*Mansonella perstans* vs co-infections) and clinical associations (symptomatic vs asymptomatic).

## Clinical & Laboratory data

The main clinical and analytical patient data are described in **Table 1**, according to the variables "Microbiological association", *Mansonella perstans* ( $n_1=308$ , 61.2%) vs. coinfections ( $n_2=195$ , 38.8%) and "Clinical association", symptomatic ( $n_3=228$ , 45.3%) vs. asymptomatic ( $n_4=275$ , 54.7%). Clinical manifestations were observed in 228 patients (45.3%). The most frequent symptom was pruritus in 190 patients (83.3%), arthralgia in 50 patients (9.9%), abdominal pain in 15 patients (3.0%) and Calabar swelling in 18 patients. Significant differences ( $p=0.010$ ) were found between the percentage of men with symptoms (41.2%) and women with symptoms (58.8%), and no significant differences in age were found ( $p=0.771$ ).

Regarding eosinophilia levels, 105 (20.9%) patients had a normal eosinophil count, 58 (11.5%) had relative eosinophilia and 340 (67.6%) migrants had absolute eosinophilia: mild eosinophilia in 159 patients (31.6%), moderate eosinophilia in 154 patients (30.6%) and severe eosinophilia in 27 patients (5.4%). Moreover, 152/228 (66.6%) symptomatic patients had absolute eosinophilia vs. 188/275 (68.3%) asymptomatic patients. Statistically significant differences were found between the percentages of symptomatic vs. asymptomatic cases and levels of eosinophilia ( $p<0.001$ ). Concerning IgE detection, 49 (10%) migrants had normal levels, and the remainder (442, 90%) had hyper-IgE: mild hyper-IgE (57, 11.6%), moderate hyper-IgE (105, 21.4%) and severe hyper-IgE (280, 55.7%). No statistically significant differences were found between asymptomatic and symptomatic patients and IgE ( $p=0.749$ ). Finally, 240 (47.7%) cases healed (**Table 1**). Among the remaining 263 (52.3%) cases, 245 (48.7%) individuals had no follow-up data and 18 (3.6%) cases were not cured.

## Microbiology data

**Table 2** describes the microorganism coinfection in patients with *Mansonella perstans* infection. Globally, 308 migrants (61.2%) had only *Mansonella perstans* infections and 195 patients (38.8%) had coinfection with other filarial nematodes. *Onchocerca volvulus* was the most frequent filarial coinfection with 138 patients (27.4%). One hundred eighty-six patients (37%) had coinfection with other helminths. *Trichuris trichiura* was the most frequent helminthic coinfection, with 137 patients (27.24%). Moreover, 73 (14.5%) migrants presented simultaneously with other filarial and helminthic coinfections. Additionally, 78 migrants (15.5%) were protozoa coinfecting, mainly with amoebas (58, 11.5%). Finally, 26 patients (5.2%) had HIV coinfection.

	Total=503 (100%) n (%)
Only <i>Mansonella perstans</i>	308 (61.2)
Filarial co-infection	195 (38.8)
<i>Onchocerca volvulus</i>	110 (21.9)
<i>Loa loa</i>	48 (9.5)
<i>Mansonella streptocerca</i>	7 (1.4)
<i>Loa Loa</i> + <i>Onchocerca volvulus</i>	12 (2.4)
<i>Onchocerca volvulus</i> + <i>Mansonella streptocerca</i>	13 (2.6)
<i>Loa Loa</i> + <i>Onchocerca volvulus</i> + <i>Mansonella streptocerca</i>	3 (0.6)
<i>Loa Loa</i> + <i>Mansonella streptocerca</i> + <i>Wuchereria bancrofti</i>	1 (0.2)
All negative*	1 (0.2)
Other helminthic co-infections	186 (37.0)
<i>Trichuris trichiura</i>	74 (14.7)
<i>Ascaris lumbricoides</i>	33 (6.6)
Hookworms	6 (1.2)
<i>Strongyloides stercoralis</i>	5 (1.0)
<i>Schistosoma</i> spp	2 (0.4)
<i>Trichuris trichiura</i> + <i>Ascaris lumbricoides</i>	44 (8.7)
<i>Trichuris trichiura</i> + Hookworms	8 (1.6)
<i>Trichuris trichiura</i> + <i>Schistosoma</i> spp	1 (0.2)
<i>Ascaris lumbricoides</i> + Hookworms	2 (0.4)
<i>Ascaris lumbricoides</i> + <i>Strongyloides stercoralis</i>	1 (0.2)
<i>Trichuris trichiura</i> + <i>Ascaris lumbricoides</i> + Hookworms	9 (1.8)
<i>Trichuris trichiura</i> + <i>Ascaris lumbricoides</i> + <i>Strongyloides stercoralis</i>	1 (0.2)
<i>M. perstans</i> + Other filarials + Other helminthics co-infection	73 (14.5)
Protozoa co-infection	78 (15.5)
Amebas	54 (10.7)
<i>Dientamoeba fragilis</i>	3 (0.6)
<i>Giardia lamblia</i>	17 (3.4)
Amebas + <i>Giardia lamblia</i>	4 (0.8)
Viruses co-infection	26 (5.2)
HIV	26 (5.2)
*All negative but presence of calabar swelling	

Table 2  
Co-infections in patients with *Mansonella perstans* infections.  
The presence of coinfections was not significantly related to gender (48.4% men vs. 51.6% women,  $p=0.627$ ). No significant differences were found between age groups ( $p=0.228$ ). By contrast, a higher percentage of patients infected only with *M. perstans* was

asymptomatic (63%) than patients coinfecting with symptoms (58.5%) ( $p < 0.001$ ). Coinfecting patients had higher absolute eosinophilia percentages ( $p < 0.001$ ), severe eosinophilia (10.8% vs 1.9%) and moderate eosinophilia (40.5% vs 24.4%). Similarly, coinfecting patients had higher hyper-IgE, severe and moderate ( $p = 0.001$ ) (see **Table 1**).

## Treatment and outcome

Four hundred thirty-seven cases (86.9%) were treated, 66 (13.1%) cases were untreated, and 292 cases (58.1%) used only one drug: mebendazole ( $n = 267$ ), ivermectin ( $n = 16$ ) and albendazole ( $n = 9$ ). By contrast, 145 cases (28.8%) used combined therapy, mainly ivermectin and mebendazole ( $n = 113$ ), as shown in **Table 3**. Therefore, the drug most used, alone or associated, was mebendazole, in 407 patients. Most of them ( $n = 382$ ) received a single dose, 24 double doses and 1 triple dose. Corticosteroid therapy was administered concurrently with the anti-filarial drug in 20 (4%) cases, and an antihistaminic drug was administered with the anti-filarial drug in 38 (7.6%) cases. **Figure 1** shows a significant decrease in eosinophilia before and after treatment ( $p < 0.001$ ).

	Anti-filarial drugs	Adverse effects	Healing
	437/503 (86.9%)	25/503 (5%)	240/503 (47.7%)
Simple treatment	292/503 (58.1)	12/292 (4.1)	154/292 (52.7)
Mebendazole	267/503 (53.1)	9/267 (3.4)	152/267 (56.9)
Ivermectine	16/503 (3.2)	2/16 (12.5)	0
Albendazole	9/503 (1.8)	1/9 (11.1)	2/9 (22.2)
Combined treatment	145/503 (28.8)	13/145 (9.0)	86/145 (59.3)
Diethylcarbamazcine + Ivermectine + Mebendazole	20/503 (4.0)	3/20 (15.0)	17/20 (85.0)
Diethylcarbamazcine + Mebendazole	3/503 (0.6)	1/3 (33.3)	3/3 (100.0)
Diethylcarbamazcine + Albendazole	5/503 (1.0)	2/5 (40.0)	4/5 (80.0)
Ivermectine + Mebendazole	113/503 (22.5)	7/113 (6.2)	61/113 (54.0)
Ivermectine + Mebendazole + Albendazole	1/503 (0.2)	0	0
Mebendazole + Albendazole	3/503 (0.6)	0	1/3 (33.3)
None/No follow-up	66/503 (13.1)	478/503 (95.0)	263/503 (52.3)

Table 3  
Treatment in patients with *Mansonella perstans*

Adverse reactions to the anti-filarial drugs were in 25 (5%) migrants: 15 had pruritus (13 associated with mebendazole and 2 with ivermectin), 3 had pruritus and skin abscess (3 mebendazole), 5 had Calabar swelling (mebendazole or albendazole associated with DEC or ivermectin), 1 had arthritis (DEC and ivermectin and mebendazole), and 1 had hepatotoxicity (DEC and albendazole).

No significant differences were found in the healing outcomes between the two most frequent treatment patterns: mebendazole (152/26, 56.9%) vs mebendazole and ivermectin (61/113, 54%). The combined anthelmintics and steroid treatment presented significantly better results; 14/20 (70%) cases healed ( $p = 0.042$ ). However, no significant results were obtained for the combined anthelmintic and antihistamine treatment; 22/38 (57.9%) cases healed ( $p = 0.191$ ). No statistical association was found between the percentage of side effects of treatment and cases healed ( $p = 0.094$ ).

## Discussion

*M. perstans* is transmitted by biting midges (Culicoides). The life cycle is similar to that of other filariae. Microfilariae are responsible for the transmission of infection because they are taken up during the blood meal of the insect vector. The epidemiology of *M. perstans* has not been clearly defined. Among the known human filarial infections, mansonellosis is probably the most frequent filariasis in sub-Saharan Africa as well as a northern part of the Amazon rainforest stretching from equatorial Brazil to the Caribbean coast of South America[9]. It has been estimated that 114 million people may be infected and as many as 581 million people in 33 countries are at risk

for *M. perstans* infection in Africa alone[1]. Many publications refer to mansonellosis as one of the most common human helminthiasis in endemic areas, and it is more prevalent and more neglected than other filarial diseases such as lymphatic filariasis, onchocerciasis, and loiasis. In endemic areas, the probability of infection increases with age, with the prevalence reaching 100% in highly endemic areas. However, the infection is the least studied and is likely one of the most neglected of all tropical diseases, subject to more neglect than schistosomiasis, taeniasis, echinococcosis, or rabies[9,10].

Most symptoms ascribed to *M. perstans* infections in modern scientific literature are based on symptoms that have been recorded in case study reports. Most of these reports have been based on the treatment of tourists and expatriate Europeans and North Americans returning home from endemic areas, and not on people who have lived all their lives in endemic areas. It is unclear whether the symptoms reported from these studies can be used to compile a clinical picture that represents all or even most infections caused by mansonellosis. The identification of these infections is often complicated by coinfection with other infective agents. Bassene et al. analyzed patients infected only with *M. perstans* and concluded that these infections had low pathogenicity because most individuals with *M. perstans* infection are asymptomatic[11,12]. When symptoms occur, they are predominantly related to the migration of the adult worms and include dermatological symptoms such as transient subcutaneous swellings similar to the Calabar swellings of *Loa loa* infection, serositis (i.e., pericarditis and pleuritis)[13,14], and ocular symptoms (granulomatous nodules in the conjunctiva, retinal lesions, and periorbital inflammation surrounding dead adult worms)[1]. Nonspecific symptoms, including pruritus, urticaria, fever, pain in bursae and/or joint synovia, enlarged lymph glands, vague abdominal symptoms and fatigue, have also been attributed to *M. perstans*. Headaches, neuropsychiatric manifestations, meningoencephalitis, and hepatitis have also been described. Nonspecific but characteristic laboratory abnormalities include high-level eosinophilia and elevated serum immunoglobulin IgE levels observed in some but not all patients with *M. perstans* infections, likely because of the body's reaction against the adult worm, rather than against microfilariae[15,16]. In our series, all the described features are represented. Moreover, recent observations have suggested that mansonellosis infections can influence the human immune system's response, which can influence the development of secondary infections, such as malaria[17].

Parasitological diagnosis is based on the detection and identification of sheathless *Mansonella* microfilariae in the skin or blood at any time of day or night. Additionally, the diagnosis is established by identifying the adult worm in tissues. Serologic tests based on crude filarial antigens are useful but do not distinguish between active or past infection and show cross-reactivity among different filarial species and with other nematode infections. Consequently, their usefulness is limited, although a negative result can exclude the possibility of infection. Given the limitations of serology, we do not apply this diagnostic technique and may underestimate the number of cases.

Among the three types of human mansonellosis, that caused by *M. perstans* is usually regarded as the most difficult to treat[1,9]. Our work shows great variability in the treatment of this disease. Therefore, human infection with *M. perstans* raises questions about treatment because of poor responses to standard antifilarial drugs and limited findings from controlled trials. In contrast to conventional anthelmintic treatments, doxycycline has proven to be excellent, effective, and safe in the treatment of *M. perstans* infections[5,18]. However, the course of treatment over 6 weeks that is necessary for this type of therapy probably makes it impractical for control programs, although it appears to be curative, making it a very desirable therapeutic for travel medicine[5,18]. *M. perstans* is relatively resistant to standard antifilarial agents, including DEC, ivermectin, albendazole, and mebendazole[2]. The usefulness of doxycycline in the treatment of *M. perstans* varies according to geographic region [5]. An ideal drug treatment for *M. perstans* infections needs to be identified that is effective, fast acting, tolerable and easy to administer. The search for new treatments may include a more meticulous quantitative assessment of the above-mentioned drugs, both alone and in various combinations[1].

Despite accumulating evidence of a high prevalence of human infections, no current large-scale filariasis control program has targeted mansonellosis. Mansonellosis is not listed among the neglected diseases of the World Health Organization, and no control strategy has been defined against this human filariasis.

The health-related impact on individuals living with these filariae remains unknown, and evidence regarding treatment strategies is scarce. Like other neglected diseases, it mainly affects poor populations living in tropical and subtropical climates and it has not been associated with a clear and distinct clinical picture[1]. It can be considered one of the most neglected tropical infectious diseases [10].

Our study had some limitations, which were caused mostly by the retrospective design. First, most patients visited our center because of symptoms or an increased eosinophil count. Thus, the proportion of symptomatic patients is not representative of the general population with *M. perstans* infection. Second, posttreatment follow up was available for only a few patients. Thus, we could not properly describe the response to treatment.

## Conclusions

In summary, a long series of *M. perstans* infections is presented in sub-Saharan immigrants whose data indicate that it should be included in the differential diagnosis in patients with pruritus or analytical alterations such as eosinophilia or hyper-IgE presentation. These patients also have a high number of coinfections with other microorganisms, the treatment of which needs to be protocolized.

## Declarations

## Ethics approval and consent to participate

This was a retrospective analysis of data obtained over a 19-year period. The data were collected anonymously to ensure impartial analysis; therefore, written informed consent was not obtained from the individual participants. The research was approved by the La Paz–Carlos III Hospital's Ethics Committee.

## Consent for publication

Not applicable

## Availability of data and materials

The dataset supporting the conclusions of this article is included within the article and its additional file.

## Competing interests

All authors declare no potential conflicts of interest and no sources of support.

## Funding

None.

## Authors' contributions

SP conceived the study

SP, MS, ISE, ML, MBG, AM designed the study protocol

SP, ISE, ML, MA, MS carried out clinical assessment

MS and BV carried out microbiological analysis and interpretation of these data

MAS analyzed and interpreted statistical data

MBG and AM drafted the manuscript

All authors contributed to writing the manuscript

SP critically revised the manuscript for intellectual content

All authors read and approved the final manuscript.

SP, MBG and AM are guarantors of the paper.

## Acknowledgements

Virginia Velasco-Tirado.

## References

1. Simonsen PE, Onapa AW, Asio SM. Mansonella perstans filariasis in Africa. *Acta Tropica* **2011**; 120 Suppl 1:S109–20.

2. Bregani ER, Rovellini A, Mbaïdoum N, Magnini MG. Comparison of different anthelmintic drug regimens against *Mansonella perstans* filariasis. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **2006**; 100:458–463.
3. Asio SM, Simonsen PE, Onapa AW. Analysis of the 24-h microfilarial periodicity of *Mansonella perstans*. *Parasitology Research* **2009**; 104:945–948.
4. Keiser PB, Coulibaly Y, Kubofcik J, et al. Molecular identification of *Wolbachia* from the filarial nematode *Mansonella perstans*. *Mol. Biochem. Parasitol.* **2008**; 160:123–128.
5. Coulibaly YI, Dembele B, Diallo AA, et al. A randomized trial of doxycycline for *Mansonella perstans* infection. *The New England journal of medicine* **2009**; 361:1448–1458.
6. Büttner DW, Wanji S, Bazzocchi C, Bain O, Fischer P. Obligatory symbiotic *Wolbachia* endobacteria are absent from *Loa loa*. *Filaria J* **2003**; 2:10.
7. Grobusch MP, Kombila M, Autenrieth I, Mehlhorn H, Kreamsner PG. No evidence of *Wolbachia* endosymbiosis with *Loa loa* and *Mansonella perstans*. *Parasitology Research* **2003**; 90:405–408.
8. ADOLPH PE, KAGAN IG, McQUAY RM. Diagnosis and treatment of *Acanthocheilonema perstans* filariasis. *American Journal of Tropical Medicine and Hygiene* **1962**; 11:76–88.
9. Ta-Tang T-H, Crainey JL, Post RJ, Luz SL, Rubio JM. Mansonellosis: current perspectives. *Res Rep Trop Med* **2018**; 9:9–24.
10. Mediannikov O, Ranque S. Mansonellosis, the most neglected human filariasis. *New Microbes New Infect* **2018**; 26:S19–S22.
11. Bassene H, Sambou M, Fenollar F, et al. High Prevalence of *Mansonella perstans* Filariasis in Rural Senegal. *The American journal of tropical medicine and hygiene* **2015**; 93:601–606.
12. JANSSENS PG. [IS D. PERSTANS PATHOGENIC FOR MAN?]. *Ann Soc Belges Med Trop Parasitol Mycol* **1964**; 44:989–998.
13. FOSTER DG. Filariasis, a rare cause of pericarditis. *The Journal of tropical medicine and hygiene* **1956**; 59:212–214.
14. Kahn JB. Pleural effusion associated with *Dipetalonema perstans* (*Acanthocheilonema perstans*). *The Journal of infectious diseases* **1983**; 147:166.
15. Holmes GK, GELFAND M, Boyt W, Mackenzie P. A study to investigate the pathogenicity of a parasite resembling *Acanthocheilonema perstans*. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **1969**; 63:479–484.
16. Bouyou-Akotet MK, Moussavou Boussougou MN, Ovono-Abessolo F, Owono-Medang M, Kombila M. Influence of *Mansonella perstans* microfilaraemia on total IgE levels in Gabonese patients co-infected with *Loa loa*. *Acta Tropica* **2014**; 131:11–15.
17. Dolo H, Coulibaly YI, Dembele B, et al. Filariasis attenuates anemia and proinflammatory responses associated with clinical malaria: a matched prospective study in children and young adults. *PLoS Neglected Tropical Diseases* **2012**; 6:e1890.
18. Medeiros JF, Crainey JL, Pessoa FAC, Luz SLB. Mansonellosis. In: *Arthropod Borne Diseases*. Cham: Springer International Publishing, 2016: 405–426.

## Figures




Figure 1

Eosinophilia before and after treatment