

Single-wall Carbon Nanotubes Improve Cell Survival Rate and Reduce Oxidative Injury in Cryopreservation of *Agapanthus praecox* Embryogenic Callus

Li Ren

Shanghai Academy of Agricultural Sciences <https://orcid.org/0000-0001-8255-7678>

Shan Deng

Shanghai Academy of Agricultural Sciences

Yunxia Chu

Shanghai Academy of Agricultural Sciences

Yiying Zhang

Shanghai Academy of Agricultural Sciences

Hong Zhao

Shanghai Academy of Agricultural Sciences

Hairong Chen (✉ saasdus@163.com)

<https://orcid.org/0000-0003-3996-5857>

Di Zhang

Shanghai Jiao Tong University

Research

Keywords: Cryopreservation, Single-wall Carbon Nanotubes, Oxidative stress, *Agapanthus praecox*

Posted Date: June 30th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-28448/v2>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Plant Methods on September 21st, 2020. See the published version at <https://doi.org/10.1186/s13007-020-00674-6>.

Abstract

Background: Cryopreservation is the best way for long-term in vitro preservation of plant germplasm resources. The preliminary studies found that reactive oxygen species (ROS) induced oxidative stress and ice-induced membrane damage are the fundamental causes of cell death in cryopreserved samples. How to improve plant cryopreservation survival rate is an important scientific issue in the cryobiology field.

Results: This study found that the survival rate was significantly improved by adding single-wall carbon nanotubes (SWCNTs) to plant vitrification solution (PVS) in cryopreservation of *Agapanthus praecox* embryogenic callus (EC), and analyzed the oxidative response of cells during the control and SWCNTs-added cryopreservation protocol. The SWCNTs entered EC at the step of dehydration, and mainly located around the cell wall and in the vesicles, and most of SWCNTs moved out of EC during dilution step. Combination with physiological index and gene quantitative expression results, SWCNTs affect ROS signal transduction and antioxidant system response during plant cryopreservation. The EC treated by SWCNTs had higher antioxidant levels, like POD, CAT and GSH than the control group EC. EC mainly depended on AsA-GSH and GPX cycle to scavenge H₂O₂ in the control cryopreservation, but depended on CAT in the SWCNTs-added cryopreservation which lead to low levels of H₂O₂ and MDA. Elevated antioxidant level in dehydration by adding SWCNTs enhanced cells resistance to injury during cryopreservation. The ROS signals of EC were balanced and stable in the SWCNTs-added cryopreservation.

Conclusions: SWCNTs regulated oxidative stress responses of EC during the process, and controlled oxidative damages by the maintenance of ROS homeostasis to achieve high survival rate after cryopreservation. This study is the first to systematically describe the role of carbon nanomaterial in the regulation of plant oxidative stress response, and provided a novel insight into the application of nanomaterials in the field of cryobiology.

Background

Long-term preservation of plant in vitro cultures of cell cultures, embryogenic materials, and endangered germplasm is a vital requirement around the world [1]. Cryopreservation, the storage of living cells, tissues, organs or whole plants in extra low temperature, such as in liquid nitrogen (LN), is thought to be the ideal method to realize the safe and cost-efficient medium- and long-term preservation of almost all economically important crops [2-4]. Vitrification-based cryopreservation has been successfully applied to long-term preservation by achieving the glassy state in cryopreservation of many plant species for its rapid and convenient procedure [5]. Plant vitrification solution (PVS), a commonly used cryoprotectant in vitrification-based cryopreservation, can replace cellular water, alter the freezing behavior, and prevent ultra-water loss [6].

Vitrification-based cryopreservation achieves storage by enhancing cell viscosity, and accompanies many stresses leading to the accumulation of reactive oxygen species (ROS) [7]. ROS-induced oxidative stress

is the fundamental trigger of cell death during cryopreservation [8]. Ren et al. [7, 9] utilized comparative transcriptomics to obtain genes response to cryoinjury and revealed that peroxidation was a key element affecting viability, and genes related to oxidative stress played important roles in *Arabidopsis thaliana* cryopreservation. Zhang et al. [10] revealed that oxidative stress and apoptosis were the major factors which injure embryogenic callus during *Agapanthus praecox* cryopreservation.

Plants have complex antioxidant systems including antioxidant enzymes, such as catalase (CAT), superoxide dismutase (SOD), glutathione reductase (GR) and ascorbate peroxidase (APX), and non-enzymatic antioxidants, like glutathione (GSH) and ascorbic acid (AsA) [11]. The high antioxidant level is related to the tolerance to cryopreservation [12]. In antioxidant enzymes, SOD removes O_2^- to H_2O_2 , and CAT catalyzes H_2O_2 to O_2 and H_2O [13, 14]. In non-enzymatic antioxidants, AsA and GSH helps keep a normal REDOX state to counteract oxidative damages [15].

Many researchers have been focusing on the improvement of cryopreservation procedure, by adding exogenous compounds to the cryoprotectant to reduce damages. Nanomaterials have a good effect to improve the cryoprotectant, due to its small particle size and large specific surface [16]. After adding nanomaterials to the cryoprotectant can enhance the thermal conductivity and relative viscosity, promote vitrification, improve the stability of solution during the rewarming process, and suppress the occurrence of devitrification [17-19]. Han et al. [20] added 0.2% Nano-diamond to the ethylene glycol cryoprotectant, and found it doubled the freeze rate and decreased the glass transition temperature significantly. In addition, Adding 0.05% hydroxyapatite to the cryoprotectant can improve the pig (*Sus domesticus*) oocytes developmental rate from 14.7% to 30.4% after cryopreservation [19, 21]. After adding Nanomaterials Rhodiola Sachanensis Polysaccharide (NRSP), the motility, acrosome integrity and the membrane integrity of boar sperm were improved significantly compared to the control groups after freezing and thawing [22]. With the addition of NRSP in the cryoprotectant, the malondialdehyde (MDA) content decreased and the activity of SOD increased [21]. Furthermore, lecithin nanoparticles can enhance the cryosurvival of caprine sperm [23], and albumin coated copper-cysteine nanozyme and Au-Ag-AFT nanozyme both can improve survival of human sperm after cryopreservation by reducing oxidative stress [24, 25]. Enrichment of semen extender with selenium nanoparticles improved fertility rate through decreasing peroxidation and injury during cryopreservation of Holstein bulls [26].

However, some nanomaterials have been reported to induce cell death [17, 18], and carbon nanomaterials (CNMs) have better biosafety and biocompatibility [27,28]. 60-h seedlings of *A. thaliana* are often used as an experimental model to evaluate the optimal effect of exogenous additives on cryopreservation of plants [7]. In our previous studies, CNMs including graphene, single-walled carbon nanotubes (SWCNTs) and graphene quantum dots have been applied in this cryopreservation model to identify their effects on cryopreservation, and 0.1 g/L SWCNTs in PVS2 was the most effective one (unpublished results). Then we applied it to the callus or protocorm cryopreservation of *lily*, *Cymbidium* and *Anoectochilu*, and it also played a positive role in each process. Chen et al. [28] investigated the efficiency of SWCNTs, Graphene, Graphene quantum dots and fullerene (C_{60}) on the cryopreservation of *Agapanthus praecox* callus by adding to PVS2. 0.3 g/L fullerene and 0.1 g/L SWCNTs were the best two CNMs. The effect on

vitrification behaviors was detected, and the glass transition temperatures (T_g) showed no significant difference on Nano-PVS2. Raman spectroscopy and transmission electron microscopy (TEM) analyses demonstrated that the SWCNTs and fullerene are able to enter the callus cells and protect the cell structure [28].

Agapanthus praecox is a perennial herbaceous plant, and sensitive to low temperature. We have successfully cryopreserved the embryonic callus (EC) of *A. praecox*. In this study, SWCNTs adding into PVS2 was evaluated in *Agapanthus praecox* EC cryopreservation. It aims to point the positive effects to oxidative stress in cryopreservation, and provides a new application of CNMs using as protectants during cryopreservation. This study is the first time to systematically investigate the effects of nanomaterial to regulate cell response to oxidative stress in plant cryopreservation applications.

Results

Effects of SWCNTs on the cell viability after cryopreservation

Based on the previous studies, CNMs have been applied in the *Arabidopsis* cryopreservation model to identify their effects on cryopreservation, and 0.1 g/L SWCNTs in PVS2 was the most effective one (unpublished results). In this study, we applied it to the EC cryopreservation of *A. praecox*, and 0.1 g/L SWCNTs have improved the relative survival rate from 53.42% to 84.57% (Fig. 1). Compare it to the effects of other compounds adding to PVS2 in previous studies including 0.08 mM GSH, 1 μ M abscisic acid (ABA), 0.1 μ M melatonin, 1 mM AsA, 10 mM betaine, 1 mM CaCl_2 , 6 mM lipoic acid (LA), 6 mM polyvinyl alcohol (PVA), or 3% polyvinyl pyrrolidone (PVP), 0.1g/L SWCNTs had a significant effect on the EC survival increasing by 58.31% higher than others (Fig. 1).

Localizations of SWCNTs in EC during cryopreservation

Transmission electron microscopy imaging provided accurate results, which displayed the localizations of SWCNTs inside the EC. The SWCNTs entered EC at the step of dehydration, and mainly located around the cell wall and in the vesicles (Fig. 2a-c). It is demonstrated that plasmolysis, protoplast concentrated, organelles damage and more intracellular vesicles were observed after dehydration (Fig. 2d). The dilution treatment caused most of SWCNTs to move out of EC, and the retained SWCNTs were mostly tube-like fragmented, mainly distributed in the cytoplasm and vesicles (Fig. 2e-h).

Effects of SWCNTs on the ROS and MDA contents during cryopreservation

In the control procedure, H_2O_2 contents highly increased in dehydration and reached the peak after rapid cooling-warming which was more the 2-fold compared with CK (Fig. 3a). Adding SWCNTs could inhibit H_2O_2 content during the dehydration step, and H_2O_2 content kept at a low level in rapid cooling-warming and decreased significantly in dilution in the improved cryopreservation. O_2^- inhibition activities were higher in all steps of the control group, and $\text{OH}\cdot$ generation activities were higher after rapid cooling-warming in the SWCNTs-added cryopreservation (Fig. 3b and c). The MDA content trend was similar to

that of H_2O_2 . MDA accumulation increased apparently during the control cryopreservation, and the EC treated with SWCNTs had less MDA than in control through the whole procedure (Fig. 3d).

Effects of SWCNTs on the antioxidant system during cryopreservation

Adding SWCNTs significantly increased the activity of antioxidant system during cryopreservation. With the addition of SWCNTs to the PVS2, the activities of enzymatic antioxidants highly increased, especially POD and CAT activities (Fig. 3f and g). The changes of SOD activity were not significant (Fig. 3e). The increases of POD activity treated by SWCNTs were much greater than those in the control process (Fig. 3f). Furthermore, the non-enzymatic antioxidant GSH in the SWCNTs group increased nearly 2-fold compared to the control group during dehydration (Fig. 3i).

Correlation analysis of oxidative physiological indices

In the control cryopreservation (Table 1), MDA contents showed a significant positive correlation with H_2O_2 contents, which revealed that peroxidation was mainly caused by H_2O_2 in EC during cryopreservation. At the same time, $OH\cdot$ generation activities had a significant positive correlation with H_2O_2 and MDA contents. SOD activities had a significant positive correlation with CAT activities and AsA contents. In the SWCNTs cryopreservation (Table 2), O_2^- inhibition activities had a significant negative correlation with POD activities, MDA and AsA contents, and $OH\cdot$ generation activities had a significant positive correlation with SOD activities and AsA contents. MDA contents had a significant positive correlation with SOD activities, and SOD activities had a significant positive correlation with AsA contents like in the control cryopreservation. In addition, there was a significant positive correlation between POD and CAT activities indicating that the two enzymes may work together to decrease ROS level during cryopreservation.

Table 1

The correlation analysis of physiological indices in the control cryopreservation

Indices	H ₂ O ₂ O ₂ ⁻ inhibition activity		OH· generation activity	MDA	SOD	POD	CAT	AsA	GSH
H ₂ O ₂	1	0.263	0.918*	0.974**	0.630	-0.141	0.510	0.702	0.830
O ₂ ⁻ inhibition activity	1		-0.076	0.146	-0.477	0.381	-0.678	-0.319	0.190
OH· generation activity			1	0.888*	0.764	-0.447	0.709	0.747	0.718
MDA				1	0.707	-0.046	0.621	0.794	0.844
SOD					1	-0.107	0.935*	0.973**	0.764
POD						1	-0.277	0.075	0.269
CAT							1	0.878	0.527
AsA								1	0.859
GSH									1
All the data were correlation coefficients, and significant levels are indicated at * P < 0.05 or ** P < 0.01.									

Table 2

The correlation analysis of physiological indices in the SWCNTs-added cryopreservation

Indices	H ₂ O ₂ O ₂ ⁻ inhibition activity		OH· generation activity	MDA	SOD	POD	CAT	AsA	GSH
H ₂ O ₂	1	-0.388	0.042	0.440	0.195	0.451	0.490	0.166	0.529
O ₂ ⁻ inhibition activity	1		-0.793	-0.904*	-0.864	-0.894*	-0.867	-0.884*	-0.437
OH· generation activity			1	0.851	0.982**	0.568	0.550	0.979**	0.124
MDA				1	0.934*	0.868	0.875	0.877	0.147
SOD					1	0.693	0.683	0.980**	0.152
POD						1	0.996**	0.647	0.175
CAT							1	0.621	0.134
AsA								1	0.304
GSH									1

All the data were correlation coefficients, and significant levels are indicated at * P < 0.05 or ** P < 0.01.

Quantitative expression analysis of genes related to oxidative stress response

Some genes related to oxidative stress response were chosen to study the molecular protection of SWCNTs during cryopreservation (Fig. 4). As a ROS signal transduction related gene, *oxidative signal-inducible 1 (OXI1)* was significantly upregulated at rapid cooling-warming stage in control group, but maintained a low expression level in SWCNTs group. *MAPK3/6* is located downstream of *OXI1*, and their expression pattern were very similar to that of *OXI1*. As a ROS signal amplification related gene, *NADPH oxidase (RbohA)* maintained a low expression level in control group, and was significantly upregulated after dehydration in the SWCNTs group.

In ROS-scavenging network, SOD and CAT catalyzed O₂⁻ by forming H₂O₂ and H₂O, respectively. The expression levels of *Cu/Zn* and *Fe SOD* were higher in control group than SWCNTs group, but those of *CAT* were quite the opposite which shown that *CAT* was upregulated in SWCNTs group. The glutathione peroxidase (GPX) and AsA-GSH cycle were also essential to scavenge H₂O₂. Glutathione reductase (GR) and GPX drive the GPX cycle, and GR, monodehydroascorbate reductase (MDHAR), and ascorbate

peroxidase (APX) regulate the AsA-GSH cycle. The expression levels of *GR* and *APX* were higher in control group than SWCNTs group. The expression patterns of *GPX1* and *MDHAR* had the similar trend. They were only significantly upregulated in the rapid cooling-warming step of the control cryopreservation, and the dilution step of the SWCNTs cryopreservation. *POD* only significantly upregulated in the dilution step of the SWCNTs cryopreservation.

Discussion

SWCNTs contribute to the better efficiency of cryoprotectant

The field of plant cryobiology seeks to modify existing techniques that allow for more efficient storage by reducing multiple stresses. Adding some exogenous compounds can improve the survival of cryopreserved cells in many species, like antioxidants [29-33], anti-stress compounds [34, 35], metabolism related compounds [36, 37] and ice inhibitors [38, 39]. Previously we found that CNMs especially SWCNTs adding to PVS2 can improve survival rate of *Arabidopsis* seedlings, callus or protocorm of *lily*, *Cymbidium* and *Anoectochilu* (unpublished results) and *A. praecox* callus [28] after cryopreservation. In this study, the relative survival rate was highly improved by adding SWCNTs to PVS2 in cryopreservation of *A. praecox* EC compared to above compounds (Fig. 1). Thus, SWCNTs are the potential and important exogenous additions of cryoprotectant.

In our previous study, the differential scanning calorimetry (DSC) analysis suggested that the glass-transition temperature of SWCNTs-PVS2 was slightly decreased [28]. Adding 0.1 g/L SWCNTs to PVS2 reduced the glass-transition temperature from -112.67 to -114.18 °C still within the reported temperature range of PVS2 (-115 °C to -112 °C). Moreover, a melting peak around -38.87 °C and an endothermic peak at -50.12 °C were observed in the DSC curve of 0.1 g/L SWCNTs-PVS2. It indicated that SWCNTs-PVS2 may be more stable than PVS2, but no significant changes on glass transition parameters were detected [28]. Since the glass-transition temperature is not changed, do SWCNTs regulate the physiological response of plant cells?

Effective antioxidant response during dehydration step improved EC survival after cryopreservation

ROS-induced oxidative stress is a major reason of low survival in samples after cryopreservation [40-48]. In many species cryopreservation, H₂O₂ is the major component of ROS leading to oxidative stress [8, 45]. Adding SWCNTs in the cryoprotectant might suppress H₂O₂ production and maintain the H₂O₂ content in the lower level than that in the control process, in which H₂O₂ content increased dramatically otherwise.

Membrane lipids are the primary target in oxidative damage [49], and MDA acting as a breakdown product of lipid peroxidation increased in cryopreserved *Oryza sativa* [47, 50], *Azadirachta indica* [51], *Arabidopsis thaliana* [8] and *Agapanthus praecox* [44], *Hancornia speciosa* [52], *Passiflora suberosa* [48]. With the addition of NRSP in the cryoprotectant, the MDA content decreased and the activity of SOD increased leading to more intact membrane and better quality of boar sperm [21]. MDA content significantly increased in the control process, and led to low survival rate of EC after cryopreservation.

Antioxidant system works to prevent plant cells from oxidative injury through cleaning ROS [11]. SWCNTs can activate related antioxidant enzymes, and improve survival after cryopreservation. In general, antioxidant enzyme activities increased after rapid cooling-warming (Fig. 3). In the SWCNTs cryopreservation, the enzyme activities rose up when cells were treated with SWCNTs-added PVS2, and maintained in a relative high level through the process. Like CAT was involved in scavenging the intracellular H_2O_2 in SWCNTs treated group, it is also found that cryopreserved of *Dendrobium* suffered serious oxidative stress because of decreasing CAT activity after cryopreservation leading to low survival [53], and high tolerance was related to high CAT activity in *Haematococcus pluvialis* cryopreservation [54]. In the study of Liu [55], the graphene-treated rice produced oxidative stress response, and the activities of SOD, POD and CAT were all increased in seedlings treated with graphene.

AsA-GSH cycle is a key way to scavenge H_2O_2 [56]. In this cycle, APX can scavenge H_2O_2 followed by a series of catalytic reactions involving GR, MDHAR and DHAR, in which GSH and AsA work as reducing substrates [57]. Adding SWCNTs increased GSH contents. Expression levels of *GPX* and *GR* in SWCNTs cryopreservation were lower than those in the control group, and the difference between these two processes became very obvious after rapid cooling-warming. This study indicated that PVS2 with SWCNTs improved the survival of *Agapanthus praecox*. SWCNTs promoted dehydration protection. This improvement is due to the scavenging of ROS and improving of antioxidative system activity, especially POD and CAT.

Effects of SWCNTs on ROS signal transduction during cryopreservation

Because of their key signaling roles (at low levels) and toxic roles (at high levels), the levels of ROS are regulated by the complex pathway including many genes [58-60]. In this pathway, plants sense ROS by three ways: (a) unknown ROS receptors; (b) redox-sensitive transcription factors; (c) direct inhibition of phosphatases [61]. The ROS signal is detected by unknown ROS receptors leading to the accumulation of Ca^{2+} signal [62-64], and the signal is transmitted to *oxidative signal-inducible 1 (serine/threonine protein kinase, OXI1)* which was significantly up-regulated in the control rapid cooling-warming with ROS level reaching the maximum which is much higher than that in the SWCNTs-added cryopreservation. OXI1 acts as a central factor in the ROS sensing, and is up-regulated in many H_2O_2 -generating stimulus [65]. *MAPK3/6* following *OXI1* [66, 67] were mainly up-regulated during the rapid cooling-warming of the control group.

The ROS signals further influenced following pathways including ROS producing and scavenging. The producing pathway has *NADPH oxidases* [68, 69], which were up-regulated in the SWCNTs treated steps. The producing pathway might be activated by ROS at low levels leading to the ROS production and amplification, and the scavenging pathway might be activated by ROS accumulation leading to the ROS suppression [61]. The interaction between producing and scavenging pathway determines the intensity of ROS signals [61]. In summary, EC in the control group suffered excessive ROS after dehydration, which broke the ROS metabolism balance. By contrast, SWCNTs both enhanced producing and scavenging pathway smoothly, and maintained ROS signals balanced and stable in EC during cryopreservation.

Regulation of physiological response of carbon nanotubes in plants

In the past decade, researchers have applied carbon nanotubes to plant studies and found that they have a certain regulatory effect on physiological response, especially on enzyme activity and gene expression. The effective impact of CNMs on plant development and growth has been studied by many research groups [70]. Giraldo et al. [71] pointed out that SWCNTs localized in the lipid envelope of *Arabidopsis* chloroplasts, promote over three times higher photosynthetic activity than that of controls, and concentrations of ROS inside chloroplasts were significantly suppressed. Industrialized MWCNTs can stimulate the growth of *Onobrychis arenaria* and enhance peroxidase activity [72]. In this study, SWCNTs contribute to the better efficiency of cryoprotectant, and also improved enzyme activities including POD and CAT. At the molecular level, SWCNTs regulated gene expression levels including *NADPH oxidase*, *CAT* and *POD*. Other studies have found similar results. MWCNTs enhanced the tobacco cell growth, and up-regulated genes related to water transport and cell division [73]. A number of genes regulated by MWCNTs were related to plant stress-signal transduction in tomato. Important stress signal pathways could be regulated in response to the uptake of carbon nanotubes [74]. For instance, MAPK was up-regulated in leaves exposed to MWCNTs, and played a positive role to promote plant development and stress response of carbon nanotubes [74]. In this study, MAPK3/6 were higher in control rapid cooling and warming than those treated with SWCNTs. Adding MWCNTs to the seeds of soybean (*Glycine max*), corn (*Zea mays*) and barley (*Hordeum vulgare*) led to the improvement of germination, and activated expression levels of aquaporins [75]. On the contrary, experiment on suspension rice cells with MWCNTs showed that it induced the accumulation of ROS leading to cell death [76]. It also decreases the dry weight and the activity of SOD in *Arabidopsis* suspension cells [77]. Whether carbon nanotubes play a positive or negative regulatory role, they can regulate a variety of biological processes in plants, like water transport, cell division, stress response, electron transfer, ROS generation and metabolism [78, 79]. However, the mechanism of how the nanotubes can regulate the physiological response especially regulate the gene expression is an unresolved issue and deserved further studies.

Conclusions

This study found that the relative survival rate was highly improved by adding SWCNTs to PVS2 in cryopreservation of *A. praecox* EC, and analyzed the oxidative response of EC at some steps in the control and SWCNTs-added cryopreservation. The SWCNTs entered EC at the step of dehydration, and mainly located around the cell wall and in the vesicles, and most of SWCNTs moved out of EC during dilution step. SWCNTs affect ROS signal transduction pathway and antioxidant system response through the physiological and gene expression results. The EC treated with SWCNTs exhibited higher antioxidant levels, including POD, CAT and GSH than the control group EC. EC mainly depended on the GPX and AsA-GSH cycle to scavenge H₂O₂ in the control cryopreservation, but depended on CAT in the SWCNTs-added cryopreservation which lead to low levels of H₂O₂ and MDA. Elevated antioxidant level in dehydration by adding SWCNTs improved cells tolerance to injury from cryopreservation procedure. The ROS signals of EC were balanced and stable in the SWCNTs-added cryopreservation. Overall, SWCNTs regulated the

oxidative stress response of EC in cryopreservation, and controlled cell oxidative injury by keeping ROS homeostasis to achieve high survival rate after cryopreservation.

Methods

Plant materials and SWCNTs

EC was induced and cultured from pedicel tissue of *A. praecox* as described by Wang et al. [80].

The SWCNTs aqueous solution (5.0 g/L, particle diameter 1 nm, length 1 μ m) was kindly provided by Prof. Yafei Zhang (Shanghai Jiao Tong University, Shanghai, China).

Cryopreservation procedure

EC was cryopreserved as described by Chen et al. [33]. The cryopreservation procedure included pre-culture (PC), osmoprotection (OP), dehydration with PVS2 (DH), rapid cooling-warming (RW), dilution (DL) and recovery (RC).

In the SWCNTs-added cryopreservation procedure, we added SWCNTs at 0.1 g/L in PVS2 at the dehydration step. The samples for the physiological experiments and qRT-PCR analysis were taken after some key steps including PC, DH, RW and DL. Experiments were performed three times individually.

Viability detection

In order to detect the survival of cryopreserved EC, the 2,3,5-triphenyltetrazolium chloride (TTC) protocol was used in this study [33]. EC (0.05 g) after 24 h-recovery was put into 2 mL TTC and kept in dark for 20 h. After rinsing 3 times with sterile water, EC was immersed in 95% ethanol and 85 °C water bathed for 1 h. EC was centrifuged at 3000 g for 5 min, and the optical density value of supernatant was tested at 485 nm. The relative survival rate was the ratio of cryopreserved and non-cryopreserved value. Each sample procedure was repeated 3 times.

Transmission Electron Microscopy

The EC was treated as described by Chen et al. [28], and observed by 120 kV biology transmission electron microscope (Tecnai G2 spirit Biotwin).

Detection of physiological indices

H₂O₂ levels, O₂⁻ inhabitation and OH[·] generation activities, MDA, SOD, CAT and POD activities, AsA and GSH contents were tested using related biological assay kits (Nanjing Jiancheng Bioengineering Institute, China) following the manufacturer's instructions with some modifications as according to Yang et al. [81].

qRT-PCR analysis

The qRT-PCR was performed as described by Chen et al. [33]. The amplifications were repeated for 3 times. The relative quantitative expression was calculated using the $2^{-\Delta\Delta CT}$ method. All primer sequences were listed in Additional file 1: Table S1.

Statistical analysis

The one-way ANOVA was used to analyze differences followed by least significant difference multiple range test using Statistics Analysis System 9.1.3 software (SAS Institute, Inc., Cary, NC, USA). Correlation analysis was calculated by Statistics Analysis System 9.1.3 software, and $P < 0.05$ was considered as significant.

Abbreviations

ABA: abscisic acid; APX: ascorbate peroxidase; AsA: ascorbic acid; CAT: catalase; DH: Dehydration; DHAR: dehydroascorbate reductase; DL: Dilution; DSC: differential scanning calorimetry; EC: embryogenic callus; GPX: glutathione peroxidase; GR: glutathione reductase; GSH: glutathione; H_2O_2 : hydrogen peroxide; LA: lipoic acid; LN: liquid nitrogen; LSD: least significant difference; MAPK: mitogen-activated protein kinase; MDA: malonaldehyde; MDHAR: monodehydroascorbate reductase; MS: Murashige and Skoog; NRSP: Nanomaterials Rhodiola Sachanensis Polysaccharide; $O_2^{\cdot -}$: superoxide anion; OH: hydroxyl radicals; OP: Osmoprotection; OXI1: oxidative signal-inducible 1; PC: Pre-culture; PVA: polyvinyl alcohol; PVP: polyvinyl pyrrolidone; PVS: plant vitrification solution; qRT-PCR: quantitative reverse-transcription polymerase chain reactions; RC: Recovery; ROS: reactive oxygen species; RW: Rapid cooling-warming; SOD: superoxide dismutase; TEM: transmission electron microscopy; T_g : glass transition temperatures; TTC: triphenyltetrazolium chloride; CNMs: carbon nanomaterials; CNTs: carbon nanotubes; SWCNTs: single-wall carbon nanotubes.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

All data generated or analysed during this study are included in this published article and its Additional file 1: Table S1.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by the National Natural Science Foundation of China [NO. 31670693, NO. 31870686 and NO. 31971705], and Academic Specialty Development Project of Shanghai Academy of Agricultural Sciences (2019-13).

Author contributions

HC and DZ conceived and designed research. LR and SD conducted experiments. YC, YZ and HZ analyzed the data. LR was a major contributor in writing the manuscript. All authors read and approved the final manuscript.

Acknowledgements

The authors thank Prof. Yafei Zhang (Shanghai Jiao Tong University, Shanghai, China) for providing the SWCNTs aqueous solution.

Authors' information

¹ Institute for Agri-food Standards and Testing Technology, Shanghai Academy of Agricultural Sciences, Shanghai 201403, China. ² Department of Landscape Science and Engineering, School of Design, Shanghai Jiao Tong University, Shanghai 200240, China.

References

1. Reed BM. Plant cryopreservation: a continuing requirement for food and ecosystem security. *In Vitro Cell Dev-Pl.* 2017;53:285-8.
2. Sakai A. Survival of the twigs of woody plants at -196°C . *Nature.* 1960;185:392-4.
3. Engelmann F. Cryopreservation: progress and prospects. *In Vitro Cell Dev-Pl.* 2004;40:427-33.
4. Wang MR, Yang W, Zhao L, Li JW, Liu K, Yu JW, Wu YF, Wang QC. Cryopreservation of virus: a novel biotechnology for long-term preservation of virus in shoot tips. *Plant Methods.* 2018;14:47.
5. Sakai A, Hirai D, Niino T. Development of PVS-based vitrification and encapsulation-vitrification protocols. In: Reed BM, editor. *Plant Cryopreservation: a practical guide.* New York: Springer-Verlag; 2008. p. 33-57.
6. Volk GM, Walters C. Plant vitrification solution 2 lowers water content and alters freezing behaviour in shoot tips during cryoprotection. *Cryobiology.* 2006;52:48-61.
7. Ren L, Zhang D, Jiang X, Gai Y, Wang W, Reed BM, Shen X. Peroxidation due to cryoprotectant step is a vital factor for cell survival in *Arabidopsis* cryopreservation. *Plant Sci.* 2013;212:37-47.

8. Benson EE. Free radical damage in stored plant germplasm. International board for plant genetic resources. Italy; 1990.
9. Ren L, Zhang D, Chen G, Reed BM, Shen X, Chen H. Transcriptomic profiling revealed the regulatory mechanism of *Arabidopsis* seedlings response to oxidative stress from cryopreservation. *Plant Cell Rep.* 2015;34:2161-78.
10. Zhang D, Ren L, Chen G, Zhang J, Reed BM, Shen X. ROS-induced oxidative stress and apoptosis-like event directly affect the cell viability of cryopreserved embryogenic callus in *Agapanthus praecox*. *Plant Cell Rep.* 2015;34:1499-513.
11. Gill SS, Tuteja N. Reactive oxygen species and antioxidant machinery in abiotic stress tolerance in crop plants. *Plant Physiol Bioch.* 2010;48:909-30.
12. Volk GM. Application of functional genomics and proteomics to plant cryopreservation. *Curr Genomics.* 2010;11:24-29.
13. Van Breusegem F, Vranová E, Dat JF, Inzé D. The role of active oxygen species in plant signal transduction. *Plant Sci.* 2001;161:405-14.
14. Scandalios JG. Molecular genetics of SOD in plants. In: Scandalios JG, editor. *Oxidative stress and the molecular biology of antioxidant defense*. Cold Spring Harbor: Cold Spring Harbor Laboratory Press; 1997. p. 527-68.
15. Meyer AJ. The integration of glutathione homeostasis and redox signaling. *J Plant Physiol.* 2008;165:1390-403.
16. Husen A, Siddiqi KS. Carbon and fullerene nanomaterials in plant system. *J Nanobiotech.* 2014;12:1-10.
17. Hao BT. Application of nanoparticles in cell cryopreservation. *J Clin Rehabilita Tiss Eng Res.* 2008;12:8140-2.
18. Derui DI, Zhizhu HE, Liu J. Advances in nano-cryopreservation of biological materials. *Ciesc J.* 2011;62:1781-9.
19. Li W, Zhou X, Lv F, Liu B. Preliminary Study on the Mechanism of Promoting the Survival Rate of Vitrified Oocytes with Nano-cryoprotectants. *J Refrigeration* 2014;35:114-8.
20. Han X, Ma HB, Wilson C, Critser JK. Effects of nanoparticles on the nucleation and devitrification temperatures of polyol cryoprotectant solutions. *Microfluidics & Nanofluidics.* 2008;4:357-61.
21. Li W, Zhou XL, Liu BL, Dai JJ, Lv FK, Zhang DF, Xu L. The Effect of Nanoparticle on Vitrification of Porcine GV-Stage Oocytes. *Chinese Journal of Biomedical Engineering.* 2013;32:601-5.
22. Han X, Dai JJ, Zhang SS, Zhang TY, Wu CF, Gu XL, Li X, Xu L, Li FF, Zhang DF. Effects of nonomaterials rhodiola sachalinensis polysaccharide on cryopreservation of boar semen. *Chinese Journal of Veterinary Science.* 2013;33:1937-42.
23. Nadri T, Towhidi A, Zeinoaldini S, Martínez-Pastor F, Mousavi M, Noei R, Tar M, Mohammadi Sangcheshmeh A. Lecithin nanoparticles enhance the cryosurvival of caprine sperm. *Theriogenology.* 2019;133:38-44.

24. Dashtestani F, Ghourchian H, Najafi A. Albumin coated copper-cysteine nanozyme for reducing oxidative stress induced during sperm cryopreservation. *Bioorg Chem.* 2018;80:621-30.
25. Dashtestani F, Ghourchian H, Najafi A. Silver-gold-apoferritin nanozyme for suppressing oxidative stress during cryopreservation. *Mat Sci Eng C-Mater.* 2019;94:831-40.
26. Khalil WA, El-Harairy MA, Zeidan AEB, Hassan MAE. Impact of selenium nano-particles in semen extender on bull sperm quality after cryopreservation. *Theriogenology.* 2019;126:121-7.
27. Zhang J, Yang K, Zhang H, Liang X. Application Status and Prospect of Carbon-Based Nanomaterials in Biomedical Field. *Prog Chem.* 2013;25:397-408.
28. Chen SM, Ren L, Zhang D, Zhang YF, Shen XH. Carbon nanomaterials enhance survival of *Agapanthus praecox* callus after cryopreservation by vitrification. *Cryo-Lett.* 2017;38:125-36.
29. Wang ZC, Deng XX. Cryopreservation of shoot-tips of citrus using vitrification: effect of reduced form of glutathione. *Cryo-Lett.* 2004;25:43-50.
30. Uchendu EE, Leonard SW, Traber MG, Reed BM. Vitamins C and E improve regrowth and reduce lipid peroxidation of blackberry shoot tips following cryopreservation. *Plant Cell Rep.* 2010;29:25-35.
31. Uchendu EE, Muminova M, Gupta S, Reed BM. Antioxidant and anti-stress compounds improve regrowth of cryopreserved *Rubus* shoot tips. *In Vitro Cell Dev Biol-Pl.* 2010;46:386-93.
32. Zhao Y, Qi LW, Wang WM, Saxena PK, Liu CZ. Melatonin improves the survival of cryopreserved callus of *Rhodiola crenulata*. *J Pineal Res.* 2011;50:8-88.
33. Chen GQ, Ren L, Zhang D, Shen XH. Glutathione improves survival of cryopreserved embryogenic calli of *Agapanthus praecox* subsp. *orientalis*. *Acta Physiol Plant.* 2016;38:250
34. Hu MJ. Cryopreservation of *Arabidopsis thaliana* suspension cells and the role of Abscisic Acid in stress signaling transduction pathways. Dissertation, Zhejiang University, 2003.
35. Lardet L, Martin F, Dessailly F, Carron MP, Montoro P. Effect of exogenous calcium on post-thaw growth recovery and subsequent plant regeneration of cryopreserved embryogenic calli of *Hevea brasiliensis* (Müll. Arg.). *Plant Cell Rep.* 2007;26:559-69.
36. Ryynanen L, Haggman H. Recovery of cryopreserved silver birch shoot tips is affected by the pre-freezing age of the cultures and ammonium substitution. *Plant Cell Rep.* 2001;20:354-60.
37. Decruse SW, Seeni S, Nair GM. Preparative procedures and culture medium affect the success of cryostorage of *Holostemma annulare* shoot tips. *Plant Cell Tiss Org.* 2004;76:179-82.
38. Kami D, Kasuga J, Arakawa K, Fujikawa S. Improved cryopreservation by diluted vitrification solution with supercooling-facilitating flavonol glycoside. *Cryobiology.* 2008;57:242-5.
39. Wang HY, Inada T, Funakoshi K, Lu SS. Inhibition of nucleation and growth of ice by poly (vinyl alcohol) in vitrification solution. *Cryobiology.* 2009;59:83-9.
40. Johnston JW, Harding K, Benson EE. Antioxidant status and genotypic tolerance of *Ribes* in vitro cultures to cryopreservation. *Plant Sci.* 2007;172:524-34.
41. Fang JY, Wetten A, Johnston J. Headspace volatile markers for sensitivity of cocoa (*Theobroma cacao* L.) somatic embryos to cryopreservation. *Plant Cell Rep.* 2008;27:453-61.

42. Whitaker C, Beckett RP, Minibayeva FV, Kranner I. Production of reactive oxygen species in excised, desiccated and cryopreserved explants of *Trichilia dregeana* Sond. S Afr J Bot. 2010;76:112-8.
43. Skyba M, Petijová L, Košuth J, Koleva DP, Ganeva TG, Kapchina-Toteva VM, Čellárová E. Oxidative stress and antioxidant response in *Hypericum perforatum* L. plants subjected to low temperature treatment. J Plant Physiol. 2012;169:955-64.
44. Zhang D, Ren L, Chen GQ, Zhang J, Reed BM, Shen XH. ROS-induced oxidative stress and apoptosis-like event directly affect the cell viability of cryopreserved embryogenic callus in *Agapanthus praecox*. Plant Cell Rep. 2015;34:1499-13.
45. Chen GQ, Ren L, Zhang J, Reed BM, Zhang D, Shen XH. Cryopreservation affects ROS-induced oxidative stress and antioxidant response in *Arabidopsis* seedlings. Cryobiology. 2015;70:38-47.
46. Jia MX, Shi Y, Di W, Jiang XR, Xu J, Liu Y. ROS-induced oxidative stress is closely related to pollen deterioration following cryopreservation. In Vitro Cell Dev Biol-Pl. 2017;53:433-9.
47. Huang B, Zhang JM, Chen XL, Xin X, Yin GK, He JJ, Lu XX, Zhou YC. Oxidative damage and antioxidative indicators in 48 h germinated rice embryos during the vitrification-cryopreservation procedure. Plant Cell Rep. 2018;37:1325-42.
48. Vianna MG, Garcia RO, Mansur E, Engelmann F, Pacheco G. Oxidative stress during the cryopreservation of *Passiflora suberosa* L. shoot tips using the V-Cryo-plate technique: determination of the critical stages of the protocol. Plant Cell Tiss Org. 2019;139:369-79.
49. Benson EE, Bremner D. Oxidative stress in the frozen plant: a free radical point of view. In: Fuller BJ, Lane N, Benson EE, editors. Life in the frozen state. Boca Raton: CRC Press; 2004. p. 205-41.
50. Benson EE, Lynch PT, Jones J. The detection of lipid peroxidation products in cryoprotected and frozen rice cells—consequences for post thaw survival. Plant Sci. 1992;85:107-14.
51. Varghese B, Naithani SC. Oxidative metabolism-related changes in cryogenically stored neem (*Azadirachta indica* A. Juss) seeds. J Plant Physiol. 2008;165:755-65.
52. Prudente DO, Paiva R, Nery FC, Paiva PDO, Alves JD, Máximo WPF, Silva LC. Compatible solutes improve regrowth, ameliorate enzymatic antioxidant systems, and reduce lipid peroxidation of cryopreserved *Hancornia speciosa* Gomes lateral buds. In Vitro Cell Dev Biol-Pl. 2017;53:352-62.
53. Poobathy R, Sinniah UR, Xavier R, Subramaniam S. Catalase and superoxide dismutase activities and the total protein content of protocorm-like bodies of *Dendrobium Sonia-28* subjected to vitrification. Appl Biochem Biotechnol. 2013;170:1066-79.
54. Fleck RA, Benson EE, Bremner DH, Day JG. A comparative study of antioxidant protection in cryopreserved unicellular algae *Euglena gracilis* and *Haematococcus pluvialis*. Cryo-lett. 2003;24:213-28.
55. Liu SJ. The effects of graphene on the germination and seedling growth in rice. Yangtze University, 2013.
56. Noctor G, Foyer CH. Ascorbate and glutathione: keeping active oxygen under control. Annu Rev plant Physiol Plant Mol Biol. 1998;49:249-79.

57. Alscher RG, Donahue JL, Cramer CL. Reactive oxygen species and antioxidants: relationships in green cells. *Physiol Plant*. 1997;100:224-33.
58. Dat JF, Vandenabeele S, Vranová E, Van Montagu M, Inzé D, Van Breusegem F. Dual action of the active oxygen species during plant stress responses. *Cell Mol Life Sci*. 2000;57:779-95.
59. Mittler R. Oxidative stress, antioxidants and stress tolerance. *Trends Plant Sci*. 2002;7:405-10.
60. Vranová E, Atichartpongkul S, Villarroel R, Van Montagu M, Inzé D, Van Camp W. Comprehensive analysis of gene expression in *Nicotiana tabacum* leaves acclimated to oxidative stress. *Proc Natl Acad Sci USA*. 2002;99:10870-5.
61. Mittler R, Vanderauwera S, Gollery M, Van Breusegem F. Reactive oxygen gene network of plants. *Trends Plant Sci*. 2004;9:490-8.
62. Bowler C, Fluhr R. The role of calcium and activated oxygens as signals for controlling cross-tolerance. *Trends Plant Sci*. 2000;5:241-6.
63. Knight H, Knight MR. Abiotic stress signalling pathways: specificity and cross-talk. *Trends Plant Sci*. 2001;6:262-7.
64. Coelho SM, Taylor AR, Ryan KP, Sousa-Pinto I, Brown MT, Brownlee C. Spatiotemporal patterning of reactive oxygen production and Ca²⁺ wave propagation in *Fucus rhizoid* cells. *Plant Cell*. 2002;14:2369-81.
65. Rentel MC, Lecourieux D, Ouaked F, Usher SL, Petersen L, Okamoto H, Knight H, Peck SC, Grierson CS, Hirt H, Knight MR. OXI1 kinase is necessary for oxidative burst-mediated signalling in *Arabidopsis*. *Nature*. 2004;427:858-61.
66. Kovtun Y, Chiu WL, Tena G, Sheen J. Functional analysis of oxidative stress activated mitogen-activated protein kinase cascade in plants. *Proc Natl Acad Sci USA*. 2000;97:2940-5.
67. Apel K, Hirt H. Reactive oxygen species: metabolism, oxidative stress, and signal transduction. *Annu Rev Plant Biol*. 2004;55:373-99.
68. Dat JF, Pellinen R, Beeckman T, Van De Cotte B, Langebartels C, Kangasjärvi J, Inzé D, Van Breusegem F. Changes in hydrogen peroxide homeostasis trigger an active cell death process in tobacco. *Plant J*. 2003;33:621-32.
69. Rizhsky L, Davletova S, Liang HJ, Mittler R. The zinc finger protein Zat12 is required for cytosolic ascorbate peroxidase 1 expression during oxidative stress in *Arabidopsis*. *J Biol Chem*. 2004;279:11736-43.
70. Zaytseva O, Neumann G. Carbon nanomaterials: production, impact on plant development, agricultural and environmental applications. *Chem Biol Technol Agric*. 2016;3:17.
71. Giraldo JP, Landry MP, Faltermeier SM, McNicholas TP, Iverson NM, Boghossian AA, Reuel NF, Hilmer AJ, Sen F, Brew JA, Strano MS. Plant nanobionics approach to augment photosynthesis and biochemical sensing. *Nat Mater*. 2014;13:400-8.
72. Elena S, Gusev A, Zaytseva O, Sheina O, Tkachev A, Kuznetsova E, Lazareva E, Onishchenko G, Feofanov A, Kirpichnikov M. Uptake and accumulation of multiwalled carbon nanotubes change the

- morphometric and biochemical characteristics of *Onobrychis arenaria* seedlings. *Fron Chem Sci Eng*. 2012;6:132-8.
73. Khodakovskaya MV, De Silva K, Biris AS, Dervishi E, Villagarcia H. Carbon nanotubes induce growth enhancement of tobacco cells. *ACS Nano*. 2012;6:2128-35.
 74. Khodakovskaya M, De Silva K, Nedosekin D, Dervishi E, Biris AS, Shashkov EV, Galanzha EI, Zharov VP. Complex genetic, photothermal, and photoacoustic analysis of nanoparticle-plant interactions. *Proc Natl Acad Sci USA*. 2011;108:1028-33.
 75. Lahiani MH, Dervishi E, Chen J, Nima Z, Gaume A, Biris AS, Khodakovskaya MV. Impact of carbon nanotube exposure to seeds of valuable crops. *ACS Appl Mater Inter*. 2013;5:7965-73.
 76. Tan X, Lin C, Fugetsu B. Studies on toxicity of multi-walled carbon nanotubes on suspension rice cells. *Carbon*. 2009;47:3479-87.
 77. Lin C, Fugetsu B, Su Y, Watari F. Studies on toxicity of multi-walled carbon nanotubes on *Arabidopsis* T87 suspension cells. *J Hazard Mater*. 2009;170:578-83.
 78. Aslani F, Bagheri S, Julkapli NM, Juraimi AS, Hashemi FSG, Baghdadi A. Effects of engineered nanomaterials on plants growth: An overview. *Sci World J*. 2014:641759.
 79. Husen A, Siddiqi KS. Carbon and fullerene nanomaterials in plant system. *J Nanobiotechnol*. 2014;12:16.
 80. Wang Y, Fan X, Zhang D, Shen X. Regeneration of *Agapanthus praecox* ssp. *orientalis* 'big blue' via somatic embryogenesis. *Propag Ornament Plants*. 2012;12:148-54.
 81. Yang Z, Sheng J, Lv K, Ren L, Zhang D. Y₂SK₂ and SK₃ type dehydrins from *Agapanthus praecox* can improve plant stress tolerance and act as multifunctional protectants. *Plant Sci*. 2019;284:143-60.

Additional File

Additional file 1: Table S1. Primers sequences of qRT-PCR

Figures




Figure 1

Effects of different exogenous substances on survival rate of *Agapanthus praecox* EC cryopreservation. Concentration of exogenous substances added are: 0.1 g/L SWCNTs, 0.08 mM GSH, 1 μ M ABA, 0.1 μ M melatonin, 1 mM AsA, 10 mM betaine, 1 mM CaCl₂, 6 mM LA, 6 mM PVA, 3% PVP. Values with different small letters are significantly different between different treatment at 0.05 level.




Figure 2

Cell ultrastructure observation of *Agapanthus praecox* during SWCNTs-added cryopreservation. (a-d) dehydration stage, (e-h) dilution stage. Vesicle was defined as VES.




Figure 3

Physiological indices detection of SWCNTs added in cryopreservation system. Physiological indices were determined with at least three repeats. Bars represent means and standard deviation over triplicate detection. Values with different lowercase letters are significantly different among samples in the control group at 0.05 level. Values with different uppercase letters are significantly different among samples in the SWCNTs group at 0.05 level. * are significantly different between different groups in the same stage at 0.05 level, and ** are highly significantly different between different groups in the same stage at 0.01 level. The CK group is the cryopreservation without adding SWCNTs, and the CNT group is the SWCNTs-added cryopreservation. CK, untreated; PC, after preculture; DH, after dehydration; DL, after dilution.




Figure 4

Real-time PCR quantitative analysis of ROS signal transduction and antioxidant system related genes between the control and the improved (added with SWCNTs) cryopreservation. Bars show means of gene expression level and standard deviation over triplicate detections. Values with different lowercase letters are significantly different among samples in the control group at 0.05 level. Values with different uppercase letters are significantly different among samples in the SWCNTs group at 0.05 level. * are significantly different between different groups in the same stage at 0.05 level, and ** are highly significantly different between different groups in the same stage at 0.01 level. The CK group is the cryopreservation without adding SWCNTs, and the CNT group is the SWCNTs-added cryopreservation. CK, untreated; PC, after preculture; DH, after dehydration; DL, after dilution.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Additionalfile1.docx](#)