

Quality of life, Social Support and Self-Efficacy in Women After A Miscarriage

Grażyna Iwanowicz-Palus

Medical University of Lublin

Mariola Mróz (✉ mariolamroz2015@gmail.com)

Obstetrics and Gynecology Department and Clinic, Cardinal S. Wyszyński Regional <https://orcid.org/0000-0002-4779-5125>

Agnieszka Bień

Medical University of Lublin

Research

Keywords: miscarriage, pregnancy loss, quality of life, social support, general self-efficacy

Posted Date: May 13th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-28515/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published on January 7th, 2021. See the published version at <https://doi.org/10.1186/s12955-020-01662-z>.

Abstract

Background Pregnancy loss is typically experienced as a traumatic, critical event, which may lead to secondary psychological health disorders. Its burden involves both the experience of loss and related medical issues, which are associated with pain, hospitalization, limitation in one's social roles, decreased sense of security, and changes in one's perceived quality of life. The purpose of the present study was to evaluate levels of quality of life (QoL), social support and self-efficacy among women who had suffered a miscarriage.

Methods The study was performed using a diagnostic survey method with questionnaires administered to 610 patients hospitalized due to spontaneous pregnancy loss in hospitals in Lublin (Poland). The instruments used were: the Berlin Social Support Scales (BSSS), the Generalized Self-Efficacy Scale (GSES), the WHOQoL-BREF questionnaire, and a standardized interview questionnaire.

Results Respondents rated their overall QoL (3.90 points) higher than their overall perceived health (3.66). In terms of social support, the highest scores were noted for perceived available instrumental support ($M = 3.78$), perceived available emotional support ($M = 3.68$) and actually received support ($M = 3.60$). The mean generalized self-efficacy score among the women after pregnancy loss was 30.29. Respondents' QoL was significantly correlated with multiple social support subscales and self-efficacy ($p < 0.05$).

Conclusions Women after a miscarriage perceive their overall QoL as better than their overall health, while reporting the poorest QoL in the psychological domain. They also have a high level of self-efficacy. Regarding the types of social support, perceived available support, both instrumental and emotional, and actually received support was rated highly. Social support and self-efficacy contributed to better perceived QoL among the respondents.

Background

According to the World Health Organization (WHO), the term "fetal death" refers to the intrauterine death of a fetus at any time during pregnancy. In some countries, the concepts of miscarriage and intrauterine fetal death/demise are distinguished, with differences in definitions mainly resulting from advances in medical science, increased survival rates for newborns with an extremely low birth weight, and economic factors [1, 2]. The Royal College of Obstetricians and Gynecologists applies the term "miscarriage" to pregnancy losses occurring until 24 weeks of gestation [2]. In Poland, a miscarriage is defined as the loss of pregnancy before 22 weeks of gestation or when the weight of the dead fetus does not exceed 500 g. The incidence of miscarriage is correlated with gestational age, and is estimated at approx. 15%, out of which 80% occur within the first trimester. Recurrent miscarriages account for 1–2% of cases [1, 2, 3, 4].

Pregnancy loss is typically experienced as a traumatic, critical event, which may lead to secondary psychological health disorders. Its burden involves both the experience of loss and related medical issues, which are associated with pain, hospitalization, limitation in one's social roles, decreased sense of security, and changes in one's perceived quality of life (QoL) [5, 6, 7, 8, 9, 10, 11].

The process of coping with difficult situations is affected by a number of factors, including self-efficacy, defined as an individual's belief about their ability to achieve their objectives in a specific life situation [12, 13]. Individuals with a low level of self-efficacy may focus on their deficiencies, and tend to experience strong emotions, anxiety or even depression. In turn, a high level of self-efficacy is positively correlated with optimism, satisfaction with life, and a healthy distance to life, as well as with increased capacity to overcome difficulties or stress, and lower stress levels [12, 13, 14]. Alongside one's personal competences, sense of control and coherence, self-efficacy is another personal resource that influences the effectiveness of support [13, 14, 15].

Social support is defined as assistance available to an individual in difficult situations. Broadly defined, it refers to issues of social integration and interpersonal relationships that have an impact on the individual, allowing them to feel surrounded by people upon whom they can rely. Being with another in a problematic situation reduces stress and makes one feel more secure and in control. Social support is a significant contributor to the maintenance of health, prevention of disease, and success of treatment. It is also a recognized factor positively affecting an individual's QoL [16].

The communication between medical personnel and a patient who has lost a pregnancy is a special kind of social interaction, involving an exchange of emotions and instruments for action. Recognizing a hospitalized patient's individuality and complexity has a positive impact on their relationship with medical personnel and contributes to greater care effectiveness and lower incidence of psycho-emotional disorders associated with one's health situation [17].

In modern medicine, the objective is not just to treat patients, but also to improve their well-being and QoL. Hence the focus on achieving extramedical objectives within the treatment process, to help patients function normally not only in the physical domain, but also in the psychological and social ones.

In medical science, research on QoL and social support provides knowledge on institutional efforts that need to be undertaken for the sake of patients, and on ways of supporting and empowering patients in their difficult circumstances [17]. This reflects a holistic approach to the subject of care and helps identify factors with the most significant impact. Such research contributes to the continued improvement of standards of care in specific conditions, and should therefore be integrated with treatment [18].

Considering the impact and complexity of pregnancy loss, with its multiple health-related, psycho-social and economic consequences, the present study aimed at evaluating the levels of QoL, social support and self-efficacy in women after a miscarriage.

Methods

Study design and population

The study was performed between August 2016 and February 2019, and included 610 patients hospitalized due to spontaneous pregnancy loss (until 22 weeks of gestation) in the following hospitals in Lublin, Poland: the Independent Public Teaching Hospital no. 1, the Independent Public Teaching Hospital no. 4, the Cardinal Stanisław Wyszyński Regional Specialist Hospital, and the John of God Independent Public Regional Hospital. Inclusion criteria were as follows: loss of a singleton pregnancy until 22 weeks, consent to participate in the study, age above 18 years, and no psycho-physical disorders. Patients undergoing psychotherapy or psychiatric treatment, and patients in a poor psychological condition were excluded.

Data collection

The survey questionnaire was administered to each patient on the last day of her hospitalization. Before communicating with each patient, information was obtained from the medical personnel regarding the stage of treatment, duration of hospitalization, and the psychological and physical condition of the patient. The study was approved by the Bioethics Committee of the Lublin Medical University (KE-0254/221/2016) and the managers and department heads in each hospital where the study was performed. Respondents were informed that their participation in the survey was anonymous and strictly voluntary, and that the results would only be used for research purposes. Out of the 645 patients recruited for the study, 610 returned fully and correctly completed questionnaires, and were included in subsequent statistical analyses. The response rate was 94.57%.

Assessments

The study used a diagnostic survey with questionnaires. The instruments used were: the WHOQoL – BREF questionnaire, the Berlin Social Support Scales (BSSS), the Generalized Self-Efficacy Scale (GSES), and a standardized interview questionnaire used to collect respondents' characteristics.

Instruments

The World Health Organization Quality of Life Test – Bref (WHOQoL – BREF) is used to determine a patient's QoL profile in the physical, psychological, social and environmental domains, as well as overall QoL and perceived health. Respondents rate 26 items on a scale from altogether negative to altogether positive. The scoring is positive, i.e. higher scores indicate a better QoL. The questionnaire's internal consistency coefficient (Cronbach's α) is 0.54–0.91 for individual domains; for the whole scale it is 0.92 in healthy individuals, and 0.95 in ill individuals [19, 20].

The Berlin Social Support Scales (BSSS) are a set of independent instruments measuring the behavioral and cognitive dimensions of social support: available and received support, protective buffering, need for support, and support seeking. Respondents rate each item on a scale of 1 to 4, where 1 indicates complete disagreement and 4 complete agreement with the statement. Higher scores indicate more social support. In the present study, results are presented as means for each scale. The Cronbach's α for the questionnaire is 0.80 [21].

The Generalized Self-Efficacy Scale (GSES) is an instrument measuring the strength of an individual's belief in their capacity to overcome difficulties and obstacles. Respondents rate 10 statements on a scale of 1 to 4 (*1 – disagree, 2 – somewhat disagree, 3 – somewhat agree, 4 – agree*). The total score, converted into standardized sten units, reflects the overall level of self-efficacy. Low self-efficacy is indicated by scores up to 24 points (sten 1–4), moderate – by scores between 25 and 29 points (sten 5–6), and high – by scores of 30 or more points (sten 7–10). Cronbach's α for the instrument is 0.85 [12, 22].

Statistical analysis

Statistical analysis of the material was performed using the IBM SPSS Statistics (v. 21) software. Quantitative variables were described using mean (M), median (Me), standard deviation (SD), and minimum (Min) and maximum (Max) values. For qualitative variables, numbers and percentages in each category were reported. A series of regression analyses were also performed using the enter method, with the physical, psychological social, and environmental domains of the WHOQOL-BREF as the dependent variables. The independent variables included: GSES score, perceived emotional support, perceived instrumental support, need for support, support seeking, and actually received support. The study used a significance threshold of $p < 0.05$.

Results

Table 1 shows respondents' characteristics. Most were women aged between 26 and 30 (32.6%), having completed college/university education (61.1%), living in province capitals (47.7%), married (80.5%), performing white-collar work (48.7%), and reporting a good socio-economic standing (60.8%). Moreover, most respondents had had children before (59.0%), and had miscarried for the first time (59.2%).

Table 1
Participants' characteristics

Participants' characteristics		<i>n</i>	%
Age	< 25 y/o	77	12.6
	26–30 y/o	199	32.6
	31–35 y/o	195	32.0
	> 35 y/o	139	22.8
Education	Other than college/university	237	38.9
	College/university	373	61.1
Residence	Urban – province capital	291	47.7
	Urban – other	116	19.0
	Rural	203	33.3
Relationship status	Married	491	80.5
	Single	119	19.5
Professional activity	Professionally inactive	135	22.1
	White-collar work	297	48.7
	Blue-collar work	178	29.2
Self-reported financial standing	Very good	93	15.2
	Good	371	60.8
	Moderate	146	24.0
Having children	No	250	41.0
	Yes	360	59.0
History of pregnancy loss	First pregnancy loss	361	59.2
	1 previous pregnancy loss	175	28.7
	≥ 2 previous pregnancy losses	74	12.1

The mean overall QoL score was 3.90 ± 0.77 , and the mean overall perceived health score was 3.66 ± 0.76 . In terms of specific domains, QoL was highest in the social domain (17.04 ± 2.54), and lowest in the psychological domain (14.91 ± 2.45). The highest scores on the social support scales were obtained for perceived available instrumental support (3.78 ± 0.43), and the lowest for protective buffering (1.89 ± 0.68). The mean generalized self-efficacy (GSES) score among the studied women after a miscarriage was 30.29 ± 4.01 (Table 2).

Table 2
Quality of life, social support and self-efficacy scores among women after a miscarriage

		M	Me	SD	Min	Max
QoL	Overall quality of life	3.90	4.00	0.77	1.00	5.00
	Perceived general health	3.66	4.00	0.76	1.00	5.00
	Physical domain	15.96	16.00	2.14	8.57	20.00
	Psychological domain	14.91	15.33	2.45	6.67	20.00
	Social relationships domain	17.04	17.33	2.54	4.00	20.00
	Environment domain	15.90	16.00	2.04	8.50	20.00
BSSS subscales	Perceived available emotional support	3.68	3.75	0.44	1.00	4.00
	Perceived available instrumental support	3.78	4.00	0.43	1.00	4.00
	Need for support	3.16	3.25	0.57	1.00	4.00
	Support seeking	3.09	3.20	0.66	1.00	4.00
	Actually received support	3.60	3.80	0.40	1.00	4.00
	Protective buffering	1.89	1.83	0.68	1.00	4.00
GSES		30.29	30.00	4.01	17.00	40.00
<i>M – mean; SD – standard deviation; Me – median; QoL – Quality of Life;</i>						
<i>BSSS – Berlin Social Support Scales; GSES – Generalized Self-Efficacy Scale</i>						

Table 3 shows the results of regression analysis for the specific QoL (WHOQOL-BREF) domains, social support (BSSS), and generalized self-efficacy (GSES). QoL in the physical domain is positively correlated with perceived available instrumental support ($\beta = 0.152, p < 0.001$) and self-efficacy ($\beta = 0.280, p < 0.001$). The model that was tested accounted for 12% of variance in the physical QoL score.

The psychological QoL domain was analyzed next. This factor was positively correlated with perceived available emotional support ($\beta = 0.235, p < 0.001$), actually received support ($\beta = 0.110, p = 0.005$), and self-efficacy ($\beta = 0.348, p < 0.001$). Negative correlations were found with need for support ($\beta = -0.084, p < 0.027$) and protective buffering ($\beta = -0.079, p = 0.031$). This model accounted for 26% of variance in the psychological QoL score.

Analyses demonstrated that: perceived available emotional support ($\beta = 0.209, p < 0.001$), actually received support ($\beta = 0.419, p < 0.001$), protective buffering ($\beta = -0.076, p = 0.029$), and self-efficacy ($\beta = 0.113, p < 0.001$) were significantly correlated with the social QoL domain. The model accounted for 35% of variance in the social QoL score.

QoL in the environmental domain was positively correlated with perceived available emotional ($\beta = 0.153, p = 0.006$) and instrumental support ($\beta = 0.128, p = 0.026$), actually received support ($\beta = 0.135, p = 0.001$), and self-efficacy ($\beta = 0.281, p < 0.001$). A negative correlation was found with need for support ($\beta = -0.081, p = 0.036$). The model accounted for 23% of variance in the environmental QoL score (Table 3).

Table 3
Regression analysis for WHOQOL-BREF, BSSS, and GSES scores in women after a miscarriage

WHOQOL-BREF Domains															
Predictor	Physical Health				Psychological				Social Relationships				Environment		
	F = 39.027; df = 2; p < 0.001				F = 42.469; df = 5; p < 0.001				F = 79.518; df = 4; p < 0.001				F = 36.715; df = 3; p < 0.001		
	B	β	t	p	B	β	t	p	B	β	t	p	B	β	p
Constant	8.599		9.712	< 0.001	2.971		2.488	0.013	1.438		1.316	< 0.001	5.143		
BSSS Subscales	Perceived available emotional support	-0.033	0.568	-0.023	0.428	1.299	0.235	5.918	< 0.001	1.199	0.209	5.727	< 0.001	0.705	0.149
	Perceived available instrumental support	0.753	0.152	3.947	< 0.001	0.027	0.632	0.020	0.385	0.100	0.058	0.077	0.386	0.604	0.149
	Need for support	-0.050	0.221	-0.050	0.879	-0.362	-0.084	-2.221	0.027	0.015	0.674	0.017	0.859	-0.290	0.149
	Support seeking	-0.068	0.095	-0.068	0.879	0.066	0.184	0.054	0.498	0.023	0.521	0.026	0.862	0.023	0.149
	Actually received support	0.018	0.683	0.017	0.758	0.672	0.110	2.791	0.005	2.655	0.419	11.367	< 0.001	0.688	0.149
	Protective buffering	-0.071	0.070	-0.074	0.955	-0.288	-0.079	-2.156	0.031	-0.285	-0.076	-2.185	0.029	-0.054	0.149
GSES	0.149	0.280	7.239	< 0.001	0.212	0.348	9.334	< 0.001	0.072	0.113	3.355	0.001	0.142	0.149	
Regression equation	Y = 8,599 + 0,149 × GSES + 0,753 × BSSSI Instr				Y = 2.971 + 0.212 × GSES + 1.299 × BSSSI Emo – 0.362 × BSSSII + 0,672 × BSSSIV – 0,288 × BSSSV				Y = 1.438 + 0.072 × GSES + 1.199 × BSSS Emo + 2.655 × BSSSIV – 0.285 × BSSSV				Y = 5.143 + 0.149 × GSES + 0.705 × BSSS Emo – 0.290 × BSSS Instr		

WHOQOL-BREF – The World Health Organization Quality of Life-Test Bref; BSSS – Berlin Social Support Scales; GSES – Generalized Self-Efficacy Scale;

BSSS I Emo – Perceived available Emotional Support; BSSS I Instr - Perceived available Instrumental Support; BSSS II - Need for Support; BSSS III - Support Seeking; BSSS IV - Actually Received Support; BSSS V - Protective Buffering Support

Discussion

In modern medicine, the objective is not just to treat patients, but also to improve their well-being and QoL, hence the increasing interest in research on social support and QoL experienced by patients with a variety of conditions. Such a comprehensive assessment is especially important in patients who have lost a pregnancy, as this experience entails a number of consequences, not only physical, but also psychological and social, and at times, even financial. Each patient has her own needs and beliefs that affect her individual perception of her health and living situation. Therefore, we undertook to evaluate the levels of QoL, social support, and self-efficacy in this patient group [17, 18].

Our findings are to some extent consistent with those reported by other researchers studying the subject. Like patients with hyperglycemia in pregnancy and those treated for polycystic ovary syndrome, the women after a miscarriage who were studied rated their overall QoL higher than their overall perceived health [23, 24].

The social relationships domain was scored the highest, as in studies by Couto et al. and Tavoli et al. regarding women after pregnancy loss [25, 26]. The high scores in the physical domain were also consistent with the report by Tavoli et al. [26]. Interestingly, in the QoL self-assessment by physically active women during uncomplicated pregnancy, the scores in the physical domain were next to lowest, while the highest scores were reported in the psychological domain [27]. In the present study, women who had lost a pregnancy reported lower QoL in the psychological domain. Respondents' psychological well-being was also rated lower than other QoL aspects in the studies by Couto et al. and Tivoli et al., although these authors put more emphasis on the limitations in the performance of social roles, associated with emotional difficulties [25, 26]. In turn, Song et al., analyzing the long-term effects of a child's death on the parents, reported much poorer QoL than in the case of non-bereaved individuals [28]. Other authors also highlighted considerable differences in QoL between pregnant subjects with no history of obstetric problems and ones after a pregnancy loss, with the latter group having lower QoL [25, 26, 29, 30].

QoL is also significantly associated with social support, which is a major contributor to a patient's health and treatment success [16]. In the group of women after pregnancy loss, perceived available support, both instrumental and emotional, and actually received support was rated highly. The present statistical

analysis results are comparable to those reported by Iwanowicz-Palus G. et al. in a group of patients with hyperglycemia in pregnancy [23].

The association between social support and satisfaction with life was previously demonstrated e.g. by Strine et al. in their study on the US population. They found that overall dissatisfaction with life increased as the level of social support decreased [31]. Gul et al. (2018), Emmanuel et al. (2012) and Shishehgar et al. (2013) also reported a positive correlation between social support and QoL in pregnant women [32, 33, 34].

A literature review demonstrates that social bonds have a beneficial impact on various aspects of an individual's psycho-physical condition, including reduced health-related stress, anxiety, and depressive symptoms [31, 34, 35, 36, 37, 38]. The results of the present study indicate an association between perceived available emotional support and actually received social support, on the one hand, and better QoL in the psychological domain on the other. The quality of social support received from various sources and the professional demeanor of medical personnel also affect the patient's psychological well-being and satisfaction with life.

Pregnancy loss is a stressful event that may prompt a psychological crisis [8, 10]. The process of coping with a difficult health-related situation is shaped by a number of factors, one of which is self-efficacy [12]. A literature review does not reveal many reports on the subject in the field of gynecology. It is, however, a highly consequential topic, as an individual's expectations and beliefs have a significant impact on their actions, as well as on their physical and psychological health [12, 14, 39].

The patients who had experienced a miscarriage and who participated in our survey had a higher level of generalized self-efficacy than that found in a standardization group for the Polish population, which was moderate (this included women with pregnancy complications, post-menopausal women, diabetic patients, patients on dialysis, and patients with a history of myocardial infarction) [12]. This suggests that the patients studied here were involved in their treatment process and actively coping with their difficult situation. Similar results were found in groups of patients after a mastectomy and pregnant women with hyperglycemia [12, 23].

Self-efficacy may affect QoL or satisfaction with life. It is also one of the cognitive factors that affect the way individuals manage stress [12, 13, 14, 20].

The available publications suggest that building self-efficacy by strengthening a patient's sense of control and ability to manage a given life situation may counteract negative emotions or exacerbations of depressive symptoms [40, 41, 42, 43]. An analysis by Nikcevic et al. demonstrated a significant association between more self-efficacy and less anxiety and depression in women who had lost a pregnancy [39]. The present analysis confirmed the impact of self-efficacy on all QoL domains in the women studied, and the strongest positive correlation was found for the psychological QoL domain.

As pregnancy loss may have negative psychological health consequences, it is worth considering strategies to enhance self-efficacy, especially among women found to have lower levels of this resource [6, 9, 10, 11]. The present findings may contribute to a better understanding of care involving not just professional medical interventions, but also the provision of adequate social support. Providing support and improving the self-efficacy of women after pregnancy loss seem extremely important for patients' psycho-social well-being and QoL, on a level with effective medical management. In practice, these aspects should be included among the priorities of medical personnel's daily work.

Strengths and limitations of the study

The strengths of our study include the sample size (610 patients) and personal communication with each respondent. As we used standardized instruments, other researchers interested in issues related to pregnancy loss will be able to compare results, continue in-depth research and monitor changes. The available studies on women after a miscarriage were typically performed weeks or months after the event. We investigated patients during hospitalization, as this is when the coping process typically begins.

One difficulty was associated with the personal administration of surveys to the respondents, as during their 3–6 days of hospitalization, they also underwent multiple intensive diagnostic and treatment procedures. Therefore, to ensure the highest possible quality and reliability of our study, information on the stage of treatment, duration of hospitalization, and the clinical condition of the patient was obtained from medical personnel before contact with each patient. The study instrument was given to each patient on the last day of her hospitalization, having ascertained that her treatment had been completed and her psycho-physical condition had been stabilized.

Our study is limited by its cross-sectional design, as it does not allow for identifying any causal relationships between quality of life, social support and self-efficacy in women after a miscarriage.

Conclusions

1. Women after a miscarriage rate their overall QoL higher than their overall perceived health. In terms of specific domains, their QoL is highest in the social domain, and lowest in the psychological domain.
2. Among the various types of social support, perceived available support, both instrumental and emotional, and actually received support was rated the highest.
3. Women after a miscarriage have a high level of generalized self-efficacy.
4. Social support and self-efficacy affects the perceived QoL of women after a miscarriage. Perceived available emotional support and actually received support was positively correlated with QoL in the environmental, social and psychological domains, while perceived available instrumental support was likewise correlated with the environmental and physical domains. More need for support was associated with poorer QoL in the psychological and

environmental domains. Protective buffering was negatively correlated with QoL in the psychological and social domains. Self-efficacy was a predictor of QoL in all the domains studied.

Abbreviations

BSSS

Berlin Social Support Scales

BSSS I Emo

Perceived available Emotional Support

BSSS I Instr

Perceived available Instrumental Support

BSSS II

Need for Support; BSSS III - Support Seeking

BSSS IV

Actually Received Support

BSSS V

Protective Buffering Support

GSES

Generalized Self-Efficacy Scale

M

mean

Me

median

QoL

Quality of Life

SD

standard deviation

WHOQOL-BREF

The World Health Organization Quality of Life-Test Bref

Declarations

Availability of data and materials

Data from this study will be shared with qualified investigators upon reasonable request for scientific purposes.

Acknowledgements

We would like to express our deepest gratitude to the heads of health care services (hospital care) that hosted the study, and to all the women who agreed to participate.

Funding

This research received no external funding.

Author information

GI-P – Chair and Department of Development in Midwifery, Faculty of Health Sciences, Medical University of Lublin, Lublin, Poland

MM – Obstetrics and Gynecology Department and Clinic, Cardinal S. Wyszyński Regional Specialist Hospital in Lublin, Poland

AB – Chair and Department of Development in Midwifery, Faculty of Health Sciences, Medical University of Lublin, Lublin, Poland

Contributions

GI-P: study design, analyse the research material, wrote the paper, performed the manuscript review. MM: data collection, analyse the research material, manuscript preparation, wrote the paper. AB: statistical analysis, interpreted the data, wrote the paper, and performed the manuscript review. All authors read and approved the final manuscript.

Corresponding author

Correspondence to Mariola Mróz

Ethics declarations

Ethics approval and consent to participate

All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee (The Bioethics Committee of the Medical University of Lublin (KE-0254/221/2016) and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Consent for publication – Not applicable

Availability of data and materials – All data generated or analysed during this study are included in this published article

Competing interests

The authors declare that they have no competing interests" in this section

Informed consent

Informed consent was obtained from all individual participants include in the study.

The authors declare that they have no competing interests in this section

References

1. Lewicka M, Sulima M, Pyć M, Stawarz B. Characteristics of miscarriage and women's rights after pregnancy loss. *Annales AMS*. 2013;59(1):123–9.
2. **The management of early pregnancy loss. Royal College of Obstetricians and Gynaecologists Guidelin No. Retrieved from** <http://www.mp.pl/artykuly/32267>.
3. Arias-Sosa LA, Acosta ID, Lucena-Quevedo E, Moreno-Ortiz H, Esteban-Pérez C, Forero-Castro M. Genetic and epigenetic variations associated with idiopathic recurrent pregnancy loss. *J Assist Reprod Genet*. 2018;35(3):355–66.
4. Klikarová J, Snajbergová K, Měchurová A, Velebil P, Feyereisl J. Intrauterine foetal death syndrome: analysis of cases from 2008 to 2012 in Institute for the care of mother and child. *Ceska Gynekol*. 2014;79(2):120–7.
5. Bhat A, Byatt N. Infertility and Perinatal Loss: When the Bough Breaks. *Curr Psychiatry Rep*. 2016;18(3):31.
6. Christiansen DM, Olf M, Elklit A. Parents bereaved by infant death: sex differences and moderation in PTSD, attachment, coping and social support. *Gen Hosp Psychiatry*. 2014;36(6):655–61.
7. Guzewicz M, Steuden S, Szymona-Pałkowska K. Changes in the perception of self-image and the sense of purpose and meaning in life, among women who lost their child before birth. *Health Psychology Report*. 2014;2(3):162–75.
8. Hunter A, Tussis L, MacBeth A. The presence of anxiety, depression and stress in women and their partners during pregnancies following perinatal loss: A meta-analysis. *J Affect Disord*. 2017;223:153–64.
9. Kersting A, Wagner B. Complicated grief after perinatal loss. *Dialogues Clin Neurosci*. 2012;14(2):187–94.
10. Kolte AM, Olsen LR, Mikkelsen EM. Depression and emotional stress is highly prevalent among women with recurrent pregnancy loss. *Hum Reprod*. 2015;30(4):777–82.
11. San Lazaro Campillo I, Meaney S, McNamara K, O'Donoghue K. Psychological and support interventions to reduce levels of stress, anxiety or depression on women's subsequent pregnancy with a history of miscarriage: an empty systematic review. *BMJ Open*. 2017;7(9):017802.
12. Schwarzer R, Jerusalem M. Skala Uogólnionej Własnej Skuteczności – GSES. In: Juczyński Z, **editor**. *Measurement Instruments In Health Promotion and Psychooncology (Narzędzia Pomiaru w Promocji i Psychologii Zdrowia)*. **Warsaw**: Pracowania Testów Psychologicznych Polskiego Towarzystwa Psychologicznego; 2012. p. 89–94.. **In**.
13. Schwarzer R, Jerusalem M. Generalized Self-Efficacy scale. In: Weinman J, Wright S, Johnston M, **editors**. *Measures in health psychology: A user's portfolio. Causal and control beliefs*. **UK**: NFER-NELSON; 1995. p. 35 – 7.. **In**.
14. Janiszewska M, Kulik T, Żońnierczuk-Kieliszek D, Drop B, Firliej E, Gajewska I. General self-efficacy level and health behaviours in women over the age of 45 years who have undergone osteoporosis treatment. *Menopause Rev*. 2017;16(3):86–95.
15. Bellhouse C, Temple-Smith MJ, Jade E, Bilardi JE. **"It's just one of those things people don't seem to talk about... women's experiences of social support following miscarriage: a qualitative study**. *BMC Women's Health*. 2018;18(1):176..
16. Lotfi-Kashani F, Vaziri S, Akbari ME, Kazemi-Zanjani N, Shamkoeyan L. Predicting posttraumatic growth based upon self-efficacy and perceived social support in cancer patients. *Iran J Cancer Prev*. 2014;7(3):115–22.
17. Tabali M, Ostermann T, Jeschke E, Dassen T, Heinze C. The Relationship between health-related quality of life and care dependency among nursing home residents in Germany: A longitudinal study. *J Gerontol Geriatr Res*. 2015;4(5):1–6.
18. Murphy S, Shevlin M, Elklit A. Psychological consequences of pregnancy loss and infant Heath in a sample of bereaved parents. *J Loss Trauma*. 2014;19(1):56–69.
19. The WHOQOL Group. Development of the World Health Organization WHOQOL-BREF quality of life assessment. *Psychol Med*. 1998;28:551–8.
20. Jaracz K, Kalfoss M, Górna K, Bączyk G. Quality of life in Polish: Psychometric properties of the Polish WHOQOL-Bref. *Scand J Caring Sci*. 2006;20:251–60.
21. Schulz U, Schwarzer R. Soziale Unterstützung bei der Krankheitsbewältigung: Die Berliner Social Support Skalen (BSSS). *Diagnostica*. 2003;49:73–82.

22. Lönnfjord V, Hagquist C. The Psychometric Properties of the Swedish Version of the General Self-Efficacy Scale: A Rasch Analysis Based on Adolescent Data. *Curr Psychol*. 2018;37(4):703–15.
23. Iwanowicz-Palus G, Zarajczyk M, Pięta B, Bień A. Quality of life, social support, acceptance of illness, and self-efficacy among pregnant women with hyperglycemia. *International Journal of Environmental Research Public Health*. 2019;16(20):3941.
24. Rzońca E, Bień A, Wdowiak A, Szymański R, Iwanowicz-Palus G. Determinants of quality of life and satisfaction with life in women with polycystic ovary syndrome. *Int J Environ Res Public Health*. 2018;15(2):376.
25. Couto ER, Couto E, Vian B, Gregório Z, Nomura ML, Zaccaria R, Passini RJr. Quality of life, depression and anxiety among pregnant women with previous adverse pregnancy outcomes. *Sao Paulo Med*. 2009;127(4):185–9.
26. Tavoli Z, Mohammadi M, Tavoli A, Moini A, Effatpanah M, Khedmat L, et al. Quality of life and psychological distress in women with recurrent miscarriage: a comparative study. *Health Qual Life Outcomes*. 2018;16(1):150.
27. Krzepota J, Sadowska D, Biernat E. Relationships between Physical Activity and Quality of Life in Pregnant Women in the Second and Third Trimester. *Int J Environ Res Public Health*. 2018;15(12):2745.
28. Song J, Floyd FJ, Seltzer MM, Greenberg JS, Hong J. Long term effects of child death on parents' health related quality of life: a dyadic analysis. *Fam Relat*. 2010;59(3):269–82.
29. Abbaspoor Z, Razmjou PS, Hekmat K. Relation between quality of life and mental health in pregnant women with prior pregnancy loss: QoL, mental health and pregnancy loss. *J Obstet Gynaecol Res*. 2016;42(10):1290–6.
30. Soma-Pillay P, Makin JD, Pattinson RC. Quality of life 1 year after a maternal near-miss event. *Int J Gynaecol Obstet*. 2018;141(1):133–8.
31. Strine TW, Chapman DP, Balluz L, Mokdad AH. Health-related quality of life and health behaviors by social and emotional support. Their relevance to psychiatry and medicine. *Soc Psychiatry Psychiatr Epidemiol*. 2008;43(2):151–9.
32. Gul B, Riaz MA, Batool N, Yasmin H, Riaz MN. Social support and health related quality of life among pregnant women. *JPMA*. 2018;68(6):872–5.
33. Emmanuel E, St. John W, Sun J. Relationship between social support and quality of life in childbearing women during the perinatal period. *J Obstet Gynecol Neonatal Nurs*. 2012;41(6):62–70.
34. Shisheghar S, Mahmoodi A, Dolatian M, Mahmoodi Z, Bakhtiari M, Alavi Majd H. The relationship of social support and quality of life with the level of stress in pregnant women using the PATH model. *Iran Red Crescent Med J*. 2013;15(7):560–5.
35. Flenady V, Boyle F, Koopmans L, Wilson T, Stones W, Cacciatore J. Meeting the needs of parents after a stillbirth or neonatal death. *BJOG. Br J Obstet Gynaecol*. 2014;121(4):137–40.
36. Leahy-Warren P, McCarthy G, Corcoran P. First-time mothers: social support, maternal parental self-efficacy and postnatal depression. *J Clin Nurs*. 2012;21(3–4):388–97.
37. Sereshti M, Nahidi F, Simbar M, Ahmadi F, Bakhtiari M, Zayeri F. Mothers' perception of quality of services from Health Centers after perinatal loss. *Electron Physician*. 2016;8(2):2006–17.
38. Gan Y, Xiong R, Song J, Xiong X, Yu F, Gao W, et al. The effect of perceived social support during early pregnancy on depressive symptoms at 6 weeks postpartum: a prospective study. *BMC Psychiatry*. 2019;19(1):232.
39. Nikcevic AV, Kuczmierczyk AR, Nicolaidis KH. Personal Coping Resources, Responsibility, Anxiety and Depression After Early Pregnancy Loss. *J Psychosom Obstet Gynaecol*. 1998;19(3):145–54.
40. Rahimparvar S, Hamzehkhani M, Geranmayeh M, Rahimi R. Effect of educational software on self-efficacy of pregnant women to cope with labor: a randomized controlled trial. *Arch Gynecol Obstet*. 2012;286(1):63–70.
41. Schwartz L, Toohill J, Creedy DK, Baird K, Gamble J, Fenwick J. Factors associated with childbirth self-efficacy in Australian childbearing women. *BMC Pregnancy Childbirth*. 2015;15:29.
42. Torrisi M, De Cola MC, Buda A, Carioti L, Scaltrito MV, Bramanti P, et al. Self-efficacy, poststroke depression, and rehabilitation outcomes: Is there a correlation? *J Stroke Cerebrovasc Dis*. 2018;27(11):3208–11.
43. Volz M, Voelkle MC, Werheid K. General self-efficacy as a driving factor of post-stroke depression: A longitudinal study. *Neuropsychol Rehabil*. 2019;29(9):1426–38.