

Exploring Cardioprotective Potential of Esculetin Against Isoproterenol Induced Myocardial Infarction in Rats: In Vivo and in Vitro Evidence

Pullaiah Chitikela P (✉ samuelpharma@gmail.com)

Central Council for Research in Siddha

Vinod K Nelson

Raghavendra Institute of Pharmaceutical Education and Research

Sushma R

Sri Padmavathi School of Pharmacy

Narasimha Kumar GV

Dr Anjali Chatterjee Regional Institute of Homeopathy

Thyagaraju K

Sri Venkateswara University

Research Article

Keywords: Isoproterenol, Oxidative stress, Esculetin, Lysosomes, Antioxidant, Cardioprotection

Posted Date: March 15th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-289373/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Pharmacology and Toxicology on July 15th, 2021. See the published version at <https://doi.org/10.1186/s40360-021-00510-0>.

Abstract

Background

Esculetin is a natural coumarin derivative from various plants with multiple pharmacological effects. Hence, the present study was undertaken to explore the cardio protective potential of esculetin against isoproterenol induced myocardial toxicity in rats.

Methods

The treatment schedule was fixed for 28 days and the rats were divided into five groups of six each. Rats of group I received the normal saline and served as normal control, group II was received ISO (100mg/kg body weight) for last two consecutive days of the study and served as disease control. Groups III and IV received esculetin 10 and 20 mg/kg body weight respectively once a day per oral for 28 days along with ISO for last two consecutive days of the study. Cardiac biomarkers such as CK-MB and LDH, membrane bound Na^+/K^+ ATPases activity, myocardial lysosomal enzymes activity and tissue antioxidants status were estimated in the heart tissue samples. The histopathological changes in the myocardium were also assessed. Further, DPPH assay was done to evaluate the free radicals scavenging potential of esculetin. Cytotoxicity assay, intracellular ROS levels by DCFDA assay and m-RNA expression of TNF- α , IL-6 and NF- κ B by quantitative RT-PCR in H9c2 cell lines.

Results

The increased levels of CK-MB, LDH, LPO, myocardial lysosomal enzymes and membrane bound Na^+/K^+ ATPase levels by ISO administration was significantly increased with concomitant decrease in tissue antioxidant enzymes such as GSH, Catalase, and SOD. Pre-treatment with esculetin for 28 days has significantly decreased the levels of cardiac bio-markers, lysosomal enzymes, membrane bound Na^+/K^+ ATPase levels as well as Lipid peroxides which is in contrary to the ISO group. Amelioration of the antioxidant levels were also found in esculetin treated groups. Histopathological examination of heart reveals that myocardial degeneration, mononuclear cell infiltration was noticed in ISO treated rats, whereas the same was restored with esculetin treatment. In H9C2 cell lines esculetin could effectively reduced intracellular ROS inhibition and m-RNA expression of pro-inflammatory cytokines including TNF- α , IL-6 and NF- κ B to prevent apoptosis or cell necrosis.

Conclusion

The study provides the evidence of cardioprotective potentials of esculetin against isoproterenol induced myocardial infarction by antioxidant and myocardial membrane stabilization along with *in vitro* protection from arsenic induced ROS cell necrosis or apoptosis in H9C2 cells.

Background

Cardiovascular diseases (CVDs) comprise disorders of the heart and blood vessels and still represent a major cause of death globally. CVDs have shown to be responsible for approximately 17.9 million deaths each year, which accounts for 31% of all deaths worldwide. Among the various CVDs, myocardial infarction (MI) is a major cause of mortality and morbidity across the world. Any blockade in the coronary artery leads to insufficient blood supply to heart, causing the heart muscle that is being supplied by the artery to get infarcted ultimately resulting in ischemic tissue necrosis in addition to other pathological and structural changes. The pathogenesis of MI includes hyperlipidemia, oxidative stress, peroxidation of membrane lipids, and loss of plasma membrane integrity [1].

Isoproterenol (ISO), a synthetic catecholamine and β -adrenergic agonist is well-known to cause severe stress in the myocardium by generating free radicals which in turn stimulates lipid peroxidation and perhaps the major contributing factor for the irreversible damage to the myocardial membrane [2]. ISO upon administration causes increase in heart rate leading to increased oxygen demand, high calcium burden and accumulation besides causing alterations in the morphology and membrane integrity of the myocardium with elevated cAMP levels in the myocardial cells [3].

Induction of myocardial infarction was previously performed by surgical procedures, but it has incidence of morbidity, mortality and animals were prone to pneumothorax infections [4]. Isoproterenol-induced myocardial infarction is simple and non-invasive model, considered as one of the most widely used experimental model to study the beneficial effects of many drugs and cardiac function [2] and it is similar to those pathophysiological changes observed in human myocardial infarction [5].

Natural products have high universal demands due to their claimed advantage in terms of both safety and efficacy against various diseases like MI. Plant based coumarins are low- molecular weight phenolic compounds that has been used for the prevention and treatment of various thromboembolism MI and stroke [6]. Esculetin (6, 7- di hydroxyl coumarin) is a natural coumarin derivative isolated from many plants such as *Artemisia capillaries*, *Citrus limonia*, *Solanum surrettence* and *Euphorbia lathyrus* [7] with multiple pharmacological & biochemical properties[8]. There are numerous treatment approaches that have been developed to attenuate the risk of myocardial infarction, but most of them fail when translated from the bench to the bedside. Therefore, there is a need to find new drugs which hostile MI as well as be used as suitable therapeutic candidate and also can be translated to clinical use in the future [9].

The present study was intended to investigate whether esculetin can provide protection against free radical generation by ISO and its associated alterations in the membrane bound enzymes and myocardial lysosomal enzyme activities in experimental rats. Further, we also tried to explore the probable mechanism of action of the esculetin for its cardio protection in H9C2 cell lines.

Experimental Procedures

Chemicals

Isoproterenol (CAS Number 5984-95-2), and Esculetin (CAS Number 305-01-1) was purchased from Sigma Aldrich. Co, St. Louis, USA. All the chemicals used in the present study were of analytical grade and indigenous.

***In vitro* studies**

2, 2-Diphenyl-1-picrylhydrazyl (DPPH) radical scavenging activity

The antioxidant activity of esculetin was measured using DPPH radical scavenger assay in triplicate. Briefly, the stock solution (1mg/mL) of esculetin was prepared in methanol. Concentration of 5, 25, 50 and 100µg/mL were used along with ascorbic acid as reference standard. The test were then incubated with 50µL of 0.1mM DPPH solution and made up to the final volume to 3mL with methanol. A blank was prepared using DPPH solution and methanol. The reaction mixture was incubated for 30 min at room temperature in the dark followed by measuring absorbance at 517nm [10].

H9C2 cell culture

H9C2 myoblast cells from rat's myocardium were acquired from National Centre for Cell Sciences, Pune, India. The myoblast cells were cultured in Dulbecco's modified Eagle's Medium (DMEM) medium supplemented with 10% FBS and 10 ml/l100 × antibiotic–antimycotic solution containing 10,000 units of penicillin and 10 mg/ml streptomycin in 0.9% normal saline in a humidified atmosphere of 95% air and 5% CO₂ at 37°C.

Measurement of cell viability by MTT assay

The cells were seeded in 96-well culture plates at a density of 7×10⁴ cells/well. When the cells reached 80% confluence, they were treated with 1‰ dimethyl sulfoxide (DMSO) solution, and 5, 25 and 50 µg/mL of esculetin at 37°C for 24 h. The cells were then incubated with 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) (0.5 mg/mL) solution for 4h, and the resulting formazan was solubilized with 150 µL of DMSO for 30 min. The absorbance of each well was measured at 570 nm, and the absorbance of control cells was considered to indicate 100% cell viability [10, 11].

Intracellular Reactive Oxygen Species (ROS) measurement

The generation of intracellular reactive oxygen species (ROS) was measured by using the ROS-sensitive fluorescence indicator called Dichlorofluorescein diacetate (DCFH-DA) as per our previous protocol. H9C2 cells grown in 96 well plates to ~75% confluence were treated in triplicate. The cells were treated with

esculetin at 0.1, 1, 5 and 10 μ M. After 12h, 10 μ M arsenic was added to the plate. The cells were incubated for an additional 30 min at 37°C and after PBS wash cells were incubated with 20 μ M DCFH-DA 30min at 37°C in the dark. After, Cells were washed, and analyzed by flow cytometer. The florescence intensity was calculated using the FAC Suite software [10].

RNA isolation, cDNA synthesis and qPCR to assess mRNA expression of TNF- α , IL-6, NF- κ b

After arsenic and esculetin treatment, the total RNA was isolated from H9C2 cells by using Trizol reagent (Thermo Fisher Scientific, Inc.). The isolated RNA was quantified by using a nano-drop spectrophotometer and complementary DNA (cDNA) was synthesized from 1 μ g of RNA was used for reverse transcription reaction using the iScript cDNA synthesis kit (Bio-Rad, Hercules, CA, USA). Reverse transcription quantitative polymerase chain reaction (RT-qPCR) was performed by using the SYBR green reagent according to the manufacturer's protocol (MilliporeSigma).

The primer sequences used for qPCR were as follows: TNF- α forward, 5' GAACTGGCAGAAGAGGCACT-3' and reverse, 5'-GGTCTGGGCCATAGAACTGA-3'; IL-6 forward, 5'-CCGGAGAGGAGACTTCACAG-3' and reverse, 5'-CAG AATTGCCATTGCACA-3'; NF- κ B forward 5'-CCCACACTATGGATTTCTACTTATGG'-3 and reverse 5' CCAGCAGCATCTTCACGTCTC-3'. RT-qPCR reactions were performed under the conditions like, 50°C for 35 min, 85°C for 12 min, followed by 60 cycles of 95°C for 23 sec and 60°C for 1.5 min. The selected gene expression level was normalized to glyceraldehyde-3-phosphate dehydrogenase (GAPDH) forward, 5' CTTTGGTATCGTGGAAGG ACTC-3' and reverse, 5' GTAGAGGCAGGGATGATGTTCT-3' as internal loading control [12, 13].

In vivo study

Animals

Male wistar rats weighed between 230-280 g were used in this study. Rats were housed under standard conditions and fed with standard pellet with drinking water *ad libitum*. The animals were kept in polypropylene cages and maintained at a room temperature of 25 \pm 2°C with 55 \pm 5% relative humidity and 12hrs light/dark cycle. The study was carried out in compliance with the ARRIVE (Animals in Research: Reporting *In Vivo* Experiments) guidelines ("Guide for the Care and Use of Laboratory Animals" (Institute of Laboratory Animal Resources, National Academic Press 1996; NIH publication number #85-23, revised 1996). All experimental procedures and methods were approved by the Institutional Animal Ethical Committee (IAEC), Sri Padmavathi School of Pharmacy, constitute as per the directions of the Committee for the Purpose of Control and Supervision of Experiments on Animals (CPCSEA), India.

Induction of myocardial infarction

Myocardial infarction was induced by dissolving isoproterenol (100 mg/kg) in normal saline and injected subcutaneously to rats for last two consecutive days of the experimental schedule [14].

Experimental schedule

The treatment schedule was fixed for 28 days and the rats were divided into five groups of six each. Rats of group I received the normal saline and served as normal control, group II was received ISO (100mg/kg body weight) for last two consecutive days of the study and served as disease control. Groups III and IV received esculetin 10 and 20 mg/kg body weight respectively, once a day orally for 28 days of the study along with ISO for last two consecutive days of the study and serve as test groups.

Blood sample collection and analysis

At the end of treatment blood was collected from retro orbital plexus by anesthetizing the rats with thiopental sodium (35 mg/kg body weight, intra peritoneal) [15] and serum was separated by centrifugation at 2000 rpm. Serum was used to analyze various biochemical parameters such as determinations of cardiac biomarkers lactate dehydrogenase (LDH), and creatinine kinase MB (CK-MB) by using commercial diagnostic kits (Agappe Pvt. Ltd, Kerala, India).

Na⁺ /K⁺ ATPase activity of myocardial membrane

The myocardial membrane Na⁺ /K⁺ ATPases activity was determined according to procedure done by Periyathambi and Ponnian 2007. The incubation mixture contained 10mM of Tris buffer, 20mM of potassium chloride, 125mM of sodium chloride, 1mM of EDTA and 3mM of ATP. To the incubation mixture, the reaction was initiated by the addition of 0.2mL of tissue homogenate and the contents were incubated at 37°C for 15 min. To stop the reaction of 10% trichloro acetic acid (TCA) was added. The tubes were centrifuged and supernatant was used for the estimation of liberated P_i. 1.0mL of supernatant was made up to 4.3mL with distilled water and added 1.0ml 3mM of ammonium molybdate reagent. The tubes were incubated at room temperature for 10 minutes, and later 0.4ml of amino naphtholsulphonic acid reagent was added to develop the color and the P_i released recorded using a standard P_i graph [16].

Preparation of lysosomal sub cellular fractions

Lysosomal subcellular fractions were isolated according to the method of Venkatachalem *et al.*, 2003. The heart tissue sample was cut open and placed in isotonic saline to remove the blood. Then the heart tissue was rinsed in ice cold 0.25 M sucrose, blotted, weighed and minced. The enzyme extracts were

prepared by homogenizing the tissue samples in 0.25 M sucrose at 4°C. The portion of the homogenate was subjected to differential centrifugation, and the different fractions were separated as follows: structural proteins, nucleus, and cell debris at 600×g for 10 min; mitochondria at 5000×g for 10 min; lysosomes at 15,000×g for 10 min. Myocardial sub-fractions were treated with Triton X-100 (final concentration 0.2% v/v) in ice for 15 min prior to the determination of enzymatic activity [17] .

The activities of the lysosomal enzymes including β -glucuronidase [18], β -glucosidase and β -galactosidase [19], and acid phosphatase [20] were determined.

Determination of tissue antioxidants

At the end of the experimentation hearts were excised from rats and homogenate in 0.1M Tris buffer (pH 7.4) and the separated homogenates were used for estimation of heart antioxidants like super oxide dismutase (SOD) [21], Reduced glutathione (GSH) [22], Catalase [23] and lipid peroxidation (LPO) [24].

Histopathological studies of heart

After removal of myocardial tissue immediately washed with ice cold saline to remove all the blood and fixed in 10% buffered neutral formalin solution. After fixation was complete, tissues were embedded in paraffin and serial sections were cut in to 0.5 μ m. Each section was stained with hematoxylin and eosin. The sections were examined under light microscope and histograms were taken.

Statistical analysis

Results were expressed as mean \pm standard error mean multiple comparisons of the significant analysis of variance (ANOVA) followed by the Dennett's test as post parametric test using computer based fitting program (Prism graph pad 5.0). A p value of <0.05 was considered as statistically significant.

Results

DPPH radical scavenging activity of esculetin

DPPH assay is the simple and sensitive choice to assess the free radical scavenging activity of any natural origin compounds. DPPH accepts an electron or hydrogen radical to become a stable diamagnetic molecule. Esculetin exhibited significant radical scavenging activity at all the working concentrations (Fig 1) nevertheless, the maximum scavenging activity was at 50 μ M compared to control.

Esculetin Inhibits Intracellular ROS production in H9c2 cell

In the study, we investigated the inhibitory effect of esculetin on arsenic induced ROS generation in H9c2 cells by using DCFH-DA a ROS sensitive non-fluorescent agent (Fig 1). The DCF stained cells were analyzed by flow cytometer. During incubation period a significant ROS generation is noticed in cells placed with arsenic alone. Whereas cells treated with esculetin at 25 and 50µg/mL concentrations exhibited dose-dependent inhibitory effects with IC₅₀ value of 125M.

Effect of esculetin on H9c2 cell viability

MTT assay was used to determine the effect of esculetin on H9C2 cell viability at 5, 25 and 50µg/mL concentration. After the effective treatment with esculetin H9C2 cells observed as viable with no significant cytotoxicity at all the concentrations.

Esculetin attenuates mRNA expression of TNF-α, IL-6, and NF-κB against arsenic induced ROS in H9C2 cells

The mRNA expression of pro-inflammatory cytokines TNF-α, IL-6 and NF-κB were measured using RT-qPCR method against arsenic induced ROS generation in H9C2 cells (Fig 2). The mRNA expression TNF-α, IL-6 and NF-κB were significantly ($p \leq 0.05$) elevated, approximately 9 folds compared to control cells. While the cells treated with esculetin at 25 and 50µg/mL concentration in the presence of arsenic attenuated the expression of mRNA at 4 to 6 folds ($p \leq 0.05$) lesser than the control. No differences were observed in the levels of these pro-inflammatory cytokines between the control and esculetin alone groups. The above results indicated that pre-treatment with esculetin could reduce the production and expression of pro-inflammatory cytokines in TNF-α, IL-6 and NF-κB in arsenic induced ROS in H9C2 cells.

Effect of esculetin on cardiac biomarkers in ISO induced myocardial toxicity

The alterations in serum heart biomarkers like CK-MB and LDH were tabulated in Table 1. The concentration of CK-MB and LDH were significantly ($p < 0.05$) increased in isoproterenol alone treated rats compared to normal control rats. Animals treated with esculetin brought these cardiac markers near normalcy ($p < 0.05$) compared to isoproterenol alone treated rats.

Table 1

Esculetin effect on CK-MB, LDH and myocardial Na⁺/K⁺ ATPase activity in ISO induced myocardial toxicity.

	CK-MB (IU/L)	LDH (IU/L)	Na ⁺ K ⁺ ATPases activity (μmol of Pi/mg/h)
Normal control	85.14± 4.374	252.0±10.68	3.285±0.2040
Disease control	336.1±23.43 [#]	449.6±26.48 [#]	1.080±0.158 [#]
Esculetin(10 mg/kg)	144.4±6.399 [*]	319.9±12.77 [*]	2.190±0.099 [*]
Esculetin (20 mg/kg)	121.6±7.470 [*]	257.9±13.33 [*]	2.542±0.303 [*]
All values are shown as mean ± SEM and n=6. Data analysis done by one way analysis of variance followed by the Tukey multiple comparison tests. [#] p<0.05, indicate disease control compared with normal. ^{**} p<0.05, indicate Esculetin compared with disease control group.			

Effect of esculetin on membrane bound Na⁺/K⁺ ATPase activity in ISO induced myocardial toxicity

The effect of Esculetin on membrane bound Na⁺/K⁺ ATPase activity has shown in Table 1, significant (p<0.05) decrease of myocardial membrane bound Na⁺/K⁺ ATPase in rats treated with control when compared with normal rats. Whereas, the same Na⁺/K⁺ ATPase significantly (p < 0.05) increased in rats pretreated with esculetin for four weeks when compared with control rats.

Effect of Esculetin on lysosomal membrane destabilization in ISO induced myocardial toxicity

Table 2, shows that alterations of activities of lysosomal hydrolases enzymes like β-glucuronidase, β-galactosidase, β-glucosidase, and acid phosphatase. Activity of these enzymes were significantly (p<0.05) increased in heart tissue homogenates of rats treated with isoproterenol alone when compared with normal rats. Animals pre-treated with esculetin at doses of 10 and 20 mg/kg body weight for 28 days brought these enzyme activities significantly (p<0.05) low and near to normal when compared with isoproterenol alone treated rats.

Table 2

Esculetin effect on myocardial lysosomal enzyme's activity in ISO induced myocardial toxicity.

	β -glucuronidase (p-nitro phenol/h/mg protein)	β -glucosidase (p-nitro phenol/h/mg protein)	β -galactosidase (p-nitro phenol/h/mg protein)	Acid phosphatase (p-nitro phenol/h/mg protein)
Normal	0.425 \pm 0.003	0.050 \pm 0.003	0.042 \pm 0.004	0.071 \pm 0.008
Disease Control	0.088 \pm 0.007 [#]	0.114 \pm 0.003 [#]	0.107 \pm 0.007 [#]	0.159 \pm 0.009 [#]
Esculetin (10mg/kg)	0.536 \pm 0.005 [*]	0.063 \pm 0.005 [*]	0.073 \pm 0.005 [*]	0.084 \pm 0.005 [*]
Esculetin (20mg/kg)	0.042 \pm 0.001 [*]	0.06 \pm 0.002 [*]	0.062 \pm 0.001 [*]	0.079 \pm 0.003 [*]
All values are shown as mean \pm SEM and n=6. Data analysis done by one way analysis of variance followed by the Tukey multiple comparison tests. [#] p<0.05, indicate disease control compared with normal. ^{**} p<0.05, indicate Esculetin compared with disease control group.				

Effect of esculetin on tissue antioxidants in ISO induced myocardial toxicity

The changes in heart antioxidants are presented in Table 3. In ISO induced rats, there was a significant (p<0.05) decrease in GSH, Catalase, SOD and a significant increase in LPO compared to normal control group and pre-treatment with esculetin at doses of 10 and 20mg/kg, respectively, brought the elevated heart antioxidants near normal.

Table 3

Esculetin effect on myocardial tissue antioxidant enzyme's activity in ISO induced myocardial toxicity.

	SOD (U/mg Protein)	CATALASE (μ M H2O2 Consumed/mg Protein)	GSH (μ g of GSH/mg Protein)	LPO (nm of MDA/mg Protein)
Normal	9.248 \pm 1.047	7.965 \pm 1.217	11.65 \pm 1.108	1.693 \pm 0.200
Disease Control	1.646 \pm 0.516 [#]	1.159 \pm 0.386 [#]	4.027 \pm 0.784 [#]	3.313 \pm 0.407 [#]
Esculetin (10 mg/kg)	5.804 \pm 0.267 [*]	6.484 \pm 1.145 [*]	7.766 \pm 0.384 [*]	2.044 \pm 0.255 [*]
Esculetin (20mg/kg)	7.241 \pm 1.070 [*]	9.249 \pm 1.856 [*]	9.719 \pm 0.332 [*]	1.661 \pm 0.365 [*]
All values are shown as mean \pm SEM and n=6. Data analysis done by one way analysis of variance followed by the Tukey multiple comparison tests. [#] p<0.05, indicate disease control compared with normal. ^{**} p<0.05, indicate Esculetin compared with disease control group.				

Histopathological observations (Fig 3) of normal control rat's heart revealed normal cardiac muscle bundles without any inflammation, whereas myocardium of rats treated with isoproterenol alone has shown a marked inflammatory signs like membrane damage and cellular infiltration along with focal myonecrosis. Rats pre-treated with esculetin at doses of 10 and 20mg/kg body weight has shown reduction in inflammatory signs and myonecrosis compared to disease control.

Discussion

The present study validates to reduce the acuteness of isoproterenol induced myocardial infarction by stabilizing the myocardial membrane integrity. The results of experiment bring a new outcome that, pre-treatment with esculetin in experimental rats protects against myocardial infarction. Reactive oxygen species generation by oxidative stress plays an important role in the development of myocardial infarction both experimental and clinically [25].

Antioxidant compounds, especially polyphenol compounds from plants, are capable of counteracting the harmful effects caused by ROS and therefore it can prevent chronic diseases related to oxidative stress [26]. In the present study reveals that, esculetin has shown that direct radical quenching effect in DPPH radical scavenging assay. In addition, it also reduced the intracellular ROS production induced by arsenic [27] in H9C2 cell lines. Treatment with esculetin in H9C2 cells, a noticeable reduction of DCFDA fluorescence intensity was observed this indicates that, esculetin may interact with cellular anti-oxide enzymes. Thus, esculetin has the ability to alleviate free radicals ions both *in vitro* and *in vivo*. This was further confirms by tissue antioxidant data of present study.

Anti-oxidant enzymes such as catalase, superoxide dismutase (SOD), glutathione peroxidase and glutathione-S transferase (GSH) are the first line of cellular defense against oxidative injury by superoxide anion radical and H_2O_2 before interacting to form the more reactive hydroxyl radical [28]. Auto-oxidation of ISO produces highly cytotoxic free radicals like quinines which in addition with superoxide anion and potent hydroxyl radicals damages the polyunsaturated fatty acids of myocardial membrane [29, 30]. The elevation of these free radicals brings an imbalance between tissue bound cellular scavenging enzymes like SOD, GSH and Catalase and lipid peroxidase [31].

The elevation of free radical including superoxide, hydroxyl ions and hydrogen peroxide ions was the reason for decreasing of tissue bound antioxidant enzymes [32, 33]. It is in line with previous studies, that tissue anti-oxidant enzymes were significantly decreased where the LPO levels were increased in the hearts of ISO alone treated rats [34, 35]. Pre-treatment with esculetin brought the elevated levels of tissue anti-oxidant enzymes like SOD, GSH and Catalase near base line. Our results are agreement with several previous findings in which esculetin is a potential naturally occurring antioxidant both *in vitro* and *in vivo* [36, 37].

Probably, the anti-oxidant effect of esculetin could by hydrogen donating ability of esculetin [38], the catechol structure of esculetin contributes the stronger antioxidant activity [36]. The available data of the present study confirms with existing literature that, esculetin has the ability to activate anti-oxidant enzymes or and might combat the excess ROS formed with in the cell [39, 40].

Arsenic one of the major component to induce intra-cellular ROS [27]. Cytokines such as IL-1 β , TNF- α and IL-6 are pro-inflammatory markers play a vital role in various inflammatory pathways like JNK, NF- κ B and NLRP [41]. NF- κ B is a multiple transcription factor that regulates transcription of various genes involved in the pathogenesis of myocardial infarction like TNF- α and IL-6 [42, 43]. These pro-inflammatory cytokines and inflammatory signaling pathways could promote the development of myocardial infarction. Intracellular ROS promotes the production and release of these pro-inflammatory cytokines from cardiomyocytes. TNF- α , IL-1b, and IL-6, are early predictors of organ dysfunction and, causes to activate cardiomyocytes apoptosis [44]. In our study we investigated for the change in the mRNA expression levels of ROS sensitive transcription factor NF- κ B along with its pro-inflammatory cytokines TNF- α and IL-6 in esculetin treated H9C2 cells and noticed a dose dependent decline in the mRNA expressions of above said markers. This denotes that esculetin could effectively inhibited the expression of mRNAs of pro-inflammatory cytokines, which are the prominent markers of inflammation and myocardial infarction, this effect may prevent the myocardium to undergo apoptosis or cell necrosis. The present findings are in line with earlier research saying esculetin could effectively down regulated the TNF- α , NF- κ B and IL-6 levels upon various stimulus including ROS [13, 45, 46].

CK-MB and LDH are indicator biomarker to diagnose the severity of myocardial infarction and number of necrotic cells [47]. These enzymes are present in cardiac muscle, upon the injury release into the blood stream [48]. The available data [49, 50], reveals that isoproterenol administration significantly elevates the CK-MB and LDH levels in the serum; in our study also the same. In the *in vitro* model [28], esculetin

improved the cell viability and reduces the release of LDH, in the same line, pre-treatment with esculetin reduce the ISO induced raise of CK-MB and LDH to protect the myocardium, and it also maintained the cell viability in H9C2 cell lines. This implying that esculetin has the potentiality to protect pathological and morphological changes in the rats in a dose-independent manner.

The functioning ability of the membrane bound enzymes are depends on the stability of plasma membrane. Under normal cell physiology Na^+ - K^+ ATPase will balances the Na^+ and K^+ ions across the membrane, upon its disruption may leads to ionic inequality and cell death. Therefore, the determination of membrane bound enzyme activities will indicate any alteration to the membrane physiology under pathological conditions [51]. ISO liberated free radicals bind to the membrane lipids to cause injure, thereby inhibits the membrane associated enzymes function including ATPases [2]. The loss of ATPase activity in the ischemic state may be responsible for causing functional damage and reversible necrotic changes in the involved myocardial cell [52].

Palanivel Karthika et al., has found that, pre-treatment with esculetin has effectively increased the Na^+ - K^+ ATPase activity, this could be due to the ability of esculetin to protect SH group of Na^+ - K^+ ATPase enzyme from oxidative damage through inhibition of peroxidation of membrane lipids [53]. It is in line that, the same circumstances were observed in the present study with pre-treatment with esculetin appreciably restored the membrane bound Na^+ - K^+ ATPase near to normalcy.

Lysosomes are surrounded by phospholipid enriched membranes, and are a potential site for free-radical attack including ISO produces quinines [54]. Therefore, radical ions causing to loss the membrane stability and leads the release of hydrolytic enzymes from its sacs to augments cell necrosis [55], [56]. Hence, significant attention has been raised on the involvement of lysosomal enzymes in myocardial damage. From our previous report it was observed that, administration of the ISO to the rats causes a significant elevation in the activities of the lysosomal enzymes [30]. In the present study same effect observed.

Stabilization of myocardial cell membranes, mainly the lysosomal membranes, may extend the viability of myocardial cells to prevent MI. Pre-treatment with esculetin could effectively inhibit the release of lysosomal enzymes from their sacs, and protects the myocardium from necrosis. This provides the first *in vivo* evidence that esculetin protects myocardial from ISO-induced injury by lysosomal membrane stabilization.

Biochemical findings that proves the cardio protective effect was further supported by histopathological examination. Myocardial tissue sections of normal rats depicted clear integrity of the myocardial cell membrane whereas the sections of hearts treated with ISO showed necrosis of muscle fibers with inflammatory cell infiltration, edema and fragmentation of muscle fibers, which indicated involvement of oxidative stress and inflammatory processes. Pre-treatment with esculetin showed the integrity to the structure of myocardium from ISO injury near to the normal myocardium denotes its cardio protective

action. The limitation of the study was a clear mechanism of action by which pathway esculetin interfering to reducing the myocardial necrosis has not done.

Conclusion

Based on the results of this study, it can be concluded that Esculetin ameliorates isoproterenol associated pathological features like attenuating the oxidative stress, stabilization of lysosomal, myocardial membrane and modulation of pro-inflammatory markers TNF- α , NF- κ B and IL-6 cytokines to ascertain its cardioprotective potentials. Understanding the molecular mechanisms involved in the lysosomal enzyme leakage may prove beneficial measures in the prevention of myocardial infarction. Thus, esculetin may consider as a better therapeutic option for treatment of myocardial infarction. Further work need to be done in order to establish specific molecular mechanisms by which esculetin prevent myocardial infarction and possibility for its clinical application.

Abbreviations

CVDs: Cardiovascular diseases; MI: Myocardial infarction; ISO: Isoproterenol; TNF- α : Tumor necrosis factor alfa; IL-6: Including interleukin-6; NF- κ B: Nuclear factor kappa B; NLRP3: Nucleotide-binding Leucine-rich repeat Receptors Protein DMEM: Dulbecco's Modified Eagle's Medium; DMSO: Dimethyl sulfoxide; ROS: Intracellular reactive oxygen species; DCFH-DA: Dichlorofluorescein diacetate; cDNA: complementary DNA; RT-qPCR: Reverse transcription quantitative polymerase chain reaction; LDH: Lactate dehydrogenase; CK-MB: Creatinine kinase MB; TCA: Trichloro acetic acid; SOD: Super oxide dismutase; GSH: Reduced glutathione; LPO: Lipid peroxidation; mRNA: Messenger ribonucleic acid.

Declarations

Acknowledgement

Authors are thankful to principal Prof Ranganayakulu D and management of Sri Padmavathi School of Pharmacy, Tirupati-517503, for providing the animal house and laboratory facilities.

Author's contribution

CPP has designed and prepared the manuscript, VKN performed all in vitro and cell line work and in the preparation of manuscript. SR performed animal activities and biochemical assays, NKGV analyzed the data and results interpretation and TK edited the complete manuscript and made ready for communication.

Funding

Not applicable

Availability of the data and materials

Corresponding author will provide the data used in present work upon the request.

Ethics approval and Declaration

All methods including animal experimental procedures were adopted in the study were performed in accordance with ARRIVE guidelines. Animal experimental protocols used in the present study were approved by Institutional Animal Ethical Committee of Sri Padmavathi School of Pharmacy (SPSP/CPCSEA/IAEC-1016/a /2015/009), constitute as per the directions of the Committee for the Purpose of Control and Supervision of Experiments on Animals (CPCSEA), India. The further methods and procedures adopted in the study are in agreement with appropriate guidelines and regulations.

Consent for publication

Not applicable

Competing of interest

The authors declare that they have no competing interest.

References

1. Zaki AA, Hashish NE, Amer MA, Lahloub MF: **Cardioprotective and antioxidant effects of oleogum resin "Olibanum" from Bos Boswellia carteri Birdw. (Bursearceae).** *Chinese journal of natural medicines* 2014, **12**:345-350.
2. Lobo RO, Chandrasekhar Sagar BK, Shenoy CK: **Bio-tea prevents membrane destabilization during Isoproterenol-induced myocardial injury.** *Journal of microscopy and ultrastructure* 2017, **5**:146-154.
3. Eltobshy SAG, Hussein AM, Elmileegy AA, Askar MH, Khater Y, Metias EF, Helal GM: **Effects of heme oxygenase-1 upregulation on isoproterenol-induced myocardial infarction.** *The Korean journal of physiology & pharmacology : official journal of the Korean Physiological Society and the Korean Society of Pharmacology* 2019, **23**:203-217.
4. Wong ZW, Thanikachalam PV, Ramamurthy S: **Molecular understanding of the protective role of natural products on isoproterenol-induced myocardial infarction: A review.** *Biomedicine & pharmacotherapy = Biomedecine & pharmacotherapie* 2017, **94**:1145-1166.

5. Rona G, Chappel CI, Balazs T, Gaudry R: **An infarct-like myocardial lesion and other toxic manifestations produced by isoproterenol in the rat.** *AMA archives of pathology* 1959, **67**:443-455.
6. Voora D, McLeod HL, Eby C, Gage BF: **The pharmacogenetics of coumarin therapy.** *Pharmacogenomics* 2005, **6**:503-513.
7. Masamoto Y, Ando H, Murata Y, Shimoishi Y, Tada M, Takahata K: **Mushroom tyrosinase inhibitory activity of esculletin isolated from seeds of Euphorbia lathyris L.** *Bioscience, biotechnology, and biochemistry* 2003, **67**:631-634.
8. Jeon YJ, Jang JY, Shim JH, Myung PK, Chae JI: **Esculetin, a Coumarin Derivative, Exhibits Anti-proliferative and Pro-apoptotic Activity in G361 Human Malignant Melanoma.** *Journal of cancer prevention* 2015, **20**:106-112.
9. Schreiber T, Salhöfer L, Quinting T, Fandrey J: **Things get broken: the hypoxia-inducible factor prolyl hydroxylases in ischemic heart disease.** *Basic research in cardiology* 2019, **114**:16.
10. Nelson VK, Ali A, Dutta N, Ghosh S, Jana M, Ganguli A, Komarov A, Paul S, Dwivedi V, Chatterjee S, et al: **Azadiradione ameliorates polyglutamine expansion disease in Drosophila by potentiating DNA binding activity of heat shock factor 1.** *Oncotarget* 2016, **7**:78281-78296.
11. Nelson VK, Sahoo NK, Sahu M, Sudhan HH, Pullaiah CP, Muralikrishna KS: **In vitro anticancer activity of Eclipta alba whole plant extract on colon cancer cell HCT-116.** *BMC complementary medicine and therapies* 2020, **20**:355.
12. Singh BK, Vatsa N, Nelson VK, Kumar V, Kumar SS, Mandal SC, Pal M, Jana NR: **Azadiradione Restores Protein Quality Control and Ameliorates the Disease Pathogenesis in a Mouse Model of Huntington's Disease.** *Molecular neurobiology* 2018, **55**:6337-6346.
13. Choi RY, Ham JR, Lee MK: **Esculetin prevents non-alcoholic fatty liver in diabetic mice fed high-fat diet.** *Chemico-biological interactions* 2016, **260**:13-21.
14. Stanely Mainzen Prince P, Priscilla H, Devika PT: **Gallic acid prevents lysosomal damage in isoproterenol induced cardiotoxicity in Wistar rats.** *European journal of pharmacology* 2009, **615**:139-143.
15. Prabhu S, Narayan S, Devi CS: **Mechanism of protective action of mangiferin on suppression of inflammatory response and lysosomal instability in rat model of myocardial infarction.** *Phytotherapy research : PTR* 2009, **23**:756-760.
16. Devika PT, Mainzen Prince PS: **(-)-Epigallocatechin gallate (EGCG) prevents isoprenaline-induced cardiac marker enzymes and membrane-bound ATPases.** *The Journal of pharmacy and pharmacology* 2008, **60**:125-133.
17. Sathish V, Ebenezar KK, Devaki T: **Synergistic effect of nicorandil and amlodipine on lysosomal hydrolases during experimental myocardial infarction in rats.** *Biomedicine & pharmacotherapy = Biomedecine & pharmacotherapie* 2003, **57**:309-313.
18. Hultberg B, Lindsten J, Sjöblad S: **Molecular forms and activities of glycosidases in cultures of amniotic-fluid cells.** *The Biochemical journal* 1976, **155**:599-605.

19. Conchie J, Levvy GA: **Inhibition of glycosidases by aldonolactones of corresponding configuration.** *The Biochemical journal* 1957, **65**:389-395.
20. Kavanagh JP, Bardsley WG: **The identity of the acid and alkaline phosphatases of human seminal plasma.** *Journal of reproduction and fertility* 1979, **57**:43-48.
21. Misra HP, Fridovich I: **The role of superoxide anion in the autoxidation of epinephrine and a simple assay for superoxide dismutase.** *The Journal of biological chemistry* 1972, **247**:3170-3175.
22. Moron MS, Depierre JW, Mannervik B: **Levels of glutathione, glutathione reductase and glutathione S-transferase activities in rat lung and liver.** *Biochimica et biophysica acta* 1979, **582**:67-78.
23. Aebi H: **Catalase in vitro.** *Methods in enzymology* 1984, **105**:121-126.
24. Kovachich GB, Mishra OP: **Lipid peroxidation in rat brain cortical slices as measured by the thiobarbituric acid test.** *Journal of neurochemistry* 1980, **35**:1449-1452.
25. D'Oria R, Schipani R, Leonardini A, Natalicchio A, Perrini S, Cignarelli A, Laviola L, Giorgino F: **The Role of Oxidative Stress in Cardiac Disease: From Physiological Response to Injury Factor.** *Oxidative medicine and cellular longevity* 2020, **2020**:5732956.
26. Witacenis A, Seito LN, da Silveira Chagas A, de Almeida LD, Jr., Luchini AC, Rodrigues-Orsi P, Cestari SH, Di Stasi LC: **Antioxidant and intestinal anti-inflammatory effects of plant-derived coumarin derivatives.** *Phytomedicine : international journal of phytotherapy and phytopharmacology* 2014, **21**:240-246.
27. Chowdhury R, Chatterjee R, Giri AK, Mandal C, Chaudhuri K: **Arsenic-induced cell proliferation is associated with enhanced ROS generation, Erk signaling and CyclinA expression.** *Toxicology letters* 2010, **198**:263-271.
28. He Y, Li C, Ma Q, Chen S: **Esculetin inhibits oxidative stress and apoptosis in H9c2 cardiomyocytes following hypoxia/reoxygenation injury.** *Biochemical and biophysical research communications* 2018, **501**:139-144.
29. Priscilla DH, Prince PS: **Cardioprotective effect of gallic acid on cardiac troponin-T, cardiac marker enzymes, lipid peroxidation products and antioxidants in experimentally induced myocardial infarction in Wistar rats.** *Chemico-biological interactions* 2009, **179**:118-124.
30. Pullaiah CP, Kumar GN, Jyothsna K, Thyagaraju K, Nelson VK, Reddy GDJOP, Medicine E: **Rosa damascena Mill. L. attenuates myocardial lysosomal membrane destabilization in isoproterenol induced oxidative stress.** 2017, **17**:373-380.
31. Shahzad S, Mateen S, Mubeena Mariyath PM, Naeem SS, Akhtar K, Rizvi W, Moin S: **Protective effect of syringaldehyde on biomolecular oxidation, inflammation and histopathological alterations in isoproterenol induced cardiotoxicity in rats.** *Biomedicine & pharmacotherapy = Biomedecine & pharmacotherapie* 2018, **108**:625-633.
32. S M, Shaik AH, E MP, Al Omar SY, Mohammad A, Kodidhela LD: **Combined cardio-protective ability of syringic acid and resveratrol against isoproterenol induced cardio-toxicity in rats via attenuating NF- κ B and TNF- α pathways.** *Scientific reports* 2020, **10**:3426.

33. Fang WJ, Wang CJ, He Y, Zhou YL, Peng XD, Liu SK: **Resveratrol alleviates diabetic cardiomyopathy in rats by improving mitochondrial function through PGC-1 α deacetylation.** *Acta pharmacologica Sinica* 2018, **39**:59-73.
34. Anandan R, Chatterjee NS, Sivakumar R, Mathew S, Asha KK, Ganesan B: **Dietary Chitosan Supplementation Ameliorates Isoproterenol-Induced Aberrations in Membrane-Bound ATPases and Mineral Status of Rat Myocardium.** *Biological trace element research* 2015, **167**:103-109.
35. Tao W, Zuo T, Ma C, Long H: **Suppressing receptor-interacting protein 140: A new sight for esculetin to treat myocardial ischemia/reperfusion injury.** *RSC Adv* 2016, **6**.
36. Kim SH, Kang KA, Zhang R, Piao MJ, Ko DO, Wang ZH, Chae SW, Kang SS, Lee KH, Kang HK, et al: **Protective effect of esculetin against oxidative stress-induced cell damage via scavenging reactive oxygen species.** *Acta pharmacologica Sinica* 2008, **29**:1319-1326.
37. Lee BC, Lee SY, Lee HJ, Sim GS, Kim JH, Kim JH, Cho YH, Lee DH, Pyo HB, Choe TB, et al: **Anti-oxidative and photo-protective effects of coumarins isolated from Fraxinus chinensis.** *Archives of pharmacal research* 2007, **30**:1293-1301.
38. Wang K, Zhang Y, Ekunwe S, Yi X, Liu X, Wang H, Pan Y-m: **Antioxidant activity and inhibition effect on the growth of human colon carcinoma (HT-29) cells of esculetin from Cortex Fraxini.** *Medicinal Chemistry Research* 2011, **20**:968-974.
39. Arora R, Sawney S, Saini V, Steffi C, Tiwari M, Saluja D: **Esculetin induces antiproliferative and apoptotic response in pancreatic cancer cells by directly binding to KEAP1.** *Molecular cancer* 2016, **15**:64.
40. Han MH, Park C, Lee DS, Hong SH, Choi IW, Kim GY, Choi SH, Shim JH, Chae JI, Yoo YH, Choi YH: **Cytoprotective effects of esculetin against oxidative stress are associated with the upregulation of Nrf2-mediated NQO1 expression via the activation of the ERK pathway.** *International journal of molecular medicine* 2017, **39**:380-386.
41. Mangali S, Bhat A, Udumula MP, Dhar I, Sriram D, Dhar A: **Inhibition of protein kinase R protects against palmitic acid-induced inflammation, oxidative stress, and apoptosis through the JNK/NF- κ B/NLRP3 pathway in cultured H9C2 cardiomyocytes.** *Journal of cellular biochemistry* 2019, **120**:3651-3663.
42. Jin JL, Lv RG, Guo J, Liu XH, Liang YW, Wei JR, Wang L: **Improvement of Left Ventricular Remodelling by Inhibition of NF- κ B in a Rat Model of Myocardial Infarction.** *Heart, lung & circulation* 2016, **25**:1007-1012.
43. Ma L, Sun P, Zhang JC, Zhang Q, Yao SL: **Proinflammatory effects of S100A8/A9 via TLR4 and RAGE signaling pathways in BV-2 microglial cells.** *International journal of molecular medicine* 2017, **40**:31-38.
44. Zhang S, Zhang Y: **Isoflurane reduces endotoxin-induced oxidative, inflammatory, and apoptotic responses in H9c2 cardiomyocytes.** *European review for medical and pharmacological sciences* 2018, **22**:3976-3987.

45. Zhu X, Gu J, Qian H: **Esculetin Attenuates the Growth of Lung Cancer by Downregulating Wnt Targeted Genes and Suppressing NF- κ B.** *Archivos de bronconeumologia* 2018, **54**:128-133.
46. Chen T, Guo Q, Wang H, Zhang H, Wang C, Zhang P, Meng S, Li Y, Ji H, Yan T: **Effects of esculetin on lipopolysaccharide (LPS)-induced acute lung injury via regulation of RhoA/Rho Kinase/NF- κ B pathways in vivo and in vitro.** *Free radical research* 2015, **49**:1459-1468.
47. Geetha A, Sankar R, Marar T, Devi CS: **Alpha-tocopherol reduces doxorubicin-induced toxicity in rats—histological and biochemical evidences.** *Indian journal of physiology and pharmacology* 1990, **34**:94-100.
48. Li Y, Feng J, Mo Y, Liu H, Yang B: **Concordance between cardio-protective effect on isoproterenol-induced acute myocardial ischemia and phenolic content of different extracts of *Curcuma aromatica*.** *Pharmaceutical biology* 2016, **54**:3226-3231.
49. Yates JC, Dhalla NS: **Induction of necrosis and failure in the isolated perfused rat heart with oxidized isoproterenol.** *Journal of molecular and cellular cardiology* 1975, **7**:807-816.
50. Rajadurai M, Prince PS: **Preventive effect of naringin on isoproterenol-induced cardiotoxicity in Wistar rats: an in vivo and in vitro study.** *Toxicology* 2007, **232**:216-225.
51. Brindha E, Rajasekapandiyam M: **Preventive effect of phytic acid on lysosomal hydrolases in normal and isoproterenol-induced myocardial infarction in Wistar rats.** *Toxicology mechanisms and methods* 2015, **25**:150-154.
52. Dhivya V, Priya LB, Chirayil HT, Sathiskumar S, Huang CY, Padma VV: **Piperine modulates isoproterenol induced myocardial ischemia through antioxidant and anti-dyslipidemic effect in male Wistar rats.** *Biomedicine & pharmacotherapy = Biomedecine & pharmacotherapie* 2017, **87**:705-713.
53. Karthika P, Rajadurai M, Ganapathy P, Kanchana GJJPR: **Preventive effect of esculetin on lipid peroxides and antioxidants in isoproterenol-induced myocardial infarction in Wistar rats.** 2012, **5**:915-918.
54. Roy SJ, Stanely Mainzen Prince P: **Protective effects of sinapic acid on lysosomal dysfunction in isoproterenol induced myocardial infarcted rats.** *Food and chemical toxicology : an international journal published for the British Industrial Biological Research Association* 2012, **50**:3984-3989.
55. George J: **Elevated serum beta-glucuronidase reflects hepatic lysosomal fragility following toxic liver injury in rats.** *Biochemistry and cell biology = Biochimie et biologie cellulaire* 2008, **86**:235-243.
56. Karthikeyan K, Sarala Bai BR, Niranjali Devaraj S: **Grape seed proanthocyanidins ameliorates isoproterenol-induced myocardial injury in rats by stabilizing mitochondrial and lysosomal enzymes: an in vivo study.** *Life sciences* 2007, **81**:1615-1621.

Figures




Figure 1

Effect of esculletin on A) DPPH radical scavenging potentials of esculletin B) Intracellular ROS alleviating effect in H9C2 cell lines. Values are presented as the mean \pm SD. All the treatments were compared with control and $P < 0.05$ has been considered as significant.




Figure 2

Effect of esculentin on mRNA expression levels against arsenic induced ROS in H9c2 cells. The mRNA levels of (A) TNF- α , (B) IL 6 and (C) NF- κ B were measured by using RT-qPCR. Values are presented as the mean \pm SD. All the treatments were compared with control and $P < 0.05$ has been considered as significant.




Figure 3

(A–D): Histopathological study of the heart. A): Normal control rat's heart showing normal texture of cardiac muscle bundles. B) ISO induced MI rat's heart showing area of infarction with inflammatory cell infiltration, cardiac necrosis and splitting of muscle bundles. C&D) pre-treatment with esculetin showing the prevention of cell infiltration, splitting of muscle bundle by ISO has been prevented.