

Hydroxychloroquine for Post-Exposure Prophylaxis of COVID-19 among naval personnel in Sri Lanka: study protocol for a randomized, controlled trial

Madunil Anuk Niriella (✉ maduniln@yahoo.co.uk)

University of Kelaniya Faculty of Medicine <https://orcid.org/0000-0002-7213-5858>

Dileepa Senajith Ediriweera

University of Kelaniya Faculty of Medicine

Arjuna Priyadarshin De Silva

University of Kelaniya Faculty of Medicine

Ranjan Premarathna

University of Kelaniya Faculty of Medicine

Priyantha Balasooriya

Naval Hospital, Welisara

K D Duminda

Army Hospital, Narahenpita

Neelika Gathsaurie Malavige

Faculty of Medical Sciences, University of Sri Jayawardenapura

Kamani Wanigasuriya

Faculty of Medical Sciences, University of Sri Jayawardenapura

Sarath Lekamwasam

University of Ruhuna Faculty of Medicine

S A Kularathne

University of Peradeniya Faculty of Medicine

Sisira Siribaddana

Rajarata University of Sri Lanka Faculty of Medicine and Allied Sciences

Hithanadura Janaka de Silva

University of Kelaniya Faculty of Medicine

Saroj Jayasinghe

University of Colombo Faculty of Medicine

Study protocol

Keywords: Randomized controlled trial, COVID-19, SARS-CoV-2, prophylaxis, post-exposure, hydroxychloroquine, HCQ, Sri Lanka

Posted Date: August 10th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-28949/v2>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published on August 27th, 2020. See the published version at <https://doi.org/10.1186/s13063-020-04659-7>.

Abstract

Background: The first case of a corona virus 2019 (COVID-19) infection in a Sri Lankan was reported on March 11, 2020. The situation in Sri Lanka changed with the rapid increase of personnel contracting COVID-19 in a Naval base camp that housed more than 4000 people. This provided a unique opportunity to study the effectiveness of hydroxychloroquine (HCQ) for post-exposure prophylaxis (PEP), while taking stringent, non-pharmacologic, public health measures to prevent spread. Our aim is to study the effectiveness and safety of HCQ for PEP among naval personnel with exposure to COVID-19 positive patients.

Methods/design: This is a placebo-controlled, randomized, clinical trial carried out in the Naval base camp and quarantine centers of the Sri Lanka Navy, Ministry of Defense, Sri Lanka. Navy personnel who are exposed to a patient with confirmed COVID-19 infection but test negative for the virus on reverse, real-time polymerase chain reaction (rRT-PCR) at recruitment will be randomized, 200 to each arm, to receive HCQ or placebo, and monitored for the development of symptoms or rRT-PCR positivity for severe acute respiratory syndrome corona virus 2 (SARS-CoV-2) virus for 14 days.

Discussion: This trial will provide high-quality evidence of the effectiveness and safety of HCQ as PEP for COVID-19. The study design is unique due to the circumstances of the outbreak in a confined area among otherwise healthy adults, at a relatively early stage of its spread.

Trial registration: This trial was registered in the Sri Lanka Clinical Trials Registry (SLCTR) – SLCTR/2020/011 on 04.05.2020 (<https://slctr.lk/trials/slctr-2020-011>)

Background

The novel coronavirus 2019 (COVID-19) epidemic is a massive threat to public health worldwide. Current estimates suggest that the Severe Acute Respiratory Syndrome Corona Virus 2 (SARS-CoV-2), responsible for COVID-19 illness, is both highly contagious (estimated basic reproductive rate, 2-3) and five to fifty-fold more lethal than seasonal influenza (estimated mortality rate, 0.5-5%) (1). Interventions to decrease the incidence and severity of COVID-19 are urgently needed.

The first case of COVID-19 in a Sri Lankan was reported on March 11, 2020, in a tour guide who was exposed to European tourists. Since then, the cumulative numbers have increased. After a critical evaluation of evidence, the use of hydroxychloroquine (HCQ) in symptomatic patients with COVID-19 was included in a circular issued by the Ministry of Health, Sri Lanka, on March 27, 2020. Since then, the possible use of HCQ for pre- or post-exposure prophylaxis (PEP) has been intensely debated within the medical profession.

HCQ is an anti-malarial medication with immunomodulatory effects. HCQ has been shown to clear COVID-19 infection status and reduce viral shedding early during the disease course (2, 3). Those against its introduction for prophylaxis cited the poor quality of the study designs and lack of reproducible results elsewhere. However, there were in-vitro studies showing that it elevated endosomal pH, slowed the entry of COVID-19 into cells, and inhibited its replication (4, 5). The drug is inexpensive, has a good safety profile - especially with short term use, and used extensively in the past to prevent malaria and currently used to treat autoimmune diseases (4, 6).

Clinical trials have been commenced in other countries to assess the efficacy of hydroxychloroquine as pre- or PEP to reduce the attack rate or severity among contacts of known or suspected COVID-19 patients (7-10). India commenced pre-exposure prophylaxis of healthcare workers in March 22, 2020 (11). In mid-April, 2020 Indian scientists commenced mass prophylaxis of slum communities in Mumbai (12).

By late April there were four studies from USA, UK, Singapore, Canada to evaluate the role of HCQ in prophylaxis, registered in ClinicalTrials.gov (13-16).

The situation in Sri Lanka changed with the rapid increase of personnel contracting COVID-19 in a Navy camp housing more than 4000 people in an area of about 52 acres. This was believed to have been the result of navy personnel being exposed to infected patients during contact tracing. This provided a unique opportunity to study the effectiveness and safety of HCQ in prophylaxis while taking stringent, non-pharmacologic, public health measures to prevent spread. Our research question is, "is HCQ effective and safe for PEP among naval personnel with exposure to COVID-19 positive patients?".

Methods

Study design

This will be a 2-arm, placebo-controlled, randomized, clinical trial with parallel group, 1:1 allocation ratio. The study will be carried out among all eligible persons meeting the inclusion/exclusion criteria at the Naval Base camp, Welisara, Sri Lanka who are directed to be quarantined in quarantine centers (QCs) of the Sri Lanka Navy, Ministry of Defense, Sri Lanka. A patient information leaflet will be provided and informed written consent will be obtained. Those willing to participate in the trial will be randomized and monitored for development of symptoms or real-time reverse polymerase chain reaction (rRT-PCR) positivity for SARS-CoV-2 virus. All eligible participants who test negative for the virus on rRT-PCR at baseline, will be randomly assigned to the study arms within 48 hours. Permission will be obtained through the Navy to commence a double-blind randomized trial of HCQ to investigate its effectiveness and safety to prevent development of COVID-19 infection in persons who are exposed but not yet infected.

Study setting

Naval Base camp, Welisara, Sri Lanka and the designated QCs maintained by the Sri Lanka Navy, Ministry of Defense, Sri Lanka.

Study population

All the Naval personnel who are currently being directed for quarantine in QCs of the Sri Lanka Navy, Ministry of Defense, Sri Lanka.

Inclusion criteria

1. All consenting male and female adult Naval personnel (18-59.9 years of age)
2. Exposure to a patient with confirmed COVID-19 infection (rRT-PCR positive for SARS-CoV-2 virus)
3. The exposure should be for >30 mins within 2 meters of the infected individual and should be within ONE WEEK at the time of inclusion in the study
4. Recruited within 48 hours of admission to these centers
5. Willing to take study drug as directed for 5 days.

Exclusion criteria

1. Pregnant and lactating females

2. Suspected or confirmed to have COVID-19 infection (i.e Navy personnel having positive rRT-PCR test for COVID-19 or presence of fever with cough, shortness of breath, sore throat, anosmia or diarrhoea at the time of recruitment)
3. Presence of pre-existing cardiovascular disease including being on medication for ischaemic heart disease
4. Presence of contraindication to the use of HCQ (on other medications predisposing to long QT interval or presence of long QT interval on baseline ECG)
5. Prior diagnosis of retinopathy, G6PD, malignancy or advanced kidney disease
6. Known sensitivity or allergy to hydroxychloroquine

Outcome measure

All participants will be monitored for the development of symptoms suggestive of COVID-19 and will undergo a rRT-PCR test for SARS-CoV-2 virus at the end of their quarantine period of 14-days.

Primary outcome measure

1. Number of participants testing positive for SARS-CoV-2 virus using rRT-PCR on day 14 at the completion of quarantine period

Secondary outcome measures

1. Number of participants symptomatic for COVID-19 (symptomatic illness will be defined as presence of fever with cough, shortness of breath, sore throat, anosmia or diarrhoea from the time of recruitment until the completion of quarantine period of 14 days).
2. Any reported adverse effects of HCQ

Sample size

Altogether 400 participants will be recruited to the study. Among them, 200 each will be allocated to the two arms: intervention and control arms.

Sample size calculation

We used a simulation-based method to estimate design power. Required sample size was determined by simulating 1000 samples for different sample sizes, and the minimum sample size which resulted in an 80% power to detect a significant intervention effect was selected. We considered below assumptions to simulate data.

Assumptions:

1. We assumed 50% of the study population have COVID-19 infections (i.e. but not diagnosed) at the time of recruitment.
2. If no intervention is adopted (i.e. control arm), we assumed 83.5% of the sample would be infected with COVID-19 infections (i.e. 66.7% increase) after 2 weeks.
3. If intervention initiated, we expect 25% reduction in the natural increase of COVID-19 infections after 2 weeks. Therefore, the intervention arm will experience 71.4% COVID-19 infections.

Our simulation-based experiment showed the required minimal sample size as 190 for each arm to achieve 80% power. We considered additional 10% (i.e. $190 / (1-0.1)$) to allow loss to follow ups and non-response bias. Therefore, the total required sample size for the study will be 211 persons for each arm.

Consent, recruitment and randomization

Recruitment for the study will be performed in the main Naval base camp in Welisara, Sri Lanka. The chief medical officer of the Naval camp base, Welisara, Sri Lanka is responsible for the briefing of the study details to potential participants, obtaining informed consent and recruitment. Those who are willing to participate will be formally recruited after obtaining informed written consent.

Permuted block randomization method will be used to allocate the participants to intervention and placebo groups. Placebo will be produced similar to HCQ tablets in both size and shape. Allocation concealment will be maintained by packing the interventional product and the placebo in sequentially numbered, opaque, sealed envelopes with similar appearance.

These envelopes will be numbered sequentially from 1 to 400 and will be handed over to the Naval base camp in Welisara, Sri Lanka without revealing which envelopes contain HCQ and placebo to ensure double blindness.

After randomization, study participants will be handed over to the concealed intervention. Thereafter, the participants will be moved to designated quarantine centres maintained by the Sri Lanka Navy. Once the participants reach the QC, they will be monitored for continuation of intervention, compliance and adverse effects by a designated Chief nursing officer in each QC. There are 45 QCs maintained by the Sri Lanka Navy, Ministry of Defense, Sri Lanka distributed throughout Sri Lanka.

- **Arm 1: Intervention arm (n=200)**

Participants will receive a loading dose of oral HCQ 400mg (2 tablets) 12 hourly Day 1 followed by oral HCQ 200mg (1 tablets) 12 hourly for the next 4 days.

- **Arm 2: Control arm (n=200)**

Participants will receive an oral matching placebo (oral tablet containing 100mg of elemental calcium) 2 tablets 12 hourly Day 1, 12 hourly on Day 1, followed by 1 tablet orally, 12 hourly for the next 4 days.

Trial steering committee (TSC)

The trial steering committee comprised several authors (MAN, JB and HJdeS). This committee will be responsible for the overseeing of the recruitment, randomization as well as monitoring the adherence to intervention, any reported adverse effects and interim analysis. The TSC will also be responsible for communication with the ERC. This committee will meet once a week to oversee the trial conduct and progress. An unblinded statistician (author DSE) will audit the trial every two weeks and provide recruitment data to TSC.

The TSC is responsible for any protocol amendments and to notify the sponsor, Ethics review committee and Sri Lanka Clinical Trial Registry. The PI will notify the quarantine centres regarding the amendments and a copy of the revised protocol will be sent to the investigator site file. Any deviations from the Protocol will be fully documented using a breach report form.

Data collection method

Two trained doctors of the Sri Lanka Navy, in personal protective equipment, will obtain informed written consent from all the participants at the Naval base camp, Welisara, Sri Lanka. Demographic details, co-morbidities and exposure history will be collected via an interviewer administered questionnaire at the recruitment. An ECG will be performed at the Navy Hospital, Welisara, Sri Lanka on all participants to assess for pre-existing prolonged QT interval. A rRT-PCR for SARS-CoV-2 virus will be performed on all participants prior to recruitment (which was done as part of the existing screening program for COVID-19 of the Sri Lanka Navy) and after the completion of the quarantine period of 14-days. This sample collection for rRT-PCR will be performed by the already available, trained SL Navy personnel designated to perform this task, with the necessary precautions including personal protective equipment.

Samples from the naso-pharynx will be obtained and transported in viral transport media and shipped at 4° C to the laboratory at The Center for Dengue Research, Faculty of Medical Sciences, University of Sri Jayawardenapura, Sri Lanka, for analyses by one of the authors (NM) who will be blinded to the randomization. Confirmation of cases of COVID-19 will be based on detection of unique nucleic acid sequences of virus RNA by rRT-PCR (17). The collected biological material once used for the present study will not be stored or used for future any ancillary studies.

All enrolled participants will be monitored for possible adverse effects of the intervention and the placebo for the period of the study. In case of development of symptoms suggestive of COVID-19 infection or any adverse events, a participant will be admitted to the Navy Base Hospital, Welisara, Sri Lanka.

Data analysis

All main analyses will be based on an intention to treat principle. Baseline characteristics will be summarized by treatment group to characterize the study sample and identify potential baseline imbalance. Post interventional prevalence of SARS-CoV-2 virus for the control and intervention group will be calculated and presented. Subgroup analysis will be done for gender and age categories. A generalized linear mixed effect model will be used to evaluate the effectiveness of HCQ as PEP. A secondary analysis include adjustment for potential confounders including body weight and metabolic co-morbidities (i.e. diabetes, hypertension and dyslipidaemia); multi-level models will address the dependence due to quarantine centres (i.e. random intercept will be introduced to for each QC). Additional analyses will include interactions to determine whether the effects differ by gender or age categories. Treatment effect will be presented using odds ratio with 95% confidence intervals and P values for the treatment effect will be presented.

If the drop-out process is missing completely at random (MCAR), unbiased estimates of the intervention effects at post-randomization will be obtained. If missing data patterns are found to depend on specific baseline covariates, inverse-probability weighted generalized estimating equations (GEE) can be used to account for this in order to test sensitivity of results to assumptions. If the models do not converge, multicollinearity of the variables and outliers will be addressed and refit the models and number of iteration will be increased and reanalysis will be conducted.

R programming language version 3.6.3 will be used for analyses.

Ethical clearance

Ethical clearance for the study was obtained from the Ethics Review Committee (ERC) of the Faculty of Medicine, University of Kelaniya, Ragama – P/22/04/2020

Trial registration

This trial was registered in the Sri Lanka Clinical Trails Registry (SLCTR) – SLCTR/2020/011 on 04.05.2020 (<https://slctr.lk/trials/slctr-2020-011>) and authorization for the trial was obtained from the National Medicinal Regulatory Authority (NMRA) – CLITRI/2020/00032. WHO Universal trial number (UTN) for the study is U1111-1251-3613.

Main ethical issues

All participants will be provided with a detailed information sheet in their preferred language. They will be allowed to provide informed written consent without undue influence. The collected biological material once used for the present study will not be stored or used for future any ancillary studies.

HCQ has been used for a long period of time for treatment of a variety of illnesses and its safety profile is well documented (6). At the dosage given in this study investigators do not envisage significant challenges with its safety. There is no anticipated harm and compensation for trial participation.

However, participants will be monitored daily, and all instances of adverse events will be documented. Furthermore, this trial will include generally fit participants (i.e. Navy personnel on active duty) who have had medical examinations. All participants will have an ECG to exclude pre-existing prolongation of the QT interval.

All serious adverse events and suspected unexpected serious adverse reactions (SUSARs) will be reported according to guidelines issued by the National Medicines Regulatory Authority, Sri Lanka (NMRA) to the NMRA and ERC. Serious adverse events and SUSARs that require expedited reporting will be reported within stipulated timelines. All adverse events will be managed as appropriate by the trial physicians.

Trial progress will be monitored by an unblinded trial statistician. The trial will be terminated if there is an unexpectedly high serious adverse events rate among trial participants and / or evidence of futility. Further, unblinded data will be sent weekly to the Ethical Review Committee (which will in effect act as a data monitoring committee for this study).

Budget

This clinical trial will be done in collaboration with the State Pharmaceutical Manufacturing Cooperation and Sri Lanka Navy, Ministry of Defense. All the necessary rRT-PCR test kits, drug and placebo for the study have been donated to the SL Navy. rRT-PCR testing will be done free of charge at the Department of Microbiology, Faculty of Medical Sciences, University of Sri Jayewardenepura, in accordance with the already existing protocol of assessment of quarantined persons.

Discussion

The study has a unique design due to the outbreak of COVID-19 infection in a confined area, among otherwise healthy adults at a relatively early stage of its spread. The space of 52 acres houses naval personnel in dormitory-type accommodation was ideal for rapid spread, and immediate measures were taken to send the initial positive cases to hospitals, transfer contacts to other QCs, and for segregation of personnel to minimize contact. Other preventive measures such as handwashing and physical distancing are also stringently followed. The communication of the completed trial results will be via a publication in a peer-reviewed, high impact journal.

Trial status

Trial protocol version 2.0 dated May 02, 2020 was granted ethical approval and regulatory registration. Recruitment began on May 04, 2020, and the approximate date on which recruitment will be completed will be May 31, 2020.

List Of Abbreviations

SARS-CoV-2	Severe acute respiratory syndrome corona virus 2
COVID-19	Novel coronavirus infection 2019
HCQ	Hydroxychloroquine
PEP	post-exposure prophylaxis
QCs	quarantine centers
rRT-PCR	real-time reverse-polymerase chain reaction
ERC	Ethic Review Committee

SLCTR Sri Lanka Clinical Trails Registry

NMRA National Medicinal Regulatory Authority

Declarations

Ethics approval and consent to participate

Ethical clearance for the study was obtained from the Ethics Review Committee (ERC) of the Faculty of Medicine, University of Kelaniya, Ragama – P/22/04/2020. Written, informed consent to participate will be obtained from all participants.

Consent for publication

Not applicable.

Availability of data and materials

Individual participant data which is used to generate results will be shared immediately following publication after de-identification (text, tables, figures and appendices). Study protocol, statistical analysis plan and analytic code will be made available. Data will be shared with investigators whose proposed use of the data has been approved by an independent review committee identified for this purpose. Data will be shared to achieve the aims in an approved proposal. Proposals should be directed to maduniln@yahoo.co.uk. To gain access, data requestors will need to sign a data access agreement. Similarly, the participant information sheet, consent form and data collection form are available from the corresponding author (MAN) on request.

Competing interests

The authors declare that they have no competing interests.

Funding and sponsor

This clinical trial will be done in collaboration with the State Pharmaceutical Manufacturing Cooperation and Sri Lanka Navy, Ministry of Defense. All the necessary rRT-PCR test kits, drug and placebo for the study have been donated to the SL Navy. rRT-PCR testing will be done free of charge at the Department of Microbiology, Faculty of Medical Sciences, University of Sri Jayewardenepura, in accordance with the already existing protocol of assessment of quarantined persons. Trial sponsors had no role in the study design; collection, management, analysis, and interpretation of data; writing of the report; and the decision to submit the report for publication.

Authors' contributions

MAN, HJdeS and SJ are the Chief Investigators; MAN, KDD and SJ she conceived the study, led the proposal and protocol development. ADeS, BHRP, DSE, PB, GNM, CW, SL, SAMK, SS contributed to study design and to development of the proposal. MAN, DSE, HJdeS, and SJ are the lead trial methodologists. All authors read, approved and agreed to the publication of the final manuscript. All named authors adhere to the authorship guidelines of Trials.

Acknowledgements

The authors wish to acknowledge Lieutenant General Shavendra Silva, Chief of Defense Staff and Head of the National Operation Center for the Prevention of COVID- 19, Sri Lanka, Vice Admiral Piyal de Silva, Commander of the Sri Lanka Navy and Dr Lalith Mendis for their support to carry out this study.

References

1. Cevik, C Bamford and A. Ho. COVID-19 pandemic – A focused review for clinicians. *Clinical Microbiology and Infection*, 2020
2. Gao, Z. Tian, and X. Yang, “Breakthrough: Chloroquine phosphate has shown apparent efficacy in treatment of COVID-19 associated pneumonia in clinical studies.,” *Biosci. Trends*, pp. 1–2, 2020.
3. Schrezenmeier and T. Dörner, “Mechanisms of action of hydroxychloroquine and chloroquine: implications for rheumatology,” *Nat. Rev. Rheumatol.*, 2020
4. Liu, J., Cao, R., Xu, M. et al. Hydroxychloroquine, a less toxic derivative of chloroquine, is effective in inhibiting SARS-CoV-2 infection in vitro. *Cell Discov* 2020;6:16. <https://doi.org/10.1038/s41421-020-0156-0>
5. Wang et al., “Remdesivir and chloroquine effectively inhibit the recently emerged novel coronavirus (2019-nCoV) in vitro,” *Cell Res.*, no. January, pp. 2019–2021, 2020.
6. Das SK, Pareek A, Mathur DS, et al. Efficacy and safety of hydroxychloroquine sulphate in rheumatoid arthritis: a randomized, double-blind, placebo controlled clinical trial—an Indian experience. *Curr Med Res Opin.* 2007;23(9):2227-2234. doi:10.1185/030079907X219634\
7. Hydroxychloroquine Post Exposure Prophylaxis for Coronavirus Disease (COVID-19) <https://clinicaltrials.gov/ct2/show/NCT04318444>
8. Safety and Efficacy of Hydroxychloroquine for at Risk Population (SHARP) Against COVID-19 (SHARP COVID-19) <https://clinicaltrials.gov/ct2/show/NCT04342156>
9. Chloroquine/ Hydroxychloroquine Prevention of Coronavirus Disease (COVID-19) in the Healthcare Setting (COPCOV). <https://clinicaltrials.gov/ct2/show/NCT04303507>
10. Post-exposure Prophylaxis / Preemptive Therapy for SARS-Coronavirus-2 (COVID-19 PEP) <https://clinicaltrials.gov/ct2/show/NCT04308668>
11. National Taskforce for COVID-19. Advisory on the use of hydroxy-chloroquine as prophylaxis for SARS-CoV-2 infection. 2020. <https://www.mohfw.gov.in/pdf/AdvisoryontheuseofHydroxychloroquinasprophylaxisforSARSCoV2infection.pdf> (accessed 7 May 2020)
12. Shelar J. BMC to try out hydroxychloroquine as prophylaxis in Dharavi, Worli Koliwada. *The Hindu.* 2020. <https://www.thehindu.com/news/cities/mumbai/bmc-to-try-out-hydroxychloroquine-as-prophylaxis-in-dharavi-worli-koliwada/article31335133.ece> (accessed 7 May 2020)
13. Hydroxychloroquine Post Exposure Prophylaxis for Coronavirus Disease (COVID-19) <https://clinicaltrials.gov/ct2/show/NCT04318444>. Accessed May 07, 2020.
14. Safety and Efficacy of Hydroxychloroquine for at Risk Population (SHARP) Against COVID-19 (SHARP COVID-19) <https://clinicaltrials.gov/ct2/show/NCT04342156>. Accessed May 07 2020.
15. Chloroquine/ Hydroxychloroquine Prevention of Coronavirus Disease (COVID-19) in the Healthcare Setting (COPCOV). <https://clinicaltrials.gov/ct2/show/NCT04303507>. Accessed May 07, 2020.
16. Post-exposure Prophylaxis / Preemptive Therapy for SARS-Coronavirus-2 (COVID-19 PEP). <https://clinicaltrials.gov/ct2/show/NCT04308668>. Accessed May 07,2020.
17. Laboratory testing for coronavirus disease (COVID-19) in suspected human cases. <https://apps.who.int/iris/rest/bitstreams/1272454/retrieve>. Accessed May 07, 2020

Figures

TIMEPOINT	STUDY PERIOD					Close-out
	Enrolment	Allocation	Post-allocation			
	$-t_1$	5 days	Week 1	Week 2	Week 3	
ENROLMENT:						
Eligibility screen	X					31 May 2020
Informed consent	X					31 May 2020
Allocation		X				31 May 2020
INTERVENTIONS:						
HCQ		X				31 May 2020
Placebo		X				31 May 2020
ASSESSMENTS:						
Symptoms of COVID-19	X		X	X		15 June 2020
rRT-PCR for COVID-19	X				X	15 June 2020

Figure 1

SPIRIT Figure for the trial - the schedule of enrolment, interventions, and assessments

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SPIRITFillablechecklist31.07.20.pdf](#)