Risk analysis of 2-year mortality in elderly male military veterans with non-thyroidal illness syndrome

Xinyu Miao

Chinese PLA General Hospital

Hongzhou Liu

Chinese PLA General Hospital

Xiaomin Fu

Chinese PLA General Hospital

Huiping Ma

Handan Central Hospital

Xuefeng Chen

First People's Hospital of Wu'an City

Zhaoyan Gu

Chinese PLA General Hospital

Banruo Sun

Chinese PLA General Hospital

Nan Li

Chinese PLA General Hospital

Chunlin Li

Chinese PLA General Hospital

Shuangtong Yan (✉ ystf@yeah.net)

Chinese PLA General Hospital <https://orcid.org/0000-0003-2344-4504>

Zhaohui Lyu

Chinese PLA General Hospital

Research article

Keywords: NTIS, mortality, old age, inpatients

Posted Date: November 6th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-29177/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background: Elderly patients with non-thyroidal illness syndrome (NTIS) have relatively severe symptoms and a poor prognosis. However, there are few studies on the correlation between NTIS and mortality among hospitalised elderly patients. This study aimed to investigate the characterization of NTIS inpatients with biochemical indicators and mortality prediction.

Methods: In the present study, 931 male veteran inpatients ≥ 60 years of age who visited our hospital from January 2012 to December 2013 were selected and divided into the NTIS group ($n = 193$) and normal thyroid function (non-NTIS) group ($n = 738$). Following propensity score matching to match the two groups according to age and body mass index, the NTIS group and non-NTIS group consisted of 192 and 660 patients, respectively. Data on biochemical indicators and mortality were collected.

Results: Patients had more primary care and more respiratory disease and chronic kidney disease in the NTIS than in the non-NTIS group. Serum total protein (TP), albumin (Alb), prealbumin (PA), haemoglobin (Hb), uric acid (UA), triglyceride (TG), and high-density lipoprotein cholesterol (HDL) levels were significantly lower, and fasting blood glucose (FBG) and urea nitrogen (UN) levels were higher, in the NTIS than in the non-NTIS group. Triiodothyronine, TP, Alb, and PA levels correlated positively with the Hb level and negatively with FBG, UN, and creatinine (Cr) levels. The free T_3 level correlated positively with TP, Alb, PA, Hb, and UA levels and negatively with FBG, UN, and Cr levels. A lower free T_3 level was associated with increased all-cause mortality after adjusting for covariates. Patients in the NTIS group had a lower survival rate at 6-month (78.65% vs. 97.73%), 1-year (68.23% vs. 96.97%) and 2-year (64.58% vs. 86.52%). Receiver operating characteristic curve (ROC) analysis showed that a cut-off free T_3 level of 3.45 pmol/L yielded the highest sensitivity and specificity for predicting all-cause mortality at 2 years.

Conclusion: Among elderly male inpatients, the survival rate was lower in the NTIS group. Serum protein levels and renal function decreased and the FBG level increased with decreasing triiodothyronine and free T_3 levels. A decreased free T_3 level in hospitalized elderly male patients was a predictor of poor prognosis.

Background

Many changes in thyroid hormones occur as a result of illness or nutritional deprivation. These changes consist of decreased levels of serum triiodothyronine (T_3) and/or thyroxine (T_4), without an increase in the thyroid-stimulating hormone (TSH) level, and usually accompanied by elevated reverse T_3 (rT_3) [1, 2]. The combination of these findings is termed non-thyroidal illness syndrome (NTIS), euthyroid sick syndrome, or low T_3 syndrome, indicating a systemic disease outside the thyroid that causes abnormal levels of thyroid hormones and is often considered a compensatory mechanism for the body. NTIS has been reported in patients with acute and chronic illnesses, including infectious diseases, cardiovascular and gastrointestinal diseases, cancer, and trauma [3], which are quite common in patients in intensive care units (ICUs) [4, 5]. In previous studies, NTIS showed a high sensitivity and specificity for predicting patient mortality in ICU patients [6]. Serum T_3 levels further decrease as the severity of disease progresses. However, among hospitalised elderly patients in the general ward, data on the correlation between NTIS and patient mortality are lacking.

Elderly patients often have multiple chronic diseases and a poor nutritional status; thus, NTIS is quite common in these patients. [Tognini](#) [7] reported that among elderly patients (≥ 65 years of age) hospitalised for acute illness, the prevalence of NTIS was 31.9% and significantly associated with acute renal failure, New York Heart Association classification IV heart failure, and metastatic cancer. We identified elderly patients who were hospitalised in our hospital. Triiodothyronine, total T_4 , free T_3 , free T_4 , thyrotropin and serum levels of total protein (TP), albumin (Alb), prealbumin (PA), urea nitrogen (UN), creatinine (Cr), uric acid (UA), fasting blood glucose (FBG), blood lipids, alanine transaminase (ALT), aspartate transaminase (AST), and haemoglobin (Hb) were analysed, and 6-month, 1-year and 2-year all-cause mortality rates based on free T_3 levels was assessed. This study investigated the characterization of NTIS patients with biochemical indicators and mortality prediction that may provide a basis for predicting disease outcome and mortality in hospitalised elderly male patients with NTIS.

Methods

Patients

The inclusion criteria for the study were as follows: 1) ≥ 60 years of age, 2) male, 3) hospitalised between January 2012 and December 2013, and 4) complete data on thyroid function obtained within 1 day of admission. The exclusion criteria were as follows: 1) diagnosed with thyroid disease such as hyperthyroidism, hypothyroidism, subclinical hyperthyroidism, subclinical hypothyroidism, or Hashimoto's thyroiditis, 2) taking amiodarone, euthyroid, glucocorticoid, dopamine, or interferon, and 3) history of thyroid surgery. Because the patients in our hospital are Chinese military veterans, there were few female patients. As such, only male patients were selected as research subjects to reduce the influence of sex bias on the results. Within 2 years, a total of 9,832 patients were admitted to the hospital, 931 of whom were enrolled in the present study. All patients were followed up for 2 years, until December 2015. The subjects were divided based on the presence of NTIS into the NTIS group and normal thyroid function (non-NTIS) group. Because the baseline age and body mass index (BMI) in the two groups differed significantly, the groups were matched by age and weight using the propensity score matching (PSM). After matching, 852 patients were included in the study: 192 in the NTIS group and 660 in the non-NTIS (normal thyroid function) group. At the end of the 2-year follow-up, 157 patients had died and 45 were lost to follow-up (5.28%). The time and cause of death were recorded for all deceased patients.

Study protocol

Venous blood samples were collected early in the morning after an overnight fast on the second day after admission. Total T_4 , triiodothyronine, free T_3 , free T_4 , TSH, TP, Alb, PA, UN, Cr, UA, ALT, AST, and Hb levels were measured. Radioimmunoassay was used to measure the serum levels of triiodothyronine, free T_3 , total T_4 , free T_4 , and TSH. The Sysmex Xt-1800 Automated Hematology Analyzer (SYSMEX Corporation, Japan) was used for routine blood tests. Biochemistry measurements were performed using i-CHROMA Reader (Boditech Med Inc. Korea). Normal thyroid hormone levels are as follows: total T_4 (55.34–160.88 nmol/L), triiodothyronine (1.01–2.95 nmol/L), free T_3 (2.76–6.3 pmol/L), free T_4 (10.42–24.32 pmol/L), and TSH (0.35–5.5 uIU/mL). Intra-batch and batch-to-batch variations were $< 3.35\%$ and $< 5.04\%$, respectively.

Diagnostic criteria for NTIS

Decreased serum triiodothyronine and/or free T₃, normal or mildly reduced total T₄ or free T₄, and normal TSH levels were the diagnostic criteria for NTIS [8].

Statistical analysis

PSM (1:4) was used to account for demographic differences in the selected participants. SPSS 22.0 was used for the statistical analysis. Normally distributed data are expressed as means ± standard deviation ($\bar{x} \pm SD$) and non-normally distributed data as quartiles. Normally distributed data were compared between the two groups using *t*-test or *t'*-test. The rank sum test was used to compare the data distribution between the two groups. All data were evaluated using two-sided tests. Correlations were determined using Pearson correlation analysis for normally distributed data and Spearman’s test for non-normally distributed data. Overall survival at 6-month, 1-year and 2-years was estimated using the Kaplan–Meier method. Bivariate and multivariate Cox proportional hazards models were used to evaluate the risk factors associated with patient mortality. Statistical significance was set at *P* < 0.05.

Results

Comparison of baseline data between the NTIS and non-NTIS groups

To minimize research bias, PSM was used to match the two groups according to two major factors that influence metabolic indicators and mortality: age and BMI. After matching, statistically significant differences in age and BMI were not observed (see Additional file 1). And after matching by age and BMI, the triiodothyronine, free T₃, total T₄ (all *P* < 0.001), and TSH levels (*P* = 0.001) were significantly lower in the NTIS group than in the non-NTIS group, moreover, the percentage of patients with primary care was much more in the NTIS group than in the non-NTIS group (*P* < 0.001), and that for patients admitted with respiratory disease and accompanied by respiratory disease and chronic kidney disease were more in the NTIS group than in the non-NTIS group (*P* < 0.001, Table 1).

Table 1 Comparison of baseline characteristics between the NTIS and non-NTIS groups ($\bar{x} \pm SD$)

	Total (n=852)	NTIS group (n=192)	Non-NTIS group (n=660)	<i>t</i> / <i>Z</i> / <i>c</i> ²	P
(y)	87.63±6.11	87.84±5.96	87.56±6.16	0.56	0.57
	23.85±3.23	23.72±3.56	23.88±3.13	−0.57	0.57
triiodothyronine (ol/L)	1.27±0.31	0.90±0.16	1.38±0.25	−25.21	<0.001
<i>T</i> ₃ (ol/L)	3.63±0.83	2.63±0.54	3.92±0.66	−24.78	<0.001
l <i>T</i> ₄ (ol/L)	86.34±16.31	75.10±16.38	89.60±14.78	−11.67	<0.001
<i>T</i> ₄ (ol/L)	15.99±2.73	15.62±2.99	16.10±2.63	−2.17	0.31
[(J/mL)	1.88(1.30, 2.88)	1.68(1.04, 2.74)	1.95(1.36, 2.93)	−3.37	0.001
single degree	60	42(21.88%)	18(2.73%)	83.30	<0.001
primary	792	150(78.12%)	642(97.27%)		
secondary					
tertiary					
reason for hospital admission					
TD	434	51(26.56%)	383 (58.03%)	111.24	<0.001
	282	110 (57.29%)	172 (26.06%)		
SD	60	13 (6.77%)	47 (7.12%)		
TD	10	10 (5.21%)	0 (0.00%)		
others	66	8 (4.17%)	58 (8.79%)		
comorbidities					
SD	282	110(57.29%)	172(26.06%)	65.51	<0.001
SD	82	24(12.50%)	58(8.79%)	2.36	0.13
SD	759	174(90.63%)	585(88.64%)	0.61	0.44
SD	112	45(23.44%)	67(10.15%)	22.99	<0.001
tumour	180	30(15.63%)	150(22.73%)	4.50	0.03

RD respiratory disease, *NSD* nervous system disease, *CVD* cardiovascular disease, *CKD* chronic kidney disease, *T*₃ triiodothyronine, *T*₄ thyroxine, *TSH* thyroid-stimulating hormone.

After matching by age and BMI, TP, Alb, PA, Hb, UA, TG, and HDLC levels were significantly lower, while UN and FBG levels were higher, in the NTIS group than in the non-NTIS group (Table 2).

Table 2 Comparison of other baseline biochemical indicators between the NTIS and non-NTIS groups ($\bar{x} \pm SD$)

	Total (n=852)	NTIS group (n=192)	Non-NTIS group (n=660)	<i>t/Z</i>	<i>P</i>
)	66.37±6.34	64.48±6.67	66.92±6.14	-4.78	<0.001
)	38.06±4.57	34.82±4.43	39.01±4.17	-12.08	<0.001
)	23.19±7.52	21.86±9.18	23.58±6.93	-2.79	0.005
/dL)	119.10±18.99	111.46±18.30	121.32±18.63	-6.48	<0.001
)	335.25±100.74	320.34±120.80	339.58±93.78	-2.34	0.02
ol/L)	5.88±1.88	6.68±2.28	5.65±1.68	6.83	<0.001
i iol/L)	3.99±0.88	4.07±1.06	3.97±0.82	1.48	0.14
iol/L)	1.20(0.87, 1.65)	1.05(0.79, 1.65)	1.21(0.90, 1.65)	-2.09	0.04
iol/L)	1.19±0.37	1.13±0.42	1.20±0.35	-2.37	0.02
.C iol/L)	2.30±0.75	2.34±0.91	2.28±0.69	0.94	0.35
.C iol/L)	13.00(10.00, 20.00)	13.00(9.00, 22.00)	13.00(10.00, 20.00)	-0.54	0.59
)	15.00(18.00, 23.00)	19.00(14.00, 26.83)	18.00(15.00, 22.00)	-0.90	0.37
)	6.40(5.20, 8.90)	7.80(5.90, 11.30)	6.20(5.10, 8.30)	-5.95	<0.001
iol/L)	82.00(69.00, 99.00)	82.00(66.25, 121.75)	82.00(70.25, 97.00)	-1.07	0.285
ol/L)					

TP total protein, *Alb* albumin, *PA* prealbumin, *Hb* haemoglobin, *UA* uric acid, *FBG* fasting blood glucose, *TC* total cholesterol, *TG* triglyceride, *HDLC* high-density lipoprotein cholesterol, *LDLC* low-density lipoprotein cholesterol, *ALT* alanine transaminase, *AST* aspartate transaminase, *UN* urea nitrogen, *Cr* creatinine

Correlations of triiodothyronine and free T₃ levels with important biochemical indicators

Correlation analysis showed that triiodothyronine was positively correlated with the levels of TP ($r = 0.15$, $P < 0.001$), Alb ($r = 0.40$, $P < 0.001$), PA ($r = 0.08$, $P = 0.01$), and Hb ($r = 0.37$, $P < 0.001$). Triiodothyronine was negatively correlated with the levels of FBG ($r = -0.28$, $P < 0.001$), UN ($r = -0.26$, $P < 0.001$), and Cr ($r = -0.12$, $P < 0.001$).

Free T₃ was positively correlated with the levels of TP ($r = 0.14$, $P < 0.001$), Alb ($r = 0.50$, $P < 0.001$), PA ($r = 0.19$, $P < 0.001$), Hb ($r = 0.40$, $P < 0.001$), and UA ($r = 0.11$, $P = 0.001$). Free T₃ was negatively correlated with the levels of FBG ($r = -0.26$, $P < 0.001$), UN ($r = -0.20$, $P < 0.001$), and Cr ($r = -0.05$, $P < 0.001$; Table 3).

Table 3 Factors associated with triiodothyronine and free T₃

	Triiodothyronine		Free T ₃	
	<i>r</i>	P	<i>r</i>	P
TP	0.15	<0.001	0.14	<0.001
Alb	0.40	<0.001	0.50	<0.001
PA	0.08	0.01	0.19	<0.001
Hb	0.37	<0.001	0.40	<0.001
UA	0.06	0.07	0.11	0.001
FBG	−0.28	<0.001	−0.26	<0.001
TC	−0.07	0.05	0.01	0.70
TG	0.03	0.46	0.06	0.09
HDLC	0.06	0.08	0.07	0.06
LDLC	0.00	0.93	0.06	0.11
ALT	0.04	0.25	0.04	0.21
AST	−0.06	0.08	−0.06	0.07
UN	−0.26	<0.001	−0.20	<0.001
Cr	−0.12	<0.001	−0.05	<0.001

TP total protein, *Alb* albumin, *PA* prealbumin, *Hb* haemoglobin, *UA* uric acid, *FBG* fasting blood glucose, *TC* total cholesterol, *TG* triglyceride, *HDLC* high-density lipoprotein cholesterol, *LDLC* low-density lipoprotein cholesterol, *ALT* alanine transaminase, *AST* aspartate transaminase, *UN* urea nitrogen, *Cr* creatinine

Association of the free T₃ level with mortality according to Cox proportional hazards models

Univariate Cox proportional hazards models showed that the free T₃ level was associated with reduced patient mortality. Among the other biochemical indicators, Alb and Hb levels were negatively, whereas FBG, total cholesterol (TC), and ALT levels were positively, correlated with patient mortality. Multivariate models adjusted for other confounding factors showed that a lower free T₃ level in elderly male patients is associated with all-cause mortality. Other indicators including Alb and Hb levels were negatively correlated and the ALT level positively correlated with patient mortality. Patients with respiratory diseases (RD), chronic kidney disease (CKD), and tumours had a higher mortality rate (Table 4). The free T₃ level and 2-year mortality rate of all the patients were also determined. According to a receiver operating characteristic (ROC) analysis of the free T₃ level and 2-year mortality rate, a cut-off free T₃ level of 3.45 pmol/L yielded the highest Youden index (0.327), with a sensitivity of 0.675 and specificity of 0.642 (Figure 1).

Table 4. Cox proportional hazards models of the associations of thyroid hormone levels, biochemical indicators, and systemic disease with mortality

Parameters	Univariate adjusted				Multivariate adjusted			
	β	HR	95% CI	P	β	HR	95% CI	P
Free T ₃ (pmol/L)	-1.30	0.27	(0.19, 0.38)	<0.001	-0.66	0.52	(0.35, 0.77)	0.001
Alb (g/L)	-1.55	0.21	(0.15, 0.29)	<0.001	-0.94	0.39	(0.27, 0.56)	<0.001
Hb (g/L)	-1.54	0.21	(0.16, 0.29)	<0.001	-0.84	0.43	(0.30, 0.62)	<0.001
UA (μ mol/L)	-0.19	0.82	(0.53, 1.29)	0.40	-0.06	0.95	(0.58, 1.53)	0.82
FBG (mmol/L)	0.62	1.86	(1.30, 2.66)	0.001	-0.01	0.99	(0.67, 1.48)	0.96
TC (mmol/L)	0.99	2.69	(1.63, 4.75)	0.001	0.23	1.26	(0.68, 2.36)	0.46
ALT (U/L)	1.12	3.07	(1.99, 4.75)	<0.001	1.18	3.26	(2.03, 5.22)	<0.001
RD	1.16	3.20	(2.33, 4.39)	<0.001	0.42	1.52	(1.06, 2.17)	0.02
NSD	-0.02	0.99	(0.58, 1.68)	0.96	0.16	1.18	(0.68, 2.04)	0.56
CVD	0.42	1.52	(0.85, 2.75)	0.16	-0.04	0.96	(0.53, 1.75)	0.89
CKD	1.08	2.95	(2.07, 4.20)	<0.001	1.00	2.72	(1.91, 3.89)	<0.001
Tumour	1.04	2.82	(2.05, 3.88)	<0.001	0.97	2.63	(0.83, 3.77)	<0.001

T₃ triiodothyronine, Alb albumin, Hb haemoglobin, UA uric acid, FBG fasting blood glucose, TC total cholesterol, ALT alanine transaminase, RD respiratory diseases, NSD nervous system disease, CVD cardiovascular disease, CKD chronic kidney disease

Figure 1 Receiver operating characteristic curve assessing the ability of the free triiodothyronine (T₃) level to predict all-cause mortality (free T₃ cut-off level, 3.445; sensitivity, 67.5%; specificity, 64.2%).

Comparison of 6-month, 1-year and 2-year survival rates between the NTIS and non-NTIS groups

During the 6-month follow-up period, 56 patients died: 41 in the NTIS group and 15 in the non-NTIS group. At 6 months, the survival rate was 78.65% in the NTIS group and 97.73% in the non-NTIS group (log-rank $\chi^2 = 95.23$, $P < 0.001$, Figure 2a). During the 1-year follow-up period, 81 patients died: 61 in the NTIS group and 20 in the non-NTIS group. At 1 year, the survival rate was 68.23% in the NTIS group and 96.97% in the non-NTIS group (log-rank $\chi^2 = 158.37$, $P < 0.001$, Figure 2b). A total of 157 patients died during the 2-year follow-up period; 68 in the NTIS group and 89 in the non-NTIS group. At 2 -years, the survival rate was 64.58% in the NTIS group and 86.52% in the non-NTIS group (log-rank $\chi^2 = 70.56$, $P < 0.001$, Figure 2c).

Figure 2 Kaplan–Meier survival curves of the NTIS and non-NTIS groups for the period 6-months (a), 1-year (b) and 2-year (c) after study enrolment.

Discussion

In the elderly population, thyroid hormone levels can help monitor health status, predict short-term and long-term clinical prognoses, predict disease severity, and assess quality of life and survival status. In previous studies, the frequency of thyroid dysfunction increased with advancing age in the hospitalised elderly patients. The prevalence of NTIS in hospitalised severely or debilitated elderly patients can be as high as 32%-62% [7, 9, 10]. In the present study, among the 931 elderly male patients hospitalised for various reasons,

there were 193 NTIS patients (20.73% prevalence rate). After matching by age and BMI, the Alb levels were significantly lower, and the renal function indices and FBG levels higher, in the NTIS group than in the non-NTIS group. In addition, the 2-year survival rate was significantly lower in the NTIS group than in the non-NTIS group. A reduced free T_3 level was strongly associated with all-cause mortality in NTIS patients, a similar finding of previous studies [11, 12].

NTIS is often associated with nutritional deficiencies or acute and chronic diseases. Protein and UA levels are indicators of nutritional status. Proteins also play an important role in the synthesis and transport of thyroid hormones. In several studies, the serum Alb level was reduced, and the free T_3 level was positively correlated with the Alb level, in patients with NITS [13, 14]. In the present study, the Alb levels were also significantly reduced in the NTIS group compared with the non-NTIS group. Correlation analysis showed that TP, Alb, and PA levels decreased as the triiodothyronine and free T_3 levels decreased. Hypothetically, decreased Alb levels leads to a decrease in the conversion of T_4 to T_3 , resulting in a decrease in T_3 levels or a decrease in T_4 binding to the protein, which accelerates the removal of thyroid hormones [15, 16]. In the present study, the free T_3 level was also positively correlated with the UA and Hb levels, further confirming that fasting and hunger can cause NTIS [5].

The T_4 level is strongly associated with CKD. NTIS is a common thyroid dysfunction in CKD patients [13, 17], and its mechanism is associated with the kidney's involvement in the synthesis, secretion, and metabolism of thyroid hormones. In kidney disease, chronic metabolic acidosis and inflammatory factors lead to the inhibition of deiodinase activity, and the conversion of T_4 to T_3 in kidney tissue and other tissue types is reduced [18]. Hypothalamic–pituitary–thyroid axis dysfunction [19] combined with loss of T_4 in the urine causes triiodothyronine and total T_4 levels to decrease. A decrease in the glomerular filtration rate (GFR) reduces iodine excretion, resulting in an iodine-blocking effect (Wolff–Chaikoff effect) [19]. Song et al. [20] retrospectively analysed 2,284 subjects with normal TSH levels and found that as the estimated GFR (eGFR) decreased in CKD patients, the prevalence of low T_3 syndrome gradually increased; the eGFR was positively correlated with the serum T_3 level and was independent of age and serum protein levels. In another study, reduced free T_3 levels predicted an increased risk of cardiovascular events in CKD patients with proteinuria [21]. In patients with chronic haemodialysis, reduced free T_3 levels were a strong predictor of all-cause mortality [22]. In the present study, among hospitalised elderly male patients, the UN levels in the NTIS group were higher than in the non-NTIS group. Correlation analysis showed triiodothyronine and free T_3 levels were negatively correlated with UN and Cr levels. After adjusting for confounding factors, all-cause mortality was significantly increased in CKD patients.

Compared with young, short-term diabetic patients, the older patients are more likely to develop NTIS. Some studies have compared the thyroid function status of diabetic patients according to age, disease course, and blood glucose control status and found lower free T_3 levels in diabetic patients than in normal controls [23, 24]; furthermore, the incidence of cardiovascular events in patients with type 2 diabetes and NTIS was significantly increased [25]. Aging, long diabetes duration, poor blood sugar control, and several complications can increase the prevalence of NTIS, especially in patients with diabetic nephropathy and ketoacidosis [26, 27]. In the present study, triiodothyronine and free T_3 levels were also negatively correlated with the FBG level.

In this study, at the end of the 2-year follow-up, a total of 157 patients had died (18.43% mortality rate). Kaplan–Meier survival analyses showed that the survival rate was significantly lower in the NTIS group (64.58%) than in the non-NTIS group (86.52%). Cox proportional hazards models showed that after removing confounding factors, reduced free T₃ levels increased the risk of all-cause death. The ROC analysis showed that when using a free T₃ cut-off level of 3.45 pmol/L, the Youden index was highest, with a sensitivity of 0.675 and specificity of 0.642, indicating that when free T₃ levels are less than 3.45 pmol/L, the risk of death increases. The cut-off level in our study was 3.45 pmol/L which was near to the low limit of the normal range (2.76 pmol/L), indicated that for older men with a free T₃ level < 3.45 pmol/L in spite of higher than 2.76 pmol/L may also have a poor prognosis and merit close clinical attention, however, it is still need to be further demonstrated by larger studies. In a recent study of 1,190 patients with acute heart failure, the survival rates were significantly lower in patients with low free T₃ compared with normal levels, and a multivariate Cox proportional hazards model showed that a low free T₃ level was an independent predictor of mortality [28]. Studies on ICU patients and hospitalised chronic patients reported the free T₃ level to be an independent predictor of all-cause mortality [11, 29]. Similar to previous studies, a lower free T₃ level in the present study was associated with a worse prognosis in elderly male patients with chronic diseases.

The present study had several limitations including failure to evaluate many factors that affect the patient prognosis. In this retrospective cohort study, thyroid hormone levels were measured at random in patients from different departments according to their doctor's discretion, not according to a standard indication. In addition, although age and BMI were matched between the two groups, the type of care received, treatment plan, treatment timing, severity of the patient's condition, and response to the treatment plan may have affected the patient's condition. Due to the small sample size, many influencing factors were difficult to quantify. The patients were not stratified according to the above-mentioned factors. In addition, only elderly male inpatients were analysed. Whether the above study results can be generalised to the general population requires further research.

Conclusion

In summary, among elderly male patients hospitalised for multiple causes, the 2-year survival rate was lower in the NTIS group than non-NTIS group. As the triiodothyronine and free T₃ levels decreased, the protein levels and renal function worsened, and the FBG level increased. A lower free T₃ level was associated with an increased risk of all-cause mortality.

Abbreviations

NTIS: non-thyroidal illness syndrome; ROC: Receiver operating characteristic curves; T₃: triiodothyronine; T₄: thyroxine; TSH thyroid-stimulating hormone; TP: total protein; Alb: albumin; PA: prealbumin; Hb: haemoglobin, UA: uric acid, TG: triglyceride; HDLC: high-density lipoprotein cholesterol; UN: urea nitrogen; FBG: fasting blood glucose; Cr: creatinine; ICU: intensive care units; ALT: alanine transaminase; AST: aspartate transaminase; BMI: body mass index; PSM: propensity score matching; TC: total cholesterol; RD: respiratory diseases; CKD: chronic kidney disease; NSD: nervous system disease; CVD: cardiovascular disease.

Declarations

Acknowledgements

We thank Meng Wei in the department 3 of orthopedics, Central Hospital of Handan City, Xiaomei Guo in the department of Endoscope Room, First People Hospital of Wu'an City, and Lin Liu in the department of Minimally Invasive Interventional Surgery, First Hospital of Handan City for providing help in data collection.

Authors' contributions

ST Y conceived, designed and developed the project. XM F, HP M and XF C assisted in data collection. HZ L, XY M and ZH L conducted data analysis. XY M and HZ L developed the initial drafts of the manuscript. ZY G, BR S, N L and CL L helped in revising subsequent drafts. All authors read and approved the final manuscript.

Funding

No funding was obtained for this study.

Availability of data and materials

The datasets used in the analyses described in this study are available from the corresponding author on reasonable request.

Ethics approval and consent to participate

The study was conducted in accordance with the ethical rules of the Helsinki Declaration. The study protocol was approved by the Ethics Committee of Chinese PLA General Hospital. The permission to access the raw data was granted by the Ethics Committee of Chinese PLA General Hospital.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹ Department of Endocrinology, The Second Medical Center & National Clinical Research Center for Geriatric Diseases, Chinese PLA General Hospital, Beijing 100853, China

² Department 2 of Pediatrics, Central Hospital of Handan City, Handan, Hebei province 056001, China

³ Department of Endocrinology, First People Hospital of Wu'an City, Wu'an, Hebei province 056300, China

⁴ Department of Endocrinology, The First Medical Center, Chinese PLA General Hospital, Beijing 100853, China

References

1. de Vries EM, Fliers E, Boelen A. The molecular basis of the non-thyroidal illness syndrome. *J Endocrinol* 2015; 225(3):R67-81.
2. Farwell AP. Nonthyroidal illness syndrome. *Curr Opin Endocrinol Diabetes Obes* 2013; 20(5):478-84.
3. Wang B, Liu S, Li L, Yao Q, Song R, Shao X et al. Non-thyroidal illness syndrome in patients with cardiovascular diseases: A systematic review and meta-analysis. *Int J Cardiol* 2017; 226:1-10.
4. Fliers E, Bianco AC, Langouche L, Boelen A. Thyroid function in critically ill patients. *Lancet Diabetes Endocrinol* 2015; 3(10):816-25.
5. Van den Berghe G. Non-thyroidal illness in the ICU: a syndrome with different faces. *Thyroid* 2014; 24(10):1456-65.
6. Wang YF, Heng JF, Yan J, Dong L. Relationship between disease severity and thyroid function in Chinese patients with euthyroid sick syndrome. *Medicine (Baltimore)* 2018; 97(31):e11756.
7. Tognini S, Marchini F, Dardano A, Polini A, Ferdeghini M, Castiglioni M et al. Non-thyroidal illness syndrome and short-term survival in a hospitalised older population. *Age Ageing* 2010; 39(1):46-50.
8. K H, M S, K S, E A. Reed larsen williams textbook of endocrinolgy, 11 edn: Canada:Saunders Elsevier; 2008.
9. Iglesias P, Munoz A, Prado F, Guerrero MT, Macias MC, Ridruejo E et al. Alterations in thyroid function tests in aged hospitalized patients: prevalence, aetiology and clinical outcome. *Clin Endocrinol (Oxf)* 2009; 70(6):961-7.
10. Bossoni S, Cossi S, Marengoni A, De Martinis M, Calabrese P, Leonardi R et al. Low T3 syndrome and outcome in elderly hospitalized geriatric patients. *J Endocrinol Invest* 2002; 25(10 Suppl):73-4.
11. Ataoglu HE, Ahbab S, Serez MK, Yamak M, Kayas D, Canbaz ET et al. Prognostic significance of high free T4 and low free T3 levels in non-thyroidal illness syndrome. *Eur J Intern Med* 2018; 57:91-5.
12. Sato Y, Yoshihisa A, Kimishima Y, Kiko T, Kanno Y, Yokokawa T et al. Low T3 Syndrome Is Associated With High Mortality in Hospitalized Patients With Heart Failure. *J Card Fail* 2019; 25(3):195-203.
13. Pan B, Du X, Zhang H, Hua X, Wan X, Cao C. Relationships of Chronic Kidney Disease and Thyroid Dysfunction in Non-Dialysis Patients: A Pilot Study. *Kidney Blood Press Res* 2019; 44(2):170-8.
14. Chavez Valencia V, Mejia Rodriguez O, Viveros Sandoval ME, Abraham Bermudez J, Gutierrez Castellanos S, Orizaga de la Cruz C et al. Prevalence of malnutrition-inflammation complex syndrome and its correlation with thyroid hormones in chronic haemodialysis patients. *Nefrologia* 2018; 38(1):57-63.
15. Flier JS, Harris M, Hollenberg AN. Leptin, nutrition, and the thyroid: the why, the wherefore, and the wiring. *J Clin Invest* 2000; 105(7):859-61.
16. De Groot LJ. Dangerous dogmas in medicine: the nonthyroidal illness syndrome. *J Clin Endocrinol Metab* 1999; 84(1):151-64.
17. Iglesias P, Bajo MA, Selgas R, Diez JJ. Thyroid dysfunction and kidney disease: An update. *Rev Endocr Metab Disord* 2017; 18(1):131-44.
18. Zoccali C, Tripepi G, Cutrupi S, Pizzini P, Mallamaci F. Low triiodothyronine: a new facet of inflammation in end-stage renal disease. *J Am Soc Nephrol* 2005; 16(9):2789-95.

19. Basu G, Mohapatra A. Interactions between thyroid disorders and kidney disease. *Indian J Endocrinol Metab* 2012; 16(2):204-13.
20. Song SH, Kwak IS, Lee DW, Kang YH, Seong EY, Park JS. The prevalence of low triiodothyronine according to the stage of chronic kidney disease in subjects with a normal thyroid-stimulating hormone. *Nephrol Dial Transplant* 2009; 24(5):1534-8.
21. Yang JW, Han ST, Song SH, Kim MK, Kim JS, Choi SO et al. Serum T3 level can predict cardiovascular events and all-cause mortality rates in CKD patients with proteinuria. *Ren Fail* 2012; 34(3):364-72.
22. Fragidis S, Sombolos K, Thodis E, Panagoutsos S, Mourvati E, Pikilidou M et al. Low T3 syndrome and long-term mortality in chronic hemodialysis patients. *World J Nephrol* 2015; 4(3):415-22.
23. Custro N, Scafidi V, Borsellino T. [Changes in the thyroid hormone picture that may be found in severely decompensated type II diabetics]. *Minerva Med* 1991; 82(1-2):9-14.
24. Kabadi UM, Premachandra BN. Low triiodothyronine and raised reverse triiodothyronine levels in patients over fifty years of age who have type II diabetes mellitus: influence of metabolic control, not age. *J Am Geriatr Soc* 1984; 32(5):375-9.
25. Moura Neto A, Parisi MC, Tambascia MA, Pavin EJ, Alegre SM, Zantut-Wittmann DE. Relationship of thyroid hormone levels and cardiovascular events in patients with type 2 diabetes. *Endocrine* 2014; 45(1):84-91.
26. Zhao W, Li X, Liu X, Lu L, Gao Z. Thyroid Function in Patients with Type 2 Diabetes Mellitus and Diabetic Nephropathy: A Single Center Study. *J Thyroid Res* 2018; 2018:9507028.
27. Mirboluk AA, Rohani F, Asadi R, Eslamian MR. Thyroid function test in diabetic ketoacidosis. *Diabetes Metab Syndr* 2017; 11 Suppl 2:S623-s5.
28. Asai K, Shirakabe A, Kiuchi K, Kobayashi N, Okazaki H, Matsushita M et al. Relation of Low Triiodothyronine Syndrome Associated With Aging and Malnutrition to Adverse Outcome in Patients With Acute Heart Failure. *Am J Cardiol* 2020; 125(3):427-35.
29. Wang F, Pan W, Wang H, Wang S, Pan S, Ge J. Relationship between thyroid function and ICU mortality: a prospective observation study. *Crit Care* 2012; 16(1):R11.

Supplementary Table

Additional file 1 Baseline patient characteristics before and after PSM

Variable	Before PSM		P	After PSM		t	P
	NTIS (n=193)	Non-NTIS (n=738)		NTIS (n=192)	Non-NTIS (n=660)		
Age	87.90±5.93	85.97±8.57	3.36 <0.001	87.84±5.96	87.56±6.10	0.56	0.57
BMI	23.68±3.62	23.97±3.12	-1.10 0.27	23.72±3.56	23.88±3.13	-0.50	0.57

PSM propensity score matching, NTIS non-thyroidal illness syndrome, BMI body mass index

Figures

Figure 1

Receiver operating characteristic curve assessing the ability of the free triiodothyronine (T3) level to predict all-cause mortality (free T3 cut-off level, 3.445; sensitivity, 67.5%; specificity, 64.2%).

Figure 2

Kaplan–Meier survival curves of the NTIS and non-NTIS groups over 24 months.