

# Attendance, activation and health profiles of participants in a regional chronic disease self-management program in Laval, Canada

Magali Girard (✉ [magali.girard.chum@ssss.gouv.qc.ca](mailto:magali.girard.chum@ssss.gouv.qc.ca))

Centre Hospitalier de L'Université de Montréal <https://orcid.org/0000-0002-2325-1676>

Janusz Kaczorowski

Université de Montréal

Marie-Thérèse Lussier

Université de Montréal

Vivianne Martin

CIUSSS du Centre-Sud-de-l'Île-de-Montréal

---

## Research article

**Keywords:** Chronic disease, Self-management programs, Health promotion, Volunteers, Health profile, Patient Activation Measure

**Posted Date:** May 20th, 2020

**DOI:** <https://doi.org/10.21203/rs.3.rs-29386/v1>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

**Version of Record:** A version of this preprint was published on March 12th, 2021. See the published version at <https://doi.org/10.1186/s12889-021-10558-6>.

1 **Title:**

2 Attendance, activation and health profiles of participants in a regional chronic  
3 disease self-management program in Laval, Canada

4

5 Magali Girard PhD

6 CRCHUM, Montreal, Canada

7

8 Janusz Kaczorowski PhD

9 Department of Family and Emergency Medicine, University of Montreal, and  
10 CRCHUM, Montreal, Canada

11

12 Marie-Thérèse Lussier MD MSc

13 Department of Family and Emergency Medicine, University of Montreal,  
14 Montreal, Canada

15

16 Vivianne Martin MSc

17 CIUSSS du Centre-Sud-de-l'Île-de-Montréal

18

19 Corresponding author:

20 Magali Girard PhD

21 CRCHUM, Tour Saint-Antoine

22 850, rue St-Denis, Office: S01.121.01

23 Montreal (Quebec) H2X 0A9

24 Telephone number: 514-890-8000, 31834

25 Fax number: 514-412-7037

26 E-mail address: Magali.girard.chum@ssss.gouv.qc.ca

27

28

29 **Abstract**

30 **Background:** Chronic diseases are responsible for over 70% of all deaths  
31 globally. While some self-management programs have been shown to be  
32 efficacious in preventing or altering trajectories for some chronic conditions,  
33 scaling-up and sustaining such programs beyond tightly-controlled study  
34 conditions remain a major challenge. The objective of the current study was to  
35 describe the profile of attendees and the level of engagement of participants in a  
36 new, region-wide cardiometabolic disease self-management program offered in  
37 Laval, Canada. **Methods:** CISSS-Laval partnered with the Cardiovascular Health  
38 Awareness Program team to co-develop Cible-santé and evaluate the first cohort  
39 of participants enrolled in the program, in order to better understand the  
40 program's implementation and scope. Trained volunteers (supervised by a nurse)  
41 assisted participants with physical measurements (blood pressure, height, weight  
42 and waist circumference), a questionnaire on healthy habits, and assessment of  
43 cardiovascular disease and type 2 diabetes risk. **Results:** The evaluation of the  
44 first cohort of Cible-Santé participants showed very low attendance. Furthermore,  
45 only 1 in 10 of enrolled participants completed the full program. The program  
46 typically attracted adults with some risk factors associated with their chronic  
47 conditions (high waist circumference, obesity), but with an already high level of  
48 knowledge, skills and confidence to participate in self-management activities.  
49 **Conclusion:** A limitation of the study is the lack of data regarding the number of  
50 professionals who referred their patients to the program. This study provides a  
51 portrait of new participants to a self-management chronic disease program,

52 which highlights the potential of supporting chronic disease patients ready to  
53 make changes but also exposes the difficulty of attracting a larger number and  
54 diversity of participants and in encouraging completion of the program.

55

56

57 **Keywords:**

58 Chronic disease

59 Self-management programs

60 Health promotion

61 Volunteers

62 Health profile

63 Patient Activation Measure

64

65

66 **Background:**

67 Chronic diseases — including cardiovascular diseases, cancers, chronic  
68 respiratory diseases and diabetes — are responsible for over 70% of all deaths  
69 globally. Modifiable risk factors that include smoking, physical inactivity and  
70 unhealthy diets are the main drivers behind the growing epidemic of chronic  
71 diseases [1]. While some health promotion and self-management programs have

72 been shown to be efficacious in preventing or altering trajectories for some  
73 chronic conditions, in modifying risk profiles and in improving satisfaction and  
74 knowledge of participants [2], scaling-up and sustaining such programs beyond  
75 tightly-controlled study conditions remain a major challenge.

76

77 Participation rates in health promotion and chronic disease self-management  
78 programs are generally low [3], despite their wide availability in most high-income  
79 countries, often not requiring referrals from a health professional and offered free  
80 of charge or at low cost [4]. People who engage in health promotion or self-  
81 management programs are more likely to be female [5], younger [4, 6, 7], more  
82 educated [6, 8], of higher socioeconomic status [9], Caucasian [5], in better  
83 physical health [3, 6, 7] and to already be adhering to healthier lifestyles [10].  
84 They are also more likely to have a better understanding of their condition than  
85 patients who do not participate [6].

86

87 Program type and program location also influence attendance. The participation  
88 rates are higher when programs are offered in a variety of time and place  
89 settings, when there is no need for referral and when participation is free or low  
90 cost [5]. Elzen and colleagues showed that living far away from a program's  
91 location and transportation constraints both represented barriers to participation  
92 [10]. Another study found that living in rural areas with limited access to

93 physicians increased utilization of a diabetes self-management education  
94 program [4].

95

96 The objective of the current study was to describe the profile of attendees and  
97 the level of engagement of participants in a new, region-wide cardiometabolic  
98 disease self-management program offered in Laval, Canada. The study took  
99 place in 2015 and looked at the first cohort of Cible-Santé participants.

100

#### 101 **Program description**

102 The Cible-Santé program was developed by the regional health authority, Centre  
103 intégré de santé et des services sociaux de Laval (CISSS-Laval) servicing over  
104 400,000 residents of Laval, the second largest city in the province of Québec,  
105 Canada. The goal of the program was to help participants modify some of their  
106 unhealthy habits in order to support the self-management of their cardiometabolic  
107 diseases. The program was led by an interdisciplinary team (nutritionists,  
108 kinesiologists and nurses) trained in motivational interviewing techniques [11,  
109 12]. The SMART<sup>1</sup> objectives developed by participants were at the heart of the  
110 change process. The 8-month program included three group workshops and five  
111 individual follow-ups.

112

---

<sup>1</sup> SMART= Specific, Measurable, Attainable, Relevant and Time-Bound.

113 CISSS-Laval partnered with the Cardiovascular Health Awareness Program team  
114 to co-develop Cible-santé and evaluate the first cohort of participants enrolled in  
115 the program, in order to better understand the program's implementation and  
116 scope. These findings might offer useful insights that can inform similar future  
117 programs.

118

119 CHAP is a patient-centred, interdisciplinary, multi-faceted, community-led,  
120 cardiovascular disease prevention and management program targeting older  
121 Canadian adults. Building on 20 years of work around CHAP [13-22], the  
122 evaluation consisted of physical measurements that included blood pressure,  
123 height, weight and waist circumference, a questionnaire on healthy habits, and  
124 assessment of cardiovascular disease and type 2 diabetes risk at the onset of  
125 the first Cible-Santé group session. The CHAP sessions were run by locally  
126 recruited and trained volunteers and supervised by public-health nurses.

127

## 128 **Method**

### 129 *Design:*

130 This was a prospective, open-cohort study with no comparison group. Our goal  
131 was to describe the attendance, health profile and level of activation of new  
132 participants in the Cible-Santé cardiometabolic diseases self-management  
133 program.

134

135 *Recruitment of participants:*

136 To be eligible for Cible-Santé program, patients had to be registered with one of  
137 Laval's interdisciplinary primary care clinics (*Groupe de médecine de famille* or  
138 GMF) and had to have at least one of the following chronic conditions: type 2  
139 diabetes, pre-diabetes, hypertension, pre-hypertension, cardiovascular disease,  
140 dyslipidemia, stroke/transient ischemic attack, heart failure, kidney failure,  
141 metabolic syndrome or peripheral vascular disease. Participants had to be  
142 identified and referred to the program by their primary care health professional  
143 practicing in one of the city's interdisciplinary primary care clinics. Referrals were  
144 sent to CISSS-Laval appointment centre, who subsequently contacted potential  
145 participants to enroll them into the program.

146

147 Cible-Santé participants were invited to attend three group-based workshops and  
148 five individual follow-up sessions with a nutritionist, kinesiologist or nurse over an  
149 eight-month period. At their first group workshop, patients were invited to  
150 participate in a CHAP evaluation session. After signing a consent form, CHAP  
151 volunteers, supervised by a Cible-Santé program nurse, had their blood  
152 pressure, waist circumference and body mass index measured, completed a  
153 questionnaire on health habits, level of activation and the risk of developing  
154 prediabetes and type 2 diabetes over the next 10 years.

155

156 *Measures:*

157 Data on demographic characteristics, health status, risk factors and patient  
158 activation were collected at baseline and were self-reported by participants.

159

160 For physical activity, patients were asked to list the type and number of minutes  
161 of physical activities for a typical week. Then, categories were created from a  
162 measure of intensity in Metabolic Equivalents (METs) multiplied by the duration  
163 of exercise. The equivalence in minutes of slow walking per day was used as a  
164 guide.

165

166 The Patient Activation Measure (PAM) assesses the knowledge, skills and  
167 confidence for self-management. [23]. Four levels of activation were categorized  
168 according to the score ranging from 0 to 100. At Stage 1, the person is  
169 considered "unaware", does not feel confident enough and is a passive recipient  
170 of care. At Stage 2, the person lacks confidence and understanding of their  
171 health care (e.g., how to take or know what their medications do). At Stage 3, the  
172 person knows the essential facts and begins to take steps to take care of their  
173 health, but may lack the confidence and skills to take complete control of their  
174 health care. At Stage 4, knowledge, skills and confidence for self-management  
175 are all present.

176

177 All respondents who reported not being diagnosed with type 2 diabetes or being  
178 unsure were asked to complete the Canadian Diabetes Risk Questionnaire  
179 (CANRISK). Developed by the Public Health Agency of Canada, this  
180 questionnaire-based tool assesses the risk of developing pre-diabetes and type 2  
181 diabetes in the next 10 years [24].

182

183 During the sessions, physical measurements were taken by trained volunteers  
184 and included waist circumference, weight and height (to calculate the body mass  
185 index), as well as participant blood pressure, using an automated measuring  
186 device, the Microlife WatchBP Office Afib, which allows dual-cuff measurement  
187 and atrial fibrillation detection. The participants with elevated blood pressure  
188 (>140/90 mm Hg) were assessed by the nurse and referred to their family  
189 physician for follow-up.

190

191 Program participation was defined as completing all three group-based  
192 workshops and five individual follow-ups. Program completion rate was  
193 calculated using the total number of participants who registered as the  
194 denominator.

195

196 *Analyses*

197 Data were analyzed with SPSS, version 24.0. Univariate descriptive statistics  
198 and frequency distributions were used to summarize the data.

199

200 **Results:**

201

202 [INSERT FIGURE 1 HERE]

203

204 The total number of patients referred by health professionals practicing in one of  
205 the GMFs in Laval has not been recorded. A total of 270 patients had registered  
206 with the Cible-Santé program. Of these, 190 attended at least one group-based  
207 workshop and 141 took part in the CHAP evaluation session that preceded the  
208 first group workshop. One in ten (10.4%, 28/270) of the enrolled participants had  
209 attended all three group workshops and five individual follow-ups.

210

211 [INSERT TABLE 1 HERE]

212

213 More than half of the participants were 55 years of age or older, and the majority  
214 were female (Table 1).

215

216 [INSERT TABLE 2 HERE]

217

218 The vast majority of participants (83.1%, 113/136) had a waist circumference of  
219 more than 88 cm for women and more than 102 cm for men, and nearly 70%  
220 (97/140) were classified as obese. Over 17% had a diagnosis of hypertension  
221 (not shown in table) and more than a third had blood pressure above >140/ 90  
222 mm Hg).

223

224 [INSERT TABLE 3 HERE]

225

226 Approximately half of participants reported to eating the recommended daily  
227 servings of fruits and vegetables (four to six servings), and a similar proportion  
228 reported little to no physical activity (16.8% and 32.1%) (Table 3). Nearly half of  
229 the participants were classified at a level 4 of activation (Table 4).

230

231 [INSERT TABLE 4 HERE]

232

233 Of all 141 participants who had participated in a CHAP evaluation session, 68  
234 (48.2%) reported being diabetic (Table 5). Those who had no diagnosis or who  
235 were unsure (n=73) completed the CANRISK. More than 60% were classified as  
236 having high or very high risk of developing pre-diabetes or diabetes.

237

238

[INSERT TABLE 5 HERE]

239

240 **Discussion:**

241 The evaluation of the first cohort of Cible-Santé participants showed very low  
242 attendance (270 adults who registered from over 200,000 patients<sup>2</sup> registered in  
243 all of Laval's GMFs). Furthermore, only 1 in 10 of enrolled participants completed  
244 the full program. While the total number of patients referred to the program was  
245 not recorded, given the prevalence of targeted conditions and risk factors in  
246 primary care, the number of patients who registered and completed the program  
247 was quite low. These results, while below expectations, are consistent with what  
248 has been reported in the literature. Although the program was offered free of  
249 charge by the local health authorities and in settings familiar to participants, there  
250 are likely several explanations for the low attendance and completion rates.

251

252 First, Cible-Santé was a new program introduced in the already large spectrum of  
253 services provided by the regional health authority. Moreover, no specific  
254 communication strategy was put forth to inform referring health professionals of  
255 this innovative program made available to their chronic disease patients. Many  
256 community health providers were thus probably unaware of the program or  
257 uncertain of its value to their patients, which might have limited referrals. Given  
258 that this was the first cohort of participants, a more comprehensive

---

<sup>2</sup> This is an estimation of the number of patients registered in one of Laval's GMFs at the time of data collection. We don't know the total number of patients who met the eligibility criteria of Cible-Santé.

259 communication strategy targeting health professionals in the region might help  
260 raise awareness about the program and its goals. Second, the evidence  
261 suggests that when programs are offered in a variety of time and place settings,  
262 participation rates are typically higher [5]. Cible-Santé was, at the time, offered to  
263 one new cohort per month and this study focused only on the first cohort. Third,  
264 the low completion rate could be explained by the long duration of the program (8  
265 months) and low frequency of interactions with health professionals (one group  
266 workshop or individual follow-up per month), which may have de-motivated some  
267 participants. Finally, limited public transportation and the need to pay for parking  
268 may have limited access to the program for some of the participants.

269

270 The profile of Cible-Santé participants suggests that they tend to engage in a  
271 relatively healthy lifestyle. It is difficult to establish to what extent these self-  
272 reported “healthy habits” represent a true reflection of reality in contrast to a  
273 response bias resulting from the pressure they may have felt to provide socially  
274 acceptable answers. Over 80% of participants had waist circumference that put  
275 them at high risk for cardiovascular disease and 69% had a BMI of 30 or over  
276 (obese to morbid obese), two risk factors associated with the development of  
277 health problems such as diabetes, heart disease and hypertension. The CHAP  
278 sessions contributed objective physical measures data as well as validated data  
279 on level of risk of developing cardiovascular disease, prediabetes or diabetes  
280 found in the Cible santé program participants. These results tend to confirm the

281 appropriateness of the healthcare provider referrals to this chronic disease self-  
282 management program.

283

284 Nearly half of participants were considered highly activated (stage 4 of the PAM).

285 Given that people self-select to participate in the Cible-Santé program (with  
286 referral from a health professional), high levels of activation are not surprising.

287 Their participation in the program is probably the result of their high level of  
288 knowledge, skills and confidence.

289

290 A limitation of the study is the lack of data regarding the number of professionals  
291 who referred their patients to the program. This study was conducted in a real  
292 care setting and in collaboration with managers responsible for the program,  
293 which explains the absence of certain data (missing denominator). This would  
294 have allowed us to know the proportion of referred patients who subsequently  
295 enrolled in the program, and thus know the number of patients who follow-up on  
296 such referrals from their health care provider.

297

298 Another limitation of this study is the lack of a control group. Having a  
299 comparison group with similar baseline characteristics, that would not have  
300 participated in this chronic disease self-management program or maybe enrolled  
301 in a different program, would have enabled us to better assess the characteristics  
302 associated with recruitment and participation.

303

304 **Conclusion:**

305 The evaluation by the CHAP team made it possible to draw a health profile of the  
306 first cohort of Cible-Santé participants. CHAP volunteers were trained to  
307 adequately measure blood pressure, body mass index and waist circumference,  
308 and to assist participants in completing their questionnaires. The results showed  
309 a very low attendance and completion rate. The program typically attracted  
310 adults with some risk factors associated with their chronic conditions (high waist  
311 circumference, obesity), but with an already high level of knowledge, skills and  
312 confidence to participate in self-management activities.

313

314 This study provides a portrait of new participants to a self-management chronic  
315 disease program, which highlights the potential of supporting chronic disease  
316 patients ready to make changes but also exposes the difficulty of attracting a  
317 larger number and diversity of participants and in encouraging completion of the  
318 program. Further studies are needed to assess the effect of the program on self-  
319 management skills and health outcomes of participants, test several recruitment  
320 strategies to increase the diversity of the target population and to compare  
321 participants' health profiles with non-participants.

322

323

324 **List of abbreviations:**

325 BMI: Body Mass Index

326 CANRISK: Canadian Diabetes Risk Questionnaire

327 CHAP: Cardiovascular Health Awareness Program

328 CISSS-Laval: Centre intégré de santé et services sociaux de Laval

329 GMF: *Groupe de médecine de famille* (Interdisciplinary primary care clinics)

330 METS: Metabolic Equivalents

331 PAM: Patient Activation Measure

332 SMART: Specific, Measurable, Attainable, Relevant, and Time-Bound

333

334

335 **Declarations**

336 *Ethics approval and consent to participate*

337 The study protocol was reviewed and approved by the scientific and research ethics  
338 committee of the Laval regional health authority (Centre intégré de santé et de services  
339 sociaux de Laval). Participants and volunteers provided written consent to participate.

340

341 *Consent for publication*

342 Not applicable.

343

344 *Availability of data and materials*

345 The datasets used and/or analysed during the current study are available from the  
346 corresponding author (MG) on reasonable request.

347

348 *Competing interests*

349 The authors declare that they have no competing interests.

350

351 *Funding*

352 This project was supported by a grant from the Canadian Institute of Health Research.

353 The funding agency had no role in study design, data collection and analysis, decision to  
354 publish, or preparation of the manuscript.

355

356 *Authors' contributions*

357 MG, JK and MTL conceptualized the study protocol and its implementation. MG and VM  
358 carried out analyses. MG drafted the initial manuscript, and reviewed and revised the  
359 manuscript. All authors assisted with data interpretation and synthesis and reviewed and  
360 revised the manuscript. MG, JK, MTL and VM read and approved the final manuscript as  
361 submitted and agreed to be accountable for all aspects of the work.

362

363 *Acknowledgements*

364 The authors would like to express a special thanks to nurses from CISSS-Laval who  
365 supervised the volunteers; to Jocelyne Gagné, Secretary, for managing the volunteers'

366 schedules; and to CISSS-Laval for recruiting participants and helping the team organize  
367 the sessions.

368

369

## 370 **References**

371 1. World Health Organisation. Noncommunicable diseases country profiles  
372 2018. Geneva: World Health Organization; 2018. Licence: CC BY-NC-SA  
373 3.0 IGO

374

375 2. Reynolds R, Dennis S, Hasan I, Slewa J, Chen W, Tian D, et al. A  
376 systematic review of chronic disease management interventions in primary  
377 care. BMC Fam Pract. 2018; doi: 10.1186/s12875-017-0692-3.

378

379 3. Dattalo M, Giovannetti ER, Scharfstein D, Boult C, Wegener S, Wolff JL, et  
380 al. Who Participates in Chronic Disease Self-Management (CDSM)  
381 Programs? Differences between Participants and Non Participants in a  
382 Population of Multi-Morbid Older Adults. Med Care. 2012;  
383 doi:10.1097/MLR.0b013e318268abe7.

384

385 4. Cauch-Dudek, K, Victor, JC, Sigmond, M, Shah BR. Disparities in  
386 attendance at diabetes self-management education programs after  
387 diagnosis in Ontario, Canada: a cohort study. BMC Public Health. 2013;  
388 doi.org/10.1186/1471-2458-13-85.

389

390 5. Bruce B, Lorig K, Laurent D. Participation in patient self-management  
391 programs. *Arthritis Rheum.* 2007;57:851–4.

392

393 6. Muntner P, Sudre P, Uldry C, Rochat T, Courteheuse C, Naef AF, et al.  
394 Predictors of participation and attendance in a new asthma patient self-  
395 management education program. 2001; 120:778-84.

396

397 7. Jong ZD, Munneke M, Jansen LM, Runday K, van Schaardenburg DJ,  
398 Brand R. Differences between participants and nonparticipants in an  
399 exercise trial for adults with rheumatoid arthritis. *Arthritis Care &*  
400 *Research.* 2004;51:593–600.

401

402 8. Santorelli ML, Ekanayake RM, Wilkerson-Leconte L. Participation in a  
403 Diabetes Self-Management Class Among Adults With Diabetes, New  
404 Jersey 2013-2015. *Prev Chronic Dis.* 2017; doi: 10.5888/pcd14.170023.

405

406 9. Hardman R, Begg S, Spelten E. What impact do chronic disease self-  
407 management support interventions have on health inequity gaps related to  
408 socioeconomic status: a systematic review. *BMC Health Serv Res.* 2020;  
409 doi: 10.1186/s12913-020-5010-4.

410

- 411 10. Elzen H1, Slaets JP, Snijders TA, Steverink N. Do older patients who refuse  
412 to participate in a self-management intervention in the Netherlands differ  
413 from older patients who agree to participate? *Aging Clinical and*  
414 *Experimental Research*. 2008;20:266–271.
- 415
- 416 11. Miller WR, Rollnick S. *Motivational Interviewing. Preparing people for*  
417 *change*. 2nd ed. New York: The Guilford Press, 2002.
- 418
- 419 12. Prochaska J, DiClemente C. Towards a comprehensive model of change.  
420 In: Miller WR, Heather N, editors. *Treating addictive behaviours: processes*  
421 *of change*. New York: Pergamon, 1986.
- 422
- 423 13. Karwalajtys T, McDonough B, Hall H, Guirguis-Younger M, Chambers  
424 LW, Kaczorowski J, et al. Development of the volunteer peer educator role  
425 in a community Cardiovascular Health Awareness Program (CHAP): a  
426 process evaluation in two communities. *J Community Health* 2009; 34: 336-  
427 45.
- 428
- 429 14. Chambers LW, Kaczorowski J, Dolovich L, Karwalajtys T, Hall HL,  
430 McDonough B, et al. A community-based program for cardiovascular health  
431 awareness. *Can J Public Health* 2005; 96: 294-8.
- 432

- 433 15. Jones C, Simpson SH, Mitchell D, Haggarty S, Campbell N, Then K, et al.  
434 Enhancing hypertension awareness and management in the elderly:  
435 lessons learned from the Airdrie Community Hypertension Awareness and  
436 Management Program (A-CHAMP). *Can J Cardiol* 2008; 24: 561-7.  
437
- 438 16. Carter, M, Karwalajtys, T, Chambers, L, Kaczorowski, J, Dolovich, L,  
439 Gierman, T, et al. Implementing a standardized community-based  
440 cardiovascular risk assessment program in 20 Ontario communities. *Health  
441 Promot Int* 2009; 24: 325-33.  
442
- 443 17. Pora VV, Farrell B, Dolovich L, Kaczorowski, J, Chambers, L. Promoting  
444 cardiovascular health among older adults: A pilot study with community  
445 pharmacists. *Canadian Pharmacy Journal* 2005;138:50-55.  
446
- 447 18. Karwalajtys T, Kaczorowski J, Chambers L, Levitt C, Dolovich L,  
448 McDonough B, et al. A randomized trial of mail vs. telephone invitation to a  
449 community-based cardiovascular health awareness program for older family  
450 practice patients. *BMC Family Practice* 2005, 6:35.  
451
- 452 19. Kaczorowski J, Chambers L, Karwalajtys T, Dolovich L, Farrell B,  
453 McDonough B, et al. Cardiovascular Health Awareness Program (CHAP): a  
454 community cluster-randomised trial among elderly Canadians. *Prev Med*  
455 2008; 46: 537-44.

456

457 20. Kaczorowski J, Chambers LW, Dolovich L, Paterson JM, Karwalajtys T,  
458 Gierman T, et al. Improving cardiovascular health at the population level: A  
459 39 community cluster-randomized trial of the Cardiovascular Health  
460 Awareness Program (CHAP). *BMJ* 2011;342:d442.

461

462 21. Goeree R, von Keyserlingk C, Burke N, He J, Kaczorowski J, Chambers L,  
463 et al. Economic appraisal of a community-wide cardiovascular health  
464 awareness program. *Value Health J Int Soc Pharmacoeconomics*  
465 *Outcomes Res.* 2013; doi:10.1016/j.jval.2012.09.002.

466

467 22. Chambers LW, Kaczorowski J, O’Rielly S, Ignagni S, Hearps SJC.  
468 Comparison of blood pressure measurements using an automated blood  
469 pressure device in community pharmacies and family physicians’ offices: a  
470 randomized controlled trial. *Can Med Assoc Open Access J.* 2013;  
471 doi:10.9778/cmajo.20130005.

472

473 23. Hibbard JH, Stockard J, Mahoney ER, Tusler M. Development of the  
474 Patient Activation Measure (PAM): Conceptualizing and Measuring  
475 Activation in Patients and Consumers. *Health Serv Res.* 2004; doi:  
476 10.1111/j.1475-6773.2004.00269.x


477

478 24. Robinson CA, Agarwal G, Nerenberg K. Validating the CANRISK prognostic  
479 model for assessing diabetes risk in Canada's multi-ethnic population.  
480 Chronic dis Inj Can. 2011;32:19-31.

481

482 **Figure 1: Algorithm for participation in the Cible-Santé**

483



484 **Table 1: Socio-demographic characteristics of participants**

	n	%
<b>Age</b> (n=140, average 59 years-old)		
18 to 45 years-old	15	10.7
45 to 54 years-old	34	24.3
55 to 64 years-old	41	29.3
65 years-old and over	50	35.7
<b>Gender</b> (n=141)		
Male	55	39.0
Female	86	61.0

485

486

487 **Table 2: Physical Measures**

	n	%
<b>Waist Circumference</b> (n=136)		
F<80cm   H<94cm (Low Risk)	7	5.1
F80-88cm   H<94-102cm (High Risk)	16	11.8
F>88cm   H>102 cm (Very High Risk)	113	83.1
<b>BMI</b> (n=140)		
Normal (<25)	9	6.4
Overweight (25-29)	34	24.3
Obese (30-34)	57	40.7
Severely Obese (35-39)	25	17.9

Morbid Obese (40 et plus)	15	10.7
---------------------------	----	------

**Systolic Blood Pressure** (n=140, average= 136.51)

Less than 120	18	12.9
120 to 139	73	52.1
140 to 159	39	27.9
160 and over	10	7.1

**Diastolic Blood Pressure** (n=140, average= 78.93)

Less than 80	78	55.7
80 to 84	30	21.4
85 to 89	15	10.7
90 and over	17	12.1

488

489

490 **Table 3: Lifestyle Habits**

	n	%
<b>Daily consumption of fruit and vegetables</b> (n=138, mean (SD) = 5)		
1 to 3 portions	41	29.7
4 to 6 portions	69	50.0
7 to 9 portions	24	17.4
10 portions and over	4	2.9
<b>Physical Activity</b> (n=131)		
Inactive	24	18.3
Very light (< 20 min walk /day)	22	16.8

Light (20 to 40 min walk /day)	42	32.1
Moderate (40 to 60 min walk /day)	27	20.6
Vigorous (> 1 h walk /day)	16	12.2

**Smoking Status (n=141)**

Smoker	13	9.2
None smoker	128	90.8

**Stress day per week (n=126)**

Rarely or never	60	47.6
Sometimes (1-2 day/week)	18	14.3
Often (3-4-5 day/week)	34	27.0
Regularly (6 -7 day/week)	14	11.1

491

492

493 **Table 4: Patient Activation Measure (n=141, average = 65.65)**

	n	%
Level 1: Disengaged and overwhelmed	9	6.4
Level 2: Becoming aware but still struggling	20	14.2
Level 3: Taking action and gaining control	43	30.5
Level 4: Maintaining behaviours and pushing further	69	48.9

494

495

496 **Table 5: Diabetes and CANRISK Score**

	n	%
<b>Self-reported diabetes diagnosis (n=141)</b>		
Diabetic	68	48.2
Non-diabetic	64	45.4
Unsure	9	6.4
<b>Score CANRISK (n=72/73 did not declare being diagnosed with diabetes or unsure)</b>		
Low Risk (score < 21)	3	4.2
Average Risk (21 to 32)	24	33.3
High Risk (33 to 42)	34	47.2
Very High Risk (> 42)	11	15.3

497

498

499

500

# Figures




Figure 1

Algorithm for participation in the Cible-Santé