

Primary Care Population Management for COVID-19 Patients

Deborah Blazey-Martin, MD MPH (✉ dblazeymartin@tuftsmedicalcenter.org)

Tufts Medical Center

Elizabeth Barnhart, FNP

Tufts Medical Center

Joseph Gillis, Jr.

Tufts Medical Center

Gabriela Andujar Vazquez, MD

Tufts Medical Center

Method Article

Keywords: COVID, Population health, primary care

Posted Date: May 19th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-29530/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Journal of General Internal Medicine on July 27th, 2020. See the published version at <https://doi.org/10.1007/s11606-020-05981-1>.

Abstract

BACKGROUND: Most patients infected with SARS-CoV-2 have mild to moderate symptoms manageable at home; however up to 20% develop severe illness requiring additional support. Primary care practices performing population management can use these tools to remotely assess and manage COVID-19 patients and identify those needing additional medical support before becoming critically ill.

AIM: We developed an innovative population management approach for managing COVID-19 patients remotely.

SETTING: Development, implementation, and evaluation took place in April 2020 within a large urban academic medical center primary care practice.

PARTICIPANTS: Our panel consists of 40,000 patients. By April 27, 2020, 305 had tested positive for SARS-CoV-2 by RT-qPCR. Outreach was performed by teams of doctors, nurse practitioners, physician assistants, and nurses.

PROGRAM DESCRIPTION: Our innovation includes an algorithm, an EMR component, and a twice daily population report for managing COVID-19 patients remotely.

PROGRAM EVALUATION: Of the 305 patients with COVID-19 in our practice at time of submission, 196 had returned to baseline; 54 were admitted to hospitals, six of these died, and 40 were discharged.

DISCUSSION: Our population management strategy helped us optimize at-home care for our COVID-19 patients and enabled us to identify those who require inpatient medical care in a timely fashion.

Introduction

Approximately 80% of laboratory confirmed COVID-19 patients have had mild to moderate disease that can be managed without hospitalization.^[1] It is known that a significant percentage also require guidance with a worsening of respiratory status in the second week of illness²⁻⁴; there are anecdotal reports of patients dying at home for failure to recognize the severity of their symptoms. It is critical for COVID-19 patients to receive remote support and monitoring so that they can recover successfully at home when possible and be advised when in-person evaluation is needed. This monitoring ensures that patients remain at home when they are safe to do so and avoids exposing health care workers and using valuable PPE unnecessarily. In addition, there are suggestions in the literature that earlier admission to the hospital results in better outcomes.⁵

Due to responsibility for a large number of patients, access to the results of SARS-CoV-2 testing, and expertise in population management, our primary care practice was able to quickly design, implement, and evaluate an algorithm, an electronic medical record (EMR) component, and a population report for managing COVID-19 patients remotely. We sought to standardize outreach to COVID-19 patients, manage

patients effectively at home when appropriate, and identify and quickly facilitate care for those who require inpatient medical support.

Setting And Participants

We designed, implemented, and evaluated our innovative approach within a large primary care practice at an urban academic medical center.

The study was deemed exempt by the Tufts Health Sciences Institutional Review Board (STUDY00000569).

Participants

Our hospital-based primary care practice staff includes 40 attending physicians, 76 internal medicine residents, five physician assistants (PAs), five nurse practitioners (NPs), and a team of registered nurses (RNs) caring for patients both in person and remotely. The practice is divided geographically by suite into 5 teams of clinicians, each consisting of 4-5 clinical full time equivalents (cFTE) of MD (up to 8 physicians), one NP, one PA and one cFTE RN. Since the start of the COVID crisis, each clinician works on site 1 day per week and remotely 4 days per week; the teams remain unchanged, and each continues to care for the same cohort of patients. Our patient panel consists of 40,000 people (~8000 per team), including patients participating in both Medicare and Medicaid accountable care organizations (ACOs), patients who are dually eligible, and patients who are commercially insured in risk contracts as well as PPOs. By April 27, 2020, 305 patients in our panel had tested positive for COVID-19 requiring ongoing evaluation and management.

Program Description

In March 2020, our practice implemented rapid changes to prepare for the COVID-19 pandemic. Among our first steps, we began shifting in-person appointments to tele-health visits, created a nursing phone triage algorithm to manage the high call volume from patients with questions about the novel coronavirus and concerns about getting tested, and launched an on-site respiratory symptoms clinic in one of our 5 suites. All of our residents were pulled to staff the inpatient and ICU services, and about half of our attendings entered a rotation to staff additional COVID and non-COVID ward services. Our Respiratory symptoms clinic was staffed by preceptors with a now available session and other MD and NP/PA volunteers from all 5 teams. Once patients started testing positive for the virus in our clinic and the hospital testing center, we identified the need to innovate a population management structure to help us manage our growing number of COVID-19 patients remotely.

Because our EMR system can receive the results of most diagnostic tests ordered throughout the hospital, we were able to identify many patients in our practice who had tested positive for SARS-CoV-2 by RT-qPCR. However, we needed a population management strategy to allow us to outreach to this cohort. We wanted a solution that would: 1) help us remotely manage COVID-19 patients recovering at

home, 2) indicate when a visit to our respiratory symptoms clinic was required to answer clinical questions (oxygen saturation, chest x-ray for viral pneumonia, white blood count, e.g.) without taxing the Emergency Department, and 3) alert us in a timely fashion to instances when additional urgent interventions were required.

Based on published data regarding the clinical progression of the disease, we knew that some patients are more likely to rapidly deteriorate within one week after illness onset. Therefore, it was essential for our outreach program to identify not only patients with risk factors for severe illness but also increase monitoring during the days when they are most likely to worsen.¹ With this knowledge of the illness, we designed a population management algorithm, an EMR component for entering important COVID-19 identifiers, and a population report. The algorithm (Figure 1) is used to guide triage decisions including how frequently to contact patients depending on symptom day and risk factors, and to identify clinical findings that indicate if the patient is safe to remain at home, the need for additional assessment, or emergency intervention. The component (Figure 2) is used to populate a twice daily report that allows us to tailor our outreach, and includes symptom date, age, comorbidities, last date of outreach, PCP, and whether a patient is currently hospitalized or had returned to baseline.

We run the population report (Figure 3) at 7:30 am each morning and distribute it by email to the five practice teams of MDs, NPs, PAs, and RNs who then reach out by phone to patients individually. Typically the primary care doctor, who knows each patient's history best, reviews their list and determines based on and their clinical schedule and patient complexity who they can outreach to themselves, with plans to turn outreach calls into telehealth visits when appropriate and the patient consents. The NP and PA on the team collaborate with all the PCP's on their team and reach out to patients they have relationships with and those the PCP's will not have time to reach. The report is run again at 3:00 pm and redistributed so that the primary teams can see who has not been contacted and the team working in the Respiratory Symptoms clinic, including RN's who have been triaging COVID patients and know some of those on the list, can supplement our calls and ensure that those who are particularly concerning are reached.

Since the COVID-19 outbreak began in Massachusetts, our practice has had daily leadership meetings and a daily email update to all working in the practice both at home and remotely. The clinical algorithm for the population management of COVID-19 patients was created by the Division Chief and Clinical Director of Nursing and reviewed by an Infectious Disease attending physician on the Infection Prevention Team. We then worked with our Manager of Applications and Analytics to build the EMR component and the population report. Once completed, we piloted the solution with our nursing team, and then rolled it out to the practice using our daily emails and our weekly practice-wide Zoom meeting to ensure all had a chance to learn the system and receive answers to their questions. The entire process, from design to implementation, took less than one week.

Program Evaluation

With a rapid rollout of our COVID-19 management program, we have applied transparent communication and a continuous improvement approach to ensure its effectiveness. Our staff is currently working predominantly from home, and attendance at our practice-wide (now virtual) meetings has skyrocketed, resulting in improved communication and collaboration. Based on feedback from the team since rolling out our new tool, we have made some adjustments. For example, initially, “Back to Baseline” was the only criterion to stop reaching out to COVID-19 patients, but we have added “Inpatient,” “ICU,” and “Deceased” to the drop-down menu. We have also added identifiers for patients diagnosed clinically or at other institutions to ensure outreach happens regardless of location of diagnosis. We expect to continue to adapt our strategy as we learn more about Covid-19 disease progression and how best to care for COVID-19 patients.

As of April 27, 2020, 305 patients in our practice have been diagnosed with COVID-19 in by SARS-CoV-2 by RT-qPCR testing performed at our Tufts Medical Center testing site, in our respiratory clinic, or performed at an outside location or empirically diagnosed and manually entered into our system. Of the 305 cases of COVID-19, 196 patients have returned to baseline, 54 were admitted to hospitals; of the hospitalized patients six have died, and 40 have been discharged.

Discussion

Our primary care response to the COVID-19 pandemic builds upon our knowledge and expertise in population management for chronic conditions with the need for proactive management and outreach in order to achieve best patient outcomes. We leveraged our experience with existing EMR tools and workflows developed for population management of patients with conditions such as diabetes and opioid addiction when we realized that responding to the COVID-19 pandemic would require a similar approach. Our new population management strategy for COVID-19 patients enables us to provide these patients with remote support and monitoring so that they can isolate and recover at home where they are most comfortable when appropriate, while also allowing us to identify those patients requiring additional assessment and support in a timely manner. Additionally, this strategy helps conserve PPE and medical staff resources both in our clinic and in our Emergency Department. It is worth noting that the robustness of remote monitoring for COVID-19 patients would be enhanced by an increased availability of at-home blood oxygen saturation monitors and/or improved accuracy of oxygen saturation smartphone apps.

Our new algorithm, EMR component, and population report for COVID-19 patients allows us to not only identify patients at high risk of disease complications, but also to focus our outreach on symptom days 5-10, when patients are most vulnerable. Frequent conversations with these patients enable us to follow their symptoms and quickly determine when a patient is deteriorating, and we have sent patients to the Emergency Department who required admission. Patients, both young and old, who are socially isolated in their homes due to the COVID-19 pandemic, have told us how much they have appreciated our calls. We anticipate using this program as long as COVID-19 is active in our community, and we believe other primary care practices could successfully implement our population strategy to help them optimize care for their own COVID-19 patients.

Declarations

Acknowledgments: *The authors wish to thank the Kathy Siranosian and Karen Freund, MD MPH for their contributions to this manuscript. We also thank the Tufts Medical Center Primary Care Boston teams for embracing this intervention and providing such expert and compassionate care to our patients.*

Corresponding Author: *Deborah Blazey-Martin, MD MPH; Chief, Division of Internal Medicine and Adult Primary Care, Tufts Medical Center, Boston, MA, USA (e-mail: dblazey-martin@tuftsmedicalcenter.org).*

Compliance with Ethical Standards:

Conflict of Interest: *The authors declare that they do not have a conflict of interest.*

Ethical Approval: *This study was given an exempt approval (STUDY00000569) by the Tufts Health Sciences Institutional Review Board (IRB).*

References

¹ Report of the WHO-China Joint Mission on coronavirus disease 2019 (COVID-19).

[https://www.who.int/publications-detail/report-of-the-who-china-joint-mission-on-coronavirus-disease-2019-\(covid-19\)](https://www.who.int/publications-detail/report-of-the-who-china-joint-mission-on-coronavirus-disease-2019-(covid-19)) Date: Feb 28, 2020; Date accessed: April 21, 2020.

² Huang, Chaolin, et al. "Clinical Features of Patients Infected with 2019 Novel Coronavirus in Wuhan, China." *The Lancet*, vol. 395, no. 10223, 2020, pp. 497–506., doi:10.1016/s0140-6736(20)30183-5.

³ Yang, Xiaobo, et al. "Clinical Course and Outcomes of Critically Ill Patients with SARS-CoV-2 Pneumonia in Wuhan, China: a Single-Centered, Retrospective, Observational Study." *The Lancet Respiratory Medicine*, 2020, doi:10.1016/s2213-2600(20)30079-5.

⁴ Zhou, Fei. "Clinical Course and Risk Factors for Mortality of Adult Inpatients with COVID-19 in Wuhan, China: a Retrospective Cohort Study." *The Lancet*, 2020, doi.org/10.1016/S0140-6736(20)30566-3.

⁵ Richardson, Safiya, et al. "Presenting Characteristics, Comorbidities, and Outcomes Among 5700 Patients Hospitalized With COVID-19 in the New York City Area." *JAMA*, 2020, doi:10.1001/jama.2020.6775.

Figures

COVID-19 OUTREACH/Population Management – As of 4/9/2020

	Day 1-5 COVID symptoms	Day 5-10 COVID Symptoms			COVID Post-Discharge Management
Risk Assessment Using COVID outreach component	<p><u>Risk assessment—</u> ✓ Low if no comorbidities, ✓ High if 1 or more Comorbidities: Age > 60, Immunocompromise, Chronic lung dz, Poorly controlled DM, CHF, Transplant, CKD 4/5 <u>Symptom assessment—</u> Fever, chills, myalgia, anosmia/ageusia, anorexia, pharyngitis <u>COVID-19 diagnosis—</u> Positive test? Clinical diagnosis? ✓ Add to problem list ✓ Document date symptoms started</p>	<p>Questions:</p> <p>1) Shortness of Breath—measure O2 sat vs subjective dyspnea Mild dyspnea but improving? Trouble going up a flight of stairs without getting winded? Pleuritic chest pain? Elevated respiratory rate >22-25? O2 sat <95%</p> <p>2) Trouble remaining hydrated? Decreased fluid intake, <50% normal Lightheaded on standing? Unable to keep fluids or meds down? ?Vomiting and/or diarrhea?</p> <p>3) Fever >100.4, >102 Responding to antipyretics or not? Myalgias and headaches?</p> <p>4) Chronic conditions well-controlled?</p>			<p>Add COVID outreach component to post discharge encounter</p> <p>Family support?</p> <p>Continue isolation for at least 7 d with mask for 14 days from start of symptoms</p> <p>Follow recommendations for Symptom Day outreach (Day 1-5 vs 5-10)</p> <p>VNA involvement?</p> <p>Comorbidity management</p>
Management	<p><u>Prevent spread-</u> 1) Health care worker or group living? Must be tested 2) Home isolation and social distancing (wash hands, own bathroom and bed, wipe surfaces) 3) Do they have VNA? Live alone? 4) Encourage hydration, take regular meds 5) Counsel about when to call 911, GMA</p>	<p>Manage at home</p> <ul style="list-style-type: none"> Breathing ok Fevers controllable Able to eat and drink Comorbidities managed Remain isolated 	<p>UC Respiratory Clinic Assessment</p> <ul style="list-style-type: none"> High risk comorbidities CXR/O2 sat would help with triage Other comorbidities need to be assessed Testing might put patient on non-COVID service Needs further assessment/support but not ED level of care 	<p>Send to ED</p> <ul style="list-style-type: none"> Immediate support needed—hydration, O2 911 vs family to drive patient Comorbidity needs acute management (chest pain, etc) 	<p>When to stop phone outreach</p> <ol style="list-style-type: none"> Breathing ok Afebrile w/o antipyretics Back to baseline and >day 10 <p>Return to work advice</p> <ul style="list-style-type: none"> 72 hours afebrile w/o antipyretics AND >7 days total from start of symptoms Home isolation and social distancing COVID symptoms have resolved Call your own Employee Health Mask for 14 days total from date of symptom onset
OTHER	<p>Low risk—call every 2 d High risk— call daily</p>	<p>Low risk call daily High risk 2x per day</p>	<p>Low risk call daily High risk 2x per day</p>	<p>Call 2x per day if not admitted</p>	<p>Call based on illness day (1-5 vs 5-10)</p>

CDC Symptom Checker <https://www.cdc.gov/coronavirus/2019-ncov/symptoms-testing/index.html>
 CDC 10 things to care for self at home <https://www.cdc.gov/coronavirus/2019-ncov/downloads/10Things.pdf>
 Produced By: Liz Barnhart, FNP, Deb Blazey-Martin, MD and Gabriela Andujar Vasquez, MD

Figure 1

COVID -19 Outreach Algorithm

CareTransCOVID: DINELI LOGICIAN

Outreach COVID-19

Patient Details

Risk Factors: CDC Page COVID Risk Factors

High Risk Factors: Age 65+, Obesity, Nursing Home/LTC, Diabetes, Chronic Lung/Asthma, CKD, Heart Conditions, Liver Disease, Immunocompromised.

Patient is High Risk: No Yes Problem List (Pop-Up)

Date Hospital Admission: Date Hospital Discharge:

Hospital Name:

Visit Details

Minutes: Include time reviewing labs, recent hospital visits, and any other reports

Patient Vitals: *(Patient supplied data, stored in alternate OBS variables.)*

BP: / on Pt supplied BP: /

BMI: on

Temperature: on Pt supplied Temp:

Respiration: on Pt supplied Resp Rate:

Pulse: on Pt supplied Pulse:

O2 Sat: on Pt supplied O2 Sat: 97%

Caregiver: Home Alone Family Friend Other

Congregate Setting:

COVID Status: Asymptomatic

Notes:

COVID-19 Review

Lab: Record outside lab results with first yellow section; record Presumed Positive with orange section.

Outside Lab: Date: 05/04/2020 Record Outside Lab

Should contain word 'positive' in the New field. Make sure to click the box to RECORD the data.

Presume Pos: Date: Record Pt Presum Pos

Use to indicate patient presumed to have COVID. Make sure to click the box to RECORD the data.

Date of Symptoms: 05/03/2020 Days since (calculated): 6

Assessment: CDC Page COVID Steps When Sick

Remember to check the box for the condition, and fill in the text area with details.

Breathing Issues or Cough

Fever

Dehydration

Myalgia or Headaches

Other

Plan:

Action:

Med Reconciliation: Done Overall Symptom Assessment: Mild Moderate Severe

Disposition:

Call Back on: 05/09/2020 Schedule Visit For:

Patient sent to the ER: Yes

Patient Resolution: (Typically = Back to Baseline)

Return To Work: OK For Date:

Add Problems
Add Medications
Add Allergies
Add Directives
Add Orders

Prev Form (Ctrl+PgUp)
Next Form (Ctrl+PgDn)
Close

Figure 2

COVID-19 Outreach EMR Component (GE Centricity)

COV120-COVID19_Registry-2020-05-09-06-55-18.xls

New Open Save Print Import Copy Paste Format Undo Redo AutoSum Sort A-Z Sort Z-A Gallery Toolbox Zoom Help

	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P
	MRN	Patient Name	Age	Tri	PCP	C19_pos	SymDa	PRis	SymptRsl	HspAdm	HspDsc	CalBckD	EstOutcr	Q	Resolution	Next Sche
9			38	6B	Tishler, MD, Julie	05/06	4	No	Mild			05/11	05/07	1		
11			73	6B	McNamara, MD, M	05/06	8	Yes	Severe			05/09	05/08	4		06/19
12			31	4A	Gao, MD, -Jenny	05/06	8	Yes	Mild			05/09	05/08	2		
13			30	Sys	Termination Insure	05/06										
14			69	4B	Ramadan, MD, Fa	05/06		Yes					05/07	1	In-Patient	
15			47			05/06										
16			62	6B	Carroll, MD, Niam	05/05	5	No	Moderate			05/10	05/08	2		
17			63	6B	McNamara, MD, M	05/05	6	Yes	Mild			05/09	05/08	3		
18			44	5_	Kaplan, MD, Lindi	05/05	7	No	Mild				05/08	2		
19			63	4B	Chu, MD, Jaclyn	05/05	24	Yes	Moderate			05/01	04/30	6	In-Patient	
20			88	4A	Huang, MD, Kristir	05/05	26					05/12	05/02	1		06/04
23			46			05/05										
25			30	Sys	PCP not in GMA, I	05/05										
27			42	4B	Naef, MD, Jana	05/04	5					05/13	05/08	1		
29			72	4A	Kopelman, MD, Ri	05/04	28	Yes	Mild				05/08	3		07/10
30			46	4A	Gao, MD, -Jenny	05/03	6		Moderate			05/15	05/08	2	In-Patient	08/10
32			32	5_	Chinoy, MD, Sami	05/03	7		Mild			05/11	05/08	3		
36			68	4A	Gao, MD, -Jenny	05/03	11	Yes	Moderate			05/09	05/08	5		07/20
37			32	4B	Lau, MD, Sandra	05/03	46	No	Mild				05/08	4		
38			31	5_	Mouli, MD, Manas	05/03		No	Mild			05/11	05/07	2		
40			66	4B	Chu, MD, Jaclyn	05/02	7	Yes				05/13	05/08	2		06/09
43			39	6A	Ahearn, MD, Dine	05/02	11	No	Mild			05/11	05/08	3		
47			46	4A	Raval, MD, Pooja	05/01	4		Moderate			05/11	05/07	2		
49			50	R2020	[-KM 4B-] Rao M,	05/01	8	No	Mild			05/11	05/08	1		
50			77	6A	Ahearn, MD, Dine	05/01	8	Yes	Moderate			05/09	05/08	3		08/11
52			68	6A	Ahearn, MD, Dine	05/01	15	Yes	Mild			05/12	05/08	2		11/05
53			66	6A	Murphy, MD, Kerri	05/01	22	Yes	Mild			05/11	05/08	7	almost back to baseline	
54			54			05/01							05/04	1	In-Patient	
57			59	R2022	[-JC 4B-] Tran M,	04/30	12	Yes	Mild			05/10	05/08	4	In Rehab or SNF	
62			63	4A	Chui, MS, NP, Ele	04/29	14						05/04	5		
70			24	6B	Schelling, MD, Kin	04/29	14	No	Mild			05/11	05/08	4		
72			73	4B	Wilder, MD, Elisab	04/29	15	Yes	Mild			05/08	05/08	5		08/19
77			64	4B	Kross, MD, Joan	04/28	14	Yes	Mild			05/12	05/08	5		07/20
78			33	4A	Sathi, MD, Kinjalik	04/28	26		Mild			05/11	05/08	4		12/21
81			55	6A	Naggar, MD, Jack	04/28	31	Yes	Moderate			05/11	05/08	10	In-Patient	07/17
83			63	5_	Kaplan, MD, Lindi	04/27	16		Mild			05/11	05/07	7		01/11
86			42	4B	Chawla, MD, Priar	04/27	19	Yes	Mild			05/08	05/08	9	discharge from hos	07/13
89			80	5_	Duncan, MD, Paul	04/25	24	Yes	Severe			04/24	04/24	7	Deceased	
92			51	Sys	PCP not in GMA, I	04/26	13					04/26		1		
96			69	4B	Chawla, MD, Priar	04/25	16					05/11	05/07	5	ICU	06/22
100			55	6A	Marturano, MD, C	04/25	19	Yes	Mild			05/09	05/08	6	almost back to bas	05/20
103			80	6B	Tishler, MD, Julie	04/25	50	Yes					05/04	3	ICU	
106			64	6B	Krause, MD, Dian	04/24	15	Yes	Moderate				05/08	9	In-Patient	07/13
109			90	4B	Ramadan, MD, Fa	04/24	16	Yes	Mild			05/08	05/07	6	In-Patient	07/29
113			41	4A	Sathi, MD, Kinjalik	04/24	19		Moderate			05/11	05/08	10		
115			64	6A	Marturano, MD, C	04/23	19	No	Severe			05/09	05/08	6	In-Patient	05/11
116			42	4B	Ramadan, MD, Fa	04/23	22	Yes	Mild			05/08	05/08	9		
129			33	6A	Marturano, MD, C	04/23	23	No	Mild			05/09	05/08	9	In-Patient	05/19
140			69	4B	Kross, MD, Joan	04/22	19	Yes	Mild			05/11	05/08	9	In-Patient	05/15
142			71	6B	Schelling, MD, Kin	04/22	22	Yes	Moderate			05/11	05/07	4	In-Patient	05/11
143			45	4B	Wilder, MD, Elisab	04/22	24	Yes	Moderate			04/29	04/29	7		
153			65	6B	Krause, MD, Dian	04/22	26	Yes	Mild				04/24	6	ICU	
157			54	4A	Sathi, MD, Kinjalik	04/21	18		Mild			05/11	05/08	11		
179			27	6B	McNamara, MD, M	04/20	19	Yes	Mild				05/04	8		02/09
199			60	5_	Chandler, MD, Da	04/20	24	Yes	Mild				05/07	9		

Figure 3

COVID-19 Daily Outreach Report