Association between short sleep duration and overweight/obesity among middle-school students: a cross-sectional study in Fuzhou, China

Guobo Li

Fujian Medical University

Shichao Wei

Fujian Provincial Hospital

Fuhao Zheng

Fujian Provincial Hospital

Xingyan Xu

Fujian Medical University

Yingying Cai

Fujian Medical University

Siyang Wu (✉ fmuwsy@163.com)

the Key Laboratory of Environment and Health among Universities and Colleges in Fujian; Fujian Medical University

Research article

Keywords: Short sleep duration, Overweight, Obesity, Adolescents, Association

Posted Date: October 22nd, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-30129/v3>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: This study was designed to investigate the prevalence of short sleep duration and explore the association between short sleep duration and overweight or obesity in adolescents from middle school in Fuzhou, China.

Methods: A total of 13,063 students (male: 6,500; female: 6,553) from 18 different schools were included in our study. Questionnaires focusing on short sleep duration and overweight or obesity related factors were collected. The sleep duration was self-reported by the subjects. The body weight and height of each subject were measured by the qualified personnel. Logistic regression analysis was used to evaluate the effects of short sleep duration on overweight or obesity in school students.

Results: The overall rates of overweight and obesity were 12.1% and 7.1%, respectively. The prevalence of short sleep duration among students was 82.8%. The majority of high school students (92.5%) suffered from short sleep duration. Compared with male children with a sleep duration of 8-10 h day, the odds ratios (95% *CI*) of overweight/obesity for those with a sleep duration of less than 6 h or 6-8 h, were 1.63 (1.25-2.13) and 1.06 (0.88-1.27) after adjusting social and demographic status, mental health and lifestyle factors. The homologous odds ratios of female children were 1.38 (0.99-1.93) and 1.04 (0.79-1.35), respectively.

Conclusions: A large number of adolescents suffered from short sleep duration. Short sleep duration is associated with higher odds of overweight/obesity in male children.

Background

Overweight and obesity, with an increased trend in the past few decades, is becoming a threat to the public health worldwide. In a recent survey, about 711.4 million individuals were diagnosed with obesity, among which 107.7 million were children [1]. Meanwhile, a large proportion of adolescents (19.4%) aged 7-18 yrs presented overweight and obesity in a national survey [2]. In China mainland, the number of obese children is very huge because of alternations in the diet compared with the previous decades [1]. Fuzhou is the provincial capital of Fujian Province localized in the southeast of China. In 2010, Lin et al reported the body development of the students in middle school in Fuzhou city [3], however, few studies focused on the incidence of overweight/obesity and the relationship between overweight/obesity and the sleep duration in these students.

Obesity is considered to be associated with the genetic and environmental factors [4, 5]. Nowadays, more and more attention has been paid to sleep duration among the individuals with obesity [6]. Short sleep duration among students is regarded to be related to increased risk of obesity, injuries, behavior problems, attention-deficit disorder, poor academic performance and psychological problems [7-9]. In adulthood, short sleep duration is a risk factor of cardiovascular diseases [10]. Short sleep duration mainly affects metabolism, endocrine function and immune systems [11]. However, there are still some controversies on the relationship between short sleep duration and overweight or obesity. Some studies proposed a U-

shaped [12, 13] or linear inverse correlation [14] between sleep duration and obesity, while others proposed no associations [15, 16]. Meanwhile, the association between short sleep duration and the overweight/obesity was somehow modulated by gender [17] and it was proposed that gender should be considered a specific trait for short sleep duration in these with overweight/obesity [18, 19]. In this study, we aimed to investigate the prevalence of short sleep duration among students in Fuzhou. In addition, we explored the relationship between short sleep duration and overweight or obesity.

Methods

Survey design

In this survey, cluster random sampling method was utilized. Firstly, five representative cities and four rural areas were selected from Fuzhou City. Secondly, the adolescents from 2 selected middle schools were sampled randomly from the selected counties. Thirdly, all students in the chosen classes were invited to participate in the study by the qualified teachers in their school after informing about the potential benefits and risks by our staff. All the subjects were invited to participate in this study at their wills. The subjects not willing to participate in this survey were excluded from this study. A total of 15025 students from 18 different schools were invited to this study, with 14,499 students participating in the survey (response rate: 96.5%). After exclusion of subjects with missing key variables (including sex, height, weight and sleep duration), 13,063 eligible subjects (male: 6500; female: 6553) were included in the final analyses. The missing data were handled in the regression analysis as shown in Table 1.

Existing investigation was modeled from the questionnaire including the Youth Risk Behavior Survey, performed by the Centers for Disease Control and Prevention [20], and the international Global School-based Student Health Survey conducted with the support of WHO [21]. Questionnaires distributed to the respondents included the demographic characteristics (e.g. age and gender), athletic activity, screen-time, nutrient supplements, as well as sleep and mental health (e.g. sadness, annoyance and loneliness). This survey was conducted between May and June in 2019. The age of the youngest participants included in the study was 12 years. Written parental consent was obtained on behalf of participants below the age of 16. The study was approved by the Ethical Committee of Fujian Medical University (approval No. 2019-117).

The sample size was calculated using the following formula: $N = deff \times \mu^2 \times P \times (1 - P) / d^2$. In addition, the 95% CI (2-sided for $\mu = 1.96$) was determined, and the measure of probability (P) was the obese rate (8.6%) of China [2]. The design effect ($deff$) value was set to 2 and the relative error (d value) was $d = r \times 0.01$ ($r = 15\%$). On this basis, the sample size was 2,680 for each stratum. After taking 4 strata (i.e. boy and girl, urban and rural area) and an assumed potential non-response rate of 10% into consideration, the final sample size was 11,929.

Definition of the variables

The key schools were defined as those supported by the local government and the education bureau, with more high quality teachers and students. The ordinary schools were those supported less by the local government and the education bureau, and lower in the quality of the teachers and students. The environment for the learning and capacity of the key schools was much better than that of the ordinary schools.

Overweight and obesity screening

The body weight and height of each subject were measured by the qualified staff annually in the routine physical examination. Body mass index (BMI) was the result of measured weight (kg) divided by the square of height in meters. The diagnostic criteria for overweight and obesity among adolescents were conducted based on the guidelines proposed by the Chinese Working Group on Obesity for Children (WGOC) [22]. Overweight was defined as increase of one standard deviation (SD) of BMI compared with the normal children of the same gender at the same age, and obesity was defined as increase of two SDs of BMI. The disease history was inquired by the staff, followed by physical examination to exclude the secondary obesity induced by severe diseases, endocrine or metabolic disorders.

Sleep duration

Sleep duration was calculated by self-administered questionnaires by recalling the average time of falling asleep and getting up in the preceding 7 days. The average time of using electronic products each day was also self-administered. Short sleep duration was defined as a sleep duration less than 8 hrs a day for teenagers aged ≥ 13 according to the recommendation of American Academy of Sleep Medicine (AASM) [23]. Sleep duration was subdivided into 3 groups: "<6h", "6-8 h", and "8-10 h", as a categorical variable in the multivariable logistic regression.

Co-variates

The questionnaire also investigated other factors related to obesity, including the gender and age, days for consuming breakfast and night snack, physical activity, homework time, high-energy snacks, fast food intake and nutrient supplements.

Statistical analysis

SPSS version 21 was used for the statistical analysis. Continuous variables were described by mean $\pm$ standard deviation. Chi square test was utilized to analyze the differences in overweight and obese rates of different groups. Logistic regression analysis was conducted to investigate the relationship between sleep duration and obesity. Adjusted odds ratios (ORs) and 95% CI were evaluated through three logistic regression models. $P < 0.05$ was considered to statistical significance. In this study, obesity and overweight was classified as one group based on the following aspects: (1) the sample size for obesity was small, which may hamper the confidence of the data. (2) The effects of short sleep duration on overweight and obesity were similar. There was a relationship between sleep duration and overweight. Short sleep duration may lead to overweight in children [24, 25]. Besides, short sleep duration may

increase the risk of obesity in children [26-28]. Meanwhile, many studies classified the overweight and obesity as the same group, in order to analyze the effects of sleep duration on overweight and obesity [13, 29-31].

In order to further validate whether the gender would modulate the effects of overweight on the short sleep duration, we analyzed the correlation between short sleep duration and overweight/obesity among the male and female adolescence. In the Logistics regression analysis, no additional variable was given in Model 0. The Model 1 of total subject involved adjustment for gender, age, region, types of school. For the Model 1 of male and female, we adjusted gender, region, and types of school. For the Model 2 of the three aspects, we additionally adjusted for whether using electronic product daily during school, electronics in bedroom, someone smoking at home, strenuous physical activity, moderate physical activity, days of consuming breakfast and night snack per week, having fried food or pastry in the past 12 months, having taken nutrient supplements in the past 12 months. Besides, they were further adjusted for feeling annoy, sad or lonely in the Model 3 of the three aspects.

Results

Respondent characteristics

Finally, 13,063 subjects (male: 6,500; female: 6,553) were included in this study. The respondents with short sleep duration had longer screen-time than that of the counterparts with short sleep duration, ($P<0.001$, Table 2). Meanwhile, compared with those with short sleep duration, they less frequently participated in physical activity ($P<0.001$) or consuming breakfast ($P<0.001$). Instead, short sleep duration was significantly linked to consuming night snack ($P<0.001$) and feeling irritated ($P<0.001$), sad ($P<0.001$) and lonely ($P<0.001$).

Overweight and obesity

The prevalence of overweight and obesity were 12.1% and 7.1%, respectively (Table 3). They were statistical differences in the rate of obesity and overweight among various regions ($P<0.001$). No statistical differences were noticed in the rate of obesity and overweight between the students in the key schools and ordinary schools ($P=0.566$). The prevalence of obesity rate of among the male children was significantly higher than that of female counterparts ($P<0.001$).

Short sleep duration

The total prevalence of short sleep duration was 82.8%. The prevalence of short sleep duration among the female children was significantly higher than that in male counterparts (86.0% vs. 79.5% $P<0.001$). There were statistical differences among the prevalence of short sleep duration in the students aged ≤ 13 yrs (67.7%), those aged 14-15 yrs (77.8%) and those aged ≥ 16 yrs (92.5%, $P<0.001$). The prevalence of short sleep duration in the students lived in the urban was significantly lower than those lived in the rural areas

(80.5% vs. 84.6%, $P < 0.001$). No statistical difference was noticed in the prevalence of short sleep duration in the students of the key schools and ordinary schools (82.9% vs. 82.6%, $P = 0.726$).

Relationship between sleep duration and overweight/obesity

Model 1 showed that, compared with all the subjects and male children who slept 8-10h per day, those who slept for less than 6 hrs a day showed a higher possibility of overweight/obesity after adjusting socio-demographic status [OR=1.32 (95%CI: 1.12-1.55) and 1.25 (95%CI: 1.02-1.54)] (Table 4 and Table 5). After additionally adjusting life-style factors in Model 2, all the subjects and male children who slept for less than 6 hrs a day showed a significantly higher association for overweight/obesity by comparing to the reference group [OR=1.55 (95%CI:1.27-1.89) and 1.65, (95%CI:1.27-2.13)]. The odds ratios were 1.53 (1.24-1.87) and 1.63 (1.25-2.13), after further adjustment for mental health such as feeling irritation, sad or lonely. Nevertheless, there were no significant differences among the female children. The homologous odds ratios (95%CI) for female children were 1.38(1.06-1.81) in Model 1, 1.41 (1.02-1.96) in Model 2 and 1.38 (0.99-1.93) in Model 3, respectively. Nevertheless, the lower CI was very close to significance.

Discussion

This study revealed the link between short sleep duration and overweight/obesity in adolescents of Fuzhou, China. After adjusting socio-demographic status, lifestyle factors, and mental health, there was a negative correlation between short sleep duration and overweight/obesity among male children.

The prevalence of overweight or obesity in our study was 19.2%, which was higher than the national average [2]. Moreover, our findings demonstrated the differences among gender, age and region, which were similar with the Report on Childhood Obesity in China [32]. The prevalence of obesity showed decline with the elevation of the ages, which may be possibly related to physical development and attention to their appearances[33, 34].

The prevalence of obesity in male children was significantly higher than that of female children. In addition, the prevalence of short sleep duration in adolescents showed strike increase with age. About 92.4% of adolescents in high school suffered from short sleep duration. Study pressure, especially pressure from the college entrance examination, may contribute to such a high prevalence.

Our study revealed that short sleep duration was negatively related to overweight/obesity which was consistent with previous study showing that sleep deficiency seemed to parallel the increased prevalence of obesity[35]. Meanwhile, it could be an independent risk factor for obesity[10, 36, 37]. One study provided causal evidence on the relationship between short sleep duration and weight gain in the population-level [38]. Moreover, Krietsch et al reported that there was a U-shaped correlation between those with short sleep duration and obesity only in the female children [39]. These differences may be related to the physiology of adolescence between female and male.

To date, little is known about the relationship between sleep and obesity. Sleep duration involved in the regulation of cerebral function such as controlling the appetite, which could lead to over-eating in an obesogenic environment [40]. The homeostatic control of appetite was achieved by complex interactions among numerous neuroendocrine hormones [41]. Many pivotal hormones (e.g. insulin, leptin, cortisol and ghrelin) [42] may involve in the correlation between sleep and obesity. Even after adjusting the BMI, sleep duration was negatively correlated with circulating leptin [43]. Eptin pathway could explain the key mechanism via modificatory effects [44]. Under some circumstances, the short sleep duration could lead to disruption of insulin, leptin, cortisol and ghrelin expression [45, 46]. After a period of sleep loss, people could experience a 24% increase in hunger with largely whetting the appetite for high carbohydrate foods [45]. In our study, students with short sleep duration spent more time on night snack than those with adequate sleep, which may lead to weight gain. Fatigue caused by short sleep duration may result in reduced physical activity, which then promoted the weight gain [47, 48]. In this study, adolescents with adequate sleep did more exercise of moderate-intensity than those with short sleep duration.

Circadian Locomotor Output Cycles Kaput (CLOCK) genes involved in regulation of diurnal rhythm, and their effects on neuroendocrine systems might have an impact on obesity [49]. The variants of *CLOCK* gene was related to sleep duration [50], as with calorie intake [51], metabolic syndrome [52], and obesity [53]. Meanwhile, methylation of *CLOCK* gene was associated with carbohydrate intake, total energy intake, insulin resistance, and BMI [54]. *REV-ERBa* rs2071570 and rs2071427 were related to BMI and sleep duration in male children, confirming the association of the *REV-ERBa* gene with human obesity, mainly in males [55]. In line with our findings, this theory supported a negative correction between sleep duration and overweight/obesity among male children [56].

Obesity is not simply related to biology and behavior, but also to the our social context [57]. Social support from friends and awareness/internalization of thinness ideals were significantly related to odds of overweight/obesity in youth. Such association varied by age and sex, and persisted after control for intra-familial factors such as overall family support/function, diet and activity specific support [58]. A cross-sectional study on societal risk factors for overweight and obesity in women in Zimbabwe showed that the key social factors associated with overweight and obesity were older age, wealthy and the use of hormonal contraception. Besides, a higher education and being Christian also increased the risk of obese and overweight, respectively [59]. Some home environment related factors may increase the incidence of overweight/obesity among children, such as sleeping too late that may lead to inadequate sleeping and increased opportunity for taking food in night [60]. Less daytime care by mothers and shorter sleep duration were associated with increased risk of becoming overweight during childhood [61].

In this study, there were no statistical differences between the short sleep duration and overweight/obesity in the female children, however, the lower CI (95% CI 0.99-1.93) was very close to significance. It appeared that this association was just smaller than what was observed for males. It may be that the stratified analysis lacked statistical power. In our subsequent study, studies of a large sample size involving more social variables are required to investigate the effects of short sleep duration on the overweight/obesity in female children.

A large and representative sample was included in the survey. Meanwhile, we analyzed a wide range of covariance to verify the relationships between sleep duration and overweight or obesity by adjusting potential confounding factors correlated with overweight or obesity. Nevertheless, there are some limitations in our study. First, causal inference will be limited in the cross-sectional design, although there are several theories supporting our findings. Second, the determination of sleep duration was reported by the students themselves, which was a limiting factor. Indeed, the facilities may contribute to the reduction of the errors to some extent, however, it was a challenge for the promotion of the facility as the sample size was too large. In the subsequent study, representative samples will be selected for the monitor of the sleep duration of the subjects including sleep quality. Third, the impact of prolonged sleep duration on overweight or obesity among adolescents was not explored in our study, as we laid emphasis on the risk of overweight or obesity induced by short sleep duration in middle-school students. Then we will focus on the effects of long sleep duration on the overweight and obesity in the students. Fourth, the psychologic status of the subjects when self-reporting may lead to some bias of the results. In a follow-up study, qualified mental scales will be required to evaluate the psychologic status including depression and sorrow, which can correct the effects of psychologic variables on overweight and obesity.

Conclusion

Adolescents suffered from short sleep duration. Short sleep duration is associated with higher odds of overweight/obesity in male children. In female children, further studies are needed to investigate the potential association between short sleep duration and overweight/obesity. In future, further researches are required to investigate the relationship between sleep duration and specific hormones in adolescents, in order to explore the potential mechanisms of overweight or obesity.

Abbreviations

Body mass index (BMI)

Chinese Working Group on Obesity for Children (WGOC)

American Academy of Sleep Medicine (AASM)

Odds ratios

Circadian Locomotor Output Cycles Kaput (CLOCK)

Declarations

Ethics approval and consent to participate

The study was approved by the Ethical Committee of Fujian Medical University.

Consent for Publication

Written parental consent was obtained on behalf of participants below the age of 16.

Availability of data and material

All the data were available upon appropriate request.

Competing interests

The author(s) declare(s) that there is no conflict of interest regarding the publication of this paper.

Funding

This work was funded by Joint Funds for the Innovation of Science and Technology, Fujian Province (2018Y9089), the Natural Science Foundation of Fujian Province (Grant No. 2019 J01315), and Professor Development Fund Project of Fujian Medical University (JS15002), and Open Project of Fujian Key Laboratory of Environmental Factors and Cancer (GWSZD-201802). These funds showed no roles in the design of the study.

Authors' contributions

Designed and modified the manuscript: WSC and WSY; Designed the research and participated in the experimental design, coordinated and drafted the manuscript: LGB, ZFH, XXY, CYY; Data collection, achievement interpretation and manuscript writing: LGB, ZFH, XXY, CYY; Analysed the data: LGB, XXY, CYY. All of the authors have given final approval of the version to be published.

Acknowledgements

Not applicable.

References

1. Afshin A, Forouzanfar MH, Reitsma MB, Sur P, Estep K, Lee A, Marczak L, Mokdad AH, Moradi-Lakeh M, Naghavi M *et al*: **Health Effects of Overweight and Obesity in 195 Countries over 25 Years**. *The New England journal of medicine* 2017, **377**(1):13-27.
2. Wang S, Dong YH, Wang ZH, Zou ZY, Ma J: **[Trends in overweight and obesity among Chinese children of 7-18 years old during 1985-2014]**. *Zhonghua yu fang yi xue za zhi [Chinese journal of preventive medicine]* 2017, **51**(4):300-305.
3. al XLQZQZe: **A Survey on the Growth and Development of primary and middle school students in Fuzhou [J]**. *Strait Journal of Preventive Medicine* 2014.
4. Wang L, Perez J, Heard-Costa N: **Integrating genetic, transcriptional, and biological information provides insights into obesity**. 2019, **43**(3):457-467.
5. Vasanth Rao VRB, Candasamy M, Bhattamisra SK: **Obesity an overview: Genetic conditions and recent developments in therapeutic interventions**. *Diabetes & metabolic syndrome* 2019, **13**(3):2112-

2120.

6. Celis-Morales C, Lyall DM, Guo Y, Steell L, Llanas D, Ward J, Mackay DF, Biello SM, Bailey ME, Pell JP *et al*: **Sleep characteristics modify the association of genetic predisposition with obesity and anthropometric measurements in 119,679 UK Biobank participants.** *The American journal of clinical nutrition* 2017, **105**(4):980-990.
7. Allott EH, Hursting SD: **Obesity and cancer: mechanistic insights from transdisciplinary studies.** *Endocrine-related cancer* 2015, **22**(6):R365-386.
8. Bragg F, Tang K, Guo Y, Iona A, Du H, Holmes MV: **Associations of General and Central Adiposity With Incident Diabetes in Chinese Men and Women.** 2018, **41**(3):494-502.
9. Lu Y, Hajifathalian K, Ezzati M, Woodward M, Rimm EB, Danaei G: **Metabolic mediators of the effects of body-mass index, overweight, and obesity on coronary heart disease and stroke: a pooled analysis of 97 prospective cohorts with 1·8 million participants.** *Lancet (London, England)* 2014, **383**(9921):970-983.
10. Miller MA, Kruisbrink M, Wallace J, Ji C, Cappuccio FP: **Sleep duration and incidence of obesity in infants, children, and adolescents: a systematic review and meta-analysis of prospective studies.** *Sleep* 2018, **41**(4).
11. Miller MA, Cappuccio FP: **Biomarkers of cardiovascular risk in sleep-deprived people.** *Journal of human hypertension* 2013, **27**(10):583-588.
12. Danielsen YS, Pallesen S, Stormark KM, Nordhus IH, Bjorvatn B: **The relationship between school day sleep duration and body mass index in Norwegian children (aged 10-12).** *International journal of pediatric obesity : IJPO : an official journal of the International Association for the Study of Obesity* 2010, **5**(3):214-220.
13. Wu J, Wu H, Wang J, Guo L, Deng X, Lu C: **Associations between Sleep Duration and Overweight/Obesity: Results from 66,817 Chinese Adolescents.** *Scientific reports* 2015, **5**:16686.
14. Lytle LA, Pasch KE, Farbakhsh K: **The relationship between sleep and weight in a sample of adolescents.** *Obesity (Silver Spring, Md)* 2011, **19**(2):324-331.
15. Calamaro CJ, Park S, Mason TB, Marcus CL, Weaver TE, Pack A, Ratcliffe SJ: **Shortened sleep duration does not predict obesity in adolescents.** *Journal of sleep research* 2010, **19**(4):559-566.
16. Hassan F, Davis MM, Chervin RD: **No independent association between insufficient sleep and childhood obesity in the National Survey of Children's Health.** *Journal of clinical sleep medicine : JCSM : official publication of the American Academy of Sleep Medicine* 2011, **7**(2):153-157.
17. Cao M, Zhu Y, Li X, Chen Y, Ma J, Jing J: **Gender-dependent association between sleep duration and overweight incidence in CHINESE school children: a national follow-up study.** *BMC public health* 2018, **18**(1):615.
18. Rosi A, Calestani MV, Parrino L, Milioli G, Palla L, Volta E, Brighenti F, Scazzina F: **Weight Status Is Related with Gender and Sleep Duration but Not with Dietary Habits and Physical Activity in Primary School Italian Children.** *Nutrients* 2017, **9**(6).

19. Cao M, Zhu Y, He B, Yang W, Chen Y, Ma J, Jing J: **Association between sleep duration and obesity is age- and gender-dependent in Chinese urban children aged 6-18 years: a cross-sectional study.** *BMC public health* 2015, **15**:1029.
20. **Centers for Disease Control and Prevention. State and local youth risk behavior survey.** 2017.
21. **World Health Organization. Global school-based students health survey (GSHS).** 2015.
22. **[Body mass index reference norm for screening overweight and obesity in Chinese children and adolescents].** *Zhonghua liu xing bing xue za zhi = Zhonghua liuxingbingxue zazhi* 2004, **25**(2):97-102.
23. Paruthi S, Brooks LJ, D'Ambrosio C, Hall WA, Kotagal S, Lloyd RM, Malow BA, Maski K, Nichols C, Quan SF *et al*: **Consensus Statement of the American Academy of Sleep Medicine on the Recommended Amount of Sleep for Healthy Children: Methodology and Discussion.** *Journal of clinical sleep medicine : JCSM : official publication of the American Academy of Sleep Medicine* 2016, **12**(11):1549-1561.
24. Tuohino T, Morales-Muñoz I, Saarenpää-Heikkilä O, Kiviruusu O, Paunio T, Hovi P, Pietiläinen KH, Paavonen EJ: **Short Sleep Duration and Later Overweight in Infants.** *The Journal of pediatrics* 2019, **212**:13-19.
25. Rangan A, Zheng M, Olsen NJ, Rohde JF, Heitmann BL: **Shorter sleep duration is associated with higher energy intake and an increase in BMI z-score in young children predisposed to overweight.** *International journal of obesity (2005)* 2018, **42**(1):59-64.
26. Sakamoto N, Gozal D, Smith DL, Yang L, Morimoto N, Wada H, Maruyama K, Ikeda A, Suzuki Y, Nakayama M *et al*: **Sleep Duration, Snoring Prevalence, Obesity, and Behavioral Problems in a Large Cohort of Primary School Students in Japan.** *Sleep* 2017, **40**(3).
27. Gileles-Hillel A, Almendros I, Khalyfa A, Nigdelioglu R, Qiao Z, Hamanaka RB, Mutlu GM, Akbarpour M, Gozal D: **Prolonged Exposures to Intermittent Hypoxia Promote Visceral White Adipose Tissue Inflammation in a Murine Model of Severe Sleep Apnea: Effect of Normoxic Recovery.** *Sleep* 2017, **40**(3).
28. Anderson SE, Andridge R, Whitaker RC: **Bedtime in Preschool-Aged Children and Risk for Adolescent Obesity.** *The Journal of pediatrics* 2016, **176**:17-22.
29. Wang F, Liu H, Wan Y, Li J, Chen Y, Zheng J, Huang T, Li D: **Sleep Duration and Overweight/Obesity in Preschool-Aged Children: A Prospective Study of up to 48,922 Children of the Jiaxing Birth Cohort.** *Sleep* 2016, **39**(11):2013-2019.
30. Krietsch KN, Armstrong B, McCrae CS, Janicke DM: **Temporal Associations Between Sleep and Physical Activity Among Overweight/Obese Youth.** *Journal of pediatric psychology* 2016, **41**(6):680-691.
31. Halal CS, Matijasevich A, Howe LD, Santos IS, Barros FC, Nunes ML: **Short Sleep Duration in the First Years of Life and Obesity/Overweight at Age 4 Years: A Birth Cohort Study.** *The Journal of pediatrics* 2016, **168**:99-103.e103.
32. M G: **the Report on Childhood Obesity in China.** *Beijing: People's Medical Publishing House* 2017.

33. Oellingrath IM, Hestetun I, Svendsen MV: **Gender-specific association of weight perception and appearance satisfaction with slimming attempts and eating patterns in a sample of young Norwegian adolescents.** *Public health nutrition* 2016, **19**(2):265-274.
34. Spencer RA, Rehman L, Kirk SF: **Understanding gender norms, nutrition, and physical activity in adolescent girls: a scoping review.** *The international journal of behavioral nutrition and physical activity* 2015, **12**:6.
35. Jean-Louis G, Williams NJ, Sarpong D, Pandey A, Youngstedt S, Zizi F, Ogedegbe G: **Associations between inadequate sleep and obesity in the US adult population: analysis of the national health interview survey (1977-2009).** *BMC public health* 2014, **14**:290.
36. Fatima Y, Doi SA, Mamun AA: **Longitudinal impact of sleep on overweight and obesity in children and adolescents: a systematic review and bias-adjusted meta-analysis.** *Obesity reviews : an official journal of the International Association for the Study of Obesity* 2015, **16**(2):137-149.
37. St-Onge MP, Shechter A: **Sleep disturbances, body fat distribution, food intake and/or energy expenditure: pathophysiological aspects.** *Hormone molecular biology and clinical investigation* 2014, **17**(1):29-37.
38. Do YK: **Causal Effect of Sleep Duration on Body Weight in Adolescents: A Population-based Study Using a Natural Experiment.** *Epidemiology (Cambridge, Mass)* 2019, **30**(6):876-884.
39. Krietsch KN, Chardon ML, Beebe DW, Janicke DM: **Sleep and weight-related factors in youth: A systematic review of recent studies.** *Sleep medicine reviews* 2019, **46**:87-96.
40. Sluggett L, Wagner SL, Harris RL: **Sleep Duration and Obesity in Children and Adolescents.** *Canadian journal of diabetes* 2019, **43**(2):146-152.
41. Morselli L, Leproult R, Balbo M, Spiegel K: **Role of sleep duration in the regulation of glucose metabolism and appetite.** *Best practice & research Clinical endocrinology & metabolism* 2010, **24**(5):687-702.
42. Li L, Fu J, Yu XT, Li G, Xu L, Yin J, Cheng H, Hou D, Zhao X, Gao S *et al*: **Sleep Duration and Cardiometabolic Risk Among Chinese School-aged Children: Do Adipokines Play a Mediating Role?** *Sleep* 2017, **40**(5).
43. Fu J, Wang Y, Li G, Han L, Li Y, Li L, Feng D, Wu Y, Xiao X, Li M: **Childhood sleep duration modifies the polygenic risk for obesity in youth through leptin pathway: the Beijing Child and Adolescent Metabolic Syndrome cohort study.** 2019, **43**(8):1556-1567.
44. Schmid SM, Hallschmid M, Jauch-Chara K, Born J, Schultes B: **A single night of sleep deprivation increases ghrelin levels and feelings of hunger in normal-weight healthy men.** *Journal of sleep research* 2008, **17**(3):331-334.
45. Jiang X, Hardy LL, Baur LA, Ding D, Wang L, Shi H: **Sleep duration, schedule and quality among urban Chinese children and adolescents: associations with routine after-school activities.** *PloS one* 2015, **10**(1):e0115326.
46. Spiegel K, Tasali E, Penev P, Van Cauter E: **Brief communication: Sleep curtailment in healthy young men is associated with decreased leptin levels, elevated ghrelin levels, and increased hunger and**

- appetite.** *Annals of internal medicine* 2004, **141**(11):846-850.
47. Allebrandt KV, Teder-Laving M, Akyol M, Pichler I, Müller-Myhsok B, Pramstaller P, Merrow M, Meitinger T, Metspalu A, Roenneberg T: **CLOCK gene variants associate with sleep duration in two independent populations.** *Biological psychiatry* 2010, **67**(11):1040-1047.
48. Mendelson M, Borowik A, Michallet AS, Perrin C, Monneret D, Faure P, Levy P, Pépin JL, Wuyam B, Flore P: **Sleep quality, sleep duration and physical activity in obese adolescents: effects of exercise training.** *Pediatric obesity* 2016, **11**(1):26-32.
49. Albrecht U: **The circadian clock, metabolism and obesity.** *Obesity reviews : an official journal of the International Association for the Study of Obesity* 2017, **18 Suppl 1**:25-33.
50. Monteleone P, Tortorella A, Docimo L, Maldonato MN, Canestrelli B, De Luca L, Maj M: **Investigation of 3111T/C polymorphism of the CLOCK gene in obese individuals with or without binge eating disorder: association with higher body mass index.** *Neuroscience letters* 2008, **435**(1):30-33.
51. Ramos-Lopez O, Samblas M, Milagro FI: **Circadian gene methylation profiles are associated with obesity, metabolic disturbances and carbohydrate intake.** 2018, **35**(7):969-981.
52. Garaulet M, Lee YC, Shen J, Parnell LD, Arnett DK, Tsai MY, Lai CQ, Ordovas JM: **Genetic variants in human CLOCK associate with total energy intake and cytokine sleep factors in overweight subjects (GOLDN population).** *European journal of human genetics : EJHG* 2010, **18**(3):364-369.
53. Scott EM, Carter AM, Grant PJ: **Association between polymorphisms in the Clock gene, obesity and the metabolic syndrome in man.** *International journal of obesity (2005)* 2008, **32**(4):658-662.
54. Nascimento Ferreira MV, Goumidi L, Carvalho HB, De Moraes ACF, Santaliestra-Pasías AM, Kafatos A, Molnar D, Lambrinou CP, De Henauw S, Gutierrez A *et al*: **Associations between REV-ERBa, sleep duration and body mass index in European adolescents.** *Sleep medicine* 2018, **46**:56-60.
55. Suglia SF, Kara S, Robinson WR: **Sleep duration and obesity among adolescents transitioning to adulthood: do results differ by sex?** *The Journal of pediatrics* 2014, **165**(4):750-754.
56. Wang H, Hu R, Du H, Fiona B, Zhong J, Yu M: **The relationship between sleep duration and obesity risk among school students: a cross-sectional study in Zhejiang, China.** *Nutrition & metabolism* 2018, **15**:48.
57. Hawkins D: **Reprint of: Obesity and social factors.** *Obesity research & clinical practice* 2016, **10**(3):355-356.
58. Bravin JI, Gutierrez AP, McCurley JL, Roesch SC, Isasi CR, Delamater AM, Perreira KM, Van Horn L, Castañeda SF, Pulgaron ER *et al*: **Extra-familial social factors and obesity in the Hispanic Community Children's Health Study/Study of Latino Youth.** *Journal of behavioral medicine* 2019, **42**(5):947-959.
59. Mangemba NT, San Sebastian M: **Societal risk factors for overweight and obesity in women in Zimbabwe: a cross-sectional study.** *BMC public health* 2020, **20**(1):103.
60. Ochoa A, Berge JM: **Home Environmental Influences on Childhood Obesity in the Latino Population: A Decade Review of Literature.** *Journal of immigrant and minority health* 2017, **19**(2):430-447.
61. Miyakoshi C, Yamamoto Y, Mishina H, Shirai C, Morioka I, Fukuhara S: **Childcare Environment and Japanese Children Who Are Overweight in Early Childhood.** *Childhood obesity (Print)* 2018, **14**(3):197-

Tables

Table 1 Proportion and handling of the missed data in the regression analysis

Variables	Total	Missed (%)	Method
Gender	13063	0 (0.00)	
Age	13063	0 (0.00)	
Region	13063	0 (0.00)	
School	13063	0 (0.00)	
Whether using electronic product daily during school	13063	0 (0.00)	
Using electronic products in bedroom	13063	0 (0.00)	Mode
Smoking at home	13063	125 (0.96)	Mode
Doing strenuous exercise in spare time	13063	283 (2.16)	Mode
Exercise at moderate intensity in spare time	13063	205 (1.57)	Mode
Days of consuming breakfast per week	13063	242(1.85)	Multiple imputation
Days of consuming night snack per week	13063	268(2.05)	Multiple imputation
Having fried food in the past 12 months	13063	71(0.54)	Mode
Having pastry in the past 12 months	13063	52(0.40)	Mode
Taken nutrient supplements in the past 12 months	13063	335(2.56)	Mode
Feeling lonely	13063	103(0.79)	Mode
Feeling sad	13063	108(0.83)	Mode
Feeling annoy	13063	262(2.01)	Mode

Table 2 Sleep duration of middle school students in Fuzhou, China

Characteristics	Total (n=13063)	<6 h (n=1762)	6-8 h (n=9051)	8-10 h (n=2250)	F/c ²	P value
Age (years)	14.44±1.64	14.94±1.65	14.53±1.64	13.66±1.36	369.68	<0.001
Gender					103.26	<0.001
Boys	6520(49.9)	802(45.5)	4381(48.4)	1337(59.4)		
Girls	6543(50.1)	960(50.1)	4670(51.6)	913(40.6)		
Types of school					3.86	0.145
Key school	7817(59.8)	1092(62.0)	5386(59.5)	1339(59.5)		
Ordinary school	5246(40.2)	670(38.0)	3665(40.5)	911(40.5)		
Place					57.41	<0.001
Urban	5924(45.3)	860(48.8)	3910(43.2)	1154(51.3)		
Rural	7139(54.7)	902(51.2)	5141(56.8)	1096(48.7)		
Screen time per week					210.20	<0.001
During school						
0-30minute	5386(41.2)	596(33.8)	3659(40.4)	1131(50.3)		
30-60minute	3949(30.2)	469(26.6)	2814(31.1)	666(29.6)		
>60minute	3728(28.5)	697(18.7)	2578(28.5)	453(20.1)		
Using electronic products	8422(64.5)	1254(71.2)	5898(65.2)	1270(56.4)	99.70	<0.001
Bedroom						
Electronic smoking	7163(55.4)	1015(58.2)	4954(55.2)	1194(53.6)	8.69	0.013
Home						
Doing strenuous exercise	8362(65.4)	1032(59.8)	5749(65.0)	1581(71.8)	64.95	<0.001
Spare time						
Exercise at moderate intensity	7057(55.8)	837(49.0)	4896(55.8)	1324(60.9)	55.43	<0.001
Spare time						
Days for consuming breakfast	6.13±1.56	5.52±1.99	6.17±1.48	6.40±1.36	163.74	<0.001
Days for consuming light snack	2.10±2.34	2.39±2.51	2.09±2.29	1.91±2.35	19.44	<0.001
Eating fried food in the past 12 months	7468(70.9)	950(70.2)	5279(71.6)	1239(68.7)	6.17	0.046
Eating pastry in the past 12 months	8342(80.6)	1035(78.5)	5880(81.1)	1427(80.3)	5.16	0.076
Supplementation	4982(51.3)	612(49.6)	3518(51.6)	852(51.7)	1.64	0.441
Supplements in the						

at 12 months							
Frequency of being lonely						316.78	<0.001
Never	4760(36.7)	443(25.3)	3255(36.2)	1062(47.7)			
Sometimes	6397(49.4)	888(50.8)	4584(51.0)	925(41.5)			
Always	1803(13.9)	417(23.9)	1145(12.7)	241(10.8)			
Frequency of being sad						245.29	<0.001
Never	4795(37.0)	489(28.0)	3291(36.7)	1015(45.5)			
Sometimes	6877(53.1)	945(54.1)	4919(54.8)	1013(45.5)			
Always	1283(9.9)	314(18.0)	768(8.6)	201(9.0)			
Frequency of being irritated						425.93	<0.001
Never	2084(16.3)	207(12.0)	1281(14.4)	596(27.1)			
Sometimes	8771(68.5)	1054(61.1)	6353(71.5)	1364(62.1)			
Always	1946(15.2)	463(26.9)	1246(14.0)	237(10.8)			

Table 3 Prevalence of overweight and obesity among different groups [n(%)]

Characteristics	Normal	Overweight/ obesity	Overweight	Obesity	Total	χ^2	P value
Gender						365.53	<0.001
Boys	4849(74.4)	1671(25.6)	1106(17.0)	565(8.7)	6520		
Girls	5704(87.2)	839(12.8)	473(7.2)	366(5.6)	6543		
Area						48.51	<0.001
Urban	4635(78.2)	1289(21.8)	789(13.3)	500(8.4)	5924		
Rural	5918(82.9)	1221(17.1)	790(11.1)	431(6.0)	7139		
Age(y)						94.38	<0.001
≤13	3866(78.3)	1071(21.7)	592(12.0)	479(9.7)	4937		
14-15	3133(81.3)	721(18.7)	470(12.2)	251(6.5)	3854		
≥16	3559(83.3)	713(16.7)	517(12.1)	196(4.6)	4272		
Types of School						1.14	0.566
Key school	6335(81.0))	1482(19.0)	939(12.0)	543(6.9)	7817		
Ordinary school	4218(80.4))	1028(19.6)	640(12.2)	388(7.4)	5246		
Total	10553(80.8)	2510(19.2)	1579(12.1)	931(7.1)	13063		

Table 4 Odds ratios of overall overweight/obese based on sleep duration

Characteristics	<6h	6-8h	8-10h
Total (n)	1762	9051	2250
Overweight/obese (n)	376	1669	465
Model 0	1.04(0.89-1.21)	0.87(0.77-1.97)	1.00
Model 1	1.32(1.12-1.55)*	1.04(0.92-1.17)	1.00
Model 2	1.55(1.27-1.89)**	1.06(0.92-1.23)	1.00
Model 3	1.53(1.24-1.87)*	1.06(0.91-1.23)	1.00

Model 0, no additional variables

Model 1, adjustment for gender, age, region, types of school.

Model 2, additionally adjusted for whether using electronic product daily during school, electronics in bedroom, someone smoking at home, strenuous physical activity, moderate physical activity, days of consuming breakfast and night snack per week, having fried food or pastry in the past 12 months, having taken nutrient supplements in the past 12 months.

Model 3, further adjusted for feeling annoy, sad or lonely

* $P < 0.05$; ** $P < 0.01$

Table 5 Odds ratios of overweight/obese based on sleep duration in different gender

Characteristics	<6 h	6-8 h	8-10 h
Female			
Total (n)	960	4670	913
Overweight/obese (n)	152	568	119
Model 0	1.25(0.97-1.63)	0.92(0.75-1.14)	1.00
Model 1	1.38(1.06-1.81)*	0.99(0.80-1.23)	1.00
Model 2	1.41(1.02-1.96)*	1.04(0.80-1.35)	1.00
Model 3	1.38(0.99-1.93)	1.04(0.79-1.35)	1.00
Male			
Total (n)	802	4381	1337
Overweight/obese (n)	224	1101	346
Model 0	1.11 (0.91-1.35)	0.96(0.84-1.11)	1.00
Model 1	1.25(1.02-1.54)*	1.07(0.92-1.23)	1.00
Model 2	1.65(1.27-2.13)**	1.06(0.89-1.27)	1.00
Model 3	1.63(1.25-2.13)**	1.06(0.88-1.27)	1.00

Model 0, no addition variables

Model 1, adjustment for age, region, types of school.

Model 2, additionally adjusted for whether using electronic product daily during school, electronics in bedroom, someone smoking at home, strenuous physical activity, moderate

physical activity, days of consuming breakfast and night snack per week, having fried food or pastry in the past 12 months, having taken nutrient supplements in the past 12 months. Model 3, further adjusted for feeling annoy, sad or lonely

* $P < 0.05$; ** $P < 0.01$