

# Quantitative Chest CT analysis in COVID-19 to predict the need for oxygenation support and intubation

EZIO LANZA (✉ [eziolanza@gmail.com](mailto:eziolanza@gmail.com))

Humanitas Clinical and Research Center - IRCCS <https://orcid.org/0000-0003-1889-8798>

**Riccardo Muglia**

Humanitas University

**Isabella Bolengo**

Humanitas University

**Orazio Giuseppe Santonocito**

Humanitas University

**Costanza Lisi**

Humanitas University

**Giovanni Angelotti**

Humanitas Clinical and Research Center - IRCCS

**Pierandrea Morandini**

Humanitas Clinical and Research Center - IRCCS

**Victor Savevski**

Humanitas Clinical and Research Center - IRCCS

**Letterio Salvatore Politi**

Humanitas Clinical and Research Center - IRCCS

**Luca Balzarini**

Humanitas Clinical and Research Center - IRCCS

---

## Research Article

**Keywords:** COVID-19, Tomography, Spiral Computed, intubation, Pulmonary Ventilation, 3D-slicer, compromised lung, coronavirus

**Posted Date:** May 22nd, 2020

**DOI:** <https://doi.org/10.21203/rs.3.rs-30481/v1>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

**Version of Record:** A version of this preprint was published at European Radiology on June 26th, 2020.  
See the published version at <https://doi.org/10.1007/s00330-020-07013-2>.

# Abstract

**OBJECTIVE:** Lombardy (Italy) was the epicentre of the COVID-19 pandemic in March 2020. The healthcare system suffered from a shortage of ICU beds and oxygenation support devices. In our Institution, most patients received chest CT at admission, only interpreted visually. Given the proven value of Quantitative CT analysis (QCT) in the setting of ARDS, we tested QCT as an outcome predictor for COVID-19.

**METHODS:** We performed a single centre retrospective study on COVID-19 patients hospitalized from January 25th, 2020 to April 28th 2020, who received CT at admission prompted by respiratory symptoms such as dyspnea or desaturation. QCT was performed using a semi-automated method (3D-Slicer). Lungs were divided by Hounsfield Unit intervals. Compromised lung (%CL) volume was the sum of poorly and non-aerated volumes (-500,100HU). We collected patient's clinical data including oxygenation support throughout hospitalization.

**RESULTS:** Two hundred twenty-two patients (163 males, median age 66, IQR 54-6) were included; 75% received oxygenation support (20% intubation rate). Compromised lung volume was the most accurate outcome predictor (logistic regression,  $p < 0.001$ ). %CL values in the 6-23% range increased risk of oxygenation support; values above 23% were at risk for intubation. %CL showed a negative correlation with PaO<sub>2</sub>/FiO<sub>2</sub> ratio ( $p < .001$ ) and was a risk factor for in-hospital mortality ( $p < .001$ )

**CONCLUSIONS:** QCT provides new metrics of COVID-19. The compromised lung volume is accurate in predicting the need for oxygenation support and intubation and is a significant risk factor for in-hospital death. QCT may serve as a tool for the triaging process of COVID-19.

## Introduction

### Key points

1. Quantitative computer-aided analysis of chest CT (QCT) provides new metrics of COVID-19
2. The compromised lung volume measured in the -500,100HU interval predicts oxygenation support, intubation and is a risk factor for in-hospital death
3. Compromised lung values in the 6-23% range prompt oxygenation therapy; values above 23% increase the need for intubation.
4. QCT may serve as a tool for the triaging process of COVID-19

In the early days of December 2019, the first pneumonia cases of a new Coronavirus named SARS-CoV-2 were identified in Wuhan, the capital city of Hubei province (China) [1]. As of April 30th, 2020, 1,112,667 cases have been reported in the EU/EEA and the UK: Spain (212,917) and Italy (203,591) had the largest contagion [2].

Since the initial spread of this new illness, known as Coronavirus Disease 2019 (COVID-19), many patients have been hospitalized with respiratory symptoms [3]. The clinical spectrum is broad, including asymptomatic infection, mild upper respiratory tract disease, and severe interstitial pneumonia with respiratory failure requiring oxygenation support or intubation [4, 5].

Computed Tomography (CT) is the most sensitive radiological technique for the diagnosis of COVID-19, showing diffuse lung alterations ranging from ground-glass opacities to parenchymal consolidations; several radiological patterns are observed at different times throughout the disease course [6, 7]. However, CT has limited specificity for distinguishing between virus etiologies [8]. Early diagnosis is hence mainly performed with nasopharyngeal swabs and virus RNA extraction with real-time reverse-transcriptase–polymerase-chain-reaction (rtRT-PCR) [9].

Recently, the visual assessment of lung damage on CT scans has been proven valuable in determining the prognostic implications [10]. Nonetheless, computer-aided quantitative analysis of the CT exam (Quantitative Computed Tomography [QCT]) can also be used for this purpose, as already demonstrated by the research on the Acute Respiratory Distress Syndrome (ARDS) [11, 12].

The main objective of our research was to explore the role of QCT analysis as an outcome predictor for COVID-19. To this end, we conducted a monocentric retrospective study on positive patients in a tertiary referral Institution in Milan (Lombardy, Italy) during the recent outbreak, where we experienced hospital overcrowding and shortage of intensive care unit (ICU) beds, mechanical ventilation devices and oxygen [13].

## Materials And Methods

### STUDY OVERSIGHT

The present is an observational retrospective study, conducted in a tertiary referral University Hospital in Milan (Lombardy, Italy). The study protocol followed the ethical guidelines established in the 1975 Declaration of Helsinki, compliant with the procedures of the local ethical committee and was approved by the Institutional Review Board. This study received no financial support.

### DATA SOURCES

We obtained laboratory, clinical, and radiological data of hospitalized patients affected by COVID-19 from the electronic medical records; the inclusion data cutoff for the analyses was January 25th, 2020, and April 28th, 2020.

COVID-19 was diagnosed based on a positive result on rtRT-PCR assay on nasal and pharyngeal swab specimens [14]. We included only laboratory-confirmed cases who received a non-contrast chest CT at admittance in the emergency department.

We analyzed patients' demographics, clinical and laboratory findings at admittance from electronic medical records. Patients demographics included age, sex, body mass index (BMI), concomitant/previous diseases, and smoking habit. Clinical and laboratory assessments consisted of body temperature, PaO<sub>2</sub>, PaCO<sub>2</sub>, and C-reactive protein. We also noted the time from symptoms onset, days of hospitalization, ICU admittance, medical therapy administered, and the most invasive level of oxygenation support provided. In particular, we distinguished between low-flow oxygenation (nasal cannula, face mask), high-flow oxygenation (Venturi mask, helmet CPAP), and mechanical ventilation through an endotracheal tube. We collected the time interval from CT and oxygenation support, as well as the first PaO<sub>2</sub>/FiO<sub>2</sub> ratio available. In-hospital deaths and healed patients' discharge dates were also noted.

The clinical features of confusion (mental test score of 8 or less), urea, respiratory rate, and blood pressure were also acquired to calculate the CURB-65, a validated score to predict the severity of Community-Acquired Pneumonia [15] that stratifies patients in groups from 1 to 3 according to the risk of mortality.

## CHEST CT AND QUANTITATIVE ANALYSIS

All patients received a standard non-contrast chest CT with a multidetector CT scanner (Philips Brilliance, Amsterdam, The Netherlands) with the following setup: collimation, 64x0.25; voltage, 120 kV; tube current, 130-200 mAs, 240 mA, pitch 1.4, slice thickness after reformat, 2.5 mm. The field of view included the whole chest and was acquired during forced inspiration, in keeping with patient compliance. The dataset was anonymized and exported to a dedicated segmentation suite for medical image computing (3D-Slicer, www.slicer.org) [16] equipped with a semi-automated segmentation algorithm (Chest Imaging Platform) [17]. This software, validated as useful in the surgical setting [18], performed a first-pass automated segmentation; then, lung volumes were manually perfected using three-dimensional tools such as spherical brushes or erasers.

As a rule, a complete segmentation included both lungs with interstitial structures, segmentary vessels, and bronchi; the main pulmonary arteries and bronchi, all mediastinal structures, and eventual pleural effusion were excluded, as well as lung masses (e.g. tumours, fungal disease).

Lung volumes, considered as percentages of the total volume, were extracted according to different Hounsfield Units (HU) intervals into Non-Aerated (%NNL, density between 100, -100 HU), Poorly Aerated (%PAL, -101,-500 HU), Normally Aerated (%NAL, -501,-900 HU), and Hyperinflated (-901,-1000 HU) [19]. The additional volume "compromised lung" (%CL) was considered as the sum of %PAL and %NNL (-500,100 HU) (Figure 1). The authors in charge of segmentation (E.L., C.L., R.M.) were unaware of the laboratory and clinical parameters or hospitalization outcomes. Conflicts were resolved in consensus. The principal investigator reviewed and confirmed all segmentations before data entry. The time needed to complete each analysis was recorded.

## OUTCOMES

The primary objective was to identify and validate the most accurate lung volume derived by QCT, to predict the two main study outcomes: the need for oxygenation support and the need for intubation in patients affected by COVID-19.

Other objectives included correlation with pulmonary dysfunction as measured by the PaO<sub>2</sub>/FiO<sub>2</sub> ratio and prediction of in-hospital death.

## **STATISTICAL ANALYSIS**

### **Development of prediction models**

All analyses were performed using Stata 13 (StataCorp LP, College Station, USA). Multiple binomial logistic regressions were performed to explore the correlation of the lung volumes, %NAL, %PAL, and %CL, over the two outcomes of interest. All clinically relevant predictors without missing data were included in the final model as covariates: age, sex, smoke habit, CPR, heart disease, chronic lung disease, cancer, diabetes, chronic kidney failure, urea levels, and CURB-65 group. Three similar models were thus developed: %NAL-model, %PAL-model, and %CL-model.

BMI was available for 161 patients and was tested in a separate model.

A Pearson's product-moment correlation was run to assess the relationship between the selected lung interval and PaO<sub>2</sub>/FiO<sub>2</sub> in 106 patients (nasal cannula = 26, Venturi mask = 28, helmet CPAP = 21, endotracheal tube = 27).

A Cox regression survival analysis was performed to explore potential predictors of mortality. All potential candidates were tested with univariate analyses; the cutoff for inclusion in the final model was set at  $p < 0.2$ .

### **Model validation**

All simulations were run using Python programming language (Python Software Foundation, <https://www.python.org/>). Categorical variables were preliminarily tested for correlation using Chi-squared tests; Wilcoxon rank-sum tests were performed on continuous covariates to inspect the probability of being sampled from the same distribution. Two separate multivariate regressions, without regularization, were performed on both outcomes over the space of covariates. Models' coefficients, confidence intervals, and their associated associations (p values) were investigated to assess whether the selected lung interval remained significant despite adjusting for possible confounders. Predictive machine learning models were built using logistic regression with regularization. To adjust for class imbalance, on both outcomes, and preserve the limited amount of available observations, the logistic regression was stacked upon a SMOTE model during training [20].

Hyperparameters were chosen by randomized selection over 1000 possible validations by means of 10-fold cross-validation each, for a total of 5000 actually trained models. The aim was to alleviate class

imbalance by maximization of class weighted F1-score, harmonic mean of precision and recall.

Cross-validated receiver operating characteristic (CV-ROC) and mean areas under the curve (CV-AUC) were calculated [21]. Different cross-validated cut-points at 90% sensitivity and at 90% specificity were estimated for both outcomes.

## Results

### DEMOGRAPHIC AND CLINICAL CHARACTERISTICS

A cohort of 222 patients (163 males, 59 females, median age 66, interquartile range [IQR] 54-6) was identified (Table 1). The median interval time between symptom onset at admission and CT was seven days (IQR, 3-8). The median body temperature during the CT exam was 37.7° C (IQR, 36.9°-38.4°), and the median PaO<sub>2</sub> was 67 mmHg (IQR, 57-90). The median hospitalization length was 11 days (IQR, 7-17, maximum 56).

During the hospital stay, 75% of patients required oxygenation support (Table 2) as follows: 29% low flow oxygenation (26% nasal cannula, 3% facial mask); 26% high flow oxygenation (16% Venturi mask, 10% helmet CPAP); and 20% mechanical ventilation with an endotracheal tube. Median PaO<sub>2</sub>/FiO<sub>2</sub> ratio, as recorded on the first day of oxygenation support, was 192.0 (IQR, 122.2-251.5). The median time interval from admission CT to oxygenation support was one day (IQR, 0-2).

During the observation, antiviral therapy with Lopinavir-Ritonavir was administered in 55% of patients and Darunavir plus Cobicistat in 36%. Hydroxychloroquine was used in 91%. All patients also received the best medical therapy tailored to the individual case, including broad-spectrum antibiotics.

Up to the reference date April 28th, 2020, out of total 222 patients, 8 (3.6%) were still hospitalized, whereas the remaining 214 (96.3%) were discharged, 150 (67.6%) of them healed and 64 (28.8%) died after a median nine days hospitalization (IQR, 5-14): their median age was 75.7 years (IQR, 71-80).

Regarding the CURB-65 score, 129 patients were categorized as low-risk (Group 1), 57 patients were medium-risk (Group2); no patients were scored as high-risk (Group 3). Death rates were 19% and 36% respectively.

### DIAGNOSTIC PERFORMANCE OF THE DIFFERENT LUNG VOLUMES

All CT scans were considered of diagnostic quality and successfully analyzed. All lung volumes were correctly extracted (Table 2). The median time for segmentation was 11 minutes (IQR, 7-14).

The need for oxygenation support was accurately predicted both by the %PAL-model (LR chi<sup>2</sup>(<sup>13</sup>), 86.05; p < 0.0001) and by the %CL-model (LR chi<sup>2</sup>(<sup>13</sup>), 86.90; p < 0.0001). In both, %PAL and %CL showed a strong predictive value (p < 0.0001) and were the only significant covariates. The %NAL-model showed worse performance (LR chi<sup>2</sup>(<sup>13</sup>), 39.18; p < 0.0002) and %NAL was not a significant predictor (p = 0.315).

The need for intubation was accurately predicted by all three models with minimal differences in LR chi2<sup>(13)</sup> (%NAL-model, 81.47; %PAL-model, 85.31, %CL-model, 89.88;  $p < 0.0001$  each) and all three lung volumes showed strong predictive value ( $p < 0.0001$ ). The compromised lung volume was considered as the best single predictor of both outcomes, and the %CL-model was chosen for validation.

A separate model was fitted adding BMI as a covariate in a subgroup of 161 patients, proving no effect on both outcomes (Oxygenation support,  $p = 0.130$ ; Intubation,  $p = 0.428$ ).

### **Validation of compromised lung volume**

For the outcome of oxygenation support, the %CL-model reached CV-AUROC 0.83 (weighted f1-score, 0.72; standard deviation [SD], 0.098). Cross-validated cut-points were identified: 6%CL at high sensitivity (90%; specificity, 43%); 13%CL at high specificity (91%; sensitivity, 53%).

For the intubation outcome, the %CL-model reached CV-AUROC 0.86 (weighted f1-score, 0.79; SD, 0.07). Cut-points were: 10%CL at high sensitivity (90%; specificity, 56%); 23%CL at high specificity (91%; sensitivity 69%).

### **COMPROMISED LUNG VOLUME AND SURVIVAL**

A moderate negative correlation between %CL and PaO<sub>2</sub>/FiO<sub>2</sub> ratio,  $r(104) = -0.39$ ,  $p < .001$  was observed, highlighting how pulmonary function worsened as %CL increased.

As a result of multivariate survival analysis, %CL was predictive of in-hospital mortality (hazard ratio [HR], 1.02; 95% CI, 1.02 to 1.05,  $p = .01$ ), together with age (HR, 1.06; 95%CI, 1.03-1.10,  $p < .0001$ ), cancer (HR, 3.27; 95% CI, 1.54-6.94), CRP (HR, 1.03, 95%CI 1.01-1.05), CKD (HR, 5.59; 95%CI, 1.97-15.86). Detailed results are reported in Table 3.

All other variables proved non-significant at univariate analyses and were excluded from the final model.

## **Discussion**

COVID-19 is a new disease outbreak reaching a pandemic level and a threat to global health [22]. Hospital overcrowding, shortness of ICU beds, oxygen, and ventilators have been a large-scale concern in Lombardy (Italy)[13]; thus accurate and rapid triaging is key to avoid a crisis of the health care network.

Chest x-ray has been proposed as a low-cost tool for detecting lung impairment in patients with suspected COVID-19[23]. However, its low sensitivity [24] makes it more appropriate for follow-up rather than early diagnosis [25]. Chest CT is instead pivotal to the early diagnosis of COVID-19 due to the ability to detect all disease characteristics [26], and its use has been recommended by the European Society of Radiology in selected patients [27], using a dedicated scanner whenever possible [28]. However, its value is currently limited to visual findings [25], whereas QCT proved to be the tool that allowed for considerable advancements in understanding the ARDS pathophysiology and establishing adequate oxygenation

support[29]. These observations led us to perform QCT analysis in all chest CT scans performed at admission on confirmed COVID-19 cases. We identified the volume of compromised lung included in the -500,100 HU interval as a predictor of oxygenation support and invasive ventilation; this was also significantly correlated with pulmonary dysfunction as measured by the PaO<sub>2</sub>/FiO<sub>2</sub> ratio and represented a risk factor for in-hospital mortality.

Typical CT appearance of COVID-19 was well described by Chung et al. in February 2020 [7] and has been confirmed by several reports [6, 7]. Based on these findings, Yuan et al. have proposed a scoring method to screen patients based on the admittance CT scan [10]. More recently, Li et al. also described a visual, quantitative analysis of lung damage, based on a “total severity score” to the degrees of parenchymal loss, correlated with a score of clinical severity [30]. However, these visual characterizations are subjective and unsuited to a systematic disease evaluation and data sharing. Conversely, we reported on a QCT method based on 3D-Slicer, which is fast, standardized, and ensures a consistently repeatable evaluation of parenchymal impairment. Some advantages are a) is a free, open-source software untied to any workstation b) has a low learning curve [16] c) documentation and internet support are easily found d) allows for rapid deployment in the ever-changing epicentres of the Coronavirus pandemic.

Colombi et al. [31] have used a similar approach to predict the outcomes of COVID-19. They reported good performance of the well-aerated volume (%WAL, -950,-700 HU) in predicting the combined outcome of ICU admission and death. Such an outcome, however, would not be informative in our cohort. In fact, 21/48 (66%) of our deaths happened outside the ICU, and 23/48 (48%) of patients successfully healed after ICU admittance, remarking that ICU admission should not be generalized as a “worse” outcome. Moreover, they reported the negative predictive value of a BMI-surrogate, the adipose tissue measured at the T7-T8 level, whereas in our study BMI was not predictive of the need for oxygenation support or intubation.

Compromised lung was significantly correlated with the PaO<sub>2</sub>/FiO<sub>2</sub> ratio and increasing pulmonary dysfunction. One reason may be that hypoxemia refractory to oxygenation support is mainly due to intrapulmonary shunt [32] that happens in poorly and non-aerated lung areas, which are well quantified in the %CL. On the contrary, aerated lung volumes such as %NAL or %WAL represent an indirect quantification of this phenomenon and may be more prone to variation during respiratory movements, or ventilation; this may also explain the poor performance of %NAL-model in our analysis.

Another approach to quantitative analysis has been proposed by Huang et al. [33], who have successfully deployed a commercial deep learning algorithm to quantify lung impairment. However, the prediction of oxygenation support was not among their reported outcomes. Despite the promising power of such an approach, the same authors also report that in 8.7% of cases, the algorithm was unable to correctly identify the lung borders without the help of the radiologist. This need is also shared by our semi-automated approach.

Regardless of the technique used, QCT is seen as the ideal tool that could be able to predict which patient will need a ventilator soon or who no longer requires one. This might help especially in situations with

limited hospital resources; in fact, in the process of triaging, the highest priority is to be accorded to those patients whose prognosis is good with intensive care, but poor without it [34]. QCT, particularly the compromised lung volume, may hence be used as a new decision tool, considering its strong correlation not only with respiratory needs but also with in-hospital survival. The need for new metrics specific to COVID-19 is remarked by the low performance of a validated score for pneumonia (CURB-65) that was tested for comparison; most patients (58%) were graded as low-mortality risk (Group 1); still, we observed a 15% death rate, much higher than the presumed 1.5% [15].

This study has several limitations. Firstly, it has a retrospective design. Secondly, we did not perform a repeat CT on the healed patients before discharge, to check whether the physiological lung volumes had been restored; this could be the subject of a future study.

Regarding the segmentation process, we included the interstitial structures in the analyzed volume (e.g. segmental arteries and bronchi). By partially falling in the same threshold as the non-aerated lung, these may have reduced precision. However, the event might have been irrelevant to the outcome. Firstly, because the poorly aerated volume alone was a strong outcome predictor; secondly, because the same method was applied to all patients thus counterbalancing the potential inaccuracy; thirdly, the added volume of the lung interstitium may be considered negligible compared to the compromised lung volume in the setting of severe pneumonia. On the contrary, this choice allowed for a fast segmentation process.

Regarding respiratory support, 20% of our patients required invasive mechanical ventilation, a higher number compared to 2.5% reported by Wu et al. and 17% by Wang and Zhou et al. These observations may be partially due to the higher age of the Italian population [35]. Indeed, the median age of our cohort (66 years) was above the ones previously reported in China by Wu et al. [36] (51 years), Wang et al. [37] and Fei Zhou et al. [4] (56 years).

The main strengths of our study are the high number of patients enrolled, the completeness of clinical data, the high statistical significance of all tests conducted, and the use of a rapidly reproducible and scalable method for QCT.

In conclusion, lung QCT provides new metrics of COVID-19 and has a promising role in predicting its clinical outcome. The percentage of lung parenchyma in the -500,100 HU interval, namely the compromised lung volume (%CL), has shown high accuracy in predicting the need for oxygenation support and mechanical ventilation and is a risk factor for in-hospital death.

We empirically identified different cut-points of compromised lung volume: patients presenting with %CL values in the 6-23% range are at risk for needing oxygenation therapy; values above 23% are at risk for intubation. These results strengthen the evidence of QCT as an ideal tool for easing the triaging process of COVID-19.

## List Of Abbreviations

COVID-19: Coronavirus Disease 2019

CT: Computed Tomography

rtRT-PCR: real-time reverse-transcriptase–polymerase-chain-reaction

QCT: Quantitative Computed Tomography

ARDS: Acute Respiratory Distress Syndrome

BMI: body mass index

HU: Hounsfield Units

IQR: interquartile range

%NNL: non-aerated lung

%PAL: poorly aerated lung

%NAL: normally aerated lung

%CL: compromised lung

## Declarations

I declare no conflicts of interests regarding this manuscript , on behalf of all authors.

The ethical committee of Humanitas Clinical and Research Center - IRCCS approved of this retrospective review.

Patients' consent was waived by the IRB due to the retrospective design of this study

## References

1. Guan W-J, Ni Z-Y, Hu Y, et al (2020) Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med*. doi: 10.1056/NEJMoa2002032
2. Situation update for the EU/EEA and the UK, as of 27 March 2020. <https://www.ecdc.europa.eu/en/cases-2019-ncov-eueea>. Accessed 28 Mar 2020
3. McMichael TM, Currie DW, Clark S, et al (2020) Epidemiology of Covid-19 in a Long-Term Care Facility in King County, Washington. *N Engl J Med*. doi: 10.1056/NEJMoa2005412
4. Zhou F, Yu T, Du R, et al (2020) Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 395:1054–1062

5. Poston JT, Patel BK, Davis AM (2020) Management of Critically Ill Adults With COVID-19. *JAMA*. doi: 10.1001/jama.2020.4914
6. Shi H, Han X, Jiang N, Cao Y, Alwalid O, Gu J, Fan Y, Zheng C (2020) Radiological findings from 81 patients with COVID-19 pneumonia in Wuhan, China: a descriptive study. *Lancet Infect Dis* 20:425–434
7. Chung M, Bernheim A, Mei X, et al (2020) CT Imaging Features of 2019 Novel Coronavirus (2019-nCoV). *Radiology* 295:202–207
8. Li Y, Xia L (2020) Coronavirus Disease 2019 (COVID-19): Role of Chest CT in Diagnosis and Management. *AJR Am J Roentgenol* 1–7
9. Lescure F-X, Bouadma L, Nguyen D, et al (2020) Clinical and virological data of the first cases of COVID-19 in Europe: a case series. *Lancet Infect Dis*. doi: 10.1016/S1473-3099(20)30200-0
10. Yuan M, Yin W, Tao Z, Tan W, Hu Y (2020) Association of radiologic findings with mortality of patients infected with 2019 novel coronavirus in Wuhan, China. *PLoS ONE* 15:e0230548
11. Nishiyama A, Kawata N, Yokota H, Sugiura T, Matsumura Y, Higashide T, Horikoshi T, Oda S, Tatsumi K, Uno T (2020) A predictive factor for patients with acute respiratory distress syndrome: CT lung volumetry of the well-aerated region as an automated method. *Eur J Radiol* 122:108748
12. Ichikado K, Suga M, Muranaka H, et al (2006) Prediction of prognosis for acute respiratory distress syndrome with thin-section CT: validation in 44 cases. *Radiology* 238:321–329
13. Nacoti M, Ciocca A, Brambillasca P, et al (2020) At the Epicenter of the Covid-19 Pandemic and Humanitarian Crises in Italy: Changing Perspectives on Preparation and Mitigation | Catalyst non-issue content. *NEJM Catalyst Innovations in Care Delivery*
14. Zhu N, Zhang D, Wang W, et al (2020) A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N Engl J Med* 382:727–733
15. Lim WS, van der Eerden MM, Laing R, Boersma WG, Karalus N, Town GI, Lewis SA, Macfarlane JT (2003) Defining community acquired pneumonia severity on presentation to hospital: an international derivation and validation study. *Thorax* 58:377–382
16. Fedorov A, Beichel R, Kalpathy-Cramer J, et al (2012) 3D Slicer as an image computing platform for the Quantitative Imaging Network. *Magn Reson Imaging* 30:1323–1341
17. Yip SSF, Parmar C, Blezek D, Estepar RSJ, Pieper S, Kim J, Aerts HJWL (2017) Application of the 3D slicer chest imaging platform segmentation algorithm for large lung nodule delineation. *PLoS ONE* 12:e0178944
18. Donadon M, Lanza E, Branciforte B, Muglia R, Lisi C, Pedicini V, Poretti D, Famularo S, Balzarini L, Torzilli G Hepatic Uptake Index in the hepatobiliary phase of Gd-EOB-DTPA-enhanced magnetic resonance imaging estimates functional liver reserve and predicts post-hepatectomy liver failure. *Surgery*
19. Cressoni M, Gallazzi E, Chiurazzi C, et al (2013) Limits of normality of quantitative thoracic CT analysis. *Crit Care* 17:R93

20. Blagus R, Lusa L (2013) SMOTE for high-dimensional class-imbalanced data. *BMC Bioinformatics* 14:106
21. Luque-Fernandez MA, Redondo-Sánchez D, Maringe C (2019) cvauroc: Command to compute cross-validated area under the curve for ROC analysis after predictive modeling for binary outcomes. *The Stata Journal* 19:615–625
22. Madjid M, Safavi-Naeini P, Solomon SD, Vardeny O (2020) Potential effects of coronaviruses on the cardiovascular system: A review. *JAMA Cardiol.* doi: 10.1001/jamacardio.2020.1286
23. Bandirali M, Sconfienza LM, Serra R, Brembilla R, Albano D, Ernesto PF, Messina C (2020) Chest X-ray findings in asymptomatic and minimally symptomatic quarantined patients in Codogno, Italy. *Radiology* 201102
24. Wong HYF, Lam HYS, Fong AH-T, et al (2019) Frequency and Distribution of Chest Radiographic Findings in COVID-19 Positive Patients. *Radiology* 201160
25. Ng M-Y, Lee EY, Yang J, et al (2020) Imaging Profile of the COVID-19 Infection: Radiologic Findings and Literature Review. *Radiology: Cardiothoracic Imaging* 2:e200034
26. Ai T, Yang Z, Hou H, Zhan C, Chen C, Lv W, Tao Q, Sun Z, Xia L (2020) Correlation of Chest CT and RT-PCR Testing in Coronavirus Disease 2019 (COVID-19) in China: A Report of 1014 Cases. *Radiology* 200642
27. Revel M-P, Parkar AP, Prosch H, Silva M, Sverzellati N, Gleeson F, Brady A (2020) COVID-19 patients and the Radiology department – advice from the European Society of Radiology (ESR) and the European Society of Thoracic Imaging (ESTI). *Eur Radiol*
28. Politi LS, Balzarini L (2020) The Radiology Department during the COVID-19 pandemic: a challenging, radical change. *Eur Radiol.* doi: 10.1007/s00330-020-06871-0
29. Gattinoni L, Cressoni M (2010) Quantitative CT in ARDS: towards a clinical tool? *Intensive Care Med* 36:1803–1804
30. Li K, Fang Y, Li W, Pan C, Qin P, Zhong Y, Liu X, Huang M, Liao Y, Li S (2020) CT image visual quantitative evaluation and clinical classification of coronavirus disease (COVID-19). *Eur Radiol.* doi: 10.1007/s00330-020-06817-6
31. Colombi D, Bordini FC, Petrini M, Maffi G, Morelli N, Milanese G, Silva M, Sverzellati N, Michieletti E (2020) Well-aerated Lung on Admitting Chest CT to Predict Adverse Outcome in COVID-19 Pneumonia. *Radiology* 201433
32. Radermacher P, Maggiore SM, Mercat A (2017) Fifty years of research in ARDS. gas exchange in acute respiratory distress syndrome. *Am J Respir Crit Care Med* 196:964–984
33. Huang L, Han R, Ai T, Yu P, Kang H, Tao Q, Xia L (2020) Serial Quantitative Chest CT Assessment of COVID-19: Deep-Learning Approach. *Radiology: Cardiothoracic Imaging* 2:e200075
34. Swiss Academy Of Medical Sciences (2020) COVID-19 pandemic: triage for intensive-care treatment under resource scarcity. *Swiss Med Wkly* 150:w20229

35. (2014) Elderly population. Organisation for Economic Co-Operation and Development (OECD). doi: 10.1787/8d805ea1-en
36. Wu C, Chen X, Cai Y, et al (2020) Risk factors associated with acute respiratory distress syndrome and death in patients with coronavirus disease 2019 pneumonia in wuhan, china. JAMA Intern Med. doi: 10.1001/jamainternmed.2020.0994
37. Wang D, Hu B, Hu C, et al (2020) Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. JAMA. doi: 10.1001/jama.2020.1585

## Tables

<b>Table 1. Patients' Characteristics</b>	
<b>Characteristic</b>	<b>Median (IQR)</b>
Age	66.4 years (53.8-75.8)
BMI	27.2 (24.0-30.1)
Symptoms onset	7 days (3-8.5)
Body Temperature	37.7°C (36.9-38.4)
pO <sub>2</sub>	65 mmHg (53.5-76)
pCO <sub>2</sub>	34 mmHg (30.5-38)
CRP	8.1 mg/L (2.7-17.9)
<b>Total (N = 222)</b>	
Male sex	163 (73%)
Smoke habit	25 (11.2%)
Cancer	18 (8.1%)
Lung disease	22 (9.9%)
Heart Disease	27 (12.16%)
Diabetes	38 (17.1%)
CKD	10 (4.5%)
<b>Medical therapy</b>	
Lopinavir-Ritonavir	122 (54.9%)
Darunavir plus Cobicistat	80 (36.0%)
Hydroxychloroquine	203 (91.4%)

CRP = C-reactive protein; CKD = Chronic kidney disease.

**Table 2. Details of oxygenation support and results of quantitative lung CT analysis****Oxygen therapy**

	No oxygenation support	Low-flow O <sub>2</sub>	High-flow O <sub>2</sub>	Intubated	Overall
Patients	56 (25%)	63 (29%)	58 (26%)	45 (20%)	222
PaO <sub>2</sub> /FIO <sub>2</sub>	..	244.4 (207.4-320)	171.43 (122.22-229.63)	128.6 (95.4-211.1)	192.0 (122.22-251.5)
Death rate	2 (1%)	15 (7%)	26 (12%)	21 (9%)	64 (29%)
Healing rate	54 (24%)	46 (21%)	30 (13%)	20 (9%)	150 (68%)

**Lung segmentation***median lung volume % (IQR)*

	No oxygenation support	Low-flow O <sub>2</sub>	High-flow O <sub>2</sub>	Intubated	Overall
Hyperinflated	15 (4.5-22)	8 (2-18)	8 (3-12)	3 (1-7)	6 (2-16)
Normal	78.5 (71-84)	77 (72-83)	77 (70-83)	61 (45-72)	76 (67-83)
Poorly aerated	5 (4-6)	8 (5-13)	11 (6-15)	22 (13-33)	9 (5-15)
Non aerated	1 (1-2)	2 (1-4)	2 (1-3)	6 (3-16)	2 (1-4)
Compromised	6 (5-9)	11 (7-16)	13.5 (7-17)	32 (15-50)	12 (7-20)
Total (cm <sup>3</sup> )	4726.3 (3835.8-5590.9)	4115.22 ( 3246.3-4915.2)	3946.51 (2826.1-4663.7)	3332.7 (2458.1-4222.6)	4057.4 (3132.3-4916.1)

**Predictive value**


% Lung volume	Any O <sub>2</sub>	Low-flow O <sub>2</sub>	High-flow O <sub>2</sub>	Intubation
Poorly aerated [median (IQR)]	9 (6-13)	8 (5-13)	11 (7-15)	22 (13-33)
P value	<0.001	0.08	0.4	<0.001
Sensitivity, Specificity	90.0%, 51.1%	..	..	52.8%, 97.1%
Accuracy	80.0%	..	..	88%
Compromised Lung [median (IQR)]	14.5 (9-25)	11 (7-16)	14.5 (10-19)	32 (15-50)
P value	<0.001	0.13	0.14	<0.001
Sensitivity, Specificity	90.0%, 51.1%	..	..	55.6%, 97.8%
Accuracy	80.0%	..	..	89.1%

**Table 3. Multivariate analyses of risk factors for oxygenation support, intubation, and in-hospital death.**

Risk factor	Oxygenation support				Intubation			Death				
	OR	95% CI	P-value	OR	95% CI	P-value	OR	P-value	95% C.I.			
Compromised lung volume	1.28	1.16	1.41	<0.001*	1.12	1.07	1.17	<0.001	1.03	0.001*	1.01	1.04
Age	1.04	1.00	1.08	0.052	0.97	0.92	1.01	0.154	1.07	<0.001*	1.03	1.10
Sex	1.59	0.63	4.04	0.329	1.04	0.33	3.24	0.952	0.93	0.836	0.47	1.83
Smoke habit	1.03	0.17	6.12	0.974	1.08	0.15	7.57	0.939	1.77	0.129	0.85	3.68
CRP	1.02	0.98	1.05	0.321	1.01	0.98	1.05	0.506	1.03	0.001*	1.01	1.05
Heart disease	2.23	0.45	11.05	0.326	0.18	0.02	1.79	0.143	1.19	0.642	0.58	2.44
Lung disease	6.87	0.70	67.95	0.099	0.65	0.07	6.14	0.703	1.05	0.904	0.47	2.33
Cancer	0.87	0.21	3.64	0.848	0.73	0.11	4.71	0.739	3.42	0.003*	1.52	7.73
Diabetes	0.88	0.31	2.52	0.809	0.87	0.23	3.23	0.833	1.88	0.052	0.99	3.55
CKD	1.53	0.13	18.18	0.738	0.69	0.07	7.01	0.752	4.14	0.003*	1.60	10.70
CURB-65 <sup>a</sup> 1	0.58	0.06	5.25	0.630	1.31	0.18	9.40	0.788	0.81	0.685	0.30	2.23
CURB-65 <sup>a</sup> 2	0.32	0.05	2.04	0.228	1.80	0.40	8.09	0.443	0.80	0.556	0.38	1.67
Urea at admission	1.60	0.49	5.21	0.437	4.93	1.39	17.47	0.014*	1.71	0.179	0.78	3.75
BMI (n=161)	1.08	0.98	1.20	0.130	0.95	0.84	1.08	0.428	..	..	..	..


OR = Odds Ratio; CI = Confidence Interval; CRP = C-reactive protein; CKD = Chronic kidney disease. <sup>a</sup> Risk group according to the CURB-65 scoring system.

## Figures




**Figure 1**

Quantitative Lung CT analysis of an 81 years old male patient affected by COVID-19. a) Non-contrast chest CT at admission showing bilateral ground-glass opacities, common findings of the novel coronavirus pneumonia b) Semi-automated segmentation using 3D-Slicer. Blue areas are normal lung parenchyma; yellow areas represent poorly aerated lung in the -500,-100 HU interval c) 3D volumetric representation of both lungs d) Comparison between normal and compromised lung volumes. This patient had 6% of compromised lung volume, required no oxygenation support and was discharged after 15 days of observation and supportive therapy.



**Figure 2**



Quantitative Lung CT analysis of a 35 years old male patient affected by COVID-19. a) Non-contrast chest CT at admission showing bilateral ground-glass opacities, interstitial thickening and consolidation in the posterior lung zones. b) Semi-automated segmentation using 3D-Slicer. Blue areas are normal lung parenchyma; yellow areas represent poorly aerated lung in the -500,-100 HU interval; red areas represent non-aerated lung and interstitium, in the -100,100 HU interval c) 3D volumetric representation of both lungs showing multiple red areas in keeping with moderate lung impairment d) Comparison between normal and compromised lung volumes. This patient had 18% of compromised lung volume, required high-flow oxygenation support through a Venturi Mask. He was discharged after 17 days of sub-intensive care.





**Figure 3**

Quantitative Lung CT analysis of a 43 years old male patient affected by COVID-19. a) Non-contrast chest CT showing extensive areas of bilateral lung consolidation, multiple ground-glass opacities and interstitial thickening and consolidation b) Semi-automated segmentation using 3D-Slicer. Blue areas are normal lung parenchyma; yellow areas represent poorly aerated lung in the -500,-100 HU interval; red areas represent non-aerated lung and interstitium, in the -100,100 HU interval c) 3D volumetric representation of both lungs showing extensive red areas of consolidation in keeping with severe pneumonia d) Comparison between normal and compromised lung volumes. This patient had 50% of compromised lung volume and required immediate intubation and mechanical ventilation. He died after 13 days of intensive care, due to multi-organ failure.

### Compromised lung volume as a predictor of oxygenation support



### Compromised lung volume as a predictor of intubation



**Figure 4**

Ten-fold Cross-Validation for Receiver Operating Characteristic (A) and Precision-Recall curves (B) showing performance of compromised 3 lung volume as a predictor of oxygenation therapy and of intubation (C and D), based on quantitative analysis of chest CT at hospital admittance.