

# Hilbert's First Problem and the New Progress of Infinity Theory

Xijia Wang (✉ [xijiawang@hunnu.edu.cn](mailto:xijiawang@hunnu.edu.cn))

Hunan Normal University <https://orcid.org/0000-0002-0302-7589>

---

## Article

**Keywords:** Hilbert's first problem, Continuum hypothesis, Grossone method, Cosmic continuum, Infinity theory

**Posted Date:** August 6th, 2021

**DOI:** <https://doi.org/10.21203/rs.3.rs-306991/v3>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# Hilbert's First Problem and the New Progress of Infinity Theory

**Xijia Wang**

College of Mathematics and Statistics, Hunan Normal University, Changsha, China, 410081

Email: [xijiawang@hunnu.edu.cn](mailto:xijiawang@hunnu.edu.cn)

ORCID: [0000-0002-0302-7589](https://orcid.org/0000-0002-0302-7589)

**Abstract:** In the 19th century, Cantor created the infinite cardinal number theory based on the "1-1 correspondence" principle. The continuum hypothesis is proposed under this theoretical framework. In 1900, Hilbert made it the first problem in his famous speech on mathematical problems, which shows the importance of this question. We know that the infinitesimal problem triggered the second mathematical crisis in the 17-18th centuries. The Infinity problem is no less important than the infinitesimal problem. In the 21st century, Sergeyev introduced the Grossone method from the principle of "whole is greater than part", and created another ruler for measuring infinite sets. At the same time, the development of the infinity theory provides new ideas for solving Hilbert's first problem, and provides a new mathematical foundation for Cosmic Continuum Theory.

**Key words:** Hilbert's first problem; Continuum hypothesis; Grossone method; Cosmic continuum; Infinity theory

## 1. Introduction

In 1874, Cantor introduced the concept of cardinal numbers based on the "1-1 correspondence" principle. Cantor proved that the cardinal number of the continuum,  $C$ , is equal to the cardinal number of the power set of the natural number set,  $2^{\aleph_0}$ , where  $\aleph_0$  is the cardinal number of the natural number set. Cantor arranges the cardinal number of infinities from small to large as  $\aleph_0, \aleph_1, \dots, \aleph_a, \dots$ . Among them,  $a$  is an arbitrary ordinal number, which means that the cardinal number of the natural number set,  $\aleph_0$ , is the smallest infinity cardinal number. Cantor conjectured:  $2^{\aleph_0} = \aleph_1$ . This is the famous Continuum hypothesis (CH). For any ordinal  $a$ ,  $2^{\aleph_a} = \aleph_{a+1}$  holds, it is called the Generalized continuum hypothesis (GCH) [1].

In 1938 Gödel proved that the CH is not contradictory to the ZFC axiom system. In 1963, Cohen proved that the CH and the ZFC axiom system are independent of each other. Therefore, the CH cannot be proved in the ZFC axiom system [2]-[3].

However, people always have doubts about infinity theory. For example, in the study of Cosmic Continuum, the existing infinity theory shows great limitations[4]-[14].

In the 21st century, Sergeyev started from "the whole is greater than the part" and introduced a new method of counting infinity and infinitesimals, called the Grossone method. The introduced methodology (that is not related to non-standard analysis) gives the possibility to use the same numeral

38 system for measuring infinite sets, working with divergent series, probability, fractals, optimization  
39 problems, numerical differentiation, ODEs, etc.[15]-[43]

40 The Grossone method introduced by Sergeev takes the number of elements in the natural number  
41 set as a total number, marked as  $\textcircled{1}$ , as the basic numeral symbol for expressing infinity and  
42 infinitesimal, in order to more accurately describe infinity and infinitesimal.

43 The Grossone method was originally proposed as a Computational Mathematics, but its significance  
44 has far exceeded the category of Computational Mathematics. In particular, the Grossone method  
45 provides a new mathematical tool for the Cosmic Continuum Theory. A new infinity theory is about to  
46 emerge.

## 47 **2. The traditional infinity paradox and the fourth mathematics crisis**

48 In the history of mathematics, there have been three mathematics crises, each of which involves the  
49 foundation of mathematics. The first time was the discovery of irrational numbers, the second time was  
50 the infinitesimal problem, and the third time was the set theory paradox[44]-[45]. However, no one  
51 dare to say that the building of the mathematical theory system has been completed, and maybe the  
52 fourth mathematical crisis will appear someday.

53 In fact, the fourth mathematics crisis is already on the way. This is the infinity problem. In 1900,  
54 Hilbert put the Cantor continuum hypothesis as the first question in his famous lecture on 23  
55 mathematics problems [46]. This will never be an impromptu work by an almighty mathematician.

56 The infinitesimal question unfold around whether the infinitesimal is zero or not. From the 1920s to  
57 the 1970s, this problem has been initially solved through the efforts of generations of mathematicians.  
58 However, there are still different opinions about the second mathematics crisis. I believe that the  
59 infinitesimal problem has not been completely solved, otherwise there would be no infinity problem.  
60 Because the infinity problem and the infinitesimal problem are actually two aspects of the same  
61 problem.

62 Let us first look at what is problem with infinity.

63 The first is the expression of infinity. Now, there are two ways to express the infinity, one is to  
64 express with infinity symbol  $\infty$ , and the other is to express with infinity cardinal number. However,  
65 neither the infinity symbol  $\infty$  nor the infinity cardinal number can effectively express infinity and  
66 infinitesimal.

67 For example: when expressed in the infinity symbol  $\infty$ , we cannot distinguish the size of the  
68 natural number set and the real number set, nor can we distinguish the size of the natural number set  
69 and the integer set, they are all  $\infty$ . When expressed in infinity cardinal number, we can distinguish the  
70 size of the natural number set and the real number set, because the cardinal number of the natural  
71 number set is  $\aleph_0$ , and the cardinal number of the real number set is  $C = 2^{\aleph_0}$ ; but it is still  
72 impossible to distinguish the size of the natural number set and the integer set, they are both  $\aleph_0$ .

73 The second is the calculation of infinity. Whether it is the infinity symbol  $\infty$  or the infinity cardinal  
74 number, it cannot play a mathematically precise role in calculations. E.g:

75 
$$\infty + 1 = \infty, \quad \infty - 1 = \infty, \quad \infty \times \infty = \infty, \quad \infty^\infty = \infty.$$

76 And  $\frac{\infty}{\infty}$ ,  $\infty - \infty$ , etc. have no meaning at all.

77 Relative to infinity symbol  $\infty$ , Cantor's infinite cardinal number is an improvement, but the  
 78 cardinal number method of infinity can only be calculated qualitatively. The theory of infinity cardinal  
 79 number is based on the principle of "1-1 correspondence". Although according to the principle of  
 80 "power set is greater than the original set", infinite cardinal number can be compared in size, but it is  
 81 only the size of classes of infinity, not the size of infinity individuals.

82 For example, according to the continuum hypothesis, the following equation holds:

$$83 \quad \aleph_0 + 1 = \aleph_0, \quad \aleph_0 + \aleph_0 = \aleph_0, \quad \aleph_0 + 2^{\aleph_0} = 2^{\aleph_0}, \quad 2^{\aleph_0} + 2^{\aleph_0} = 2^{\aleph_0}.$$

84 This obviously violates the calculation rules of finite numbers and does not meet the uniformity  
 85 requirements of mathematical theory.

86 The reason for the infinity paradox in mathematical expressions and mathematical calculations is  
 87 that the existing infinity theory does not need to follow the principle of "the whole is greater than the  
 88 part", and this principle needs to be followed in the finite number theory. In this way, there is a problem  
 89 of using different calculation rules in the same calculation formula.

90 Since there is an infinite problem, how can there be no infinitesimal problem?

91 For example: because the infinity and the infinitesimals are reciprocal of each other (when the  
 92 infinitesimal is not zero), the following equation holds:

$$93 \quad \frac{1}{\infty + 1} = \frac{1}{\infty}, \quad \frac{1}{\infty - 1} = \frac{1}{\infty}, \quad \frac{1}{\infty \times \infty} = \frac{1}{\infty}, \quad \frac{1}{\infty^\infty} = \frac{1}{\infty};$$

$$94 \quad \frac{1}{\aleph_0 + 1} = \frac{1}{\aleph_0}, \quad \frac{1}{\aleph_0 + \aleph_0} = \frac{1}{\aleph_0}, \quad \frac{1}{\aleph_0 + 2^{\aleph_0}} = \frac{1}{2^{\aleph_0}}, \quad \frac{1}{2^{\aleph_0} + 2^{\aleph_0}} = \frac{1}{2^{\aleph_0}}.$$

95 Obviously, in these equations, although mathematical calculations can also be performed, the  
 96 mathematical accuracy is lost. At the same time, treating zero as a special infinitesimal is inconsistent  
 97 with the concept of infinitesimals. Because in modern mathematics, the infinitesimal is not a number  
 98 but a variable, and zero is a specific number, which is inconsistent with the definition of infinitesimal.

99 It can be seen that the problem of infinity involves many basic mathematics problems, and the  
 100 mathematics crisis caused by it is no less than the previous three mathematics crises. No wonder  
 101 Hilbert listed the continuum problem as the top of the 23 mathematical problems.

### 102 3. Grossone method and quantitative calculation of infinity

103 Sergeyev used Grossone  $\textcircled{1}$  to represent the number of elements in set of natural numbers, which is  
 104 similar to Kantor's cardinal number method. Kantor's cardinal number and Sergeyev's Grossone  $\textcircled{1}$   
 105 are superficially the same thing. Both represent the size of the set of natural numbers, but they are two  
 106 completely different concepts.

107 The cardinal number represents the size of a type of set that satisfies the principle of "1-1  
 108 correspondence". For a finite set, the cardinal number is the "number" of elements, but for an infinite  
 109 set, the cardinal number is not the "number" of elements. Is the size of a class of infinite sets that are  
 110 equivalent to each other. And Grossone  $\textcircled{1}$  represents the "number" of elements in a natural number  
 111 set, just like any finite set. Using this as a ruler, you can measure every infinity and infinitesimal.

112 In Grossone theory, infinity and infinitesimal are not variables, but definite quantities. Infinity and  
 113 infinitesimal are the reciprocal of each other. For example, the number of elements  $\textcircled{1}$  of the natural

114 number set is an infinity, and its reciprocal  $\frac{1}{\textcircled{1}}$  is an infinitesimal. Obviously, zero is not an

115 infinitesimal.

116 Let us see how numbers are expressed. The decimal numeral we generally use now are:  
117 1,2,3,4,5,6,7,8,9,0. Among these 10 numeral, the largest numeral is 9, but we can use them to express  
118 all finite numbers, whether it is ten thousand digits, billion digits, or larger numbers.

119 As the number of elements in the natural number set, Grossone, together with 1,2,3,4,5,6,7,8,9,0, can  
120 express any finite number and infinity.

121 For example, according to the principle of "whole is greater than part", we can get:

122  $\textcircled{1} + 1 > \textcircled{1}$ ,  $\textcircled{1} + \textcircled{1} = 2\textcircled{1}$ ,  $\textcircled{1} + 2^{\textcircled{1}} > 2^{\textcircled{1}}$ ,  $2^{\textcircled{1}} + 2^{\textcircled{1}} = 2 \times 2^{\textcircled{1}}$

123 The Grossone method can not only accurately express infinity, but also accurately express  
124 infinitesimal. E.g:

125  $\frac{1}{2\textcircled{1}}$ ,  $\frac{2}{3\textcircled{1}^2}$ ,  $\frac{3}{2^{\textcircled{1}}}$

126 For example, infinity can be operated like a finite number:

127  $0 \cdot \textcircled{1} = \textcircled{1} \cdot 0 = 0$ ,  $\textcircled{1} - \textcircled{1} = 0$ ,  $\frac{\textcircled{1}}{\textcircled{1}} = 1$ ,  $\textcircled{1}^0 = 1$ ,  $1^{\textcircled{1}} = 1$ ,  $0^{\textcircled{1}} = 0$

128  $\lim_{x \rightarrow \textcircled{1}} \frac{1}{x} = \frac{1}{\textcircled{1}}$ ,  $\lim_{x \rightarrow 2^{\textcircled{1}}} \frac{1}{x} = \frac{1}{2^{\textcircled{1}}}$ ,  $\lim_{x \rightarrow \frac{1}{\textcircled{1}}} x^3 = \frac{1}{\textcircled{1}^3}$

129  $\int_0^{\textcircled{1}} x^2 dx = \frac{1}{3} \textcircled{1}^3$ ,  $\int_{\textcircled{1}}^{\textcircled{1}^2} x^2 dx = \frac{1}{3} (\textcircled{1}^6 - \textcircled{1}^3)$ ,  $\int_0^{2^{\textcircled{1}}} x^2 dx = \frac{1}{3} \cdot 2^{3\textcircled{1}}$

130 More importantly, the Grossone method solves the calculation problems of  $\frac{\infty}{\infty}$ ,  $\infty - \infty$ , etc. that  
131 cannot be performed in the infinity theory.

132 For example, the following calculations are possible:

133  $\frac{\textcircled{1}}{2\textcircled{1}} = \frac{1}{2}$ ,  $\frac{2\textcircled{1}}{3\textcircled{1}^3} = \frac{2}{3\textcircled{1}^2}$ ,  $3\textcircled{1} - \textcircled{1} = 2\textcircled{1}$

134 It can be seen that the Grossone method meets the requirements of the unity of mathematical theory.  
135 From the above discussion, we can see that the cardinal method uses the "1-1 correspondence"  
136 principle but violates the "whole is greater than the part" principle, while the Grossone method uses the  
137 "whole is greater than the part" principle, but does not violate the "1-1 correspondence" principle.

138 Therefore, the new infinity theory can integrate the infinity cardinal number method with the  
139 Grossone method. But when using the infinity cardinal number theory to calculate, we should not use  
140 the "=" symbol, but can use "≡" to indicate that it is equivalent under the "1-1 correspondence"  
141 principle. E.g:

142  $\aleph_0 + 1 \equiv \aleph_0$ ,  $\aleph_0 + \aleph_0 \equiv \aleph_0$ ,  $\aleph_0 + 2^{\aleph_0} \equiv 2^{\aleph_0}$ ,  $2^{\aleph_0} + 2^{\aleph_0} \equiv 2^{\aleph_0}$ ;

143 
$$\frac{1}{\aleph_0+1} \equiv \frac{1}{\aleph_0}, \frac{1}{\aleph_0+\aleph_0} \equiv \frac{1}{\aleph_0}, \frac{1}{\aleph_0+2^{\aleph_0}} \equiv \frac{1}{2^{\aleph_0}}, \frac{1}{2^{\aleph_0}+2^{\aleph_0}} \equiv \frac{1}{2^{\aleph_0}};$$

144 
$$\textcircled{1}+1 \equiv \aleph_0, \textcircled{1}+\textcircled{1} \equiv \aleph_0, \textcircled{1}+2^{\textcircled{1}} \equiv 2^{\aleph_0}, 2^{\textcircled{1}}+2^{\textcircled{1}} \equiv 2^{\aleph_0};$$

145 
$$\frac{1}{\textcircled{1}+1} \equiv \frac{1}{\aleph_0}, \frac{1}{\textcircled{1}+\textcircled{1}} \equiv \frac{1}{\aleph_0}, \frac{1}{\textcircled{1}+2^{\textcircled{1}}} \equiv \frac{1}{2^{\aleph_0}}, \frac{1}{2^{\textcircled{1}}+2^{\textcircled{1}}} \equiv \frac{1}{2^{\aleph_0}}.$$

146 However, things are not so simple. Sergeyev also encountered a mathematical problem, which is the  
 147 "maximal number paradox." Just imagine, if  $\textcircled{1}$  represents the number of elements in a set of natural  
 148 numbers, is  $\textcircled{1}+1$  a natural number? If  $\textcircled{1}+1$  is a natural number, because of  $\textcircled{1}+1 > \textcircled{1}$ , then  
 149 the number of elements in the natural number set is not  $\textcircled{1}$ .

150 Sergeyev thought  $\textcircled{1}+1 \notin N$ , and the number greater than  $\textcircled{1}$  is called an extended number  
 151 [40]. But this is hard to make sense, because  $\textcircled{1}+1$  fully conforms to the definition of natural  
 152 numbers, and the extended natural numbers are still natural numbers. We will discuss this issue later.

153 **4. Grossone is a number-like symbol used for calculations**

154 In Cantor's infinite cardinal theory, the cardinal number of the natural number set,  $\aleph_0$ , is the  
 155 smallest infinite cardinal number. Using Grossone method, the set of natural numbers can also be  
 156 decomposed into smaller sets of infinity. For example: the natural numbers set  $N$  can be divided into  
 157 two infinite sets, the odd set and the even set. Let  $O$  be the odd set and  $E$  be the even set. Then  
 158 there are:

159 
$$O = \{1,3,5,\dots, \textcircled{1}-3, \textcircled{1}-1\}, E = \{2,4,6,\dots, \textcircled{1}-2, \textcircled{1}\}$$

160 
$$N = O \cup E = \{1,2,3,\dots, \textcircled{1}-3, \textcircled{1}-2, \textcircled{1}-1, \textcircled{1}\}$$

161 Obviously, the number of elements in the odd number set and the even number set is  $\frac{\textcircled{1}}{2}$ , which is  
 162 less than the number of elements  $\textcircled{1}$  in the natural number set.

163 Sergeyev also created a method of constructing an infinite subset of the natural number set [40]. He  
 164 uses  $N_{k,n}$  ( $1 \leq k \leq n$ ,  $n \in N$ ,  $n$  is a finite number) to indicate a set that the first number is  $k$ ,

165 and equal difference is  $n$ , and the size of the set is  $\frac{\textcircled{1}}{n}$ .

166 
$$N_{k,n} = \{k, k+n, k+2n, k+3n, \dots\}$$

167 
$$N = \bigcup_{k=1}^n N_{k,n}$$

168 For example:

169 
$$N_{1,2} = \{1,3,5,\dots\} = O, \quad N_{2,2} = \{2,4,6,\dots\} = E$$

170 
$$N = N_{1,2} \cup N_{2,2} = O \cup E$$

171 Or:

172 
$$N_{1,3} = \{1,4,7,\dots\}, \quad N_{2,3} = \{2,5,8,\dots\}, \quad N_{3,3} = \{3,6,9,\dots\}$$

173 
$$N = N_{1,3} \cup N_{2,3} \cup N_{3,3}$$

174 Grossone  $\textcircled{1}$  is a numeral symbol that represents the number of elements in natural numbers set.

175 However, the set of integers and real numbers are larger than the set of natural numbers. According to  
 176 the principle of "the whole is greater than the part", does it mean that there are integers and real  
 177 numbers greater than  $\textcircled{1}$  ?

178 Below we use Grossone method to examine the integer set  $Z$  and real number set  $R$ .

179 
$$Z = \{-\textcircled{1}, -\textcircled{1} + 1, \dots, 2, 1, 0, 1, 2, \dots, \textcircled{1} - 1, \textcircled{1}\}$$

180 
$$R = [-\textcircled{1}, -\textcircled{1} + 1) \cup \dots \cup [1, 0) \cup \{0\} \cup (0, 1] \cup \dots \cup (\textcircled{1} - 1, \textcircled{1}]$$

181 It is easy to see that there are no integers and real numbers exceeding  $\textcircled{1}$  in both the integer set  
 182 and the real number set.

183 The number of elements in the integer set is  $2\textcircled{1} + 1$ ; because the number of elements in  $(0, 1]$  is  
 184  $10^{\textcircled{1}}$ , the number of elements in the real number set is  $C = 2\textcircled{1} \cdot 10^{\textcircled{1}} + 1$ . It can be seen that the set  
 185 of real numbers is not the power set of the set of natural numbers. Obviously, Integer set and real  
 186 number set the number of elements in are all greater than  $\textcircled{1}$ .

187 The integer set and real number set are larger than the natural number set, which refers to the number  
 188 of elements, rather than the existence of numbers exceeding  $\textcircled{1}$  in the integer set and real number set.

189 In fact,  $\textcircled{1}$  is not a number, but infinity. No number can exceed infinity, and  $\textcircled{1}$  is a symbol for  
 190 infinity.

191 Looking back at the problem of the "maximum number paradox" now, it is not difficult to solve it.

192 The problem lies in the qualitative aspect of A. In fact, A is just a number-like symbol used for  
 193 infinity calculations, and is a ruler used to measure all infinity sets.

194 Take  $\textcircled{1} + 1$  as an example. First,  $\textcircled{1} + 1$ , like  $\textcircled{1}$ , is infinity, not a numeral. Second,

195  $\textcircled{1} + 1 > \textcircled{1}$ , indicating that this infinite set exceeds a single Grossone  $\textcircled{1}$ . Exceeding does not mean  
 196 that it cannot be expressed. It is like measuring an object with a ruler. It does not matter if the object  
 197 exceeds the ruler. You can measure a few more times.  $\textcircled{1}$  is the ruler for measuring the infinite set.  
 198 An infinite set is 1 more than this ruler. You can measure it more. After the measurement is accurate,  
 199 mark it as  $\textcircled{1} + 1$ .

200 Let  $A$  be an infinite set of  $\textcircled{1} + 1$  elements, then  $A$  can be written as:

201 
$$A = N \cup \{1\} = \{1, 2, \dots, \textcircled{1} - 1, \textcircled{1}, 1\}$$

202 Or:

203 
$$A = N \cup \{\textcircled{1} + 1\} = \{1, 2, \dots, \textcircled{1} - 1, \textcircled{1}, \textcircled{1} + 1\}$$

204 Or:

205 
$$A = \{a_1, a_2, \dots, a_{\textcircled{1}-1}, a_{\textcircled{1}}, a_{\textcircled{1}+1}\}$$

206 It can be seen that the so-called "maximum number paradox" does not exist for Grossone method.

207 **5. The size of the point is not zero and the continuity of the set**

208 The continuum originally refers to the real numbers set. Since the real number corresponds to the  
 209 point 1-1 on the straight line, the straight line is intuitively composed of continuous and unbroken  
 210 points, so the real number set is called the continuum. In the number sequence, the set that satisfies the  
 211 "1-1 correspondence" relationship with the interval (0, 1) is called the continuum.

212 Traditional mathematics has an axiom: a point has no size. This can be proved by continuum theory.

213 Suppose that the interval (0,1] on the number axis is the continuum, and the points on the number  
 214 axis are "dense and no holes", so the distance between points is zero. Since the cardinal number of the  
 215 continuum is  $c$ , the interval (0,1] point size is:

216 
$$s = \lim_{c \rightarrow \infty} \frac{1}{c} = 0$$

217 However, according to the Grossone method, because the number of elements in (0,1] is  $10^{\textcircled{1}}$ , the  
 218 size of the point in the interval (0,1] is:

219 
$$s = \frac{1}{10^{\textcircled{1}}} > 0$$


220 Not only does the dot have a size, but the size of the dot is related to the decimal or binary system of  
 221 the number on the number axis. For example, when using binary system, the number of elements in  
 222 (0,1] is  $2^{\textcircled{1}}$ , and the size of the point in the interval (0,1) is:

223 
$$s = \frac{1}{2^{\textcircled{1}}}$$

224 Imagine that one-dimensional straight lines, two-dimensional planes, three-dimensional and  
 225 multi-dimensional spaces, etc. are all composed of points. If the size of a point is zero, how to form a

226 straight line, plane and space with size? The Grossone method solves this infinitesimal puzzle.

227 We use a probability problem exemplified by Sergeev to illustrate [40].



228

229 As shown in the figure above, suppose the radius of the disc in the figure is  $r$ , and the disc is  
 230 rotating. We want to ask a probabilistic event  $E$ : What is the probability that point A on the disk stops  
 231 just in front of the fixed arrow on the right? According to the traditional calculation method, point A  
 232 has no size, so the probability of occurrence of  $E$  is:

233 
$$P(E) = \lim_{h \rightarrow 0} \frac{h}{2\pi r} = 0$$

234 This is obviously contrary to experience and common sense. And if the size of the point is solved,  
 235 such as  $s = \frac{1}{10} \text{①}$ , then you can get:

236 
$$P(E) = \frac{1}{2\pi r \cdot 10 \text{①}}$$

237 This is the logical result.

238 Although point has a size, point is isolated, so its dimension is zero. We know that straight lines and  
 239 planes are one-dimensional and two-dimensional continuums, corresponding to real numbers and  
 240 complex numbers, respectively. There are also three-dimensional and multi-dimensional space  
 241 continuums, such as Euclidean space, Minkowski space, Riemann space, etc. In this sense, a point can  
 242 also be regarded as a special continuum, that is, a zero-dimensional continuum.

243 In Sergeev's view, the "dense and no-hole" continuum is an absolute continuity. Correspondingly,  
 244 he proposed a relative continuity theory [40]. In order to distinguish between the two continuums, I call  
 245 the traditional continuum collectively the absolute continuum, and the set of relative continuity as the  
 246 relative continuum.

247 Sergeev established the relative continuity on the function  $f(x)$ . The point that stipulates the  
 248 range of the independent variable  $[a, b]_S$  of  $f(x)$  can be a finite number or an infinity, but the set  
 249  $[a, b]_S$  is always discrete, where S represents a certain numeral system. In this way, for any point  
 250  $x \in [a, b]_S$ , its nearest left and right neighbors can always be determined:

251 
$$x^+ = \min\{z : z \in [a, b]_S, z > x\}, \quad x^- = \max\{z : z \in [a, b]_S, z < x\}$$

252 Suppose a set  $X = [a, b]_S = \{x_0, x_1, \dots, x_{n-1}, x_n\}_S$ , where  $a = x_0$ ,  $b = x_n$ , and the numeral  
 253 system S allow a certain unit of measure  $\mu$  to be used to calculate the coordinates of the elements in

254 the set. If for any  $x \in (a, b)_s$ ,  $x^+ - x$  and  $x - x^-$  are infinitesimal, then the set  $X$  is said to be  
 255 continuous in the unit of measure  $\mu$ . Otherwise, set  $X$  is said to be discrete in the unit of measure  
 256  $\mu$ .

257 For example, if the unit of measure  $\mu$  is used to calculate that the position difference between  
 258 adjacent elements of set  $X$  is equal to  $\textcircled{1}^{-1}$ , then set  $X$  is continuous in the unit of measure  $\mu$ ;  
 259 but if the unit of measure  $\nu = \mu \cdot \textcircled{1}^{-3}$  is used instead, calculate that the position difference between  
 260 adjacent elements of the set  $X$  is equal to  $\textcircled{1}^2$ , then the set  $X$  is discrete in the unit of measure  
 261  $\nu$ . Therefore, whether the set  $X$  is continuous or discrete depends on the size of the unit of measure  
 262  $\mu$ .

263 Function  $f(x)$  is continuous in the unit of measure at some point  $x \in (a, b)_s$  in  $[a, b]_s$ , if  
 264  $f(x^+) - f(x)$  and  $f(x) - f(x^-)$  are both infinitesimal. If only one is infinitesimal, it can be  
 265 called left continuous or right continuous. If function  $f(x)$  is continuous in the unit of measure  $\mu$   
 266 at each point of  $[a, b]_s$ , then  $f(x)$  is said to be continuous in the unit of measure  $\mu$  on set  
 267  $X = [a, b]_s$ .

268 Relative continuum theory provides a new way to solve the problem of continuity of sets other than  
 269 absolute continuum.

## 270 **6. The mathematical continuum is the foundation of the cosmic continuum**

271 The cosmic continuum is a scientific foundation theory based on the mathematical continuum  
 272 [4]-[14]. The cosmic continuum theory believes that there are three basic entities in the universe: mass  
 273 body, energy body and dark mass body. Their smallest units are elementary particle  $m_{\min}$ , elementary  
 274 quantum  $q_{\min}$ , and elementary dark particle  $d_{\min}$ .

275 In this way, we get three countably-infinite sets in the universe: elementary particle set  $M$ ,  
 276 elementary quantum set  $Q$ , and elementary dark particle set  $D$ . Suppose  $m_i$  is a elementary  
 277 particle,  $q_i$  is a elementary quantum,  $d_i$  is a elementary dark particle,  $i$  is a natural number, then:

$$278 \quad M = \{m_1, m_2, \dots, m_i, \dots\}, Q = \{q_1, q_2, \dots, q_i, \dots\}, D = \{d_1, d_2, \dots, d_i, \dots\}$$

279 Thus, we obtain the following basic existence set  $E$ :

$$280 \quad E = M \cup Q \cup D = \{m_1, m_2, \dots, m_i, \dots\} \cup \{q_1, q_2, \dots, q_i, \dots\} \cup \{d_1, d_2, \dots, d_i, \dots\}$$

281 According to cosmic continuum theory, the coupling energy quantum connects all entities to form

282 the universe as a whole, and any changes in the universe affect the whole. Due to the effect of the  
 283 coupling energy quantum, the basic existence set  $E$  will form the following power set, namely the  
 284 existence continuum  $P(E)$  :

$$285 \quad P(E) = 2^E = \{e | e \subseteq E\}$$

286 Since space is the existence dimension of the mass body, time is the existence dimension of the  
 287 energy body, and dark space is the existence dimension of the dark mass body, correspondingly, we  
 288 obtain the other three countable infinite sets in the universe: basic space set  $S$  , basic time set  $T$  and  
 289 basic dark space set  $G$  . Suppose  $s_i$  is the elementary space,  $t_i$  is the elementary time,  $g_i$  is the  
 290 elementary dark space,  $i$  is a natural number, then:

$$291 \quad S = \{s_1, s_2, \dots, s_i, \dots\}, T = \{t_1, t_2, \dots, t_i, \dots\}, G = \{g_1, g_2, \dots, g_i, \dots\}$$

292 Thus, we obtain the following basic dimension set  $W$  :

$$293 \quad W = S \cup T \cup G = \{s_1, s_2, \dots, s_i, \dots\} \cup \{t_1, t_2, \dots, t_i, \dots\} \cup \{g_1, g_2, \dots, g_i, \dots\}$$

294 The basic dimension set  $W$  will correspondingly form the following power set, namely the  
 295 dimension continuum  $P(W)$  :

$$296 \quad P(W) = 2^W = \{w | w \subseteq W\}$$

297 According to Cosmic continuum hypothesis: the universe is a continuum consisting of an existence  
 298 continuum and an existing dimension continuum. The existence continuum is composed of mass bodies,  
 299 energy bodies and dark mass bodies. The existing dimension continuum is composed of space, time  
 300 and dark space.

301 Let the cosmic continuum be  $U$  , then:

$$302 \quad U = P(E) \cup P(W) = 2^E \cup 2^W = \{e | e \subseteq E\} \cup \{w | w \subseteq W\}$$

303 Therefore, the cosmic continuum is actually a multi-dimensional continuum composed of the power  
 304 set of the basic existence set and the power set of the basic dimensional set. It is an absolute continuum  
 305 that conforms to the Cantor cardinal number method.

306 In Cosmic Continuum Theory, the dimensional continuum  $P(W)$  is the mirror image of the  
 307 existence continuum  $P(E)$  .

308 For example, in the history of physics, three fundamental theories of physics, classical mechanics,  
 309 relativity, and quantum mechanics have appeared successively. The corresponding Euclidean space,  
 310 Minkowski space, Riemann space, matrix space, and probability space are equivalent to Space-time  
 311 mirroring of physics events.

312 In this sense, the cosmic continuum is actually a continuum composed of all physical events in the  
 313 universe. In other words, every element of the cosmic continuum is a physical event.

314 Next we examine how to describe physical events in the cosmic continuum.

315 According to the change axiom of the cosmic continuum: the change of energy is the cause of the

316 state change of all cosmic systems.

317 Suppose the energy change in a universe system is  $\Delta E$ , and the corresponding change in the state  
318 of the universe system is  $\Delta x$ , then  $\Delta E \equiv \Delta x$ . In the universe continuum, energy is the independent  
319 variable, and the state of the universe system is the dependent variable.

320 Relative continuum provides us with a mathematical tool for describing physical events.

321 The change axiom tells us that the essence of all physical events is that energy changes cause the  
322 state of the universe to change. Suppose the dependent variable  $f(x)$  is the state of the universe

323 system, the independent variable  $x$  is the energy, and  $[a, b]_s$  is the range of the independent  
324 variable that can be described by a certain digital system S, then any physical event, including all  
325 interactions and quantum phenomena, can be described by the above-mentioned relative continuum .

326 Example 1, the physical event of classical mechanics: car action. The dependent variable  $f(x)$  is  
327 the car coordinates, the independent variable  $x$  is gasoline, and  $x \in (a, b)_s$  represents the range of  
328 gasoline amount.

329 Example 2: Relativistic physical event: "deflection of light" under the action of gravity. The  
330 dependent variable  $f(x)$  is the ray coordinate, the independent variable  $x$  is the gravitational field,  
331 and  $x \in (a, b)_s$  represents the range of the gravitational field.

332 Example 3: Quantum mechanics physical event: photoelectric effect. The dependent variable  $f(x)$   
333 is the electronic state, the independent variable  $x$  is the photon frequency, and  $x \in (a, b)_s$   
334 represents the photon frequency range.

### 335 **7. Conclusion**

336 This article discusses several aspects of the development of infinity theory.

337 (1) Qualitative calculation and quantitative calculation. Cantor used the cardinal number method to  
338 solve the problem of comparing infinity. Sergeyev used Grossone method to solve the problem of  
339 unifying the calculation rules of infinity and finite numbers.

340 (2) Absolute continuum and relative continuum. The continuum in traditional mathematics refers to  
341 a collection of "dense and no holes", the relative continuum is a continuum that changes with the  
342 change of measurement units.

343 (3) Cosmic continuum and continuum theory. The universe continuum is a mathematical model of  
344 existence and its dimensions in the universe. The development of continuum theory provides a new  
345 mathematical foundation for the Cosmic Continuum Theory.

346 The discussion in this article shows that:

347 (1) The theory of infinity will usher in a new era of development. Grossone method is a scientific  
348 infinity theory like the cardinal number method; in the new infinity theory, infinity and infinity can be  
349 mathematically calculated like finite numbers.

350 (2) New developments in the theory of infinity will promote the development of fundamental  
351 scientific theories. The basic theories of mathematics and physics have always been intertwined and

352 developed, such as classical mechanics and calculus, relativity and non-Euclidean geometry, etc.,  
353 which are all good stories in the history of science. The relative continuum theory provides a new path  
354 for the study of the cosmic continuum.

355 (3) The essence of Hilbert's first problem "continuum hypothesis" is that the infinity theory is not yet  
356 mature. The development of Grossone theory makes this problem self-explanatory. According to the  
357 principles of "power set is greater than original set" and "whole is greater than part", there is neither the  
358 largest infinity and infinitesimal, nor the smallest infinity and infinitesimal.

359

360 Competing interests statement: I declare that there is no competing Interest.

361

## 362 **References**

363 [1] Cantor, Contributions to the founding of the theory of transfinitive numbers, translated, and  
364 provided with an introduction and notes, by Philip E. B. Jourdain, Dover Publications, Inc., New  
365 York, N. Y., 1952. Zbl 0046.05102 MR 45635

366 [2] P. J. Cohen, Set Theory and the Continuum Hypothesis, Benjamin, New York, 1966. Zbl  
367 0182.01301 MR 232676

368 [3] K. Gödel, The Consistency of the Continuum-Hypothesis, Princeton University Press, Prince  
369 ton, 1940. Zbl 0061.00902 MR 2514

370 [4] Wang, X.J. Scientific poverty and outlet, Science and Management, 4(1990), 28–30.

371 [5] Wang, X.J. and Wu, J.X. Unification. Haitian Publishing House, Shenzhen, 1992

372 [6] Wang, X.J. and Wu, J.X. Unification - Deciphering the mysteries of the universe, Tech wave, 11  
373 (1993), 24–26. Xinhua Digest, 1(1994), 180–182.

374 [7] Wang, X.J. Modern Interpretation of Classic of Changes System. Social Science, 9(1997), 57–59.

375 [8] Wang, X.J. The Sublimation of Thinking Experiment. Invention and Innovation, 6(1997), 8-9.

376 [9] Wang, X.J. and Wu, J.X. To solve the mystery of scientific unity. Hunan Science and Technology  
377 Press, Changsha, 2001

378 [10] Wang, X.J. Unification Theory: Challenging Traditional Scientific Norms. Invention and  
379 Innovation, 4(2003), 32-33.

380 [11] Wang, X.J. Axiomatization of the Symbols System of Classic of Changes: The Marriage of  
381 Oriental Mysticism and Western Scientific Tradition. Foundations of Science, 25(2020) , 315–325.  
382 <https://doi.org/10.1007/s10699-019-09624-5>

383 [12] Wang, X.J. Cosmic Continuum Theory: A New Idea on Hilbert's Sixth Problem. Journal of  
384 Modern Physics, 9(2018), 1250-1270. <https://doi.org/10.4236/jmp.2018.96074>

385 [13] Wang, X.J. New Discovery on Planck Units and Physical Dimension in Cosmic Continuum  
386 Theory. Journal of Modern Physics, 9(2018), 2391-2401. <https://doi.org/10.4236/jmp.2018.914153>

387 [14] Wang, X.J. New Explanation on Essence of Quantum Phenomena and Interactions and the  
388 Gravitational Action in Cosmic Continuum Theory" SSRG International Journal of Applied Physics  
389 7(3)(2020), 88-96. <https://doi.org/10.14445/23500301/IJAP-V7I3P114>

390 [15] R. De Leone, G. Fasano, and Ya. D. Sergeyev, Planar methods and grossone for the Conjugate  
391 Gradient breakdown in nonlinear programming, to appear in Comput. Optim. Appl.

392 [16] D. I. Iudin, Ya. D. Sergeyev, and M. Hayakawa, Interpretation of percolation in terms of  
393 infinity computations, Appl. Math. Comput., 218 (2012), no. 16, 8099–8111. Zbl 1252.82059 MR  
394 2912732

395 [17] D. I. Iudin, Ya. D. Sergeyev, and M. Hayakawa, Infinity computations in cellular automaton

396 forest-fire model, *Comm. Nonlinear Sci. Numer.*, 20 (2015), no. 3, 861–870.

397 [18] F. Mazzia, Ya. D. Sergeyev, F. Iavernaro, P. Amodio, and M. S. Mukhametzhano, Numerical  
398 methods for solving ODEs on the Infinity Computer, in *Proc. of the 2nd Intern. Conf. “Numerical  
399 Computations: Theory and Algorithms”*, Ya. D. Sergeyev et al. (eds.), 1776, 090033, AIP Publishing,  
400 New York, 2016.

401 [19] Ya. D. Sergeyev, R. G. Strongin, and D. Lera, *Introduction to Global Optimization Exploiting  
402 Space-Filling Curves*, Springer, New York, 2013. Zbl 1278.90005 MR3113120

403 [20] Ya. D. Sergeyev, *Arithmetic of Infinity*, Edizioni Orizzonti Meridionali, CS, 2003, 2nd ed. 2013.  
404 Zbl 1076.03048 MR2050876

405 [21] Ya. D. Sergeyev, Blinking fractals and their quantitative analysis using infinite and  
406 infinitesimal numbers, *Chaos, Solitons & Fractals*, 33 (2007), no. 1, 50–75.

407 [22] Ya. D. Sergeyev, A new applied approach for executing computations with infinite and  
408 infinitesimal quantities, *Informatica*, 19 (2008), no. 4, 567–596. Zbl 1178.68018 MR2589840

409 [23] Ya. D. Sergeyev, Evaluating the exact infinitesimal values of area of Sierpinski’s carpet and  
410 volume of Menger’s sponge, *Chaos, Solitons & Fractals*, 42 (2009), no. 5, 3042–3046.

411 [24] Ya. D. Sergeyev, Numerical computations and mathematical modelling with infinite and  
412 infinitesimal numbers, *J. Appl. Math. Comput.*, 29 (2009), 177–195. Zbl 1193.68260 MR  
413 2472104

414 [25] Ya. D. Sergeyev, Numerical point of view on Calculus for functions assuming finite, infinite,  
415 and infinitesimal values over finite, infinite, and infinitesimal domains, *Nonlinear Anal.*, 71  
416 (2009), no. 12, e1688–e1707. Zbl 1238.28013 MR2671948

417 [26] Ya. D. Sergeyev, Computer system for storing infinite, infinitesimal, and finite quantities  
418 and executing arithmetical operations with them, USA patent 7,860,914, 2010.

419 [27] Ya. D. Sergeyev, Counting systems and the First Hilbert problem, *Nonlinear Anal.*, 72 (2010),  
420 no. 3-4, 1701–1708. Zbl 1191.03038 MR2577570

421 [28] Ya. D. Sergeyev, Lagrange Lecture: Methodology of numerical computations with infinities  
422 and infinitesimals, *Rend. Semin. Mat. Univ. Politec. Torino*, 68 (2010), no. 2, 95–113. MR  
423 2790165

424 [29] Ya. D. Sergeyev, Higher order numerical differentiation on the Infinity Computer, *Optim.  
425 Lett.*, 5 (2011), no. 4, 575–585. Zbl 1230.65028 MR2836038

426 [30] Ya. D. Sergeyev, On accuracy of mathematical languages used to deal with the Riemann zeta  
427 function and the Dirichlet eta function, *p-Adic Numbers Ultrametric Anal. Appl.*, 3 (2011), no. 2,  
428 129–148. Zbl 1268.11114 MR2802036

429 [31] Ya. D. Sergeyev, Using blinking fractals for mathematical modelling of processes of growth in  
430 biological systems, *Informatica*, 22 (2011), no. 4, 559–576. Zbl 1268.37092 MR2885687

431 [32] Ya. D. Sergeyev, Solving ordinary differential equations by working with infinitesimals  
432 numerically on the Infinity Computer, *Appl. Math. Comput.*, 219 (2013), no. 22, 10668–10681.  
433 Zbl 1303.65061 MR3064573

434 [33] Ya. D. Sergeyev, Computations with grossone-based infinities, in *Unconventional  
435 Computation and Natural Computation: Proc. of the 14th International Conference UCNC 2015*, C.  
436 S. Calude and M.J. Dinneen (eds.), LNCS 9252, 89–106. Springer, New York, 2015. Zbl 06481801  
437 MR3447459

438 [34] Ya. D. Sergeyev, The Olympic medals ranks, lexicographic ordering, and numerical infinities,  
439 *Math. Intelligencer*, 37 (2015), no. 2, 4–8. Zbl 1329.90074 MR3356109

440 [35] Ya. D. Sergeyev, Un semplice modo per trattare le grandezze infinite ed infinitesime, *Mat.*  
441 *Soc. Cult. Riv. Unione Mat. Ital. (I)*, 8 (2015), no. 1, 111–147. MR3364901

442 [36] Ya. D. Sergeyev, The exact (up to infinitesimals) infinite perimeter of the Koch snowflake  
443 and its finite area, *Commun. Nonlinear Sci. Numer. Simul.*, 31 (2016), no. 1–3, 21–29. MR  
444 3392467

445 [37] Ya. D. Sergeyev and A. Garro, Observability of Turing machines: A refinement of the theory  
446 of computation, *Informatica*, 21 (2010), no. 3, 425–454. Zbl 1209.68255 MR2742193

447 [38] Ya. D. Sergeyev and A. Garro, Single-tape and multi-tape Turing machines through the lens of  
448 the Grossone methodology, *J. Supercomput.*, 65 (2013), no. 2, 645–663.

449 [39] Ya. D. Sergeyev, M. S. Mukhametzhano, F. Mazzia, F. Iavernaro, and P. Amodio, Numerical  
450 methods for solving initial value problems on the Infinity Computer, *Internat. J. Unconventional*  
451 *Comput.*, 12 (2016), no. 1, 3–23.

452 [40] Yaroslav D. Sergeyev, Numerical infinities and infinitesimals: Methodology, applications,  
453 and repercussions on two Hilbert problems, *EMSSurv. Math. Sci.* 4 (2017), 219–320 DOI  
454 10.4171/EMSS/4-2-3

455 [41] Davide Rizza, A Study of Mathematical Determination through Bertrand's  
456 Paradox, *Philosophia Mathematica*. 26 (3)(2018):375-395, Advance Access Publication on  
457 December 16, 2017. DOI 10.1093/phimat/nkx035

458 [42] Cristian S. Calude and Monica Dumitrescu, Infinitesimal Probabilities Based on Grossone, *SN*  
459 *Computer Science* (2020) 1:36 <https://doi.org/10.1007/s42979-019-0042-8>

460 [43] LORENZO FIASCHI AND MARCO COCCIONI, Numerical Asymptotic Results in Game Theory  
461 Using Sergeyev's Infinity Computing, *Int. Journ. of Unconventional Computing*, Vol. 14(2018), pp.  
462 1–25


463 [44] Kline, M. *Mathematical Thought from Ancient to Modern Times*. Oxford University Press,  
464 Oxford, 1972

465 [45] L. Corry, *A brief history of numbers*, Oxford University Press, Oxford, 2015. Zbl 1335.01001  
466 MR3408089

467 [46] D. Hilbert, Mathematical problems: Lecture delivered before the International Congress of  
468 Mathematicians at Paris in 1900, *American M. S. Bull.* (2), 8 (1902), 437–479. Zbl 33.0976.07 MR  
469 1557926

470

## Figures



**Figure 1**

Imagine that one-dimensional straight lines, two-dimensional planes, three-dimensional and multi-dimensional spaces, etc. are all composed of points. If the size of a point is zero, how to form a straight line, plane and space with size? The Grossone method solves this infinitesimal puzzle. We use a probability problem exemplified by Sergeyev to illustrate.