

Doppler Ultrasound Evaluation of Patients with Popliteal Vascular Entrapment Syndrome

Hai Zhong

Second Hospital of Shandong University

Guangrui Shao

Second Hospital of Shandong University

Hengtao Qi

Shandong Medical Imaging Research Institute

Yuan Zhao

Shandong University of Traditional Chinese Medicine

Maohua Wang

Shandong Provincial Hospital

wubo Yang (✉ 404958638@qq.com)

Second Hospital of Shandong University <https://orcid.org/0000-0002-5622-8595>

Research

Keywords: Popliteal vascular entrapment syndrome, Doppler ultrasound, Provocative maneuvers

Posted Date: May 29th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-30724/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background

To retrospectively evaluate Doppler ultrasound findings for patients with popliteal vascular entrapment syndrome (PVES).

Methods

Twenty-four patients (30 legs) who underwent surgical intervention for PVES were included in this study. The popliteal artery, popliteal vein, and surrounding musculotendinous structures were explored in all symptomatic lower extremities using ultrasound. Intraoperative findings served as the gold standard. In the absence of popliteal artery occlusion, popliteal arteries were examined using Doppler ultrasound at rest and during a provocative maneuver.

Results

Ultrasonography in 25 cases (25/30, 83%) showed that gastrocnemius medial head or popliteal vascular anomalies were consistent with surgical photography results. According to Whelan and Rich classification, the classic type I was found in five limbs, type II in twelve limbs, type III in seven limbs, and type V in one limb. In the remaining five cases (5/30, 17%), no anatomical abnormalities of the popliteal fossa were discovered by the ultrasound examination. Among these, two cases (2/30, 7%) were misdiagnosed as lower extremity atherosclerosis and popliteal vascular depression was not found. In addition, increasing blood flow velocity was found in three cases (3/30, 10%) of compressed popliteal arteries using Duplex scanning during active plantar flexion, with decreasing flow at the distal end. Color Doppler ultrasound of the affected popliteal vessels showed varied degrees of vascular structure pathology, including arterial stenosis in ten limbs, occlusion in fifteen limbs, and aneurysm in two limbs.

Conclusion

Doppler ultrasound may have a high diagnostic rate in PVES. Due to its simplicity, repeatability, functional evaluation of blood flow, and non-invasiveness, it can be used as a primary screening examination modality in PVES. Provocative maneuvers could help clinicians diagnose PVES.

Introduction

The popliteal artery and popliteal vein normally run above the popliteal muscle between the medial and lateral heads of the gastrocnemius muscle. Popliteal vascular entrapment syndrome (PVES) due to an abnormal anatomical relationship between popliteal vessels and neighboring myofascial structures results in repeated vessel compression during exercise. Young adults and active adolescents are most

often affected by the syndrome. Claudication is the predominant clinical feature of PVES, which is characterized by calf pain after exercise [1]. After repeated irritation and compression for a long time, eventually irreversible trauma to the popliteal vessels can result in arterial stenosis, occlusion, thrombosis, post-stenotic aneurysms, and popliteal vein thrombosis [2] .

Therefore, early diagnosis and treatment are crucial for decreasing morbidity and preventing complications. Imaging modalities can provide information that might allow to identify entrapment [3]. The present study described the popliteal vessel and adjacent musculotendinous structure findings by Doppler ultrasound in patients with PVES.

Methods

Patients

The local institutional ethics committee approved this retrospective study (KYLL-2019(LW)025), and the need for informed consent was waived. The present study included twenty-four patients who underwent surgical intervention for PVES between May 2011 and April 2019 in The Second Hospital of Shandong University and Shandong Provincial Hospital (Jinan, China). Six patients were affected bilaterally. A total of nineteen males and five females, ranging in age from 16 to 58 years old (mean, 28 years old) were evaluated using Doppler sonography. All patients had a history of progressive claudication. Paresthesia, calf swelling, and foot coldness were also observed in some individuals. Patient clinical profiles are summarized in Table 1. All patients were surgically treated.

Table 1
Clinical profiles for patients with PVES

	Number/Total number (%)
Location	
Right	10/24 (42%)
Left	8/24 (33%)
Both	6/24 (25%)
Symptoms	
Claudication	30/30 (100%)
Paresthesia	21/30 (70%)
Calf swelling	7/30 (23%)
Foot coldness	8/30 (27%)
Risk factor	
Exercise	5/24 (21%)
Smoker	3/24 (13%)
High blood pressure	2/24 (8%)
N, number of limbs.	

Doppler ultrasound scanning

All subjects were analyzed with linear (7–10 MHz) transducers connected to Philips IU 22, (Philips, Amsterdam, the Netherlands) or GE Vivid E9 (General Electric Healthcare, Milwaukee, USA). Graded dynamic assessment combined gray-scale imaging with range-gated pulsed-Doppler analysis of blood flow at each point during imaging. Subjects with relevant limbs were evaluated in a prone position by two radiologists specializing in vascular ultrasonography. Each patient was placed on an examination couch with the knee of the examined leg slightly flexed to ensure that the gastrocnemius muscle was fully relaxed. Color Doppler ultrasound of the popliteal vessel was first placed with the foot in a neutral position and then actively plantar flexed to maximum the active extension of the knee, while an assistant added resistance to the plantar flexion using manual pressure on the sole of the foot. Duplex ultrasound during this modality of plantar flexion was designated as a provocative maneuver test [4], which was performed in patients without popliteal vessel occlusion.

The neutral position was used to visualize artery occlusion, aneurysm, popliteal vessels anomalies, and presence of collateral arteries. Repeated plantar flexion until the individual becomes symptomatic may be necessary to demonstrate occlusion [5]. Popliteal artery with a peak systolic velocity ratio exceeding 200% indicates a positive result. Careful assessment of both lower limbs was required to determine if the

abnormality was bilateral. Ultrasound PVES imaging was divided into six types according to Whelan and Rich classification (Table 2) [1, 6, 7].

Table 2
Types of patients with PVES

Type	Description
Type I	Aberrant course of the PA medial to a normal MHG
Type II	Abnormal lateral insertion of the MHG, no deviation of PA
Type III	Abnormal muscle bundle from MHG surrounding the PA
Type IV	Abnormal location of PA, deep in the popliteus muscle or a fibrous band in the popliteal fossa
Type V	Any form of the entrapment that involves both PA and popliteal vein
Type VI	PA is normally positioned and entrapped by a normally positioned hypertrophied gastrocnemius
MHG indicates medial head of gastrocnemius muscle; PA, popliteal artery.	

Results

All twenty-four patients (30 legs) underwent a successful surgical treatment. Popliteal vascular entrapment was presented unilaterally in eighteen patients and bilaterally in six patients. All cases had popliteal vessel migration or gastrocnemius medial head anomalies.

Doppler ultrasound was performed with dynamic testing in all patients. It was found that 25 cases (25/30) of limb popliteal artery and popliteal vein were separated. Ultrasound images show the popliteal artery and popliteal vein moving from a normal forward and backward orientation into a left and right orientation and the medial head of the gastrocnemius muscle or an abnormal bundle between the vessels (Fig. 1–4). Each case was classified grounded on the findings of Doppler ultrasound. Five limbs with medial deviation of the popliteal artery were classified as type I. Twelve limbs with the medial head of the gastrocnemius origin shifting laterally were classified as type II (Fig. 5). Seven limbs were type III with an anomalous accessory band of the medial head of gastrocnemius (Fig. 4). One limb was type V with any type of entrapment that includes the popliteal vein as well as artery. All ultrasound features and classifications were confirmed by surgical photography. In the remaining five cases (5/30), no anatomical abnormalities of the popliteal fossa were found by ultrasound examination. Two cases were diagnosed as lower extremity atherosclerosis. In addition, three cases were normal in the neutral position, whereas the three popliteal arteries were narrowed or occluded in the plantar flexed position (Fig. 6). Duplex imaging demonstrated popliteal vessel compression occlusion with blood flow interruption and significant popliteal vessel compression of the lumen. Doppler spectrum showed that the peak systolic

blood flow velocity increased significantly, more than two times the value before the provocative maneuvers (Table 3). Doppler blood flow velocity was decreased near the tibiofibular stem.

Table 3
Doppler ultrasound findings for three limbs with PVES

Case	At rest	Provocative test
1	Normal flow	Popliteal arterial stenosis Peak systolic blood velocity increased significantly in popliteal artery
2	Normal flow	Popliteal arterial occlusion
3	Normal flow	Popliteal arterial stenosis Peak systolic blood velocity increased significantly in popliteal artery Tibiofibular stem Doppler blood velocity was decreased
At rest, in a neutral position; Provocative test, in plantar flexed position		

When the feet were placed in the neutral position, ten of the 30 limbs with popliteal artery were stenotic (Fig. 3c). Arterial occlusion was demonstrated in fifteen of the 30 limbs (Figs. 2 and 4). Occlusion was characterized by hypoechoic or medially echoic thrombosis filling the lumen, no blood flow during color Doppler flow imaging, decreased blood flow velocity in the distal artery, and one-way spectrum. Partial lumen thinning was even more difficult to identify and distal segment of the popliteal artery had abundant collateral vessels. Doppler scan images also showed a post-stenotic aneurysm in one limb and a thrombosed aneurysm in another limb. With the feet placed in the plantar flexed position, unoccluded popliteal vascular flow tended to decrease or disappear. These ultrasound findings are shown in Table 4.

Table 4
Doppler ultrasound findings for patients with PVES in neutral position

Combined anomaly	Vascular stenosis (N)	Vascular occlusion (N)	Post-stenotic aneurysm (N)	Normal flow (N)	Total (N)
Gastrocnemius medial head anomaly	10	13	2		25
No anatomical abnormalities		2		3	5
Total (N)	10	15	2	3	30
N, number of limbs.					

Discussion

PVES is a rare cause of exercise-induced leg pain. Active young people and athletes without risk factors, such as smoking, heart disease, or systemic atherosclerosis, are affected. The symptoms are correlated with the degree of compression of the popliteal vessels. Popliteal vascular structure is normal in the early stage and the disease becomes evident only after vascular compression between the contracting gastrocnemius muscles occurs [8]. Repeated extrinsic vascular compression causes trauma to the vascular wall, leading to premature intrinsic atherosclerosis and thrombus formation. The pathology of PVES is considered to be progressive and resulting stenosis and turbulence can lead to post-stenotic aneurysm. In addition, acute ischemia can occur if there is an occluded artery or aneurysm with thrombosis [2, 9]. Early diagnosis is crucial as PVES is a progressive disorder and early intervention is recommended to prevent serious complications [8, 9].

Color Doppler ultrasound has been widely used to diagnose peripheral vascular diseases [10–13]. This non-invasive examination does not require contrast enhancement, patient preparation before study, or radiation exposure. The present study suggests that ultrasonography is able to show when popliteal artery and popliteal vein are separated, when normal forward and backward alignment of popliteal artery and vein becomes aligned in left and right directions, and medial head of gastrocnemius muscle or abnormal muscle bundle between the blood vessels, which are crucial signs for the diagnosis of PVES. The present study found that the popliteal artery and popliteal vein were separated in 83% (25/30) of the cases, which was consistent with surgical results, suggesting a high ultrasound sensitivity. This included six patients with bilateral popliteal vessel involvement. Collins et al. [14] reviewed about 22–67% of the bilateral cases. Duplex scanning examination associated with pulsed Doppler offers a more precise scope of the popliteal fossa. This modality makes it possible to identify PVES by showing a popliteal artery may take abnormal course or an anatomical aberrant composition between the popliteal artery and vein that increases the distance separating the two vessels [15]. Separation of the arteries and veins can be shown on the ultrasound because the popliteal artery and popliteal vein move in a left-right direction, even if there is an arterial occlusion. Popliteal vascular ultrasonography on the affected side shows varied degrees of vascular structure pathology, including thickening, lumen stenosis, and occlusion. When the popliteal artery is complicated by aneurysm, ultrasound examination can clearly find local thickening and aneurysmal dilation of the artery lumen. Even if the aneurysm is full of thrombosis, the aneurysm diagnosis can be confirmed. Duplex ultrasound therefore only shows the morphological structure of popliteal vessels, but also provides hemodynamic information [16]. It is a non-invasive method that might prove to be of great value in the diagnosis of PVES.

Provocative maneuvers can provide important information for early diagnosis of PVES. This type of examination allows real-time, dynamic observation of changes in popliteal vessel blood flow before and after the experiment, as well as demonstrates that compression was not detected in the neutral position [4, 17]. The presence of entrapment should be highly suspected when the popliteal artery or vein are compressed. The popliteal vessels usually show signs of narrowing or lumen occlusion and have positive peak systolic velocity results. Therefore, close attention must be paid to the examination of plantar flexion. Three examined cases had no abnormality in the neutral position, but popliteal vessels appeared to be narrowed or occluded during the provocative maneuvers, which was helpful for clinicians to

consider the possibility of PVES. It is worth noting that for patients with complete popliteal artery occlusion, there is no blood flow signal in the popliteal artery during color Doppler examination. Thus, provocative maneuver test is applicable to patients without popliteal artery occlusion.

However, in two cases no popliteal fossa malformations were detected by ultrasound and the imaging was not specific. In this group, there were two cases with lower extremity atherosclerosis, with one 58-year-old male with history of left lower limb paresthesia for three years and claudication for four months. Ultrasound imaging found left lower limb artery atherosclerotic plaques and popliteal artery thrombosis. This patient underwent computed tomographic angiography, which showed that the left popliteal artery was entrapped by the medial head of the gastrocnemius which rose more laterally than the normal. Another case included a 55-year-old male who complained about symptoms of lower limb ischemia, such as swelling of the right leg and cold right foot for six months. Ultrasound suggested atherosclerosis of right lower extremities. The popliteal artery and tibiofibular stem Doppler blood flow velocity was decreased. The operation demonstrated compression of the right popliteal artery with a fibrous band that attached to the intercondylar notch after crossing the popliteal artery. Ultrasound examination in the above two patients only diagnosed lower limb atherosclerosis and failed to find popliteal vascular depression. Thus, even though ultrasound scanning can show arterial lesions in older patients affected by atherosclerosis, pathological and etiological analysis still needs to combine clinical data and other imaging examinations.

Currently, it is believed that patients with PVES should undergo a surgical treatment as soon as they are diagnosed. The principle of surgery is that [18, 19] when the popliteal artery is not injured, the medial head of the gastrocnemius muscle causing popliteal artery constriction or abnormal accessory muscle bundles and tendons can be released. In case of popliteal artery injury, endarterectomy and venous patch repair can be used. When popliteal artery is completely occluded, great saphenous vein autograft or artificial vascular bypass can be performed [20]. Doppler ultrasound can reveal locations of the saphenous vein and provide accurate positioning for surgery (Figs. 3d and 3e). In addition, patients with popliteal artery and saphenous vein patency during postoperative follow-up can be evaluated using ultrasonography [3]. For some patients with popliteal artery compression and proximal side aneurysmal dilatation, ultrasound images can indicate the scope of aneurysms and mark them on the body surface, facilitating clinical selection of surgical methods.

Yang et al. suggested that Doppler ultrasound is a simple, rapid, non-invasive procedure that should be used routinely in the evaluation of individuals with suspected PVES [3]. However, the use of Doppler sonography is limited by its small window, operator proficiency, probe pressure, and lack of classification accuracy. Therefore, further imaging studies are required for patients with suspicion of entrapment, even if Doppler results are normal.

Limitations

The major limitations of our study were the relatively smaller sample of patients and the lack of control group. The results would be more accurate for a larger sample size, but these specific ultrasound findings provide direct evidence, we can still recommend ultrasound as a primary screening examination modality in PVES. In the absence of a healthy control group, the application value of provocative maneuvers in PVES may not be adequately evaluated. In future study, we will continue to collect more case and control study, and further in-depth discussion and improvement.

Conclusion

Doppler ultrasound may have a high diagnostic rate of depicting clinical characteristics in patients with abnormal peripheral artery disease. PVES can be diagnosed by ultrasonography if the distance between the popliteal artery and vein is widened and medial displacement of the popliteal artery and abnormal muscle tissue between the blood vessels are present. It is a non-invasive and simple peripheral vascular examination tool. Provocative maneuvers allow direct observation of the popliteal vessels while they become entrapped. Simultaneous monitoring of the effects on vascular blood flow may help diagnose the syndrome. In conclusion, these results suggest that Doppler ultrasound can be used as a primary screening examination modality in PVES.

Declarations

Ethics approval and consent to participate

The retrospective study has been approved by the institutional ethics committee of the Second Hospital of Shandong University (KYLL-2019(LW)025), and the need for informed consent was waived.

Consent for publication

Consent for publication was obtained from all the authors.

Availability of data and materials

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by the youth talent fund of the second hospital of Shandong University (2018YT04) and the Natural Science Foundation of Shandong University of Traditional Chinese Medicine (2018yq08).

Authors' contributions

Zhong H and Shao GR : study conception and design, data collection; Zhong H: data collection, data analysis, interpretation of data, drafting of the manuscript, approval of article; Zhao Y: data collection, critical revision of the manuscript, approval of article. Qi HT and Yang WB: interpretation of data, critical revision of the manuscript. Wang MH: data collection, critical revision of the manuscript, approval of article.

Acknowledgements

We thank International Science Editing (<http://www.internationalscienceediting.com>) for editing this manuscript.

Conflict of interest:

none declared

References

1. Rich NM, Collins GJ, McDonald PT, Kozloff L, Clagett GP, Collins JT. Popliteal vascular entrapment. Its increasing interest. *Arch Surg.* 1979;114:1377–84.
2. Levien LJ, Veller MG. Popliteal artery entrapment syndrome: more common than previously recognized. *J Vasc Surg.* 1999;30:587–98.
3. Yang L, Yan S, Xiangyu H, Qianxiang K, Yishi Z, Junxue W, et al. Imaging Diagnosis and Surgical Treatment of Popliteal Artery Entrapment Syndrome: A Single-Center Experience. *ANN Vasc Surg.* 2014;28:330–7.
4. Altintas U, Helgstrand UV, Hansen MA, Stentzer KF, Schroeder TV, Eiberg JP. Popliteal artery entrapment syndrome: ultrasound imaging, intraoperative findings, and clinical outcome. *Vasc Endovasc Surg.* 2013;47:513–8.
5. Williams C, Kennedy D, Bastian-Jordan M, Hislop M, Cramp B, Dhupelia S. A new diagnostic approach to popliteal artery entrapment syndrome. *J Med Radiat Sci.* 2015;62:226–9.
6. Whelan TJ, Haimovivi H. *Vascular surgery: principles and techniques.* 2nd ed. New York: McGraw-Hill; 1984. 557 – 67.

7. Gerkin TM, Beebe HG, Williams DM, Bloom JR, Wakefield TW. Popliteal vein entrapment presenting as deep venous thrombosis and chronic venous insufficiency. *J Vasc Surg.* 1993;18:760–6.
8. Andrikopoulos V, Papacharalambous G, Antoniou I, Panoussis P. The popliteal artery entrapment syndrome. *Vasc Endovasc Surg.* 1999;33:357–65.
9. Hai Z, Guangrui S, Yuan Z, et al. CT angiography and MRI in patients with popliteal artery entrapment syndrome. *AJR Am J Roentgenol.* 2008;191:1760–6.
10. Karacagil S, Lofberg AM, Granbo A, et al. Value of duplex scanning in evaluation of crural and foot arteries in limbs with severe lower limb ischaemia: a prospective comparison with angiography. *Eur J Vasc Endovasc Surg.* 1996;12:300–3.
11. Flanigan DP, Ballard JL, Robinson D, Galliano M, Blecker G, Harward TR. Duplex ultrasound of the superficial femoral artery is a better screening tool than ankle-brachial index to identify at risk patients with lower extremity atherosclerosis. *J Vasc Surg.* 2008;47:789–92.
12. Mohler ER 3rd, Bundens W, Denenberg J, Medenilla E, Hiatt WR, Criqui MH. Progression of asymptomatic peripheral artery disease over 1 year. *Vasc Med.* 2012;17:10–6.
13. Miao Z, Chuang C, Qianyi Q, Changjun W. Ultrasound in diagnosis of anatomical variation of anterior and posterior tibial arteries. *Med Ultrason.* 2016;18:64–9.
14. Collins PS, McDonald PT, Lim RC. Popliteal artery entrapment: an evolving syndrome. *J Vasc Surg.* 1989;10:484–90.
15. Rosset E, Hartung O, Brunet C, Roche PH, Magnan PE, Mathieu JP, et al. Popliteal artery entrapment syndrome: anatomic and embryologic bases, diagnostic and therapeutic considerations following a series of 15 cases with a review of the literature. *Surg Radiol Anat.* 1995;17:161–9.
16. Crawford JD, Perrone KH, Jung E, Mitchell EL, Landry GJ, Moneta GL. Arterial duplex for diagnosis of peripheral arterial emboli. *J Vasc Surg.* 2016;64:1351–6.
17. Alvarez Rey I, Alvarez Rey G, Alvero Cruz JR, Jimenez Diaz JF, Alvarez Bustos G. Popliteal artery entrapment syndrome in an elite rower: sonographic appearances. *J Ultrasound Med.* 2004;23:1667–74.
18. di Marzo L, Cavallaro A, Sciacca V, Mingoli A, Tamburelli A. Surgical treatment of popliteal artery entrapment syndrome: a ten-year experience. *Eur J Vasc Surg.* 1991;5:59–64.
19. Radonić V, Koplić S, Giunio L, Bozić I, Masković J, Buća A. Popliteal artery entrapment syndrome: diagnosis and management, with report of three cases. *Tex Heart Inst J.* 2000;27:3–13.
20. Lavingia KS, Dua A, Rothenberg KA, Fredericson M, Lee JT. Surgical management of functional popliteal entrapment syndrome in athletes. *J Vasc Surg.* 2019;70:1555–62.

Figures




Figure 1

Normal popliteal artery, popliteal vein, and tibial nerve alignment. a Sagittal ultrasound position shows that popliteal artery (red) and popliteal vein (blue) are in normal anteroposterior relationship. b Transverse position shows sequence of anterior and posterior movements of tibial nerve (arrow), popliteal vein (POV), and popliteal artery (POA). c Drawings illustrate normal popliteal vascular relationship (posterior view): front and back arrangement, followed by the tibial nerve (green), popliteal vein (blue), and popliteal artery (red) from back to front




Figure 2

Left popliteal vascular entrapment syndrome in 22-year-old man who presented with intermittent claudication of left calf for three months. a Color Doppler images for the popliteal artery (POA) and popliteal vein (POV) widening and moving from normal forward and backward orientations to left and right orientations. Left popliteal artery was occluded with hypoechoic thrombosis and no flow signal filling in the lumen (arrow). b Proximal diameter of the left popliteal artery is within the normal range and lumen diameter of the middle and far popliteal arteries is significantly thinner (arrow).




Figure 3

Left popliteal vascular entrapment syndrome in a 17-year-old man who presented with intermittent claudication and paresthesia of left lower extremity after exercise can be relieved after rest (PVES type II). a Color Doppler images show widening of distance between the popliteal artery (POA) and popliteal vein (POV), changing from normal forward and backward alignment to abnormal left and right alignment. b Proximal part of popliteal artery blood flow signal shows fullness and integrity, where a branch (arrow) of the popliteal artery blood vessel is detected. c Lumen diameter of the popliteal artery in middle and far sections is significantly thinner and there is no blood flow signal in the lumen. d and e Ultrasonic images show that blood flow signal in the left great saphenous vein (LGSV) is complete and without obvious abnormalities. f Operation photography of left knee shows that the medial head of left gastrocnemius

muscle (star) shifts outwards and downwards and lies between the popliteal artery (1) and popliteal vein (2).




Figure 4

Left popliteal artery entrapment syndrome in 43-year-old man who presented with intermittent claudication of left calf and left foot coldness for three months (PVES type III). a Ultrasound images show left the popliteal artery (LPOA) and left popliteal vein (LOPV) widening. Left popliteal artery is entrapped by aberrant accessory muscle slip (arrow) of the medial head of left gastrocnemius muscle, which passes between the popliteal artery and vein. b Left popliteal artery occlusion due to thrombosis. Color Doppler image shows that lumen is filled with hypoechoic thrombosis (arrow) and that popliteal artery was not filled with blood. c Operation photography demonstrates popliteal artery (1) is entrapped by aberrant accessory muscle slip (dotted line) of the medial head of gastrocnemius muscle (star) that passes between the popliteal artery (1) and vein (2).




Figure 5

Right popliteal vascular entrapment syndrome in a 34-year-old man who presented with intermittent claudication of right calf for six months (PVES type II). a Ultrasonic longitudinal section image shows the right popliteal artery (arrow) is entrapped by the laterally originated medial head of gastrocnemius muscle (star). b Color Doppler image shows segmental stenosis of right popliteal artery (arrow) not filling with blood. c Note that due to thrombosis (arrow), lumen is filled with medially echoic thrombosis and middle and distal segments of popliteal artery are occluded. d Operation photography reveals the right popliteal artery (1) is entrapped by medial head of gastrocnemius muscle (star) that passes between the popliteal artery (1) and vein (2).




Figure 6

Right popliteal artery entrapment syndrome in 18-year-old man who presented with calf swelling and foot coldness in right lower extremity after physical activity. a Color Doppler image of right foot in neutral position reveals right popliteal artery with normal flow. b Color Doppler image of right foot in plantar flexion position shows segmental stenosis of right popliteal artery. Right popliteal artery is entrapped by medial head of gastrocnemius muscle (star). c Doppler spectrum shows that peak systolic blood velocity increased significantly and more than doubled compared to its value prior to provocative maneuvers.