

# Women's Experiences of Safety Apps for Sexualized Violence: A Narrative Scoping Review

Nicole Doria (✉ [nicole.doria@dal.ca](mailto:nicole.doria@dal.ca))

Dalhousie University <https://orcid.org/0000-0003-4786-997X>

Christine Ausman

Dalhousie University

Susan Wilson

Dalhousie University

Annalisa Consalvo

Dalhousie University

Jad Sinno

Dalhousie University

Matthew Numer

Dalhousie University

---

## Research article

**Keywords:** Sexualized Violence, Safety Apps, Women's Health, Health Promotion, Global Health, Narrative Scoping Review, Qualitative Research

**Posted Date:** June 2nd, 2020

**DOI:** <https://doi.org/10.21203/rs.3.rs-30829/v1>

**License:** © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

---

**Version of Record:** A version of this preprint was published at BMC Public Health on December 1st, 2021. See the published version at <https://doi.org/10.1186/s12889-021-12292-5>.

## Abstract

## Background

Sexualized violence against women is a significant human rights problem worldwide. Safety apps have the capacity to provide women with resources to prevent or respond to experiences of sexualized violence. The aim of the following study was to review the scope of the literature on women's experiences of safety apps related to sexualized violence and identify if there is a current gap in this literature.

## Methods

This scoping review employed a systematic methodology guided by the Arksey and O'Malley framework. The search for this scoping review was conducted in January 2020 and four electronic databases were searched: Embase, MEDLINE, PsycINFO, and Scopus. Reference lists of systematic reviews, scoping reviews, and literature reviews that were found through the search of databases were also checked to ensure all relevant studies had been screened. Last, the table of contents for the last five years (2015–2020) were hand searched in four key journals (Journal of Technology in Human Services, Violence Against Women, BMC Public Health, and Health Promotion International) to identify any articles that may have been missed in the database search. Covidence was used to complete the screening. All data that met inclusion criteria was charted, extracted and synthesized.

## Results

Across the four databases searched, a total of 389 studies were identified; 127 duplicates were removed, resulting in 262 studies screened. In total, 6 studies were included from the search of databases and 1 study was included from hand-searching, for a total of 7 included studies. Thematic analysis identified the following themes in the literature, which are summarized narratively: (1) security; (2) accessibility; and (3) knowledge. Recommendations and barriers found in relation to each theme are also presented.

## Conclusions

This review confirms that the qualitative literature on women's experiences of sexualized violence safety apps is scarce and exposes that a gap in the literature on this topic. The gaps in the literature were identified and implications and recommendations for future research are discussed.

## Background

The high rate of sexualized violence against women is an urgent public health issue and a significant human rights problem (1). Although this is well known, global prevalence of sexualized violence is difficult to determine as there is not a universal understanding of what constitutes sexualized violence (2). Sexualized violence is a broad term that encompasses much more than sexual assault and rape. Sexualized violence includes any unwanted sexual contact that targets sexuality and is physical and/or psychological in nature, including but not limited to: sexual abuse, sexual assault, rape, sexual harassment, stalking, indecent/sexualized exposure, degrading sexual imagery, sharing sexual photographs without permission, and/or unwanted comments or jokes (3, 4). Further complicating our understanding of sexualized violence is underreporting. For example, only an estimated 32–35% of incidents of sexualized violence in the United States are reported to police (5, 6). Canadian data suggests even lower reporting rates of sexual assault, with only 5% reported to police (7, 8).

Although anyone can experience sexualized violence, women are primarily the victims and are most likely to experience more severe forms of violence, whereas men are typically the perpetrators (9, 10). Studies suggest that individuals who are young, female, Indigenous North American, and/or a gender or sexual minority are also more likely to experience sexualized violence (7, 11, 12). Further, a 2018 review of the literature on technology-facilitated sexualized violence reported that women, as well as sexual and gender minorities, are at an increased risk of being the targets of online abuse (13). These findings suggest that prevention and intervention efforts must target women, as well as marginalized populations, but further research is needed in this area (13, 14).

## Health Outcomes And Traditional Resources

The negative health outcomes associated with physical sexual assault are well known and include poorer self-rated health, exacerbated psychological distress, and post-traumatic stress disorder (PTSD) (15–20). Literature that focuses on the health outcomes associated with other forms of sexualized violence, however, is scarce (21–23). A recent Australian study found that women's experiences of sexualized violence were most commonly public harassment, flashing, unwanted groping, or being coerced into consenting to sex (10). These women were at an elevated risk for having anxiety, depression, and PTSD as compared to women who did not experience sexualized violence (10, 24). Negative mental health outcomes have been found to interfere with the victim's physical functioning and interpersonal relationships (25–28).

A variety of prevention initiatives, supports, and resources exist that aim to increase safety for women, reduce incidents of sexualized violence, and help offset worsened health outcomes. Some examples include educational campaigns, medico-legal services, specific training for healthcare professionals, prophylaxis for HIV infection, and centres that provide trauma-informed care (29). Unfortunately, these initiatives and resources are often limited to high-income and resource-rich settings (4). There is reason to believe that current measures are insufficient given that rates of sexualized violence have remained constant since the early 2000s (7, 30). In the majority of countries with available data, less than 40% of women who experienced violence sought help of any sort (31). Among women who did seek help, most reached out to family and friends instead of formal institutions, and less than 10% of women sought help from the

police (31). These statistics demonstrate a need for safe, accessible, affordable, and comprehensive resources that aim to prevent sexualized violence against women and support women who have experienced sexualized violence.

## Mobile Technology And Safety Apps

In today's digital age, mobile technology presents a unique opportunity to deliver improved sexualized violence interventions and resources (32). Research shows that 95% of adults between the ages of 18 and 34 in the United States own a mobile phone (33). Further, mobile technologies as sophisticated as smartphones are widely used around the world: 95% in South Korea, 88% in Israel, 81% in Australia, 80% in Spain, with a median worldwide ownership of smartphone technology of 76% (34). The extant literature has primarily focused on the ways that mobile apps can be used for health and fitness (35–37). There has recently, however, been interest in discovering the potential of using smartphone technology to help enhance women's safety.

A recent review on the role of emerging technologies regarding women's safety found that smartphones were the most commonly discussed protective technology in the literature (37.2% mobile phones, 18.8% apps) (38). Smartphones can be used to call for help in emergencies, film violence incidents/injuries, retain threatening text messages, and use apps to contact support networks (38). Another study found that 62.9% of college students would consider downloading a personal safety app, and even more students would consider doing so if the app contained a tracking feature (39). Women were also found to be more likely than men to download a safety app with tracking features to increase their sense of security and reassurance (39). Overall, the proliferation of technological solutions for sexualized violence such as signal-/alarm-emitting wearables and apps have surged in recent years (40).

Specific to sexualized violence, qualitative evidence shows that technology can provide access to information and services, strengthen support networks, and support sexualized violence victims in their safety and escape planning (41). An Australian review found that safety apps for public stranger violence focused mainly on location-based services, personal alarms, and crowd-sourced data (42). This review identified *MySafetyPin*, *Saven*, and *My Keeper* as useful safety apps allowing users to identify dangerous areas, and the *Women Safety Totem SOS* app for providing information to help reduce the likelihood of being a target for violence and tips on how to handle violence (e.g., self-defence information) (42). Although there are several studies that have explored safety apps regarding sexualized violence, the findings regarding women's experiences and perceptions of these apps has not been comprehensively reviewed. Therefore, the research question for this scoping review was: What is the scope of the literature on women's experiences of safety apps in relation to sexualized violence?

## Method

A scoping review method was chosen to comprehensively review the published literature on women's experiences using safety apps related to sexualized violence (43). The aim was to identify if the evidence about women's experiences using these apps presents an existing gap in the literature. This scoping review employed a systematic methodology guided by the Arksey and O'Malley (2005) framework(43). This framework includes identifying the research question, identifying relevant studies, selecting relevant studies, charting the data, and collating and summarizing the findings(43).

## Identifying Relevant Studies

The literature search for this scoping review was conducted in January 2020. The following electronic databases were searched: Embase, MEDLINE, PsycINFO, and Scopus. Our search strategy was developed in consultation with a medical librarian at the Maritime SPOR SUPPORT Unit and can be seen in Table 1. The final search strategy was first used on the MEDLINE database and then applied to the other databases (Embase, PsycINFO, and Scopus). Reference lists of systematic reviews, scoping reviews, and literature reviews that were found through the search of databases were also checked to ensure all relevant studies had been screened. Last, four key journals were identified for hand-searching (*Journal of Technology in Human Services*, *Violence Against Women*, *BMC Public Health*, and *Health Promotion International*). The table of contents for the last five years (2015–2020) were searched for each journal to identify any articles that may have been missed in the database search.

Table 1  
Search Strategy Used to Search Databases

1	exp Intimate Partner Violence/
2	exp Sex Offenses/
3	(anti-abuse or anti-assault or anti-harassment or anti-rape or anti-victim* or anti-violence).ti,ab,kw,kf.
4	((dating or domestic or gender* or partner* or relationship* or wom#n or sex*) adj2 (abuse* or assault* or violence)).ti,ab,kw,kf.
5	(intimate partner violence or ipv).ti,ab,kw,kf.
6	(rape or rapes or raped or rapist* or raping).ti,ab,kw,kf.
7	(sex* adj2 coerc*).ti,ab,kw,kf.
8	(sex* adj2 (force* or unwanted or unwelcome)).ti,ab,kw,kf.
9	(sex* adj2 harass*).ti,ab,kw,kf.
10	(sex* adj2 victimi*).ti,ab,kw,kf.
11	(unwanted pursuit or unwanted online pursuit).ti,ab,kw,kf.
12	(wom#n adj4 (safety or security)).ti,ab,kw,kf.
13	or/1-12
14	Mobile Applications/
15	(app or apps).ti,ab,kw,kf.
16	(application* adj4 (android or cell* or iphone* or mobile or smart phone* or smartphone*)).ti,ab,kw,kf.
17	or/14-16
18	13 and 17

## Selecting Relevant Studies

All of the peer-reviewed articles from the database search were imported into Covidence (an online software tool for review management) for organization and screening. Inclusion and exclusion criteria were established for selecting relevant studies prior to screening began (Table 2). Our definition of sexualized violence used was broad to include any unwanted sexual contact that targets sexuality and is physical and/or psychological in nature, including but not limited to: sexual abuse, sexual assault, rape, sexual harassment, stalking, indecent/sexualized exposure, degrading sexual imagery, sharing sexual photographs without permission, and/or unwanted comments or jokes (3, 4).

Table 2  
Inclusion and Exclusion Criteria for Selected Articles

Inclusion Criteria	Exclusion Criteria
<ul style="list-style-type: none"> <li>· Peer-reviewed, primary research.</li> <li>· Safety apps related to sexualized violence: any unwanted sexual contact that targets sexuality and is physical and/or psychological in nature.</li> <li>· Qualitative findings (including qualitative components of mixed methods studies).</li> <li>· Published in English or French.</li> <li>· Focused on adult women's (&gt;18 y/o) experiences of sexualized violence focused safety apps.</li> <li>· All dates and all countries.</li> </ul>	<ul style="list-style-type: none"> <li>· Not peer-reviewed, primary research.</li> <li>· Safety apps not related to sexualized violence.</li> <li>· Quantitative findings.</li> <li>· Not published in English or French.</li> <li>· Data that did not focus on adult women's (&lt;18 y/o) experiences (only men or not able to extract data on women).</li> <li>· Literature that did not include empirical data (commentaries, editorials, book reviews).</li> <li>· Literature/systematic reviews.</li> <li>· Conference proceedings.</li> </ul>

We sought peer-reviewed studies that collected primary qualitative data as we were only interested in identifying women's experiences for the purposes of this scoping review, which cannot be well captured by quantitative data (44). Literature reviews, systematic reviews, conference proceedings, and literature that did not include empirical data (commentaries, editorials, book reviews) were excluded. Mixed-methods studies were reviewed, however, only the qualitative components were considered and included if relevant. Sexualized violence interventions and resources that are meant for children or teenage girls are typically designed to meet the unique needs of minors, which differ from the needs of adults (45). Given that our research question was interested in understanding the experiences of adult women, inclusion criteria was limited to 18 years of age or older; if an age range was not included in the study or the age range included any participants that were 17 years of age or younger, the study was excluded. Studies that had male participants were included as long as the data on

women could be extracted. Studies conducted worldwide that were published in English or French were also included and all publication dates were considered for inclusion.

Four reviewers (AC, CA, ND, SW) independently screened all articles at the title/abstract stage and full-text stage in accordance with the inclusion/exclusion criteria (each study in Covidence requires the vote of two reviewers). If voting conflicts could not be resolved by the original two reviewers, a final decision was made in collaboration with the full team. If relevance of a study could not be determined at the title/abstract stage, it was voted forward to be reviewed at the full-text stage. Included full-text articles were obtained through online access, Dalhousie University library services, or Dalhousie University document delivery service. If full-text articles could not be obtained through these means, they were excluded. The same process was followed for hand searching.

## **Data Charting, Extraction And Synthesis**

The following data was extracted and charted from each of the included studies by ND: author/year, title, country, purpose, participants, research method, sexualized violence focus, safety app, key findings. To ensure rigour and accuracy, a second reviewer (CA) reviewed and confirmed all extracted data. Thematic analysis guided by Braun and Clarke (2006) was employed to identify emerging themes. Each reviewer (AC, CA, ND, SW) familiarized themselves with the included studies, generated initial codes, and searched for themes. Coding and preliminary themes were compared and discussed by all reviewers, and final themes were defined and named.

## **Results**

Across the four databases searched, a total of 389 studies were identified; 127 duplicates were removed, resulting in 262 studies screened. Screening at the title and abstract level resulted in the exclusion of 182 studies. There were 80 studies screened at the full-text stage, with 74 studies being excluded – the majority (60) for wrong study design. In total, 6 studies were included from the search of databases and 1 study was included from hand-searching, for a total of 7 included studies (see Fig. 1 for PRISMA diagram). Reference lists of systematic reviews, scoping reviews, and literature reviews that were found through the search of databases were also checked to ensure all relevant studies had been screened. This process did not identify any new studies.

## **Narrative Summary Of Themes**

Using thematic analysis, three themes emerged that impacted women's experiences of using sexualized violence safety apps: (1) security; (2) accessibility; and (3) knowledge. Recommendations and barriers found in relation to each theme are presented as subthemes. Table 4 offers a summary of each theme.

Table 3  
Summary of Included Studies

Author, Year	Title	Country	Purpose	Participants	Research Method	Sexualized Violence Focus	Safety App
Alhusen et al., 2015	Development of the MyPlan safety decision app with friends of college women in abusive dating relationships	United States	To explore the perceptions of friends of dating violence survivors regarding the benefits of a safety decision aid, deployed through a smart phone application prototype, for friends of female survivors of dating violence.	Thirty-one college students who self-reported having a friend who had experienced dating violence while in college. Participants were English-speaking male and female college students, aged 18-24 ( <i>Mean</i> = 20.84). The majority of participants were female ( <i>n</i> = 25) and the remainder were male. Most of the participants were White ( <i>n</i> = 16), followed by African American ( <i>n</i> = 8); the remaining participants were from a variety of ethno-racial identities.	Qualitative - Focus groups/interview. Each of the focus group discussions lasted 60–90min and was cofacilitated by two trained research assistants. The individual in-depth interviews lasted approximately 60min. The procedures implemented were consistent across the focus groups and individual interviews. At the beginning of each focus group/interview, participants were instructed to progress through the app prototype preloaded onto an iPod touch. Participants also had access to the app throughout the focus group/interview. The emphasis of the interviews was on the friend's assessment of the app. The audio-taped interviews were digitally recorded and then transcribed.	Dating violence (DV)	MyPlan – a prototype smart phone application (app) that is a safety decision aid designed to assist college women (age 18–24) experiencing dating violence/survivors of dating violence and their friends who wish to learn more about how to help them. A collaborative, multistate research team partnered with the One Love Foundation, a national relationship violence prevention advocacy organization, to develop the app.
Blayney et al., 2018	Enlisting friends to reduce sexual victimization risk: There's an app for that... but nobody uses it	United States	To collect feasibility and acceptability information on the Co6 app among college women who drink alcohol, a group at greater risk for sexualized violence, to shed light on the Co6 app and the challenges associated with app-based prevention in real-world contexts.	Forty-four college women. Women had to 18–24 years of age ( <i>Mean</i> = 20.11, <i>SD</i> = 1.33), be enrolled in college, own a smart phone, and drink alcohol at least once per week in the last 6 months. A majority of participants were White ( <i>n</i> = 23), followed by African American ( <i>n</i> = 7), Asian ( <i>n</i> = 6), Hispanic ( <i>n</i> = 3), and other ( <i>n</i> = 5). Almost all of the participants identified as heterosexual ( <i>n</i> = 41)	Mixed Methods – Participants completed questionnaires, used the Co6 app for 2 months, and returned to report their experiences. For the qualitative component, participants were interviewed individually in a semi-structured format about what they liked and did not like about the app. Follow up interviews were approximately 1 hour long and were audio-recorded and transcribed verbatim.	Sexual victimization	Circle of Six (Co6 app) - centralizes both personal and community resources to reduce SV risk. Specifically, the app calls for users to program the contact information of six trusted individuals, who are then identified as part of the user's safety network (i.e., circle of six).
Bloom et al., 2016	Developing an App for College Women in Abusive Same-Sex	United States	To establish initial content validity, feasibility, appropriateness, understandability,	Thirteen participants participated in interviews. These included eight	Qualitative - Interviews using a semi-structured interview guide with questions regarding understandability,	Same-sex dating violence	An interactive, personalized safety decision aid smartphone application (app) developed by a collaborative, multistate team. The app was

	Relationships and Their Friends		and usability of a smartphone-based safety decision aid app for college women in same-sex relationships and their friends.	college students (four female survivors, three female friends, and one male friend), five of whom were White and three African American, with a mean age 22.0 years, SD 1.9. Five college staff who worked directly with LGBT survivors on campus also participated (four female, one male; all White, mean age 28.2, SD 3.6) in the study.	appropriateness, comprehensiveness, and usefulness of the app for women in same-sex relationships and their friends. Interviews were audio-recorded.		intended to allow abused college-aged women and their friends to privately and safely assess violence severity in an abusive relationship, clarify their areas of decisional conflict, (e.g., advantages/disadvantages of the relationship) and identify their safety priorities and link to national resources (e.g., national hotline).
Gilmore et al., 2019	Usability testing of a mobile health intervention to address acute care needs after sexual assault	United States	To test the usability of a mobile health intervention targeting alcohol and drug misuse, suicide prevention, posttraumatic stress symptoms, coping skills, and referral to formal assistance for individuals who have experienced sexual assault.	Thirteen participants ( <i>Mage</i> = 28.00) who experienced sexual assault and received a sexual assault medical forensic examination. Most participants identified as white ( <i>n</i> = 13), female ( <i>n</i> = 11), and were single ( <i>n</i> = 7). One person identified as male and another as "other". Approximately two-thirds of participants were not in college ( <i>n</i> = 10) and had medical insurance ( <i>n</i> = 10). The average length of time since the sexual assault was 12.09 months. The assaults were perpetrated by an acquaintance ( <i>n</i> = 7), stranger ( <i>n</i> = 5), and partner ( <i>n</i> = 1).	Mixed Methods- The qualitative component consisted of individual interviews that were conducted in-person or through teleconferencing, according to participant preference, and lasted 45 to 60 minutes.	Sexual Assault	<i>SC-Safe</i> - a resource designed for individuals over the age of 18 residing in South Carolina who have experienced sexual assault. It was designed by the first and second authors to address a gap in clinical services after recent sexual assault.
Lindsay	Survivor	United	For young	Thirty-four	Qualitative- Ten	Dating	A prototype smart phone

et al., 2013	feedback on a safety decision aid smartphone application for college-age women in abusive relationships	States	women who had previously experienced dating violence to evaluate a mobile phone safety decision aid prototype, which was designed for use by college-age women experiencing dating violence.	English-speaking female college students, ages 18–25 ( <i>Mage</i> = 21.26, <i>SD</i> = 1.86), who reported that they experienced dating violence while in college. Self-identified ethno-racial background were 52.6% White, 23.7% Hispanic, 7.9% African America, 13.2% Multiracial, and 2.6% other. Nearly 16% of participants reported being in a previous relationship with an abusive female partner.	focus group (ranging from 2 to 7 participants) discussions, each lasting approximately 90 min, were cofacilitated by two trained research assistants in a campus or community location. Individual interviews took place in a setting of the participant's choosing, were approximately 60 min, and were conducted by one trained research assistant. Procedures were consistent across the focus group sessions and individual interviews and a semi-structured interview guide was utilized.	Violence/Intimate Partner Violence	application (app) that is a safety decision aid designed to assist college women (age 18–24) experiencing dating violence/survivors of dating violence and their friends who wish to learn more about how to help them. A collaborative, multistate research team partnered with the One Love Foundation, a national relationship violence prevention advocacy organization, to develop the app.
Ragavan et al., 2019	Thrive: A Novel Health Education Mobile Application for Mothers Who Have Experienced Intimate Partner Violence	United States	Describe the development and formative evaluation of a trauma-informed, user-friendly Smartphone-based mobile application to address the unmet health needs and improve the well-being of mothers who have experienced intimate partner violence (IPV).	Eight IPV survivors and 16 hospital-based staff (nine health care providers, four social workers, one mental health provider, and three IPV advocates; hereafter called providers).	Formative Evaluation - Participants were instructed to use <i>Thrive</i> on a study Smartphone for 10 to 20 minutes and then provide feedback about its content, design, safety features, and applicability via a structured interview.	Intimate Partner Violence	Thrive - a trauma-informed, user-friendly Smartphone based mobile application (app) to address the unmet health needs and improve the well-being of mothers who have experienced IPV. A multidisciplinary team of IPV experts developed the app in partnership with software developers.
Tarzia et al., 2017	"Technology Doesn't Judge You": Young Australian Women's Views on Using the Internet and Smartphones to Address Intimate Partner Violence	Australia	To confirm the hypothesis that technology has a potential role in responding to IPV, and to ascertain what factors might encourage or discourage women from using an IPV website or app.	Nineteen women between 20 and 25 years of age. All participants were residing in Victoria, Australia at the time of the study, and all had self-reported experiencing fear of a partner in the previous 6 months. None of the women	Qualitative- Four focus groups were held at The University of Melbourne between April and August 2014. The sessions were informal and semi structured in nature, and facilitated by a trained researcher. An additional note taker was present but did not take part in the conversation. The discussions lasted approximately 60	Intimate Partner Violence	Not specified – general exploration of safety apps

were married at the time of participation, and most were tertiary educated.

min each and were audio recorded and later transcribed verbatim by members of the research team.

Table 4  
Summary of the Themes

Themes	Specific Aspects	Sources	Sample Quotations
Security	Privacy, judgement, stigma	Blayney et al. (2018); Bloom et al. (2016); Gilmore et al. (2019); Lindsay et al. (2013); Ragavan et al. (2019); Tarzia et al. (2017)	<i>The option of ... being able to maybe correspond with people anonymously, especially if you're scared of being judged or found out ... that'd be really good.</i> (Tarzia et al., 2017: p. 205)
Accessibility	Ease and usefulness	Alhusen et al. (2015); Blayney et al. (2018); Bloom et al. (2016); Gilmore et al. (2019); Lindsay et al. (2013); Ragavan et al. (2019); Tarzia et al. (2017)	<i>It helps you organize your mind because when you're in the situation, you don't really know how to feel ... there's so much going on, you don't really know how to categorize things. When [the app does] it for you, it just helps you put yourself in order, and have more control on your life. When something's happening to you like that, you feel like you're out of control and you can't—you don't know where to go. You don't know what to do. You're just so confused, so I think it helps.</i> (Lindsay et al., 2013; p. 378)
Knowledge	Information, awareness, validation, myth debunking	Alhusen et al. (2015); Bloom et al. (2016); Gilmore et al. (2018); Lindsay et al. (2013); Ragavan et al. (2019); Tarzia et al. (2017)	<i>I liked the concrete advice...it didn't just say "talk to your friend"—rather "here are five different things you can say." I feel like everyone knows you have to talk to the person but people don't know what to say and how to say it, and that was very, very useful.</i> (Alhusen et al., 2015; p. 274)

## Security

Security was found to be a prevalent theme across studies that influenced women's experiences of using safety apps (46–52). Specific aspects of security that were discussed included privacy, judgement, and stigma. All of the studies identified that using safety apps to obtain information provided more privacy or anonymity than obtaining information from in-person health services (46–52). Using a safety app was also found to provide additional privacy because individuals could access them discreetly and apps were designed to be ambiguous to others (49, 51, 52). For example, using basic colours for the app's design so that it would not be recognized as a sexualized violence related app by another person who might see the individual's phone.

Four studies noted that participants experienced a greater sense of security because they faced less judgment regarding being a victim of sexualized violence and, therefore, avoided stigma when using a safety app (47–49, 51). Using a safety app was found to be more objective and unbiased than accessing traditional health services or speaking with friends and family (48, 49, 51). Five of the seven studies also acknowledged that individuals found it was easier to interact with an app than having to discuss their situation with health professionals or their social network (46–49, 51). Women perceived less stigma about having experienced sexualized violence as a result of using these apps (47–49, 51). One participant stated:

With suicide already being stigmatized the way it is, and communication about suicide being the way it is, I would want to know that like, it's okay to talk about this and it's okay if this is what you're feeling like (Gilmore et al., 2019, p. 10). (51)

Many participants mentioned the option of not having to discuss sexualized violence in a traditional way such as "face-to-face" as a benefit (46–49, 51). One participant noted, "it [app] gives you a privacy and accessibility. ... the fact that I don't have to go to Student Health Center to get help and not have to worry about being judged" (Alhusen et al., 2015, p. 276) (46). Another participant discussed the benefit of accessing help via an app instead of going to a counseling center on campus, noting: "I feel judged to go. ... just knowing that I'm going to see them every day since I live there [on campus] I would feel a little uncomfortable" (Alhusen et al., 2015, p. 276) (46). Interestingly, several studies identified that it would be beneficial to be able to engage with others through

the app (48, 49, 52), stating that “the option of ... being able to maybe correspond with people anonymously, especially if you’re scared of being judged or found out ... that’d be really good” (Tarzia, Iyer, et al., 2017, p. 205) (49).

**Barriers and recommendations.** The greatest barrier that emerged in relation to security was privacy, including the potential for partner monitoring or surveillance which may limit use of the app (47–49, 51). For example, one participant commented: “If somebody’s in a relationship that is abusive, and someone’s already checking their phone and checking everything they’re doing, and they have an app about this on their phone, it might cause issues” (Lindsay et al., 2013, p. 382) (48). Recommendations to address privacy concerns, some of which were already part of the apps studied, included providing password protection for the app (46, 48, 49, 52), an emergency exit on each screen (47, 52), use of an innocuous name for the app that does not refer to relationships or safety (46–49, 52), and even allowing the user to rename the program or change the icon themselves (47).

Bloom et al. (2016) suggested the ability for users to print or e-mail the contents of the app to themselves or another person and then delete the app or the answers as a useful alternative(47). This would circumvent the need for the resources on the app to be exclusively on a computer or smartphone(47). Other studies specifically recommended educating users about healthy relationship boundaries regarding technology and sharing passwords with partners and friends to better instruct users how to hide the app (e.g., bury the app in smartphone folders), and how to safely use the app if a partner monitors their phone (47, 48). Women who reviewed the Circle of 6 (Co6) app specifically identified being uncomfortable with the group messaging feature, which is customized to send messages to only select individuals that you identify as your “circle of 6” (50).

## Accessibility

The importance of accessibility emerged as a key theme that influenced experiences that women had when using sexualized violence safety apps. Accessibility included the ease and usefulness of the safety apps (46–52). Women in all studies reported that they found the safety apps to be user friendly, easy to use, and easily accessible (46–52). In six studies, the usefulness of the app was directly related to the fact that it could be used anywhere, was comprehensive in content, and all the information needed was in one place (46–49, 51, 52). For example, one participant commented:

It helps you organize your mind because when you’re in the situation, you don’t really know how to feel ... there’s so much going on, you don’t really know how to categorize things. When [the app does] it for you, it just helps you put yourself in order, and have more control on your life. When something’s happening to you like that, you feel like you’re out of control and you can’t—you don’t know where to go. You don’t know what to do. You’re just so confused, so I think it helps (Lindsay et al., 2013, p. 378). (48)

In many studies, participants found that the app could be customized or personalized to the specific user, which improved its accessibility and overall usability (46, 48, 50). Examples of personalized or customized content included safety planning suggestions (46), messages that will be sent to friends (50), and what information is presented, such as a specific risk assessment and safety plan for the user (48). Overall, most (n = 6) studies found that the accessibility of safety apps had the potential to improve safety and decrease risk (47–52).

**Barriers and recommendations.** Several studies noted suggestions to improve the accessibility of sexualized violence safety apps. Making the apps more personalized and interactive was recommended by women in five studies (47–50, 52). Examples included providing written scripts to choose from (47, 50, 51) or the ability to connect directly with a counsellor through the app (48, 49, 52). Other recommendations included making the navigation functions standard across the app and clearly indicating the purpose of each icon (51). Celebrity endorsement was also suggested as a way to promote sexualized violence safety apps, which in turn would increase a sense of accessibility for more women (49).

Alhusen et al. (2015) noted that if individuals are not ready to address the violence, then the app may be unnecessary and useless (46). One participant noted “If they’re not ready they’re not ready... don’t talk about them behind their back and don’t talk about them with others [on an app] (Alhusen et al., 2015, p. 276)” (46). Further, women in the Blayney et al. (2018) study noted the app had limited contexts for use and did not provide anything more than a mobile phone could offer, seeing the app as unnecessary(50). Feedback included:

It just generally seemed like you could do the same things without the app, because iPhones nowadays are so intricate. Like, you could click details on your messages and press ‘send location’ and type a short message. I feel like that wouldn’t take nearly as long as opening the app, clicking the button, sending the messages... It’s not really an easy way to contact friends, I think personally for me, it would just be easier to call or text them. Like it wasn’t any easier to do that [use the app] (Blayney et al., 2018, p. 771). (50)

## Knowledge

Six studies discussed the importance of knowledge in the experiences of women when using sexualized violence safety apps (46–49, 51, 52). How women experienced the knowledge provided on the safety apps was dependent on the information provided, and if the information increased awareness, validation, and myth debunking. In all six studies, participants found the safety apps to be helpful in raising awareness of sexualized violence and recognizing violent behaviour through various ways such as myth debunking (46–49, 51, 52). Additionally, six studies found the apps to be an easy way to acquire information that validated women’s experiences of violence and indicated that participants found the app provided assurance that they are not alone (47–52).

Further, six studies found that women believed the apps to contain relevant information that was credible, evidence based, and/or trauma informed (46–49, 51, 52). In relation to providing knowledgeable and credible information, one participant commented:

I liked the concrete advice...it didn’t just say “talk to your friend”— rather “here are five different things you can say.” I feel like everyone knows you have to talk to the person, but people don’t know what to say and how to say it (Alhusen et al., 2015, p. 274). (46)

Similarly, another participant commented that the app “arms you with ideas as to how to go about it [conversations] properly” (Alhusen et al., 2015; p. 276) (46).

The information, options, choices, and safety planning strategies provided on the safety apps were also found to increase a feeling of empowerment (46–49). For example, one participant commented: “[A young woman] should feel relieved. Like she is equipped to know what to do, and not lost and drowning her sorrows and burden by herself. Like someone is there to help her” (Tarzia, Iyer, et al., 2017 p. 209) (49).

**Barriers and recommendations.** Several studies noted recommendations for improving the knowledge provided on sexualized violence safety apps. Several women thought that personal anecdotes, rather than statistics about dating violence, might better help young women recognize the violence in their own relationships (47, 52). Gilmore et al. (2019) (51) and Lindsay et al. (2013) (48) found that some young women noted the desire for more information throughout the app about emotional abuse, as illustrated by the following participant: “It would be really cool if there was more stuff about emotional abuse and control because I think that is also really important” (Lindsay et al., 2013, p. 383) (48). Survivors thought information about what the police can and cannot do to assist the survivor would also be helpful because “talking to police can be kinda scary” (Lindsay et al., 2013, p. 383) (48). Last, several studies believed that sexualized violence safety apps should do more than just provide information (47–49, 51). In addition to providing information, the safety apps need to expand the ability to gain knowledge by including information for appropriate resources (47, 49, 51) and incorporating further educational modules and learning opportunities (47, 51).

## Discussion

This scoping review was conducted to explore the nature and extent of literature on women’s experiences of safety apps related to sexualized violence. To our knowledge, this is the first scoping review to explore this topic. While most safety apps are studied through quantitative measures to determine prevalence of use and downloads, as well as content creation (e.g., (39, 53), quantitative data is limited in its ability to capture the lived experiences of participants (44). The current scoping review therefore was designed to focus solely on qualitative studies. Within the literature reviewed, three common themes emerged that influenced women’s experiences of sexualized violence safety apps: security, accessibility, and knowledge. This review, however, confirms that the qualitative literature on women’s experiences of sexualized violence safety apps is scarce and exposes that a gap in the literature on this topic does indeed exist.

Of the seven studies included, the majority of the samples comprised of Caucasian/White females, and there was an under-representation of other ethno-racial groups (46–48, 50, 51). Moreover, only one study disclosed the sexual orientation of their participants, which comprised of predominantly heterosexual females (50). There was one other study that included a participant that identified as neither female nor male (51), which is insufficient to capture the experiences of gender-non-conforming and trans persons. Given the widespread evidence that members from diverse ethno-racial identities and the sexual and gender diverse community are often victims of unique forms of sexualized violence (7, 11, 13), our review has highlighted a significant gap in this literature. Future research should aim to recruit samples that are entirely, or predominantly, comprised of members of diverse communities to better understand their unique experiences and needs regarding sexualized violence safety-apps.

From the included studies in this review, it became evident that feelings of security and privacy were of utmost importance to the women using the apps (47–52). Some women felt an increased sense of privacy while using the app, which allowed them to feel more comfortable and secure, as well as less judged and stigmatized by others. While certain apps were praised for their discrete icons and layouts (47, 51), one article specifically discussed improvements in this area through increased password protection and a “quick exit” button or feature to disguise the app’s purpose to an onlooker (52). The concept of privacy on apps has become a key concern for many app users, often in regards to sharing of sensitive user data (54, 55), or where data will be stored (56). For health apps specifically, a fear surrounding privacy lies in the potential of unauthorised use or disclosure of health information, which could lead to social stigma and discrimination (57). A quantitative study with a focus on how privacy is valued by app users across a variety of apps (e.g., care sharing, diabetes app, companion and security app, and mood adjustment app) found that premium (i.e. at a cost) privacy features were more favourable than others, such as premium functions and personalization (58). As demonstrated through the literature, most privacy concerns were related to sharing personal information, whereas the studies in this review discussed privacy in the context of shielding their use of the app from a partner or from social networks.

Our review found that accessibility, inclusive of how easy and useful the app was perceived to be, was integral to the user experience. App content and design has been directly related to accessibility and the rate of use in other literature. The connection between ease of use and a user’s experience has been widely discussed, especially among health apps (e.g., (59–62). For example, a qualitative study exploring user experiences of mobile health apps found that an apps ease of use led to more desire to use the app (59). Features that made the app easy to use were automation of in-app functions that reduced time in performing tasks, and convenience such as having information for a person’s self-management plan in one location (59). In a qualitative study on barriers and facilitators of medical mobile app use, app features such as information content, accessibility of the information, and ease of use of the app, were highlighted as facilitators that promoted use of the app (62). Another study, which explored factors that influence use of a mobile app for reporting adverse drug reactions and receiving safety information, found that use of the app was influenced by ease of use and the security of the app (60). Overall, it is apparent that an apps ease of use is an integral part of the user experience.

The reviewed literature found that increased information, awareness, validation, and myth debunking were positive knowledge features on sexualized violence safety apps (46–49, 51, 52). It was also found that users appreciated information that was trauma informed, credible and evidence based (46–49, 51, 52). Similar to our findings, a study examining help-seeking for domestic violence victims found that apps were a particularly good avenue for accessing information (63). Other literature has reported that mobile apps with high quality education materials can significantly increase the knowledge of users (64) and directly influence their attitudes and behaviours (65, 66). It is important to note that sexualized violence safety apps can also be used to inform individuals who are not victims or perpetrators, such as bystanders. Shaw and Janulis (2016) found that bystander education increased sexualized violence knowledge, decreased the likelihood of endorsing rape myths, and increased a sense of efficacy for intervening as a bystander (67). Last, apps provide a

convenient and affordable way to access information that is often interactive and are an environmentally friendly alternative to information that is often conveyed on paper (68).

The use of technology to increase safety is not a novel concept. For example, ride-share apps such as Uber offer GPS tracking that can be shared with family and friends in real time, as well as a distress alarm available on the app that can signal the ride-sharing service for help (69). Increased safety through technology can also be in the form of delivering safety information, such as the use of mobile technology for delivery of safety awareness in the workplace (70) and communicating emergency safety information through mobile text alerts (71). Safety in relation to technology and sexualized violence, however, has primarily been discussed negatively. The use of social media, personal tracking applications, and smartphone technology in general have been found to facilitate forms of sexualized violence, such as harassment, stalking, violence, and dating abuse (13). Interestingly, despite these challenges, a study by Finn & Atkinson (2009) found that women who experience sexualized violence through technology still feel a sense of independence when using technology (53). This aligns with the findings presented in this review, which found that the women who had experienced sexualized violence felt that safety apps provided them with increased knowledge, anonymity, validation, and a sense of empowerment (47–49).

## Limitations

Quality appraisal is not a compulsory component of the Arksey and O'Malley (2005) scoping review framework, and the included studies in this scoping review were not critically appraised (43). Although this scoping review did not seek to assess quality of evidence, it consequently cannot determine whether the included studies provide robust or generalizable findings or if the research itself is of poor quality. Literature that was not included in this review included reviews, commentaries, editorials, and conference proceedings. Existing networks and relevant organizations were also not contacted. While the search was broad, some relevant studies may have been omitted. In addition, the review only included literature published in English and French and, therefore, relevant literature may have been omitted if published in other languages. Although many apps are currently being created and developed, they may not be at the user testing phase or are not explicitly looking at users' experiences beyond interface evaluations. Despite these limitations, the review does provide important understandings of women's experiences of sexualized violence safety-apps.

## Conclusion

This scoping review provides a comprehensive summary of the qualitative research findings in relation to women's experiences of sexualized violence safety apps. This review has highlighted that there is limited research conducted in this area. Victims of sexualized violence are in a unique position to provide insight to app developers on their priorities and specific needs, which can potentially change the way women utilize app technology for their safety. Additional research that focuses on the experiences of women users will help to better inform quality app development that is secure, informative, useful, and wanted by the user. Overall, the reviewed literature in this study found safety apps to be a private all-in-one resource for support, information, and emergency planning that were useful and easy to use. Further, women believed that sexualized violence safety apps had the potential to decrease the overall risk of experiencing sexualized violence.

## Declarations

### Ethics approval and consent to participate:

Not applicable – this was a scoping review of the literature.

### Consent for publication:

Not applicable.

### Availability of data and materials:

Available upon request.

### Competing interests:

None to disclose.

### Funding:

This project did not receive any funding.

### Authors' contributions:

All authors (ND, CA, SW, AC, JS, MN) have contributed to the manuscript scholastically and have approved the final version.

# Acknowledgements:

Leah Boulos, Evidence Synthesis Coordinator, Maritime SPOR Support Unit

## References

1. Benoit C, Shumka L, Phillips R, Kennedy MC, Belle-Isle L. Issue Brief: Sexual Violence Against Women in Canada – 2015 [Internet]. 2016 [cited 2020 May 18]. Available from: <https://www.princeedwardisland.ca/en/publication/issue-brief-sexual-violence-against-women-canada-2015>.
2. Halstead V, Williams JR, Gonzalez-Guarda R. Sexual violence in the college population: a systematic review of disclosure and campus resources and services. *J Clin Nurs*. 2017;26(15–16):2137–53.
3. Victoria Sexual Assault Centre. What is Sexualized Violence? [Internet]. Sexualized Violence is a Community Issue. [cited 2020 May 18]. Available from: <https://vsac.ca/sexualized-violence/>.
4. World Health Organization. Violence against women [Internet]. 2018 [cited 2020 May 18]. Available from: <https://www.who.int/news-room/fact-sheets/detail/violence-against-women>.
5. Truman J, Langton L. Criminal, Victimization. 2013 (Revised) [Internet]. Bureau of Justice Statistics, U.S. Department of Justice; 2014 [cited 2020 May 18]. Report No.: NCJ 247648. Available from: <https://www.bjs.gov/index.cfm?ty=pbdetail&iid=5111>.
6. Truman J, Morgan R. Criminal Victimization. 2015 [Internet]. Bureau of Justice Statistics, U.S. Department of Justice; 2016 Oct [cited 2020 May 18]. Report No.: NCJ 250180. Available from: <https://www.bjs.gov/index.cfm?ty=pbdetail&iid=5804>.
7. Conroy S, Cotter A. Self-reported sexual assault in Canada, 2014 [Internet]. Ottawa: Statistics Canada; 2017 Jul [cited 2020 May 18]. Report No.: 85–002–X. Available from: <https://search.proquest.com/openview/7412b493c403481e8d75275442c40d7f/1?pq-origsite=gscholar&cbl=44168>.
8. Cotter A, Savage L. Gender-based violence and unwanted sexual behaviour in Canada, 2018: Initial findings from the Survey of Safety in Public and Private Spaces [Internet]. Ottawa: Statistics Canada; 2019 Nov [cited 2020 May 18]. Report No.: 1209–6393. Available from: <https://search.proquest.com/openview/238b418d1887ebc3004e767fa0679d2b/1?pq-origsite=gscholar&cbl=44168>.
9. Breiding MJ, Black MC. Intimate Partner Violence in the United States – 2010 [Internet]. Atlanta, GA: National Center for Injury Prevention and Control, Centers for Disease Control and Prevention; 2014 [cited 2020 May 18]. Available from: <https://stacks.cdc.gov/view/cdc/21961>.
10. Tarzia L, Maxwell S, Valpied J, Novy K, Quake R, Hegarty K. Sexual violence associated with poor mental health in women attending Australian general practices. *Aust N Z J Public Health*. 2017;41(5):518–23.
11. Langenderfer-Magruder L, Walls NE, Kattari SK, Whitfield DL, Ramos D. Sexual Victimization and Subsequent Police Reporting by Gender Identity Among Lesbian, Gay, Bisexual, Transgender, and Queer Adults. *Violence Vict*. 2016 Jan 1;31(2):320–31.
12. Powell A, Scott AJ, Henry N. Digital harassment and abuse: Experiences of sexuality and gender minority adults. *Eur J Criminol*. 2020 Mar;17(2)(1):199–223.
13. Henry N, Powell A. Technology-Facilitated Sexual Violence: A Literature Review of Empirical Research. *Trauma Violence Abuse*. 2018 Apr 1;19(2):195–208.
14. Mortimer S, Powell A, Sandy L. ‘Typical scripts’ and their silences: exploring myths about sexual violence and LGBTQ people from the perspectives of support workers. *Curr Issues Crim Justice*. 2019 Jul 3;31(3):333–48.
15. Amstadter AB, McCauley JL, Ruggiero KJ, Resnick HS, Kilpatrick DG. Self-Rated Health in Relation to Rape and Mental Health Disorders in a National Sample of Women. *Am J Orthopsychiatry*. 2011;81(2):202–10.
16. Campbell R, Dworkin E, Cabral G. An Ecological Model of the Impact of Sexual Assault On Women’s Mental Health. *Trauma Violence Abuse*. 2009 Jul 1;10(3):225–46.
17. Koss MP, Figueredo AJ, Prince RJ. Cognitive mediation of rape’s mental, physical and social health impact: Tests of four models in cross-sectional data. *J Consult Clin Psychol*. 2002;70(4):926–41.
18. Resick PA. The Psychological Impact of Rape. *J Interpers Violence*. 1993 Jun;1(2):223–55. 8{.
19. Spohn R, Wright EM, Peterson JC. Rape and Mental Health Outcomes Among Women: Examining the Moderating Effects of “Healthy” Fear Levels. *Violence Women*. 2017 Aug 1;23(9):1055–75.
20. Zinzow HM, Resnick HS, Amstadter AB, McCauley JL, Ruggiero KJ, Kilpatrick DG. Drug- or Alcohol-Facilitated, Incapacitated, and Forcible Rape in Relationship to Mental Health Among a National Sample of Women. *J Interpers Violence*. 2010 Dec 1;25(12):2217–36.
21. Pegram SE, Abbey A. Associations Between Sexual Assault Severity and Psychological and Physical Health Outcomes: Similarities and Differences Among African American and Caucasian Survivors. *J Interpers Violence*. 2019 Oct;34(19)(1):4020–40.
22. Scott KM, Koenen KC, King A, Petukhova MV, Alonso J, Bromet EJ, et al. Post-traumatic stress disorder associated with sexual assault among women in the WHO World Mental Health Surveys. *Psychol Med*. 2018 Jan;48(1):155–67.
23. Stoner JE, Cramer RJ. Sexual Violence Victimization Among College Females: A Systematic Review of Rates, Barriers, and Facilitators of Health Service Utilization on Campus. *Trauma Violence Abuse*. 2019 Oct;20(4)(1):520–33.
24. 10.3389/fpsy.2018.00311/full  
Millon EM, Chang HYM, Shors TJ. Stressful Life Memories Relate to Ruminative Thoughts in Women With Sexual Violence History, Irrespective of PTSD. *Front Psychiatry* [Internet]. 2018 [cited 2020 May 18];9. Available from: <https://www.frontiersin.org/articles/10.3389/fpsy.2018.00311/full>.

25. Beck JG, Grant DM, Clapp JD, Palyo SA. Understanding the interpersonal impact of trauma: Contributions of PTSD and depression. *J Anxiety Disord.* 2009 May 1;23(4):443–50.
26. Bedard-Gilligan M, Kaysen D, Desai S, Lee CM. Alcohol-involved assault: Associations with posttrauma alcohol use, consequences, and expectancies. *Addict Behav.* 2011 Nov;36(11)(1):1076–82.
27. Forbes D, Fletcher S, Parslow R, Phelps A, O'Donnell M, Bryant RA, et al. Trauma at the hands of another: Longitudinal study of differences in the posttraumatic stress disorder symptom profile following interpersonal compared with noninterpersonal trauma. *J Clin Psychiatry.* 2012;73(3):372–6.
28. Ullman SE, Brecklin LR. Sexual Assault History and Suicidal Behavior in a National Sample of Women. *Suicide Life Threat Behav.* 2002 Jun 1;32(2):117–30.
29. World report on violence and health [Internet]. World Health Organization; 2002 [cited 2020 May 18]. Available from: [https://www.who.int/violence\\_injury\\_prevention/violence/world\\_report/en/](https://www.who.int/violence_injury_prevention/violence/world_report/en/).
30. Facts and figures. Ending violence against women [Internet]. UN Women. [cited 2020 May 18]. Available from: <https://www.unwomen.org/en/what-we-do/ending-violence-against-women/facts-and-figures>.
31. The World's Women. 2015, Trends and Statistics [Internet]. United Nations Economic and Social Affairs; 2015 [cited 2020 May 18]. Available from: [https://public.tableau.com/views/populationstory/YouthEducation?:embed=y&:showVizHome=no&:host\\_url=https%3A%2F%2Fpublic.tableau.com%2F&t](https://public.tableau.com/views/populationstory/YouthEducation?:embed=y&:showVizHome=no&:host_url=https%3A%2F%2Fpublic.tableau.com%2F&t)
32. Glass N, Eden KB, Bloom T, Perrin N. Computerized Aid Improves Safety Decision Process for Survivors of Intimate Partner Violence. *J Interpers Violence.* 2010 Nov 1;25(11):1947–64.
33. Smith A, Rainie L, Zickuhr K. College students and technology. Pew Research Center: Internet, Science & Tech [Internet]. 2011 Jul 19 [cited 2020 May 18]; Available from: <https://www.pewresearch.org/internet/2011/07/19/college-students-and-technology/>.
34. Silver L. Smartphone Ownership Is Growing Rapidly Around the World, but Not Always Equally. Pew Research Center's Global Attitudes Project [Internet]. 2019 Feb 5 [cited 2020 May 18]; Available from: <https://www.pewresearch.org/global/2019/02/05/smartphone-ownership-is-growing-rapidly-around-the-world-but-not-always-equally/>.
35. Abrams LC, Padmanabhan N, Thaweethai L, Phillips T. iPhone Apps for Smoking Cessation: A Content Analysis. *Am J Prev Med.* 2011 Mar 1;40(3):279–85.
36. Breland JY, Yeh VM, Yu J. Adherence to evidence-based guidelines among diabetes self-management apps. *Transl Behav Med.* 2013 Sep 1;3(3):277–86.
37. Breton ER, Fuemmeler BF, Abrams LC. Weight loss—there is an app for that! But does it adhere to evidence-informed practices? *Transl Behav Med.* 2011 Dec;1(4)(1):523–9.
38. Cardoso LF, Sorenson SB, Webb O, Landers S. Recent and emerging technologies: Implications for women's safety. *Technol Soc.* 2019 Aug 1;58:101108.
39. McCarthy OT, Caulfield B, O'Mahony M. How transport users perceive personal safety apps. *Transp Res Part F Traffic Psychol Behav.* 2016 Nov;1:43:166–82.
40. White D, McMillan L. Innovating the Problem Away? A Critical Study of Anti-Rape Technologies. *Violence Women.* 2019 Jul 21;1077801219856115.
41. Zaidi AU, Fernando S, Ammar N. An exploratory study of the impact of information communication technology (ICT) or computer mediated communication (CMC) on the level of violence and access to service among intimate partner violence (IPV) survivors in Canada. *Technol Soc.* 2015 May;1:41:91–7.
42. Maxwell L, Sanders A, Skues J, Wise L. A Content Analysis of Personal Safety Apps: Are They Keeping Us Safe or Making Us More Vulnerable? *Violence Women.* 2020 Feb 1;26(2):233–48.
43. Arksey H, O'Malley L. Scoping studies: towards a methodological framework. *Int J Soc Res Methodol.* 2005 Feb;1(1):19–32. 8(.
44. Finn J. A Survey of Online Harassment at a University Campus. *J Interpers Violence.* 2004 Apr 1;19(4):468–83.
45. Hébert M, Moreau C, Blais M, Lavoie F, Guerrier M. Child Sexual Abuse as a Risk Factor for Teen Dating Violence: Findings from a Representative Sample of Quebec Youth. *J Child Adolesc Trauma.* 2017 Mar 1;10(1):51–61.
46. Alhusen J, Bloom T, Clough A, Glass N. Development of the MyPlan Safety Decision App with Friends of College Women in Abusive Dating Relationships. *J Technol Hum Serv.* 2015 Jul;3(3):263–82. 33(.
47. Bloom T, Gielen A, Glass N. Developing an App for College Women in Abusive Same-Sex Relationships and Their Friends. *J Homosex.* 2016 Jun 2;63(6):855–74.
48. Lindsay M, Messing JT, Thaller J, Baldwin A, Clough A, Bloom T, et al. Survivor Feedback on a Safety Decision Aid Smartphone Application for College-Age Women in Abusive Relationships. *J Technol Hum Serv.* 2013 Oct 1;31(4):368–88.
49. Tarzia L, Iyer D, Thrower E, Hegarty K. "Technology Doesn't Judge You": Young Australian Women's Views on Using the Internet and Smartphones to Address Intimate Partner Violence. *J Technol Hum Serv.* 2017 Jul 3;35(3):199–218.
50. 50.
51. Gilmore AK, Davidson TM, Leone RM, Wray LB, Oesterle DW, Hahn CK, et al. Usability Testing of a Mobile Health Intervention to Address Acute Care Needs after Sexual Assault. *Int J Environ Res Public Health.* 2019 Jan;16(17):3088.
52. Ragavan MI, Ferre V, Bair-Merritt M, Thrive: A Novel Health Education Mobile Application for Mothers Who Have Experienced Intimate Partner Violence. *Health Promot Pract.* 2020 Mar 1;21(2):160–4.
53. Finn J, Atkinson T. Promoting the Safe and Strategic Use of Technology for Victims of Intimate Partner Violence: Evaluation of the Technology Safety Project. *J Fam Violence.* 2009 Jan;24(1)(1):53–9.
54. Avancha S, Baxi A, Kotz D. Privacy in mobile technology for personal healthcare. *ACM Comput Surv.* 2012 Dec 7;45(1):3:1–3:54.

55. 10.1007/978-3-642-39498-0\_10  
Egelman S, Felt AP, Wagner D. Choice Architecture and Smartphone Privacy: There's a Price for That. In: Böhme R, editor. *The Economics of Information Security and Privacy* [Internet]. Berlin, Heidelberg: Springer; 2013 [cited 2020 May 18]. p. 211–36. Available from: [https://doi.org/10.1007/978-3-642-39498-0\\_10](https://doi.org/10.1007/978-3-642-39498-0_10).
56. He D, Naveed M, Gunter CA, Nahrstedt K. Security Concerns in Android mHealth Apps. *AMIA Annu Symp Proc*. 2014 Nov 14;2014:645–54.
57. James G. Hodge J, Gostin LO, Jacobson PD. Legal Issues Concerning Electronic Health Information: Privacy, Quality, and Liability. *JAMA*. 1999 Oct;282(15):1466–71. 282(.
58. Dogruel L, Joeckel S, Vitak J. The valuation of privacy premium features for smartphone apps: The influence of defaults and expert recommendations. *Comput Hum Behav*. 2017 Dec;1:77:230–9.
59. Anderson K, Burford O, Emmerton L. Mobile Health Apps to Facilitate Self-Care: A Qualitative Study of User Experiences. *PLOS ONE*. 2016 May;23(5):e0156164. 11(.
60. de Vries ST, Wong L, Sutcliffe A, Houyez F, Ruiz CL, Mol PGM, et al. Factors Influencing the Use of a Mobile App for Reporting Adverse Drug Reactions and Receiving Safety Information: A Qualitative Study. *Drug Saf*. 2017 May 1;40(5):443–55.
61. Scheper H, Derogee R, Mahdad R, van der Wal RJP, Nelissen RGHH, Visser LG, et al. A mobile app for postoperative wound care after arthroplasty: Ease of use and perceived usefulness. *Int J Med Inf*. 2019 Sep;1:129:75–80.
62. Velu AV, Beukering MD van, Schaafsma FG, Frings-Dresen MH, Mol BW, Post JA van der, et al. Barriers and Facilitators for the Use of a Medical Mobile App to Prevent Work-Related Risks in Pregnancy: A Qualitative Analysis. *JMIR Res Protoc*. 2017;6(8):e163.
63. Cotten SR, Gupta SS. Characteristics of online and offline health information seekers and factors that discriminate between them. *Soc Sci Med*. 2004 Nov;59(9(1):1795–806.
64. Ismail S, Rangka J, Rasdi I, Rahman U, Samah M. Mobile Apps Application to Improve Safety and Health Knowledge, Attitude and Practice among University Students. *Malays J Med Health Sci* [Internet]. 2018 [cited 2020 May 18];14(101). Available from: [https://www.academia.edu/39764557/Mobile\\_Apps\\_Application\\_to\\_Improve\\_Safety\\_and\\_Health\\_Knowledge\\_Attitude\\_and\\_Practice\\_among\\_University\\_St](https://www.academia.edu/39764557/Mobile_Apps_Application_to_Improve_Safety_and_Health_Knowledge_Attitude_and_Practice_among_University_St)
65. Shin D-H, Shin Y-J, Choo H, Beom K. Smartphones as smart pedagogical tools: Implications for smartphones as u-learning devices. *Comput Hum Behav*. 2011 Nov;27(6(1):2207–14.
66. Kumar Basak S, Wotto M, Bélanger P. E-learning, M-learning and D-learning: Conceptual definition and comparative analysis. *E-Learn Digit Media*. 2018 Jul 1;15(4):191–216.
67. Shaw J, Janulis P. Re-Evaluating Sexual Violence Prevention Through Bystander Education: A Latent Growth Curve Approach. *J Interpers Violence*. 2016 Oct 1;31(16):2729–50.
68. Kumar SE. and M-Learning: A comparative study. *International Journal on New Trends in Education Their Implications*. 2013;4(3):65–78.
69. Chaudhry B, Yasar A-U-H, El-Amine S, Shakshuki E. Passenger Safety in Ride-Sharing Services. *Procedia Comput Sci*. 2018 Jan 1;130:1044–50.
70. Rodriguez A, Hagevoort GR, Leal D, Pompeii L, Douphrate DI. Using mobile technology to increase safety awareness among dairy workers in the United States. *J Agromedicine*. 2018 Oct 2;23(4):315–26.
71. Wong DJ, Jones E, Rubin GJ. Mobile text alerts are an effective way of communicating emergency information to adolescents: Results from focus groups with 12- to 18-year-olds. *J Contingencies Crisis Manag*. 2018;26(1):183–92.

## Figures




Figure 1

Safety App Prisma