Characteristics and outcomes of Stanford type A aortic dissection patients with post-operation severe hyperbilirubinemia: a retrospective cohort study

Xiaolan Chen

Xijing Hospital

Ming Bai (✉ mingbai1983@126.com)

Xijing Hospital

Lijuan Zhao

Xijing Hospital

Yangping Li

Xijing Hospital

Yan Yu

Xijing Hospital

Wei Zhang

Xijing Hospital

Feng Ma

Xijing hospital

Shiren Sun (✉ sunshiren@medmail.com.cn)

Xijing Hospital

Xiangmei Chen (✉ xmchen301@126.com)

Xijing Hospital

Research article

Keywords: Stanford type A aortic dissection, hyperbilirubinemia, acute kidney injury, continuous renal replacement therapy

Posted Date: May 26th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-30855/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published on July 28th, 2020. See the published version at <https://doi.org/10.1186/s13019-020-01243-7>.

Abstract

Background

Hyperbilirubinemia is one of the common complications after cardiac surgery and is associated with increased mortality. However, to the best of our knowledge, the report on clinical significance of postoperative severe hyperbilirubinemia in Stanford type A aortic dissection (AAD) patients is limited.

Methods

Patients who underwent surgical treatment for AAD in our center between January 2015 and December 2018 were retrospectively screened. In-hospital mortality, long-term mortality, acute kidney injury (AKI), and the requirement of continuous renal replacement therapy (CRRT) were assessed as endpoints. Univariate and multivariate regression models were employed to identify the risk factors of these endpoints.

Results

271 (12.3%) patients were included. Of the included patients, 222 (81.9%) experienced postoperative AKI, and 50 (18.5%) received CRRT. In-hospital mortality was 30.3%. The 1-year, 2-year, and 3-year cumulative mortality were 32.9%, 33.9%, and 35.3%, respectively. Multivariate Logistic regression analysis indicated that age ($P < 0.033$), AKI stage 3 ($P < 0.001$), the amount of blood transfusion after surgery ($P = 0.019$), mean arterial pressure (MAP) in the first postoperative day ($P = 0.012$), the use of extracorporeal membrane oxygenation (ECMO) ($P = 0.02$), and the peak total bilirubin (TB) concentration ($P = 0.023$) were independent risk factors of in-hospital mortality. The optimal cut-off value of peak TB on predicting in-hospital mortality was 121.2 $\mu\text{mol/L}$. Survival analysis showed significantly decreased survival for patients who developed severe, rather than mild, hyperbilirubinemia.

Conclusions

Post-operation severe hyperbilirubinemia is a common clinical presentation in AAD surgery patients. Post-operation severe hyperbilirubinemia AAD patients with older age, lower MAP, increased blood transfusion, stage 3 AKI, the use of ECMO, and the increased peak TB had higher risk of in-hospital mortality.

1. Introduction

Aortic dissection (AD) is an acute life-threatening condition with a prevalence of about 3/100,000 per year. The International Registry of AD revealed that 67% of AD patients presented with Stanford type A aortic dissection (AAD), which was characteristic as the involvement of the ascending aorta. And, approximately 86% of AAD patients required swift open cardiac surgery to avoid fatal complications such as aortic rupture and cardiac tamponade.[1–3] In spite of the improvement in medical management and surgical technique, AAD surgery was associated with early mortality as high as 20%.[4–8]

Hyperbilirubinemia is a common severe complication after cardiac operation.[9] The reported incidence of post-operation hyperbilirubinemia varied widely (10–57%) and related to the severity of cardiac diseases and the type of cardiac surgery. Although hyperbilirubinemia was reported to be associated with in-hospital mortality and mortality after discharge, the effects of hyperbilirubinemia on patient prognosis were heterogeneous.[10–14] For patients undergoing cardiac surgery with cardiopulmonary bypass (CPB), recent studies suggested that severe hyperbilirubinemia (5 times the normal upper

limit) instead of mild bilirubin significantly increased patient mortality.[15] Additionally, the occurrence of acute kidney injury (AKI) and the acceptance of continuous renal replacement therapy (CRRT) after cardiac surgery were suggested to be associated with the decrease of patient survival proportion. As we know, the pathogenesis, disease severity classification, prognosis, and operation method are different between different cardiac diseases. The inclusion of all kinds of cardiac surgery in one study most likely would reduce the specificity and repeatability of the conclusions. Up to now, the reports on the characteristics and outcomes of AAD surgery patients who had post-operation severe hyperbilirubinemia are limited.

Therefore, the purpose of our present study is to describe the clinical characteristics and to investigate outcomes of AAD surgery patients who had post-operation severe hyperbilirubinemia. The results of our present study might be useful for decision makers regarding their treatment strategy and for patients and their families regarding the expected progression.

2. Patients And Method

2.1. Study design and patients selection

Our present study was retrospectively designed. Consecutive patients who underwent surgery for AD in our center between January 2015 and December 2018 were screened. AD was proven by enhanced computed tomography and defined as type A or type B according to the Stanford classification. Postoperative severe hyperbilirubinemia was defined as occurrence of a serum TB concentration of more than 85.5 $\mu\text{mol/l}$ (5 times the normal upper limit) in any measurement during the hospital staying after AAD surgery. Patients were excluded if they had any of the following conditions: (1) serum TB concentration < 85.5 $\mu\text{mol/l}$, (2) age < 18 years; (3) Stanford type B aortic dissection; (4) the occurrence of severe hyperbilirubinemia before surgery; (5) severe hyperbilirubinemia caused by the reoperation during hospitalization. The local Institutional Review Board of the hospital approved this retrospective study and waived the requirement of informed consent for the use of patients' medical data.

2.2. Surgical procedure

The use of surgical techniques has developed during the study period. However, the basic principles for repair of AAD have no change, including (1) prompt establishment of CPB, (2) resection of the primary entry site by open distal anastomosis under deep hypothermic circulatory arrest, (3) preservation of the aortic valve whenever possible, and (4) aortic arch replacement in patients with an entry site located in or extending into the aortic arch. Aortic root replacement with a composite prosthesis and reimplantation of the coronary arteries by the modified Bentall technique was performed in patients with conspicuous dilatation of the aortic root.

2.3. Data collection

Demographic data, comorbidities, and operation details were retrieved from our hospital's electronic medical record system. All the routine laboratory data was recorded before operation (the nearest to the time of surgery) and in the post-operation period. The severity of illness before and after surgery was assessed by using the acute physiology and chronic health evaluation II (APACHE II), sequential organ failure assessment (SOFA) score, and model for end-stage liver disease (MELD) score. Urine output was recorded every day after the surgery.

2.4. Outcomes and definition

Post-operation outcomes including the amount of blood transfusion, mechanical ventilation time, the use of extracorporeal membrane oxygenation (ECMO), intra-aortic balloon pump (IABP), AKI, the usage of CRRT, bilirubin adsorption (BA) or plasma exchange (PE), and vasoactive agent, the duration of hospitalization, ICU stay time, and in-hospital mortality. For those patients who were alive on discharge, telephone survey was performed to obtain the patients long-term outcome.

Kidney Disease Improving Global Outcomes (KDIGO) criteria[16] based on SCr or urine output was employed to diagnose and grade AKI. The latest SCr concentration before surgery was defined as preoperative SCr concentration. The decision to start CRRT was made at the discretion of the attending nephrologist. Main indications for starting CRRT were progressive AKI, fluid overload, hyperkalemia, and severe metabolic acidosis.[16]

2.5. Statistical analysis

Data were analyzed using SPSS version 22.0 software (SPSS, Inc, Chicago, IL, USA). Continuous variables are presented as mean $\pm$ standard deviation. Categorical variables are presented as frequencies with percentages. To evaluate the differences between groups, the independent sample *t*-test was used for continuous variables, whereas the chi-square test or Fisher's exact test was used for categorical variables. Factors significantly associated with these endpoints in univariate analysis were included in the multivariate logistic regression analysis or Cox proportional hazard analysis to identify the independent risk factors. Accumulated survival proportion was estimated with the Kaplan-Meier method, and the between-group differences of survival proportion were assessed using the log-rank test. Area under the receiver operating characteristic curve (AUC-ROC) was calculated to assess peak TB concentration on the ability to detect in-hospital mortality. Youden index was used for assessment of optimal cut-off values. For all analyses, all statistical tests were 2-sided, and a *P*-value < 0.05 was considered as statistically significant.

3. Result

3.1. Patient characteristics

Of the 2210 screened patients, 1931, 2, 3, and 3 were excluded because of TB concentration $< 85.5 \mu\text{mol/l}$, Stanford type B aortic dissection, preoperative TB $\geq 85.5 \mu\text{mol/l}$, and post-operation severe hyperbilirubinemia caused by reoperation during this hospitalization, respectively. Ultimately, 271 AAD surgery patients with post-operation severe hyperbilirubinemia were included in our present study (Fig. 1).

The baseline characteristics of the included patients were summarized in Table 1. There were 223 male and 48 female, and the mean age of the patients was 49.1 ± 11.0 years. AAD surgery involved the aortic valve in 83.8% patients, aortic arch in 86% patients, and coronary artery in 17% patients. Preoperative TB concentration was $25.0 \pm 15.0 \mu\text{mol/l}$. The mean onset time of severe hyperbilirubinemia was 2.8 ± 1.3 days after AAD surgery. The mean peak serum TB concentration was $150.9 \pm 93.0 \mu\text{mol/l}$ and the averaged time to peak TB concentration was 3.8 ± 3.0 days after AAD surgery. The change of TB concentration during the seven days after surgery was showed in Fig. 2. The median follow-up time was 18.63 (0.5–55.9) months. And, 3 patients were lost to follow-up. The time of lost to follow-up was 16-, 95-, and 365-day after the surgery, respectively.

3.2. Postoperative AKI

Of the 271 included patients, 222 patients (82.1%) had AKI after AAD surgery, of which 102 (40.2%), 34 (12.5%), and 84 (29.2%) were stage 1, stage 2, and stage 3 AKI (Table 2), respectively. The results of univariate and multivariate logistic regression are presented in Table 3. In the univariate analysis, age, hypertension, preoperative SCr concentration, APACHE II score, and MELD score, operation time, and CPB time were associated with the occurrence of postoperative AKI. Multivariate analysis indicated that increased age (OR 1.056, 95%CI 1.026–1.088, $P < 0.001$), higher preoperative SCr concentration (OR 1.026, 95%CI 1.010–1.043, $P = 0.002$), and prolonged CPB time (OR 1.007, 95%CI 1.000–1.013, $P = 0.042$) were identified as independent risk factors of AKI in AAD surgery patients with post-operation severe hyperbilirubinemia.

3.3. Postoperative CRRT

Of the included patients, 50 (18.5%) patients received CRRT after AAD surgery. Univariate analysis indicated that male, preoperative SCr concentration, APEACHE II score, MELD score, and aortic cross-clamp (ACC) time were associated with the need for CRRT. In the multivariate logistic regression analysis, only preoperative SCr concentration (OR 1.011, 95%CI 1.002–1.016, $P=0.003$) was identified as an independent predictor of the acceptance of CRRT (Table 3).

3.4. In-hospital mortality

There were 82 (30.3%) in-hospital deaths. Univariate analysis revealed that 20 pre-operation, intra-operation, and post-operation factors were associated with in-hospital mortality. After the adjustment of the important clinical parameters, independent risk factors of in-hospital mortality identified by multivariate logistic analysis included age (OR 1.005, 95%CI 1.004–1.009, $P=0.033$), the amount of blood transfusion (OR 1.018, 95%CI 1.003–1.033, $P=0.019$), stage 3 AKI (OR 46.134, 95%CI 5.436–391.525, $P<0.001$), the use of ECMO (OR 20.795, 95%CI 1.620–266.917, $P=0.02$), and the peak TB concentration (OR 1.017, 95%CI 1.002–1.032, $P=0.023$). Postoperative MAP tended to be a protective factor (OR 0.955, 95%CI 0.922–0.990, $P=0.012$, Table 4, model 1).

ROC analysis (Fig. 3) identified that peak TB concentration was associated with increased mortality, and the optimal cut-off value identified by the Youden index was 121.2 $\mu\text{mol/l}$ (sensitivity: 72%, specificity: 60%). And, the multivariate analysis revealed that patients with peak TB concentration $\geq 121.2 \mu\text{mol/l}$ had a significantly higher risk of in-hospital mortality (OR = 2.681, 95%CI 1.119–6.425, $P=0.027$, Table 4, model 2), compared with the patients with peak TB concentration $< 121.2 \mu\text{mol/l}$.

3.5. Long-term mortality

Of the 12 patients who died after discharge, 4 had non-AKI, 3 had stage 1 AKI, 3 had stage 2 AKI, and 2 had stage 3 AKI during their hospital stay, respectively. The accumulated 1-year, 2-year, and 3-year mortality proportions were 32.9%, 33.9%, and 35.3%, respectively (Fig. 4A). The risk factors of long-term mortality were presented in Table 5. Univariate analysis revealed 20 factors, including the post-operation AKI, the acceptance of CRRT and ECMO, and the peak TB concentration $\geq 121.2 \mu\text{mol/l}$ (Fig. 4B-E), were significantly related to patient long-term mortality. Multivariate COX regression analysis revealed that stage 3 AKI (HR 12.604, 95%CI 5.002–31.762, $P<0.001$) significantly increased long-term mortality, compared with patients without AKI. The use of ECMO (HR 12.167, 95%CI 4.588–32.264, $P<0.001$) was identified as an independent predictor of long-term mortality as well. In contrast, postoperative MAP (HR 0.979, 95%CI 0.962–0.995, $P=0.012$) was identified as an independent protective factor of long-term mortality.

4. Discussion

It has been reported that severe hyperbilirubinemia was associated with increased mortality in patients underwent cardiac surgery.[11, 15, 17] Up to now, limited information was available on the risk factors associated with in-hospital and long-term mortality in AAD patients with post-operation severe hyperbilirubinemia. Our presented study had several findings. Firstly, the occurrence of AKI, the requirement for CRRT, and the in-hospital mortality were higher than previous studies of patients undergoing AAD surgery without severe hyperbilirubinemia. Secondly, age, preoperative SCr concentration, and CPB time were independent risk factors for postoperative AKI, and preoperative SCr concentration was an independent risk factor for post-operation CRRT as well. Finally, the peak TB concentration, post-operation stage 3 AKI, the total amount of blood transfusion after AAD surgery, the use of ECMO, and low MAP after surgery were significantly associated with mortality.

4.1. AAD patients with severe hyperbilirubinemia were associated with worse prognosis

The analysis of our present cohort of AAD patients with post-operation severe hyperbilirubinemia showed an overall incidence of post-operation AKI of 81.9%, a requirement for CRRT of 18.5%, and the in-hospital mortality rate of 30.3%. In previous studies of AAD patients underwent surgical treatment without post-operation severe hyperbilirubinemia, the reported AKI incidences were ranged from 40–78%, [5–7, 18–23] the reported incidences of requirement for CRRT were ranged from 3% to 8%, [5, 7] and the reported in-hospital mortalities were ranged from 15–26%. [4, 5, 7, 19, 24] The discrepancy most likely attributed to the fact that all of the included AAD patients in our study developed severe hyperbilirubinemia. The development of severe hyperbilirubinemia after AAD surgery related to the severity of the AAD and the severity of injury during AAD operation. In animal model, hyperbilirubinemia was proved to have pro-apoptotic effects and aggravate renal ischemia-reperfusion injury. [25] Additionally, high concentration of bilirubin could lead to inflammatory response and cell apoptosis of the brain, [26] which might be one of the potential mechanism of the high mortality of our present cohort.

4.2. Risk factors of postoperative AKI and CRRT

In our present cohort, older age, high preoperative SCr concentration, and prolonged CPB time were identified as independent risk factors for AKI. Meanwhile, preoperative SCr concentration was also associated with the requirement of CRRT after AAD surgery. A recent meta-analysis of patients underwent AAD surgery [18] showed that older age was identified as an independent risk factor of AKI as well. Older age-mediated adverse renal structural and functional changes might contribute to the high development of postoperative AKI after AAD surgery in older patients. The aforementioned meta-analysis showed that preoperative SCr concentration did not correlate with postoperative AKI with significant heterogeneity ($I^2 = 72.8\%$). In our opinion, pre-operation elevated SCr concentration might indicate structural kidney damage or hemodynamic derangements in AAD patients, which further aggravated the development of post-operation AKI and the requirement of CRRT. Longer CPB time can lead to more hemolysis, and a longer time on the circuit can lead to changes in perfusion to the viscera and to more inflammatory action. A CPB itself might also induce hypoperfusion of abdominal organs, hypoxia or an inflammatory reaction, which causes liver and kidney damage. Therefore, surgeons most likely could reduce the AKI risk of AAD patients by the improvement of their operation strategies and the reduction of CPB time.

4.3. Risk factors of in-hospital and long-term mortality

In previous studies, older age had been identified as an independent risk factor of mortality for patients with hyperbilirubinemia after cardiac surgery. [27] In our present study, older age was established a risk factor for in-hospital mortality for patients with hyperbilirubinemia after AAD surgery as well. Nevertheless, aging indicated diminished functional capacity of the liver and added to the cumulative burden in the case of developed severe hyperbilirubinemia. Additionally, different from the a recent study outlining the increased risk of mortality for patients with hyperbilirubinemia after cardiac surgery associated with time to peak bilirubin, [15] our study indicated that peak TB concentration was an independent driver of in-hospital mortality for patients with severe hyperbilirubinemia after AAD surgery. And, we found out that the optimal cut-off value of peak TB on predicting in-hospital mortality was 121.2 $\mu\text{mol/L}$. This difference might be attributed to the existence of severe hyperbilirubinemia after AAD surgery and the limitation of small sample size. The high concentration of bilirubin leads to inflammatory response or cell apoptosis in the brain. [26] Cui et al. [28] showed that bilirubin induced lung edema and injury by inducing the apoptosis of alveolar epithelial cells. From the above, severe hyperbilirubinemia induces oxidative stress and apoptosis in many organs, which is associated with poor outcomes in patients with AAD surgery.

In a study for patients with jaundice after open heart surgery, Chu *et al.* [29] reported patients with more severe hyperbilirubinemia or delayed serum peak TB concentration might be significantly associated with more transfused blood and hypotension. Increased amount of blood transfusions contributed to hemolysis, and hypotension would reduce hepatic perfusion [30], which both caused an increased bilirubin load. This might be one of the potential mechanisms that the number of transfused blood units and lower MAP increased mortality in our present study. Previous studies suggested

that mechanisms underlying the development of post-operation late peak TB concentration differed from those of the early peak bilirubin concentration. The immediate development of post-operation peak TB concentration and rapid decline thereafter reflected the transient damaging effects by CPB surgery, whereas late development of post-operation peak TB concentration was a consequence of hepatic dysfunction caused by persistent cardiac failure or sepsis.[9, 12] Therefore, attention should be paid to the monitoring of heart failure and optimizing hemodynamics after AAD surgery to prevent further deterioration. Furthermore, identifying and implementing effective risk reduction strategies is needed. Molecular adsorbent recirculation system, prometheus therapy, fractionated plasma separation and adsorption have shown promise in reducing bilirubin concentration,[27, 31] but further study in this area is needed.

AKI had been incorporated into a risk tool to predict early mortality for patients underwent AAD surgery.[18] Our results added to these findings and demonstrated stage 3 AKI markedly increased both in-hospital and long-term mortality. Additionally, our study also showed in-hospital and long-term mortality was independently predicted by the use of ECMO. Hemolysis is a common complication in ECMO support.[32] Pump head or oxygenator thrombosis and excessive pump speed (typically greater than 3000 rpm) which may destroy blood red cells,[33] which in turn is important factor in the development of hyperbilirubinemia. Therefore, more careful postoperative management is needed to improve prognosis.

4.4. Study limitation

Our present study was a retrospective clinical research from a single institution and had some limitations. First, SCr concentration on admission was regard as preoperative renal function. However, some patients might already have AKI on admission. As a result, the number of patients with AKI may be underestimated. Second, the number of major adverse events including the use of ECMO, plasma exchange and bilirubin adsorption was small, which will be likely to reduce the statistical power for risk factor analysis. Finally, the renal prognosis was not regular followed up after the hospital discharge, which is important for the evaluation of the renal outcome. Therefore, further prospective multicenter studies with larger samples are needed to obtain stronger evidences.

5. Conclusion

Post-operation severe hyperbilirubinemia is a common clinical presentation in AAD surgery patients. Post-operation severe hyperbilirubinemia AAD patients with older age, lower MAP, increased blood transfusion, stage 3 AKI, the use of ECMO, and the increased peak TB had higher risk of in-hospital mortality. These patients most likely need more intensive monitoring.

Abbreviations

AAD	Stanford type A aortic dissection
AKI	Acute kidney injury
ACC	Aortic cross clamp
APEACHEII	Acute physiology and chronic health evaluation II
BA	Bilirubin adsorption
CPB	Cardiopulmonary bypass
CRRT	Continuous renal replacement therapy
ECMO	Extracorporeal membrane oxygenation
IABP	Intra-aortic balloon pump
ICU	Intensive care unit;
MAP	Mean arterial pressure
MELD	Model for end-stage liver disease
PE	Plasma exchange
SCr	Serum creatinine
SOFA	Sequential organ failure assessment
TB	Total bilirubin

Declarations

Acknowledgments

Not applicable.

Funding Source

This work was supported by the National Natural Science Foundation of China (81700584) and Discipline Promotion Project of Xijing Hospital (XJZT18ML16).

Availability of data and materials

The datasets used or analyzed during the current study are available from the corresponding author on reasonable request.

Contributions

Chen, Bai, Zhao, and Sun analyzed and interpreted the data. Li, Yu, Zhang, Ma and Chen performed the statistical analyses. All authors contributed to the study design and interpretation of the study results. All authors read and approved the final manuscript.

Corresponding author

Correspondence to Ming Bai, Shiren Sun¹, Xiangmei Chen.

Ethics declarations

Ethics approval and consent to participate

The local Institutional Review Board of the hospital approved this retrospective study and waived the requirement of informed consent for the use of patients' medical data, and all experimental methods were performed in accordance with the relevant guidelines and regulations.

Consent for publication

Not applicable.

Competing interests

The authors have declared that no interest.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

1. Nienaber CA, Clough RE. Management of acute aortic dissection. *The Lancet*. 2015;385:800–11.
2. Nienaber CA, Clough RE, Sakalihasan N, et al. Aortic dissection. *Nat Rev Dis Primers*. 2016;2:16053.
3. Evangelista A, Isselbacher EM, Bossone E, et al. Insights From the International Registry of Acute Aortic Dissection: A 20-Year Experience of Collaborative Clinical Research. *Circulation*. 2018;137:1846–60.
4. Tsai HS, Tsai FC, Chen YC, et al. Impact of acute kidney injury on one-year survival after surgery for aortic dissection. *Ann Thorac Surg*. 2012;94:1407–12.
5. Sasabuchi Y, Kimura N, Shiotsuka J, et al. Long-Term Survival in Patients With Acute Kidney Injury After Acute Type A Aortic Dissection Repair. *Ann Thorac Surg*. 2016;102:2003–9.
6. Roh GU, Lee JW, Nam SB, et al. Incidence and risk factors of acute kidney injury after thoracic aortic surgery for acute dissection. *Ann Thorac Surg*. 2012;94:766–71.
7. Ko T, Higashitani M, Sato A, et al. Impact of Acute Kidney Injury on Early to Long-Term Outcomes in Patients Who Underwent Surgery for Type A Acute Aortic Dissection. *Am J Cardiol*. 2015;116:463–8.
8. Schoenrath F, Laber R, Maralushaj M, et al. Survival, Neurologic Injury, and Kidney Function after Surgery for Acute Type A Aortic Dissection. *Thorac Cardiovasc Surg*. 2016;64:100–7.
9. Wang MJ, Chao A, Huang CH, et al. Hyperbilirubinemia after cardiac operation. Incidence, risk factors, and clinical significance. *J Thorac Cardiovasc Surg*. 1994;108:429–36.
10. Collins JD, Bassendine MF, Ferner R, et al. Incidence and prognostic importance of jaundice after cardiopulmonary bypass surgery. *Lancet*. 1983;1:1119–23.
11. Sharma P, Ananthanarayanan C, Vaidhya N, et al. Hyperbilirubinemia after cardiac surgery: An observational study. *Asian Cardiovasc Thorac Ann*. 2015;23:1039–43.
12. Hsu RB, Lin FY, Chen RJ, et al. Incidence, risk factors, and prognosis of postoperative hyperbilirubinemia after heart transplantation. *Eur J Cardiothorac Surg*. 2007;32:917–22.
13. Lockey E, McIntyre N, Ross DN, et al. Early jaundice after open-heart surgery. *Thorax*. 1967;22:165–9.
14. Nishi H, Sakaguchi T, Miyagawa S, et al. Frequency, risk factors and prognosis of postoperative hyperbilirubinemia after heart valve surgery. *Cardiology*. 2012;122:12–9.

15. Farag M, Veres G, Szabo G, et al. Hyperbilirubinaemia after cardiac surgery: the point of no return. *ESC heart failure*. 2019;6:694–700.
16. Group KDIGO KAKI W. KDIGO Clinical Practice Guideline for Acute Kidney Injury. *Kidney Int Suppl*. 2012;1–138.
17. Michalopoulos A, Alivizatos P, Geroulanos S. Hepatic dysfunction following cardiac surgery: determinants and consequences. *Hepato-gastroenterology*. 1997;44:779–83.
18. Wang J, Yu W, Zhai G, et al. Independent risk factors for postoperative AKI and the impact of the AKI on 30-day postoperative outcomes in patients with type A acute aortic dissection: an updated meta-analysis and meta-regression. *Journal of thoracic disease*. 2018;10:2590–8.
19. Zhao H, Pan X, Gong Z, et al. Risk factors for acute kidney injury in overweight patients with acute type A aortic dissection: a retrospective study. *Journal of thoracic disease*. 2015;7:1385–90.
20. Zhou H, Wang G, Yang L, et al. Acute Kidney Injury After Total Arch Replacement Combined With Frozen Elephant Trunk Implantation: Incidence, Risk Factors, and Outcome. *J Cardiothorac Vasc Anesth*. 2018;32:2210–7.
21. Qiu Z, Chen L, Cao H, et al. Analysis of risk factors for acute kidney injury after ascending aortic replacement combined with open placement of triple-branched stent graft in type A aortic dissection: a new technique versus the traditional method in a single Chinese center. *Medical science monitor: international medical journal of experimental clinical research*. 2015;21:674–80.
22. Luo HY, Hu KJ, Liu ZY, et al. [Analysis of the risk factors of postoperative renal failure of type A aortic dissection]. *Zhonghua wai ke za zhi [Chinese journal of surgery]*. 2008;46:1070–2.
23. Xu S, Liu J, Li L, et al. Cardiopulmonary bypass time is an independent risk factor for acute kidney injury in emergent thoracic aortic surgery: a retrospective cohort study. *J Cardiothorac Surg*. 2019;14:90.
24. Anil Z. Apaydin SB, Posacioglu H. Fatih Islamoglu, Tanzer Calkavur, Tahir Yagdi, Mustafa Ozbaran and, Yuksel M. Perioperative Risk Factors for Mortality in Patients With Acute Type A Aortic Dissection. *Ann Thorac Surg*. 2002;74.
25. Yuan L, Liao PP, Song HC, et al. Hyperbilirubinemia Induces Pro-Apoptotic Effects and Aggravates Renal Ischemia Reperfusion Injury. *Nephron*. 2019;142:40–50.
26. Barateiro A, Domingues HS, Fernandes A, et al. Rat cerebellar slice cultures exposed to bilirubin evidence reactive gliosis, excitotoxicity and impaired myelinogenesis that is prevented by AMPA and TNF-alpha inhibitors. *Mol Neurobiol*. 2014;49:424–39.
27. Komardina E, Yaroustovsky M, Abramyan M, et al. Prometheus therapy for the treatment of acute liver failure in patients after cardiac surgery. *Kardiochir Torakochirurgia Pol*. 2017;14:230–5.
28. Cui J, Zhao H, Yi B, et al. Dexmedetomidine Attenuates Bilirubin-Induced Lung Alveolar Epithelial Cell Death In Vitro and In Vivo. *Crit Care Med*. 2015;43:e356-68.
29. Chu CM, Chang CH, Liaw YF, et al. Jaundice after open heart surgery: a prospective study. *Thorax*. 1984;39:52–6.
30. Mundth ED, Keller AR, Austen WG. Progressive hepatic and renal failure associated with low cardiac output following open-heart surgery. *J Thorac Cardiovasc Surg*. 1967;53:275–84.
31. Mitzner SR, Stange J, Klammt S, et al. Improvement of hepatorenal syndrome with extracorporeal albumin dialysis MARS: results of a prospective, randomized, controlled clinical trial. *Liver transplantation: official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society*. 2000;6:277–86.
32. Pan KC, McKenzie DP, Pellegrino V, et al. The meaning of a high plasma free haemoglobin: retrospective review of the prevalence of haemolysis and circuit thrombosis in an adult ECMO centre over 5 years. *Perfusion*. 2016;31:223–31.
33. Toomasian JM, Bartlett RH. Hemolysis and ECMO pumps in the 21st Century. *Perfusion*. 2011;26:5–6.

Tables

Table 1. Baseline characteristics of the included patients

Variables	Value
Preoperative	
Age, mean $\pm$ SD (years)	49.1 $\pm$ 11.0
Male, n (%)	223 (82.3%)
Co-morbidity	
Hypertension, n (%)	158 (58.3%)
Diabetes, n (%)	3 (1.1%)
Cerebrovascular disease, n (%)	16 (5.9%)
Previous cardiac surgery, n (%)	8 (2.9%)
APECHEII score, mean $\pm$ SD	9.2 $\pm$ 3.2
MELD score, mean $\pm$ SD	9.2 $\pm$ 4.1
SOFA score, mean $\pm$ SD	2.1 $\pm$ 1.5
MAP, mean $\pm$ SD (mm/Hg)	90.5 $\pm$ 12.9
TB, mean $\pm$ SD (μ mol/l)	25.0 $\pm$ 15.0
CB, mean $\pm$ SD (μ mol/l)	7.8 $\pm$ 8.1
WBC, mean $\pm$ SD (10^9 /l)	11.7 $\pm$ 5.0
Hb, mean $\pm$ SD (g/l)	136.1 $\pm$ 19.9
PLT, mean $\pm$ SD (10^9 /l)	157.2 $\pm$ 68.5
SCr, mean $\pm$ SD (μ mol/l)	111.1 $\pm$ 38.3
PT, mean $\pm$ SD (s)	12.0 $\pm$ 2.2
Intraoperative	
Type of operation	
Aortic valve, n (%)	227 (83.8%)
Aortic arch, n (%)	233 (86%)
Coronary artery, n (%)	46 (17%)
Operation duration, mean $\pm$ SD (h)	6.8 $\pm$ 1.6
CPB time, mean $\pm$ SD (min)	226.5 $\pm$ 60.9
ACC time, mean $\pm$ SD (min)	102.2 $\pm$ 28.0
The amount of blood transfusion (U)	18.8 $\pm$ 10.6
Postoperative	
APECHE II score	16.9 $\pm$ 2.5
SOFA score	12.1 $\pm$ 2.6
Meld score	18.6 $\pm$ 4.4
AST, mean $\pm$ SD (U /L)	347.6 $\pm$ 1387.2
ALT, mean $\pm$ SD (U/L)	164.4 $\pm$ 526.6
TB, mean $\pm$ SD (μ mol /l)	189.1 $\pm$ 74.6
CB, mean $\pm$ SD (μ mol /l)	110.7 $\pm$ 37.9
WBC, mean $\pm$ SD (10^9 /l)	12.9 $\pm$ 4.8
Hb, mean $\pm$ SD (g/l)	112.9 $\pm$ 16.6
PLT, mean $\pm$ SD (10^9 /l)	99.8 $\pm$ 49.5

SCr, mean± SD (μmol/l)	162.8±71.8
PT, mean± SD (s)	13.6±2.9
Peak TB level, mean ± SD (μmol/l)	150.9 ± 93.0
Peak TB level > 171μmol/l, n (%)	55 (20.4%)
Peak TB level > 340μmol/l, n (%)	12 (4.4%)
Time to peak TB level, mean ± SD (d)	3.8 ± 3.0
Time to peak TB level > 5 d, n (%)	47 (17.4%)

ACC, Aortic cross clamp; APECHEII, acute physiology and chronic health evaluation II; CB, conjugated bilirubin; CPB, cardiopulmonary bypass; Hb, [hemoglobin](#); MAP, [mean arterial pressure](#); MELD, model for end-stage liver disease; min, [minute](#); PLT, platelet; PT, prothrombin time; SCr, [serum creatinine](#); SD, [standard deviation](#); SOFA, sequential organ failure assessment; TB, [total bilirubin](#); WBC, white blood cell.

Table 2. The major outcomes of the included patients

Variables	Value
In-hospital mortality, n (%)	82 (30.3%)
Reoperation, n (%)	17 (6.3%)
AKI, n (%)	222 (81.9%)
Stage of AKI	
Stage 1, n (%)	102 (40.2%)
Stage 2, n (%)	34 (12.5%)
Stage 3, n (%)	84 (29.2%)
Use of CRRT, n (%)	50 (18.5%)
Use of PE/BA, n (%)	11 (14.1%)
Use of IABP, n (%)	1 (0.4%)
Use of ECMO, n (%)	7 (2.6%)
Use of vasoactive agent, n (%)	173 (63.8%)
Mechanical ventilation time, mean ± SD (days)	3.3 ± 5.7
The amount of blood transfusion, mean ± SD (U)	34.8 ± 43.3
In hospital time, mean ± SD (days)	19.8 ± 10.7
ICU stay time, mean ± SD (days)	6.6 ± 6.3
Onset of severe hyperbilirubinemia, mean ± SD (days)	2.8 ± 1.3

AKI, acute kidney injury; BA, bilirubin adsorption; CRRT, continuous renal replacement therapy; ECMO, extracorporeal membrane oxygenation; IABP, intra-aortic balloon pump; ICU, intensive care unit; PE, plasma exchange.

Table 3. Logistic regression analysis for postoperative AKI and CRRT

Characteristic	Univariate logistic regression		Multivariate logistic regression	
	OR (95%CI)	P value	OR (95%CI)	P value
Postoperative AKI				
Age	1.061 (1.032 - 1.092)	<0.001	1.056 (1.026 - 1.088)	<0.001
Hypertension (yes/no)	2.766 (1.469 - 5.208)	0.002		
Preoperative				
SCr	1.030 (1.014 - 1.047)	<0.001	1.026 (1.010 - 1.043)	0.002
APECHE II score	1.399 (1.213 - 1.615)	<0.001		
MELD score	1.130 (1.035 - 1.233)	0.007		
Operation duration	1.346 (1.089 - 1.633)	0.006		
CPB time	1.008 (1.001 - 1.014)	0.015	1.007 (1.000 - 1.013)	0.042
Postoperative CRRT				
Male (yes/no)	4.006 (1.192 - 13.462)	0.025		
Preoperative				
SCr	1.011 (1.003 - 1.018)	0.004	1.009 (1.002 - 1.016)	0.017
APECHE II score	1.333 (1.033 - 1.244)	0.008		
MELD score	1.083 (1.007 - 1.164)	0.032		
ACC time	1.011 (1.000 - 1.022)	0.049		

ACC, aortic cross clamp; AKI, acute kidney injury; APECHEII, acute physiology and chronic health evaluation II; CPB, cardiopulmonary bypass; CRRT, continuous renal replacement therapy; MELD, model for end-stage liver disease; SCr, [serum creatinine](#)

Table 4. Logistic regression analysis for in-hospital mortality

In-hospital mortality	Univariate logistic regression		Multivariate logistic regression (model 1)		Multivariate logistic regression (model 2)	
Characteristic	OR (95%CI)	P value	OR (95%CI)	P value	OR (95%CI)	P value
Age (yes/no)	1.035 (1.009 - 1.062)	0.008	1.005 (1.004 - 1.099)	0.033	1.046 (1.002 - 1.093)	0.040
Male (yes/no)	2.483 (1.106 - 5.575)	0.027				
Preoperative						
SCr	1.009 (1.002 - 1.016)	0.008				
APECHE II score	1.178 (1.082 - 1.282)	<0.001				
Intraoperative						
CPB time	1.007 (1.003 - 1.011)	0.002				
ACC time	1.012 (1.002 - 1.021)	0.014				
Postoperative						
MAP	0.948 (0.926 - 0.971)	<0.001	0.955 (0.922 - 0.990)	0.012	0.963 (0.930 - 0.977)	0.031
Re-operation (yes/no)	3.611 (1.324 - 9.852)	0.012				
The total amount of blood transfusion	1.046 (1.032 - 1.060)	<0.001	1.018 (1.003 - 1.033)	0.019	1.020 (1.005 - 1.034)	0.007
Mechanical ventilation time	1.533 (1.341 - 1.798)	<0.001				
AKI (yes/no)	28.350 (3.842 - 209.203)	0.001				
Stage of AKI		<0.001		<0.001		<0.001
Stage 1	5.859 (0.735 - 46.720)	0.095	2.424 (0.267 - 21.990)	0.431	2.218 (0.253 - 19.426)	0.472
Stage 2	10.138 (1.162 - 88.457)	0.036	4.560 (0.478 - 43.545)	0.187	3.698 (0.385 - 35.548)	0.257
Stage 3	165.053 (21.354 - 1275.729)	<0.001	46.134 (5.436 - 391.525)	<0.001	46.318 (5.666 - 378.663)	46.318
Use of CRRT (yes/no)	23.756 (10.359 - 54.481)	0.001				
Use of PE/BA (yes/no)	11.527 (2.432 - 54.631)	0.002				
Use of ECMO (yes/no)	14.842 (1.757 - 125.357)	0.013	20.795 (1.620 - 266.917)	0.02	10.130 (0.900 - 113.958)	0.061
Peak TB concentration	1.009 (1.005 - 1.012)	<0.001	1.017 (1.002 - 1.032)	0.023		
Peak TB ≥121μmol/l, (yes/no)	3.899 (2.220 - 6.847)	<0.001			2.681 (1.119 - 6.425)	0.027
Peak TB >171μmol/l, (yes/no)	3.785 (2.045 - 7.005)	<0.001				
Time to peak TB concentration	1.441 (1.277 - 1.626)	<0.001				
ICU stay time	1.268 (1.172 - 1.371)	<0.001				

Model 1 was established by analysis of peak TB concentration; Model 2 was established by subgroup analysis of peak bilirubin concentration greater or less than 121.2 μmol/l

ACC, aortic cross clamp; AKI, acute kidney injury; APECHEII, acute physiology and chronic health evaluation II; BA, bilirubin adsorption; CPB, cardiopulmonary bypass; CRRT, continuous renal replacement therapy; ECMO, extracorporeal membrane oxygenation; ICU, intensive care unit; MAP, mean arterial pressure; PE, plasma exchange; SCr, serum creatinine; TB, total bilirubin

Figures

Figure 1

Patient inclusion flow chart

Figure 2

The averaged serum total bilirubin concentration during the postoperative seven days

Figure 3

Receiver operator curve (ROC) of peak TB concentration predicting in-hospital mortality (area under the curve: 0.68, 95% CI: 0.614 - 0.758)

Figure 4

Long-term survival results of (A) all patients, (B) patients without AKI and those with stage 1, 2, or 3 AKI, (C) patients without the use of CRRT and those with CRRT, (D) patients without the use of ECMO and those with ECMO, (E) patients with post-operation peak TB $\geq 121.2 \mu\text{mol/l}$ and those with post-operation peak TB $< 121.2 \mu\text{mol/l}$.