

Prevalence and Associated Factors Of Preoperative Abnormal Electrocardiography Among Older Surgical Patients In Southern Ethiopia: Multicenter Cross-Sectional Study

Addisu Mossie

Hawassa University

Hailemariam Getachew (✉ hailemariamgetachew23@gmail.com)

Dilla University

Timsel Girma

Dilla University

Hailemariam Mulugeta

Dilla University

Belete Destaw

Dilla University

Aschalew Besha

Hawassa University

Adanech Shiferaw

Hawassa University

Research Article

Keywords: Abnormal ECG, older, preoperative, surgical

Posted Date: July 17th, 2023

DOI: <https://doi.org/10.21203/rs.3.rs-3086228/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: The prevalence of abnormal electrocardiography (ECG) increases with aging, and these abnormalities may have an impact on anesthesia management.

Objective: The aim of this study was to determine the prevalence and associated factors of preoperative abnormal electrocardiography among older surgical patients at selected teaching hospitals in southern Ethiopia, from February 15 to June 15, 2022.

Methodology: A multicenter cross-sectional study was conducted on 246 elderly surgical patients. A binary logistic regression model was used, and a P-value < 0.05 was considered statistically significant.

Result: In the current study, 120 (48.78%) of older surgical patients had abnormal preoperative ECGs. In terms of severity, 44.16% were with major ECG abnormalities. The most common ECG abnormalities were left axis deviation, left ventricular hypertrophy, and ST segment changes. The presence of comorbidity (AOR=3.44, P =0.001), age \geq 70 years (AOR=2.5, P=0.011), history of angina (AOR=5.9, P=0.011), history of smoking (AOR=5.07, P=0.024) and urban residency (AOR=1.89, P=0.039) were associated factors for an abnormal ECG.

Conclusion and recommendation: Our study showed that the prevalence of preoperative abnormal ECG increased with age, even in asymptomatic patients with no risk factors and younger than 65 years. Therefore, a preoperative ECG is advisable in all older patients.

Background

Aging is a universal and progressive physiological process characterized by changes in organ and tissue structure as well as functional status(1). As people get older, their hearts and arteries go through a variety of anatomical and functional changes. Moreover, the prevalence of cardiovascular diseases such as systemic hypertension, atherosclerosis, acute myocardial infarction, and congestive heart failure has increased, further limiting the cardiovascular system's function(2).

Surgical interventions in elderly patients are associated with increased rates of morbidity and mortality, possibly due to an increased prevalence of comorbidities, decreased physiological reserve, or both (3). The aim of the preoperative assessment is to reduce perioperative surgical and anesthetic complications and to optimize them as early as possible through history, clinical examination, and investigations(4, 5).

According to the American Society of Anesthesiologists (ASA) task force on preanesthesia evaluation preoperative investigations are classified as routine and indicated. Routine tests are those performed without any special clinical rationale or aim including a panel of blood, urine tests, and chest x-ray, and ECG) and Indicated tests are those performed for a specific clinical indication or goal, such as confirming a clinical diagnosis, assessing the severity and the progression of disease, or determining the efficacy of medication(6).

Of all investigations, ECG was commonly ordered investigation for the elderly due to the prevalence of age-related cardiovascular changes. ECG is a sophisticated galvanometer and sensitive electromagnet that can detect and record changes in electromagnetic potentials. The ECG (12-lead) is the primary clinical tool for noninvasive assessment of cardiac electrical function and is one of the most widely used, inexpensive, and convenient assessment modalities used to screen for cardiovascular disease(7, 8).

Due to their age and the prevalence of multiple comorbidities, elderly surgical patients are at increased risk of preoperative ECG abnormalities. Elderly patients are more likely to have cardiovascular disease, cerebrovascular disease, and diabetes mellitus, all of which increase the overall risk of the cardiovascular system(3).

Abnormal electrocardiograms become more common as people get older. Evidence showed that the prevalence of abnormal ECG is high in the elderly population. Studies conducted in the United States, China, Brazil, and Nigeria showed that the prevalence of abnormal ECGs was 75.2%, 75.2%, 88.85%, and 70%, respectively (9–12).

According to the American College of Cardiology/American Heart Association (ACC/AHA) guideline, Surgical procedures are classified as low (< 1%), intermediate (1–5%), or high (> 5%) risk for the development Perioperative Cardiac Events (PCE) within 30 days after surgery. Preoperative ECG is not recommended for patients undergoing low-risk surgery but is indicated for patients with risk factors such as known heart disease, peripheral arterial disease, cerebrovascular disease, or other significant structural heart disease (13).

Previous studies in this area have had inconsistent results on the prevalence and associated factors of abnormal ECG. In addition, no consensus on the lowest age for routine preoperative ECG in elderly patients presenting for surgery, particularly in patients without specific risks. The present study aimed to evaluate the prevalence and contributing factors of preoperative ECG abnormalities in elderly surgical patients aged 50 years and older.

Methodology

A multi-center cross-sectional study was conducted from February 15, 2022 to June 15, 2022. Ethical clearance was obtained from the institutional review board. Patients aged over 50 years scheduled for elective surgery during study period who have the willingness to participate in this study at selected hospitals in southern Ethiopia were included.

Patients presenting for minor surgery requiring only local anesthesia, older patients who were mentally impaired and unable to give information, patients undergoing emergency surgery, and patients who underwent two or more operations if the ECG was assessed during the first surgery were excluded from the study.

Sample Size and sampling procedure

The sample size was calculated using software Epiinfo version 7 based on the objectives of the study. A previous study published in 2014, and conducted in, Ibadan, Nigeria on elderly patients revealed that, the prevalence of abnormal ECG was 70%(12). By taking proportion 70%, 95% confidence interval, and margin of error = 5%. Finally, the sample size becomes 323.

Since the total source population was less than 10,000(number of older surgical patients undergoing surgery at selected hospitals (N = 850) by making adjustment:
$$n_f = n (1 + (n/N)) = 323 / (1 + (323/850)) = 234.05 \approx 234$$
 patients. Adding 5% of the non-response rate gives a final sample size of 246 patients. From the situational analysis, the total number of older surgical patients in the last year at HUCSH, DURH, and WSUCSH Universities were 406, 140, and 304 respectively.

The study populations were taken from each hospital with a proportion allocation formula by dividing the number surgery in each hospital by the total number of surgery at three hospitals multiplied by the sample size ($n = 246$). $H = n_1/N * n$, $D = n_2/N * n$, and $W = n_3/N * n$ where n_1 , n_2 , and n_3 were the total number of surgery in older patients in the last one Year at HUCSH, DURH, and WSUCSH respectively. So the final study populations selected from three hospitals were 118, 40, and 88 at HUCSH, DURH, and WSUCSH respectively.

Data collection procedures

The data was collected from patient's history, medical charts, and preoperative anesthesia record through interviewer administered standardized questionnaire adopted from previous literatures (14–17). All ECGs for older patients planned for elective surgery who fulfill the inclusion criteria at selected hospitals was reviewed preoperatively by the data collectors after the assigned anesthetist do his preoperative anesthesia evaluation.

The result was interpreted in terms of normal and abnormal by the staff internist. To minimize bias the investigator and data collectors were not involved in the interpretation of patients ECG results at all three sites and ECG findings were classified as normal and abnormal according to the Minnesota Code classification(16). Further, the ECG was classified as major and minor abnormalities.

Operational definition

Older surgical patient: Patients aged ≥ 50 years undergoing surgery(18).

Normal ECG: Absence of any alteration of the ECG.

Abnormal ECG: Any ECG change beyond normal sinus rhythm (ST-segment changes, T-wave change, AV nodal block, BBB, ventricular hypertrophy, arrhythmias, prolonged QT, and others).

Minor ECG abnormalities: Include minor isolated Q-QS waves, minor isolated ST-T abnormalities, high R waves, incomplete RBBB, minor QT prolongation (QT interval $> 112\%$), short PR interval, LAD, RAD, frequent PVC, and other minor abnormalities(19).

Major ECG abnormalities: Included major ventricular conduction defect; definite MI (defined as the presence of major Q-wave abnormalities); possible MI (defined as the presence of minor Q-QS wave plus major ST-T abnormalities); major isolated ST abnormalities; LVH plus major ST-T abnormalities; major Atrioventricular(AV) conduction abnormalities; and major QT prolongation (QT interval > 116% or if QRS interval > 120 ms), and other major arrhythmias(19).

Data analysis procedure

Once data was collected it was entered with Epidata version 4.6 then exported and analyzed by STATA version 16. The data analysis was carried out using both descriptive and inferential statistics, describing qualitative variables as frequencies and percentages, and quantitative variables as mean and standard deviation (SD). Normality was checked using Kolmogorov–Smirnov test and outliers were checked by both graphical and non-graphical methods. Bivariate analysis was done by binary logistic regression.

Variables that showed an association with preoperative abnormal ECG in bivariate analysis (p-value < 0.2) were entered into a multivariate logistic regression model to identify preoperative abnormal ECG independent factors. A backward stepwise elimination technique was used to build the logistic regression model. The model fitness was evaluated using the Hosmer-Lemeshow goodness of fit test. Multicollinearity between independent variables was checked using variance inflation factor (VIF) before entering the multivariable model, and the mean VIF was = 1.19. To determine statistical significance, a p-value of less than 0.05 was used. For both the crude and adjusted odds ratios, a 95% confidence interval was provided. The data was presented using numbers, frequencies, tables, charts, and figures as appropriate.

Results

A total of 246 elderly surgical patients, aged 50 years and older, appointed for elective surgery and who had a preoperative ECG were enrolled in this study. The minimum age of the patients was 50 years and the maximum age was 98 years. The median age was 61.5 (IQR; 56, 70), with the majority of 102 (41.46%) being in the second (60–69) age group. The majority of patients 167(67.89%) were males, 196(80%) patients have normal weight, 142(57.72%)resided rurally and 91(36.99%) were farmers

Preoperative baseline characteristics and health conditions of study participants

Regarding the preoperative baseline characteristics, the majority of the study participants (139 (56.5%) were ASA II, followed by ASA I (92 (37.3%), 200 (81.3%) of the patients have a MET Equivalent of 4–10, 55 (22.36%) of the elderly patients were frail and on the base of surgical complexity nearly half 126 (51.22%) of the patients had SHAPE III surgery. Additionally, 16 (6.5%) patients were former smokers, 41 (16.4%) patients had a history of alcohol, 20 (8.13%) had a history of angina, and 28 (11.38%) had a family history of CAD.

Majority of patients were underwent urologic surgery followed by general surgery and orthopedic surgery respectively (Fig. 2)

In terms of preoperative health status, 70 (28.46%) of study participants had a history of comorbid conditions, with hypertension being the most common (present in 34 (48.57%) patients, followed by diabetes mellitus (20 (29.11%)). When we see the distribution of cases according to the surgical specialty the majority of cases (87.37%) were referred from urology, followed by general surgery (63.11%) and orthopedics (42.07%) (Fig. 1)

Figure 2: *Distribution of cases according to surgical specialty and gender among older surgical patients operated at selected teaching hospitals of southern Ethiopia, February15-June 15, 2022(n = 246).*

Prevalence of preoperative ECG abnormality

In the current study, the prevalence of preoperative abnormal ECG in elderly surgical patients was 48.78% (Fig. 3), with males accounting for 33.33%. In terms of severity, 67 (55.3%) of all abnormal ECGs were classified as minor, while 53 (44.16%) were classified as major ECG abnormalities.

Left axis deviation (12.18%), LVH (11.67%), and ST segment changes (9.13%) were the most common ECG abnormalities observed in descending order of frequency. Forty-three (35.8%) patients had more than one ECG abnormality (Table 1).

In addition, the first age group (50–59 years) had a greater prevalence of ST abnormalities (4.06%), LVH (2.53%), and T-wave inversion (3.03%), while the second (60–69 years) and third age groups (70–79 years) had a higher prevalence of LAD (6.08%), RAD (3.03%) and right BBB (4.56%). The fourth age group (80 years and older) has a higher prevalence of poor R-wave progression (1.01%) and fascicular block (1.01%) (Table 1).

Table 1

Type and number of preoperative ECG abnormality based on age groups among older surgical patients operated at selected teaching hospitals in southern Ethiopia – 2022GC.

Common ECG abnormalities	Total number of abnormalities	50–59 years	60–69 years	70–79 years	≥ 80 years
LAD	24(12.18%)	3(1.52%)	9(4.56%)	7(3.55)	5(2.53%)
LVH	23(11.67%)	5(2.53%)	8(4.06%)	7(3.55)	4(2.03%)
ST segment changes	18(9.13%)	8(4.06%)	7(3.55)	3(1.52%)	0
Sinus bradycardia	17(8.62%)	2(1.01%)	6(3.04%)	7(3.55)	2(1.01%)
RBBB	13(6.59%)	4(2.03%)	6(3.04%)	3(1.52%)	0
T wave inversion	13(6.59%)	6(3.03%)	3(1.52%)	0	4(2.03%)
RAD	9(4.56%)	1(0.5%)	5(2.53%)	3(1.52%)	0
Sinus Tachycardia	9(4.56%)	5(2.53%)	1(0.5%)	3(1.52%)	0
RVH	8(4.06%)	3(1.52%)	4(2.03%)	1(0.5%)	0
Poor R-wave progression	7(3.55%)	1(0.5%)	1(0.5%)	4(2.03%)	2(1.01%)
Fascicular block	6(3.04%)	0	1(0.5%)	3(1.52%)	2(1.01%)
LBBB	6(3.04%)	1(0.5%)	2(1.01%)	3(1.52%)	0
AV block	4(2.03%)	0	2(1.01%)	1(0.5%)	1(0.5%)
Low voltage ECG	4(2.03%)	2(1.01%)	0	1(0.5%)	1(0.5%)
Others	36(18.24)	11(5.58%)	16(8.12%)	6(3.04%)	3(1.52%)

Associated Factors of preoperative abnormal ECG

The bivariate analysis for this study revealed that age greater or equal to 70 years, place of residence, ASA III-IV, MET equivalent less than four, frailty, history of comorbid disease, duration of comorbid disease greater than 5 years, smoking history, history of alcohol intake, history of angina and family history of CAD were significantly associated with preoperative abnormal ECG.

A Multivariate logistic regression analysis found that history of comorbid disease, Age \geq 70 years, History of angina, History of smoking, and urban residence were independently associated with preoperative abnormal ECG (Table 2.)

Table 2

Multivariate logistic regression-showing factors associated with preoperative abnormal ECG among older surgical patients at selected teaching hospitals in Southern Ethiopia, February 15 – June 15, 2022, (n = 246).

Variable	Category	COR (95%CI)	AOR(95% CI)	P-value
Age	50–69 years	1	1	0.011**
	≥ 70 years	2.67(1.15,3.49)	2.5(1.23-5.0)	
Place of residency	Rural	1	1	0.039**
	Urban	2.14(1.28–3.58)	1.89(1.03–3.48)	
History of comorbidity	No	1	1	0.001**
	Yes	5.07(2.71,9.48)	3.44(1.65–7.18)	
History of smoking	No	1	1	0.024**
	Yes	5.98(1.38,17.94)	5.07(1.24–20.7)	
History of angina	No	1	1	0.011**
	Yes	6.78(1.93–23.73)	5.9(1.49–23.47)	

Key: **: significant in the Multivariate logistic regression (p-value < 0.05), 1: Reference, AOR adjusted odds ratio, COR crude odds ratio, CI confidence interval.

Discussion

Abnormal ECGs become more common as people grow older due to physiological changes, the prevalence of comorbid disease, or both. The present study was conducted to determine the prevalence and associated factors of preoperative abnormal ECG among older surgical patients.

Prevalence of ECG Abnormalities

Our findings revealed that the prevalence of preoperative abnormal ECG in older surgical patients was 48.78%(CI 42–56%), which is lower than the findings of Friedman, J et al. in New York (70.1%)(20), Souza et al. in Brazil (88.85%)(11), and Awana EE, et al. in Nigeria (70%)(12).

On the other hand, our finding is higher than a study conducted in India by Prabal Bharali et al., who reported a prevalence of abnormal ECG among elderly patients of 14.45% (21) and Ghimire et al. of 25% (22). The discrepancy could be due to differences in the socio-economic status, age, selection criteria, study design, sample size, environmental and genetic variations.

Type of ECG abnormality

The prevalence of LAD in our study was 12.18%. It was also observed that the prevalence increased with age, with 1.52% in the first age group (50–59 years), 4.56.21%, and 3.55% in the second and third age groups, respectively. Our findings are consistent with those of an Indian study by Khane et al(23).

In addition, the current study revealed that the prevalence of LVH was 11.67%, which was greater in the second (60–69 years) and third age groups (70–79 years) 4.56%, and 3.55%, respectively than in the first age group (2.53%). The result of our study is in line with the Spanish study by Santos et al.(24) which reports LVH as the second most prevalent abnormal ECG finding present in 11.8% of patients and increased with age. The increasing prevalence of LVH with age can be attributed to cardiac muscle hypertrophy(1).

Furthermore, this study found that the prevalence of ST segment changes was 9.13% and the majority of subjects were from the second age group (60–69 years) 4.06%. The result of the current study is almost comparable with the study done in Turkey which reports the prevalence of ST-segment abnormalities was 12%(25).

Medical history of study participants

Patient histories in this study revealed that 28.46% of study participants had a history of comorbid disease and were taking medication. The two most common comorbidities observed in elderly surgical patients were hypertension (48.57%) and diabetes mellitus (29.11%), which is comparable to the Nigerian study by Awana EE, et al(12) and the Spanish study by Santos et al (24).

Factors associated with preoperative abnormal ECG

According to this study, smokers were 5 times more likely to have a preoperative ECG abnormality than non-smokers (AOR = 5.07). Smoking was found to be an important predictor of preoperative ECG abnormalities, and our study's finding was supported by studies conducted in India(26) and china(27). This is due to cardiac effects of nicotine a conduction system and electrophysiology which contributes to induce arrhythmias and other ECG changes(28).

Preoperative ECG abnormalities were associated with an older age of 70 years or more in our study. Patients 70 years and older were 2.5 times more likely to develop an ECG abnormality than patients 50 to 69 years of age (AOR = 2.5 95% (1.23-5.0)). Our result agrees with a study by Correll et al. which reports that older age is an independent predictor of an abnormal ECG(29).

In terms of comorbid illnesses, our study revealed patients who have a history of comorbid disease were 3.4 times more likely to have preoperative abnormal ECG than those who do not have(AOR = .3.44 & 95%CI (1.65–7.18) other studies done in China by Yu et al.(27), in Nigeria by Awana EE, et al(12). and Brazil by Gutiérrez, et al.(30) also confirmed that patients who have coexisting medical illness have a higher chance of having abnormal ECG. This finding is not surprising given that comorbid conditions like hypertension and diabetes mellitus are known to have long-term effects on cardiac function. Furthermore, drugs used in the treatment of comorbid illnesses can cause various types of ECG changes(2).

Our finding also indicated that patients who had a previous history of angina had an odd ratio of a 5.9 times higher risk of abnormal ECG than those who had no history of angina (AOR = 5.9 95% CI (1.49–23.47)). This is in agreement with a study conducted in the USA by Linda et al.(9), which reports a prior history of angina had a significant impact on preoperative abnormal ECG.

Regarding place of residence, our findings revealed that older patients in urban areas were 1.89 times more likely than those living in rural areas to have a preoperative abnormal ECG (AOR = 1.89 95% CI (1.03–3.48)). In contrast to our study a Chinese study found that living in rural areas was significantly associated with major arrhythmias(27). This could be due to differences in the study population's lifestyles and socioeconomic status of patients living in the urban and rural areas.

Strength of the study

The first strength is that we included elderly surgical patients from a variety of surgical specialties in three different teaching hospitals in southern Ethiopia which increases generalizability. Second, since this is the first study of its kind in Ethiopia, it will serve as a baseline for future valuable research in the area of interest.

Limitation of the study

Our study is not without limitations, the following are the probable weaknesses of this study.

First, the professionals involved in reading and interpreting ECG results at all three sites were different, and the ECGs were performed on various electrocardiograph devices. This may have an impact on the interpretation of ECG results, resulting in bias even if we try to control this effect by utilizing a validated tool such as the Minnesota code of ECG classification.

Second, the hospital-based data collection limits the generalizability of our findings to the general population.

Third, since its cross-sectional study there was follow-up time to assess the outcome of patients with preoperative abnormal ECG.

Conclusion

The present study revealed that the prevalence of preoperative abnormal ECG increases with age. Nearly half (48.78%) of elderly patients appointed for elective surgery have an abnormal ECG. In terms of severity, more than half of (55.3%) abnormalities were classified as minor ECG abnormalities. Left axis deviation, LVH, ST segment alterations, and sinus bradycardia were the most common ECG abnormalities observed in descending order of frequency.

Furthermore, our findings show that history of comorbid disease, age ≥ 70 years, angina history, smoking history, alcoholic history, and urban residence are the independent risk factors of preoperative abnormal ECG. This strengthens the importance of preoperative ECG in patients younger than 65 years, especially

in those with risk factors like diabetes, hypertension, history of angina, and cardiovascular disease before any elective surgery.

Recommendation

Preoperative ECG is advisable in all older patients, including those under 65 years of age without risk factors because preoperative ECG allows for the detection of asymptomatic significant cardiac abnormalities. Further prospective cohort studies with a comparator group, is demanded to assess the perioperative implications and outcome of patients with preoperative abnormal ECG with an adequate follow-up time.

Declarations

Provenance and peer review

Not commissioned, externally peer-reviewed

Ethical consideration and consent to participate

Ethical approval was obtained from Dilla University's Institutional Review Board with Ref. No: duchm/irb/048/2022 and protocol unique number of: duirb/048/22-02.

All methods were carried out in accordance with Dilla University research policy and guideline

Informed consent was obtained prior to beginning data collection, with the aim and methods of the research made clear to all participants.

Confidentiality and anonymity were maintained throughout the research process. It was made clear to each study participant that they have a right of withdrawal at any time.

Consent for publication:

'Not applicable'

Availability of data and materials

The datasets used and/or analysed during the current study available from the corresponding author on reasonable request

Competing interests:

No conflict of interest

Funding

No Funding

Authors' contributions

"All investigators are engaged in Conception and design of the study, Data analysis and interpretation, Critical revision and intellectual input of the study".

Acknowledgements

Our heartfelt appreciation also goes to our data collectors, supervisors, and employees at all locations for their genuine support and collaboration.

References

1. Tonner PH, Kampen J, Scholz J. Pathophysiological changes in the elderly. *Best Pract Res Clin Anaesthesiol.* 2003;17(2):163–77.
2. Giovanardi P, Vernia C, Tincani E, Giberti C, Silipo F, Fabbo A. Combined effects of age and comorbidities on electrocardiographic parameters in a large non-selected population. *J Clin Med.* 2022;11(13):3737.
3. Loran DB, Hyde BR, Zwischenberger JB. Perioperative management of special populations: the geriatric patient. *Surg Clin.* 2005;85(6):1259–66.
4. Garcia-Miguel F, Serrano-Aguilar P, Lopez-Bastida J. Preoperative assessment. *The Lancet.* 2003;362(9397):1749–57.
5. Oresanya LB, Lyons WL, Finlayson E. Preoperative assessment of the older patient: a narrative review. *JAMA.* 2014;311(20):2110–20.
6. Pasternak LR, Arens JF, Caplan RA, Connis RT, Fleisher LA, Flowerdew R, et al. Practice advisory for preanesthesia evaluation: a report by the American Society of Anesthesiologists Task Force on Preanesthesia Evaluation. *Anesthesiology.* 2002;96(2):485–96.
7. Perez MV, Dewey FE, Tan SY, Myers J, Froelicher VF. Added value of a resting ECG neural network that predicts cardiovascular mortality. *Ann Noninvasive Electrocardiol.* 2009;14(1):26–34.
8. Rajni R, Kaur I. Electrocardiogram signal analysis-an overview. *Int J Comput Appl.* 2013;84(7):22–5.
9. Liu LL, Dzankic S, Leung JM. Preoperative electrocardiogram abnormalities do not predict postoperative cardiac complications in geriatric surgical patients. *J Am Geriatr Soc.* 2002;50(7):1186–91.
10. Xu L, Yu C, Jiang J, Zheng H, Yao S, Pei L, et al. Major adverse cardiac events in elderly patients with coronary artery disease undergoing noncardiac surgery: a multicenter prospective study in China. *Arch Gerontol Geriatr.* 2015;61(3):503–9.
11. Souza FSd, Pedro JRP, Vieira JE, Segurado AVR, Botelho MPF, Mathias LAdST. The validity of the electrocardiogram accomplishment in the elderly surgical patient preoperative evaluation. *Revista brasileira de anesthesiologia.* 2005;55:59–71.

12. Awana EE, Adigun TA. ECG Changes in the Elderly Urology Patients during Pre-Operative Assessment in Ibadan Nigeria. *Anesth Clin Res*. 2014;5(11):1–3.
13. Fleisher LA, Fleischmann KE, Auerbach AD, Barnason SA, Beckman JA, Bozkurt B, et al. 2014 ACC/AHA guideline on perioperative cardiovascular evaluation and management of patients undergoing noncardiac surgery: a report of the American College of Cardiology/American Heart Association Task Force on practice guidelines. *J Am Coll Cardiol*. 2014;64(22):e77–e137.
14. Dzankic S, Pastor D, Gonzalez C, Leung JM. The prevalence and predictive value of abnormal preoperative laboratory tests in elderly surgical patients. *Anesth Analgesia*. 2001;93(2):301–8.
15. Davenport DL, Ferraris VA, Hosokawa P, Henderson WG, Khuri SF, Mentzer RM Jr. Multivariable predictors of postoperative cardiac adverse events after general and vascular surgery: results from the patient safety in surgery study. *J Am Coll Surg*. 2007;204(6):1199–210.
16. Prineas RJ, Crow RS, Zhang Z-M. The Minnesota code manual of electrocardiographic findings. Springer Science & Business Media; 2009.
17. Chassot PG, Delabays A, Spahn D. Preoperative evaluation of patients with, or at risk of, coronary artery disease undergoing non-cardiac surgery. *Br J Anaesth*. 2002;89(5):747–59.
18. Kowal P, Dowd JE. Definition of an older person. Proposed working definition of an older person in Africa for the MDS Project. Volume 10. Geneva: World Health Organization; 2001. p. 51889286. 2.1.
19. Denes P, Garside DB, Lloyd-Jones D, Gouskova N, Soliman EZ, Ostfeld R, et al. Major and minor electrocardiographic abnormalities and their association with underlying cardiovascular disease and risk factors in Hispanics/Latinos (from the Hispanic Community Health Study/Study of Latinos). *Am J Cardiol*. 2013;112(10):1667–75.
20. Friedman A, Chudow J, Merritt Z, Shulman E, Fisher J, Ferrick K, et al. Electrocardiogram abnormalities in older individuals by race and ethnicity. *J Electrocardiol*. 2020;63:91–3.
21. Bharali P, Taye MK. Prevalence of ECG Abnormalities during Routine Preoperative Evaluation-A Retrospective Single Institute Study.
22. Ghimire A, Rahman T, Bhattarai B, Koirala S, Khatiwada S, Dhital D. Routine preoperative electrocardiogram in patients over 40 years of age: a hearty need or a customary practice? *Health Renaissance*. 2011;9(2):73–7.
23. Khane RS, Surdi AD, Bhatkar RS. Changes in ECG pattern with advancing age. 2011.
24. Santos JPA, Ribeiro ALP, Andrade-Junior D, Marcolino MS. Prevalence of electrocardiographic abnormalities in primary care patients according to sex and age group. A retrospective observational study. *Sao Paulo Med J*. 2017;136:20–8.
25. Biteker M, Duman D, Tekkeşin A. Predictive value of preoperative electrocardiography for perioperative cardiovascular outcomes in patients undergoing noncardiac, nonvascular surgery. *Clin Cardiol*. 2012;35(8):494–9.
26. Trikha S, Singh N. The Role of Routine ECG Testing in Preoperative Evaluation Prior to Non Cardiac Surgery. *Heart India*. 2015;3(3):72.

27. Yu L, Ye X, Yang Z, Yang W, Zhang B. Prevalences and associated factors of electrocardiographic abnormalities in Chinese adults: a cross-sectional study. BMC Cardiovasc Disord. 2020;20(1):1–11.
28. Swathi K, Garimella S. Study of ECG Effects in Smokers and Normals. J Pharm Sci Res. 2015;7(3):163.
29. Correll DJ, Hepner DL, Chang C, Tsen L, Hevelone ND, Bader AM. Preoperative electrocardiograms: patient factors predictive of abnormalities. J Am Soc Anesthesiologists. 2009;110(6):1217–22.
30. Martínez DG, Jiménez-Méndez C, Hernández RM, Hernández-Aceituno A, Roca AP, Torres RA. Incidence of electrocardiographic alterations in the preoperative period of non-cardiac surgery. Revista Española de Anestesiología y Reanimación (English Edition). 2021;68(5):252–7.

Figures




Figure 1

Type of medical illness present in older surgical patients operated at selected hospitals of southern Ethiopia, February15-June 15, 2022(n=246).




Figure 2

Distribution of cases according to surgical specialty and gender among older surgical patients operated at selected teaching hospitals of southern Ethiopia, February15-June 15, 2022(n=246).




Figure 3

Prevalence of preoperative abnormal ECG among older surgical patients at three teaching hospitals of southern Ethiopia, February 15-June 15, 2022(n=246).