Psychological distress and somatization in immigrants in primary health care practices.

Rosa García-Sierra

Primary Care Research Institute Jordi Gol

María Isabel Fernández-Cano (✉ mariaisabel.fernandezc@uab.cat)

Universitat Autònoma de Barcelona <https://orcid.org/0000-0002-4267-2345>

Josep Maria Manresa-Domínguez

Primary Care Research Institut Jordi Gol

María Feijoo-Cid

Universitat Autònoma de Barcelona

Eduard Moreno Gabriel

Primary Care Research Institut Jordi Gol

Antonia Arreciado-Marañón

Universitat Autònoma de Barcelona Facultat de Veterinària

Francesc Ramos-Roure

Institut Català de la Salut

Jordi Segura

Ramon Llull University, Faculty of Psychology

Pere Torán-Monserrat

Primary Care Research Institut Jordi Gol

Research article

Keywords: Emigrants and Immigrants, Somatization disorders, Stress, Psychological, Primary Health Care, Vulnerable populations

Posted Date: June 10th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-30980/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at Healthcare on December 13th, 2020. See the published version at <https://doi.org/10.3390/healthcare8040557>.

Abstract

Background

The process of international migration causes a situation of vulnerability in people's health and greater difficulty in coping with disease. Furthermore, the adversities suffered during migration can trigger reactive signs of stress and cause anxious, depressive, confusional and somatic symptoms. This article studies the relationships between psychosocial risk, psychological distress and somatization in immigrants from 4 communities: Maghrebis, sub-Saharanans, centro-South Americans and Hindustanis.

Methods

A cross-sectional study was carried out with questionnaires on 602 immigrants who were surveyed in the Primary Care Centres of an urban area of Catalonia. The instruments used were the Demographic Psychosocial Inventory (DPSI), the Brief Symptom Inventory (BSI) and the Somatic Symptom Inventory (SSI).

Results

The average psychosocial risk obtained was 0.35, with the highest values in the sub-Saharan community. Psychological distress showed a mean value of 0.66, with the sub-Saharan community scoring the lowest in all dimensions except depression. The average somatization values were 1.65, with the sub-Saharan community scoring the least. The female gender is a risk factor for somatization and psychological distress.

Conclusions

Perceived psychosocial risk is a predictor of psychological distress, but not somatization, suggesting that the use of more adaptive coping strategies could minimize the effect of the migration process on somatizations.

Background

The concept of migration is defined as "The movement of persons away from their place of usual residence, either across an international border or within a State" (1). The number of international migrants has increased in the last 50 years and it is estimated that 272 million people live in a country different from that of their birth, triple that of 1970, with 48% of them being women (2). In Spain, resident foreigners represent 10% of the population (3), the most recent being those from the migratory movement between 2000 and 2007 motivated by the demand for labour in the construction and domestic service sector mainly and which was later slowed by the economic crisis (INE, 2008). In 2016, there was a new rise in the number of registered resident foreigners caused by conflicts in their countries of origin and also by the reunification of families of immigrants already established in Spain. In Catalonia, population growth in recent years has been due to the immigration of the foreign population, which in 2019 represented 14% of the population (3).

The migratory process is a human experience that affects people's health and is an emerging public health problem that needs a response. Migrating involves high levels of emotional stress that affect health and that can manifest themselves physically and psychically. It is comprised of several phases, and in each one of them the health of immigrants is conditioned by different determinants. In the pre-migration phase, epidemiological profiles of the origin, environmental policies, and personal exposure to conflicts or human rights violations are decisive in defining the state of health. In the migratory movement phase, the duration, circumstances and conditions of the trip are the factors that most affect health. Finally, in the arrival phase, the migration policies of the destination country, access to health systems and the sensitivity of health professionals towards cultural and linguistic differences act as decisive factors on the health of migrants (4). Added to the determinants of the above-mentioned migratory process are others such as loneliness, insecurity, stress suffered during the migratory trip and cultural shock upon

arrival in the new country, creating a situation in which the person will have difficulties anticipating and facing the impact of disease, thereby increasing their vulnerability (5). On the other hand, and even with having access to the health system, some groups of migrants may find it difficult to express their symptoms or understand therapeutic instructions, not only because of language barriers but also because the causal mechanisms of the disease may have different cultural constructs, especially the processes related to mental health. Furthermore, it should not be forgotten that having to move through the different healthcare levels is an added difficulty for people who come from systems of disorder (6). A review of reviews published in 2016 suggests that first generation migrants are at increased risk of mental ill-health, including common mental disorders and psychotic disorders(7).

All the afore-mentioned adversities can affect the person's homeostasis and the functioning of the hypothalamic-pituitary-adrenal medullary axis, causing symptoms of psychological distress in the area of depression (sadness and crying), in the area of anxiety (irritability or insomnia), in the confusional area due to an increase in cortisol, as well as somatizations manifested by fatigue, osteoarticular discomfort or headache, among other symptoms. Somatic symptoms may appear in response to chronic stress and are also known as *functional somatic symptoms (FSS)*, *medically unexplained symptoms (MUS)*, *bodily distress syndromes (BDS)*, or *somatic symptom disorder (SSD)*. These names describe persistent physical symptoms that do not have an explanation. In relation to the manifestations associated with the migratory phenomenon, the *Syndrome of Extreme Migratory Mourning* was described in 1994, also called *Ulysses Syndrome* (8) in situations in which the chronic stress associated with the migration is extreme. Health professionals respond to somatic symptoms in a polarized way: they trivialize the symptoms due to the lack of objective evidence to explain them, due to ignorance of the syndrome or lack of cultural sensitivity, or they do not adequately diagnose the condition, classifying it as a depressive or psychotic illness, instead of a reactive picture of stress, thus elevating symptoms to the field of psychiatric pathology and adding a new stressor to the person (8), (9).

Consultations for recurrent symptoms of this type are especially relevant in primary care (10), since the deterioration in the quality of life of people who suffer from them leads to 60% more use of primary care services after hours (in primary Out-of-Hours care) than people who do not have these diagnoses (11). A recent systematic review argues that immigrants with somatization disorders perceive a greater need for the use of health services and greater difficulties in their daily lives than those who do not have them. However, there are differences in the prevalence of somatization disorders between different groups of immigrants, depending on culture, exposure to stress, explanatory models of the disease, coping and other individual variables (12).

The IOM (International Organization for Migration), the WHO and the Government of Spain organized a World Consultation on the Health of Migrants in 2010 (6). Four priorities were defined in it: monitoring the health of migrants, monitoring their equitable access to health services, developing health policies and programmes which are sensitive to the migrant population, and strengthening coordination and alliances between countries. Furthermore, it should be taken into account that migrants are not a homogeneous group and that increasing knowledge about their mental health, symptoms of psychological distress and psychosomatic manifestations, while attending separately to the different communities of origin would serve as a basis to better guide the healthcare services.

The research hypotheses were:

Psychosocial risk factors, psychological distress and somatizations show differences depending on the cultural community to which one belongs.

There are differences in the perception of psychosocial risk, psychological distress and somatizations which are related to gender.

Risk factors for psychological distress will consist of female gender, older age, less time residing in the host country, less academic education and psychosocial risk before and after migration.

Risk factors for somatization will consist of female gender, older age, less time residing in the host country, less academic education and psychosocial risk before and after migration.

Methods

Aim

The objective of the study was to assess psychosocial risk, psychological distress and somatizations, as well as their main risk factors in four immigrant communities residing in Catalonia (Spain): Maghrebis, sub-Saharan, South Americans and Hindustanis.

Data and design

A cross-sectional study was designed, whose protocol has been previously published, with questionnaires administered by trained personnel with the collaboration of cultural mediators when necessary (13).

The study population were immigrants who came to visit the Primary Care Centres of an urban area of Barcelona (Spain) and who were offered the opportunity, while waiting to be attended by their doctor or nurse, to participate consecutively between November 2011 and April 2013. The inclusion criteria were that they had lived in Europe for a period of between six months and ten years, that they were from countries in North Africa, sub-Saharan Africa, Central or South America or Asia and that they agreed to participate voluntarily. An exclusion criterion was presenting some type of communicative difficulty, which despite the presence of a cultural mediator, did not allow for obtaining clear and understandable information.

Instruments

A form was prepared that collected sociodemographic data related to gender, marital status, educational level and length of residence in Spain. Psychosocial risk factors were collected through the 'Demographic Psychosocial Inventory' (DPSI) (14), an 85-item inventory comprised of two scales, one referring to pre-migration status and the other to post-migration status, and with three general indexes called global index, family structure and health problems, and five dimensions that act as risk factors that promote psychological distress in immigrants. All the scales score from 0 to 1, where the highest scores represent states of poorer health or more psychosocial risk factors depending on the scale in question. The values that suggest clinical risk correspond to 0.44 for men and 0.48 for women.

Psychological distress was recorded using the 'Brief Symptom Inventory' (BSI) (15). In this investigation, the version adapted to Spanish (16) was used, consisting of 46 Likert-type response items from 0 to 4, according to the manifestation of symptoms in the previous 30 days, where a higher score indicates poorer mental health. The version used has six scales to encompass the primary dimensions of psycho-pathological symptoms: depression, phobic anxiety, paranoid ideation, obsession-compulsion, somatization and hostility/aggressiveness, as well as an overall score that would be a measure of psychological distress.

To detect somatic symptoms, the 'Somatic Symptom Inventory' (SSI) was used(17), based on the DSM-III criteria for somatization disorders. It consists of 26 Likert-type response items from 0 to 4, with higher scores representing greater intensity of symptoms. According to Tamayo et al.(18), scores ≤ 52 represent minimal somatic discomfort and > 52 moderate to severe discomfort.

Ethical approval

The confidentiality of the data obtained was absolute and scrupulous using no data which might identify the participants of the study. The data were filed in databases with totally anonymous information technology support and were wholly analysed by persons other than the interviewers. The interviews were carried out in an isolated room with a relaxed atmosphere favouring the free and sincere narrative of each interviewee after first having obtained written informed consent. If necessary, cultural mediators/facilitators were required to obtain such consent. The protocol of investigation has been reviewed and approved by the Ethical and Clinical Investigation Committee of the Primary Care Research Institute IDIAP Jordi Gol (Barcelona, Spain) on 21st July 2010.

Data analysis

A univariate analysis was performed calculating absolute frequency and percentage in the categorical variables and mean and standard deviation in the quantitative ones. A bivariate analysis was performed using the ANOVA test and the T test on the contrasts between variables. The predictive variables of psychosomatic symptomatology were evaluated with a multiple hierarchical regression, introducing the sociodemographic variables in the first step and the pre-migration and post-migration scales and the DPSI global indices in the second step using the PIN(0.05) and POUT(0.10) enter method. The multicollinearity of the data was analysed using the [Variance Inflation Factor \(VIF\)](#) and tolerance (19). VIF values less than 10 in regressions and tolerance values greater than 0.10 indicate that the explanatory variables do not show multicollinearity in the chosen model. The level of significance used was $p \leq 0.05$.

The analyses were performed with the SPSS statistical package for Windows, version 22.0.

Results

A total of 602 people participated, 50% women, belonging to four communities of different origins, 20% of Andean origin (Bolivia, Ecuador, Colombia), 35% of Hindustani origin (Pakistan, Afghanistan, Bangladesh), 21% of North African origin (Morocco, Tunisia, Algeria) and 24% of sub-Saharan origin (Senegal, Gambia, Mali). Table 1 shows an exhaustive description of the sociodemographic characteristics of the participants according to their community of origin.

[Table 1 near here]

Regarding psychosocial risk factors, the global DPSI index was 0.35, showing differences between the four communities studied with people of sub-Saharan origin perceiving greater psychosocial risk according to the a posteriori contrasts carried out (see Table 2).

Regarding the symptoms of psychological distress, Table 2 shows a description of the different scales of the BSI, as well as the overall score differentiated by communities, highlighting that sub-Saharan immigrants have lower levels of psychological distress on the somatization, hostility, paranoid ideation and anxiety scales. On the depression scale the lowest values were obtained by the Andean community.

Psychosomatic symptoms evaluated with a single SSI score show differences between communities, with values ranging from 36 to 47, below the cut-off point indicating moderate somatic discomfort. Sub-Saharan immigrants score less than the other communities on somatization.

[Table 2 near here]

When comparing the pre-migration scales and the post-migration scales according to gender, differences between the two genders were observed with women having the worst results. However, the overall psychosocial risk score does not vary.

Regarding the symptoms of psychological distress and psychosomatic symptomatology, women showed higher scores on all scales (see Table 3).

[Table 3 near here]

The regression with the psychological distress outcome variable measured with the global BSI score provides a model that explains 15.9% of the variance of psychological distress with the variables of sex, age, marital status, post-migration scales, the global DPSI index and the number of years since migration with negative β (see Table 4).

[Table 4 near here]

A second hierarchical regression was performed with a psychosomatic symptomatology outcome variable measured with the SSI. The chosen model presents a corrected R^2 of 0.288 with significant changes with respect to the first model resulting from introducing only the sociodemographic variables that explained a 17.7% variance in psychosomatic symptomatology as predictor

variables. Predictor variables are sex, age, pre-migration scales, post-migration scales and number of years since migration with negative β . The global DPSI index is no longer a predictor of somatization (see Table 4).

Discussion

The participants in this research come mainly from nine countries (Bolivia, Ecuador, Colombia, Morocco, Tunisia, Algeria, Senegal, Gambia, Mali) grouped into four different communities: Andean, sub-Saharan, Maghrebi and Hindustani. The cultural differences between these groups are important, although the most distinctive fact is the language: only the Andean community has the same language (Spanish) as the host country.

The first hypothesis that was posed was that the fact of belonging to different cultural communities conditioned the perception of psychosocial risk, psychological distress and somatic symptoms. The results obtained would allow us to accept this hypothesis since there are significant differences in the perception of psychosocial risk, as well as in all the scales of psychological distress (BSI) and in the somatization scale (SSI). In addition, the a posteriori contrasts allow us to affirm a clearly differentiated behaviour of sub-Saharan immigrants compared with the other communities since they perceive a greater psychosocial risk, although they exhibit fewer symptoms of psychological and psychosomatic distress.

The second hypothesis that was posed was the existence of differences in the perception of psychosocial risk, psychological distress and somatizations due to gender. The results partially support this hypothesis since in all the evaluation scales, women scored significantly higher than men, showing greater pre-migration and post-migration psychosocial risk, worse psychological health and more somatoform disorders. These results are in the same line as the findings of other investigations carried out with the immigrant population in which the female gender presents greater severity of somatic symptoms (Aragona et al., 2012; (10); Morawa et al., 2017). Female gender is also shown as a significant variable in anxiety, mood disorders, eating disorders and somatization, not only in immigrant communities but also in the indigenous population(10). However, the score on the global psychosocial risk scale does not show differences between men and women. This was an unexpected result as the pre-migration and post-migration scales are part of the global scale. This difference is determined to be because the global scale adds five single items under the construct "conflict reaction", which act to moderate psychosocial risk and as a result turn out to be more adaptive coping strategies than those used by men, thereby equalizing the global perception of psychosocial risk.

The third hypothesis supposed that the risk factors for greater psychological distress are female gender, older age, less time residing in the host country, less academic education and psychosocial risk before and after migration. The results partially support the hypothesis since education is not significant. The literature shows contradictory results in relation to the effect of educational level of people who emigrate. Having a university degree was a predictor of worse general health among refugee adults in Australia (22) while it was a protective factor in refugees, where having higher education predicted fewer medical conditions (23) and less psychological distress(24). The pre-migration conditions that in some studies had appeared as a strong predictor of depression and other psychological disorders were not supported in this hypothesis(25). The study by Jamil et al. (2015)(23) highlights that the pre-migration situation behaved differently in refugees and immigrants, with refugees being the main risk factor for having worse self-perceived health, whereas it did not influence the self-perceived health of the immigrant population. Regarding pre-migration conditions, Li and Anderson(26) demonstrated that the influence of the pre-migration situation on psychological symptoms was mediated by perceived discrimination, since immigrants with traumatic experiences perceive the world as a dangerous place. In other contexts, these negative beliefs about the lack of benevolence in the world lead to increased levels of vigilance, which in turn increases stress(27).

The last hypothesis proposed as somatization risk factors the female gender, older age, less time residing in the destination country, lesser academic education and the psychosocial risk before and after migration. This hypothesis is partially supported since academic education is not significant. Evidence shows that academic education behaves ambivalently, the lower the formal education, the higher the prevalence of somatization (28), or with greater levels of education than high school, greater somatization in Russian and Hispanic immigrants (29). In relation to the effect of marital status, which does not appear as a predictor in the somatization of immigrants, the literature also shows contradictory results, (30) found a higher risk of somatization in married people, while (31) found less somatization in married people.

The global DPSI scale is not significant for the somatization outcome variable, taking into account that the other dimensions of the perception of psychosocial risk variable, if they are significant, would be indicative that coping strategies would influence the impact that psychosocial risk may have in the onset of somatization disorders. In line with this argument, in 2008, Sachs et al. explored coping strategies and stress in Tibetan refugees in India who had been exposed to traumatic situations, reporting that coping strategies acted as mediators between lived traumatic experiences and decreased somatic symptoms.

Strengths and limitations of the study

This study is the first approach to the study of somatization disorders and psychological distress of immigrants of different origins carried out in Spain. The main strength of the research is found in the composition of the sample, not only due to its high number but also due to being composed of people from 12 countries that contributed, in the year in which the research was carried out, 42% of immigrants to Catalonia(32).

There are some limitations in the study, the main one being the language barrier for some of the participants, which was resolved with the presence of cultural mediators, but which could influence some of the results. Furthermore, due to the death of the main researcher, there were changes to the research team and consequently the writing of the article took longer than desirable, although a lack of evidence in this area reduces the impact of this limitation.

Conclusions

Despite the limitations, this research contributes to the limited literature on somatization disorders and psychological distress in different immigrant communities.

Sub-Saharan immigrants perceive a greater psychosocial risk than other cultural groups, although they show lower levels of psychological distress and lower levels of somatization.

Being a woman and being older are risk factors for presenting greater psychological distress and more psychosomatic symptoms. In contrast, the time that has passed since migration and having a stable partner are protective factors.

The fact that perception of psychosocial risk is a predictor of psychological distress, but is not a predictor of somatic symptoms, could indicate that the coping strategies that are included in the global DPSI scale may have an influence on reducing the development of somatization disorders. Working on the coping strategies of people who have recently migrated (newcomers) might improve the presentation of psychosomatic manifestations.

List Of Abbreviations

DPSI: Demographic Psychosocial Inventory

BSI: Brief Symptom Inventory

SSI: Somatic Symptom Inventory

FSS: Functional Somatic Symptoms

MUS: Medically Unexplained Symptoms

BDS: Bodily Distress Syndromes

SSD: Somatic Symptom Disorder

IOM: International Organization for Migration

WHO: World Health Organization

DPSI: Demographic Psychosocial Inventory

BSI: Brief Symptom Inventory

SSI: Somatic Symptom Inventory

VIF: Variance Inflation Factor

Declarations

Ethics approval and consent to participate

The study was approved by the Ethical and Clinical Investigation Committee of the Primary Care Research Institute IDIAP Jordi Gol (Barcelona, Spain). There was an informed consent attached to the questionnaire that the participants signed.

Consent for publication

Not applicable.

Availability of data and materials

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Funding

This project has received a grant ‘Gonçal Calvo’, from the Acadèmia de Ciències Mèdiques i de la Salut de Catalunya i Balears (Mataró, Spain) 2010.

Honourable mention of the 6th 2010 Award on Primary Health Research of the Barcelona Health Region.

Authors' contributions

Contributed to the original idea: JMMD, MIFC, MFC, FRR, JSB, PTM

Designed de questionnaire: JMMD, MIFC

Conducted Research: JMMD, MIFC, MFC, FRR, JS, PTM

Conducted the initial analysis: JMMD, MIFC, RGS

Wrote the first draft of the manuscript: RGS

Made critical comments during the process of writing: All authors

Approved the final version of the manuscript: All authors

Acknowledgements

The authors would like to thank all of the study participants for their contributions to this research.

References

1. International Organization for Migration. Glossary on Migration. [Internet]. 2019. [cited 2020 May 16]. Available from: <https://www.iom.int/glossary-migration-2019>
2. IOM Online Bookstore. World Migration Report 2020 [Internet]. 2020. Available from: <https://publications.iom.int/books/world-migration-report-2020>
3. INE. España en cifras 2019 [Internet]. Madrid; 2019. Available from: http://www.ine.es/prodyser/espa_cifras
4. Gushulak B, Weekers J, MacPherson D. Migrants in a globalized world — health threats, risks and challenges: an evidence-based framework. *Emerg Health Threats J*. 2008;2.
5. Grabovschi C, Loignon C, Fortin M. Mapping the concept of vulnerability related to health care disparities: a scoping review [Internet]. 2013. Available from: <http://www.biomedcentral.com/1472-6963/13/94>
6. World Health Organization IO for M y G de E. Health of migrants: the way forward - report of a global consultation. Madrid, Spain. 2010. 119 p.
7. Close C, Kouvonen A, Bosqui T, Patel K, O'Reilly D, Donnelly M. The mental health and wellbeing of first generation migrants: A systematic-narrative review of reviews. *Global Health* [Internet]. 2016;12(1):1–13. Available from: <http://dx.doi.org/10.1186/s12992-016-0187-3>
8. Achotegui Loizate J, López A, Morales Suárez-Varela MM, Espeso D, Achotegui A. Social and demographic study on immigrants with the Ulysses syndrome: Study on 1.110 immigrants attended at SAPPIR in Barcelona. *Norte Salud Ment* [Internet]. 2015;13(52):70–8. Available from: <https://dialnet.unirioja.es/servlet/articulo?codigo=5191741&info=resumen&idioma=SPA>
9. Greco M. Pragmatics of explanation: Creative accountability in the care of 'medically unexplained symptoms'. *Sociol Rev*. 2017;65(2_suppl):110–29.
10. Salinero-Fort MA, Jiménez-García R, de Burgos-Lunar C, Chico-Moraleja RM, Gómez-Campelo P. Common mental disorders in primary health care: differences between Latin American-born and Spanish-born residents in Madrid, Spain. *Soc Psychiatry Psychiatr Epidemiol*. 2015 Mar 1;50(3):429–43.
11. Leutgeb R, Berger S, Szecsenyi J, Laux G. Patients with somatoform disorders: More frequent attendance and higher utilization in primary out-of-hours care? *PLoS One*. 2018 Aug 1;13(8).
12. Lanzara R, Scipioni M, Conti C. A clinical-psychological perspective on somatization among immigrants: A systematic review. *Front Psychol*. 2019;9(JAN).
13. Torán-Monserrat P, Cebrià-Andreu J, Arnau-Figueras J, Segura-Bernal J, Ibars-Verdaguer A, Massons-Cirera J, et al. Level of distress, somatisation and beliefs on health-disease in newly arrived immigrant patients attended in primary care centres in Catalonia and definition of professional competences for their most effective management: PROMISE Project. *BMC Fam Pract* [Internet]. 2013 May 4 [cited 2018 Nov 28];14:54. Available from: <https://doi.org/10.1186/1471-2296-14-54>
14. Ritsner M, Rabinowitz J, Slyuzberg M. The Demographic Psychosocial Inventory: A New Instrument to Measure Risk Factors for Adjustment Problems Among Immigrants. *Refuge*. 1995;14(9):8–16.
15. Derogatis Leonard R. Inventario Breve de 18 Sintomas (BSI-18) [Internet]. Madrid; 2013. Available from: <https://www.cop.es/uploads/PDF/2015/BSI-Pearson.pdf>
16. Ruipérez MÁ, Ibáñez MI, Lorente E, Moro M, Ortet G. Psychometric Properties of the Spanish Version of the BSI. *Eur J Psychol Assess* [Internet]. 2001 Sep [cited 2018 Nov 28];17(3):241–50. Available from: <https://econtent.hogrefe.com/doi/10.1027//1015-5759.17.3.241>
17. Escobar JI, Rubio-Stipec M, Canino G, Karno M. Somatic symptom index (SSI): a new and abridged somatization construct. Prevalence and epidemiological correlates in two large community samples. *J Nerv Ment Dis* [Internet]. 1989 Mar;177(3):140–6. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/2918297>
18. Tamayo JM, Román K, Fumero JJ, Rivas M. The level of recognition of physical symptoms in patients with a major depression episode in the outpatient psychiatric practice in Puerto Rico: An observational study. *BMC Psychiatry* [Internet]. 2005 Dec 20;5(1):28. Available from: <https://doi.org/10.1186/1471-244X-5-28>
19. Belsley DA, Kuh E, Welsch RE. Regression diagnostics: identifying influential data and sources of collinearity. John Wiley, editor. New York, USA: Wiley; 2004. 292 p.

20. Aragona M, Rovetta E, Pucci D, Spoto J, Villa AM. Somatization in a primary care service for immigrants. *Ethn Heal*. 2012 Oct 1;17(5):477–91.
21. Morawa E, Dragano N, Jöckel KH, Moebus S, Brand T, Erim Y. Somatization among persons with Turkish origin: Results of the pretest of the German National Cohort Study. *J Psychosom Res*. 2017 May 1;96:1–9.
22. Dowling A, Enticott J, Kunin M, Russell G. The association of migration experiences on the self-rated health status among adult humanitarian refugees to Australia: An analysis of a longitudinal cohort study. *Int J Equity Health*. 2019 Aug 22;18(1).
23. Jamil H, Barkho E, Broadbridge CL, Ventimiglia M, Arnetz JE, Lami F, et al. Self-rated Health and Medical Conditions in Refugees and Immigrants from the Same Country of Origin. *Iraqi J Med Sci [Internet]*. 2015;13(2):108–19. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26644795><http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC4669974>
24. Myhrvold T, Småstuen MC. The mental healthcare needs of undocumented migrants: an exploratory analysis of psychological distress and living conditions among undocumented migrants in Norway. *J Clin Nurs*. 2017;26(5–6):825–39.
25. Fortuna LR, Porche M V, Alegria M. Political violence, psychosocial trauma, and the context of mental health services use among immigrant Latinos in the United States. *Ethn Heal*. 2008;13(5):435–63.
26. Li M, Anderson JG. Pre-migration Trauma Exposure and Psychological Distress for Asian American Immigrants: Linking the Pre- and Post-migration Contexts. *J Immigr Minor Heal*. 2016;18(4):728–39.
27. Lilly MM, Valdez CE, Graham-Bermann SA. The mediating effect of world assumptions on the relationship between trauma exposure and depression. *J Interpers Violence*. 2011;26(12):2499–516.
28. Mak WWS, Zane NWS. The phenomenon of somatization among community Chinese Americans. *Soc Psychiatry Psychiatr Epidemiol*. 2004;39(12):967–74.
29. Shiroma PR, Alarcon RD. Time for healing: somatization among chronically mentally immigrants. *J Cult Divers*. 2011;18(1):3–7.
30. Aragona M, Monteduro MD, Colosimo F, Maisano B, Geraci S. Effect of Gender And Marital Status on Somatization Symptoms of Immigrants From Various Ethnic Groups Attending a Primary Care Service. *Ger J Psychiatry · [Internet]*. 2008; Available from: <http://www.gjpsy.uni-goettingen.xn--de-1ea>
31. Ritsner M, Ponizovsky A, Kurs R, Modai I. Somatization in an Immigrant Population in Israel: A Community Survey of Prevalence, Risk Factors, and Help-Seeking Behavior [Internet]. Vol. 157, *Am J Psychiatry*. 2000 [cited 2018 Nov 28]. Available from: <https://ajp.psychiatryonline.org/doi/pdf/10.1176/appi.ajp.157.3.385>
32. Idescat. Població estrangera per països. Catalunya [Internet]. 2019 [cited 2020 May 16]. Available from: <https://www.idescat.cat/poblacioestrangera/?b=12>

Tables

	Andean n=122	Hindustani n=212	Maghreb n=125	Sub- Saharan n=143	Total n=602
Age	39.8 (11.4)	36.2 (10.0)	33.1 (10.6)	31.8 (7.6)	35.2 (10.3)
Men	58 (47.5%)	120 (56.6%)	68 (54.4%)	55 (38.5%)	301 (50.0%)
Religion					
Atheists	6 (4.9%)	0 (0%)	0 (0%)	3 (2.1%)	9 (1.5%)
Muslims	0 (0%)	190 (89.6%)	125 (100%)	125 (88.0%)	440 (73.2%)
Christians	116 (95.1%)	0 (0%)	0 (0%)	14 (9.9%)	130 (21.6%)
Eastern	0 (0.0%)	22 (10.4%)	0 (0%)	0 (0%)	22 (3.7%)
Level of studies					
Without studies	1 (0.9%)	28 (13.2%)	24 (19.2%)	41 (28.7%)	94 (15.6%)
Primary	15 (12.3%)	123 (58.0%)	28 (22.4%)	45 (31.5%)	211 (35.0%)
Secondary	29 (23.8%)	44 (20.8%)	26 (20.8%)	32 (22.4%)	131 (21.8%)
High school	38 (31.1%)	13 (6.1%)	19 (15.2%)	6 (4.2%)	76 (12.6%)
Job training	17 (13.9%)	0 (0.0%)	16 (12.8%)	9 (6.3%)	42 (7.0%)
University	22 (18.0%)	4 (1.9%)	12 (9.6%)	10 (7.0%)	48 (8.0%)
Civil status					
Single	34 (27.9%)	58 (27.4%)	52 (41.6%)	45 (31.5%)	189 (31.4%)
With partner	22 (18%)	0 (0%)	3 (2.4%)	0 (0%)	25 (4.2%)
Married	46 (37.7%)	144 (67.9%)	55 (44%)	90 (62.9%)	335 (55.6%)
Separate	9 (7.4%)	4 (1.9%)	1 (0.8%)	3 (2.1%)	17 (2.8%)
Divorced	10 (8.2%)	5 (2.4%)	12 (9.6%)	5 (3.5%)	32 (5.3%)
Widow/er	1 (0.8%)	1 (0.5%)	2 (1.6%)	0 (0%)	4 (0.7%)
Employment situation					
Jobless	45 (36.9%)	93 (43.9%)	53 (42.4%)	76 (53.1%)	267 (44.4%)
Street trading	0 (0.0%)	0 (0.0%)	0 (0.0%)	15 (10.5%)	15 (2.5%)
Student	4 (3.3%)	0 (0.0%)	13 (10.4%)	0 (0.0%)	17 (2.8%)
Housewife	7 (5.7%)	45 (21.2%)	14 (11.2%)	9 (6.3%)	75 (12.5%)
Active worker	63 (51.6%)	72 (34.0%)	44 (35.2%)	43 (30.1%)	222 (36.9%)
Pensioner	2 (1.6%)	2 (0.9%)	1 (0.8%)	0 (0.0%)	5 (0.8%)
Others	1 (0.8%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	1 (0.2%)
Spanish knowledge					
Speak and write correctly	117 (97.5%)	17 (8.1%)	54 (43.5%)	41 (28.9%)	229 (38.4%)
Speaks and writes with difficulty	2 (1.7%)	73 (34.6%)	28 (22.6%)	43 (30.3%)	146 (24.5%)
Speak but not write	1 (0.8%)	54 (25.6%)	34 (27.4%)	43 (30.3%)	132 (22.1%)
Not speak or write it	0 (0%)	67 (31.8%)	8 (6.5%)	15 (10.6%)	90 (15.1%)
Years since migration	9.4 (3.3)	5.2 (3.0)	6.6 (3.0)	5.8 (2.6)	6.5 (3.4)
Number of children					
None	31 (25.4%)	86 (40.6%)	65 (52.0%)	51 (35.7%)	233 (38.7%)
1-2	68 (55.7%)	54 (25.5%)	29 (23.2%)	48 (33.6%)	199 (33.1%)
3-4	18 (14.8%)	41 (19.3%)	23 (18.4%)	35 (24.5%)	117 (19.4%)
5 or more	5 (4.1%)	31 (14.6%)	8 (6.4%)	9 (6.3%)	53 (8.8%)

Table 1. Sociodemographic characteristics of the participants by community.

	M/item (SD/item)					F	Sig.
	Andean	Hindustani	Maghreb	Sub-Saharans	Total		
DPSI*							
Premigration scales	0.10(0.19)	0.16 (0.23)	0.13 (0.22)	0.15 (0.23)	0.14 (0.22)	2.30	0.077
Postmigration scales	0.43 (0.35)	0.34 (0.33)	0.35 (0.36)	0.32 (0.33)	0.36 (0.35)	2.73	0.043
Index Global DPSI	.36 (0.09)	0.33 (0.09)	0.34 (0.07)	0.38 (0.07)	0.35 (0.09)	11.38	<.001
BSI**							
Depression	0.78 (74)	0.89 (.95)	1.09(.76)	1.09 (.73)	0.95	4.79	0.003
Phobic anxiety	0.30 (35)	0.43 (.71)	0.35 (.53)	0.17(.24)	0.32	6.69	<.001
Paranoid Ideation	0.72 (63)	0.88 (.97)	0.87(.69)	0.52 (.42)	0.76	7.60	<.001
Obsession-compulsion	0.65(53)	0.82 (.80)	0.93(.68)	0.67(.39)	0.77	5.32	.001
Somatization	0.72 (69)	0.75 (1)	0.72 (.69)	0.36 (.35)	0.64	8.91	<.001
Hostility/aggressivity	0.39 (58)	0.36 (.69)	0.23(.45)	0.13(.25)	0.29	7.13	<.001
General Severity	0.62 (50)	0.72 (.80)	0.75(.54)	0.53(.29)	0.66	4.05	0.007
Index SSI***	1.68 (.63)	1.82 (1.04)	1.63(.52)	1.39 (.30)	1.65 (.75)	9.99	<.001

Note: M/item: Mean divided by the number of items; SD/item: Standard deviation divided by the number of items

*Score from 0 to 1

** Score from 0 to 4

*** Score from 0 to 4

Table 2. Questionnaires' description by community. N=602

	M/item (SD/item)		F	Sig.
	Female	Male		
DPSI*				
Premigration scales	0.20 (0.25)	0.07(0.18)	54.62	<.001
Postmigration scales	0.45(0.33)	0.26(0.34)	50.32	<.001
Index Global DPSI	0.35(0.1)	0.36(0.08)	1.49	0.223
BSI**				
Depression	1.15(.90)	0.76(.7)	34.90	<.001
Phobic anxiety	0.43(.62)	0.22(.41)	22.76	<.001
Paranoid Ideation	0.87(.82)	0.66(.68)	11.62	0.001
Obsession-compulsion	0.90(.72)	0.64(.55)	24.71	<.001
Somatization	0.86(.9)	0.43(.57)	49.02	<.001
Hostility/aggressivity	0.38(.63)	0.20(.44)	16.70	<.001
General Severity Index	0.80 (0.68)	0.52(0.48)	34.01	<.001
SSI***	1.89(0.88)	1.42(0.48)	64.24	<.001

Note: M/item. mean divided by the number of items; SD/item. standard deviation divided by the number of items

*Score from 0 to 1

** Score from 0 to 4

*** Score from 0 to 4

Table 3. Questionnaires' description by gender.

<i>Outcome variable BSI</i>					<i>Outcome variable SSI</i>				
	<i>B</i>	<i>p</i>	Collinearity statistics			<i>B</i>	<i>p</i>	Collinearity statistics	
			Tolerance	FIV				Tolerance	FIV
Gender (woman)	0.135	0.002	0.744	1.344	Gender (woman)	0.165	<.001	0.745	1.343
Age	0.103	0.041	0.565	1.770	Age	0.170	<.001	0.565	1.770
Estado civil	0.111	0.017	0.668	1.497	Years after migration	-0.103	.010	0.768	1.302
Years after migration	-0.086	0.047	0.768	1.303	Premigration scales	0.245	<.001	0.719	1.391
Postmigration scales	0.151	0.001	0.689	1.452	Postmigration scales	0.173	<.001	0.689	1.452
Index Global DPSI	0.126	0.004	0.749	1.336					
R²	0.159				R²	0.288			

Table 4. Hierarchical lineal regressions (Enter).