

# Hypertension prevalence, associated factors, treatment, and control in rural Cameroon: a cross-sectional study

**Larissa Pone Simo**

Faculty of Health Sciences, the University of Bamenda, Cameroon

**Valirie Ndip Agbor** (✉ [nvagbor@gmail.com](mailto:nvagbor@gmail.com))

Nuffield Department of Population Health, University of Oxford, United Kingdom

<https://orcid.org/0000-0002-6708-6852>

**Jean Jacques Noubiap**

Centre for Heart Rhythm Disorders, University of Adelaide and Royal Adelaide Hospital, Australia

**Orlin Pagnol Nana**

Faculty of Health Sciences, the University of Bamenda, Cameroon

**Pride Swiri-Muya Nkosu**

Faculty of Health Sciences, the University of Bamenda, Cameroon

**Arnold Forlemu Asaah Anouboweh**

Faculty of Health Sciences, the University of Bamenda, Cameroon

**Jude Nfor Ndi**

Faculty of Health Sciences, the University of Bamenda, Cameroon

**Jacques Nguend Mbock**

Faculty of Health Sciences, the University of Bamenda, Cameroon

**Noel Fils Bakari**

Faculty of Health Sciences, the University of Bamenda, Cameroon

**Harold Giovanni Guifo Tambou**

Faculty of Health Sciences, the University of Bamenda, Cameroon

---

## Research Article

**Keywords:** Hypertension, public health, epidemiology, associated factors, determinants, awareness, treatment, control, Cameroon.

**Posted Date:** May 29th, 2020

**DOI:** <https://doi.org/10.21203/rs.3.rs-31780/v1>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

**Version of Record:** A version of this preprint was published at BMJ Open on September 1st, 2020. See the published version at <https://doi.org/10.1136/bmjopen-2020-040981>.

# Abstract

**Introduction:** Sub-Saharan Africa is experiencing a surge in the burden of hypertension, and rural communities seem to be increasingly affected by the epidemic.

**Objectives:** We aimed to determine the prevalence of hypertension, its associated factors, as well as its awareness, treatment, and control rates in rural communities of the Baham Health District (BHD), Cameroon.

**Design:** A community-based cross-sectional study.

**Setting:** Participants from five health areas in the BHD were recruited from August to October 2018.

**Participants:** Consenting participants aged 18 years or older were included.

**Results:** We included 526 participants in this study. The median age of the participants was 53.0 (IQR = 35 – 65) years and 67.1% were female. The prevalence of hypertension was 40.9% (95% confidence interval [CI] = 36.7 – 45.1) were hypertensive with no gender disparity. The overall age-standardised prevalence of hypertension was 23.9% (95% CI = 20.3 – 27.5). Five-year increase in age (adjusted odd's ratio [AOR] = 1.34; 95% CI = 1.23 – 1.44), family history of hypertension (AOR = 2.22; 95% CI = 1.37 – 3.60), and obesity were associated with higher odds of hypertension (AOR = 2.57; 95% CI = 1.40 – 4.69).

The rates of awareness, treatment, and control of hypertension were 37.2% (95% CI = 31.0 – 43.9), 20.9% (95% CI = 16.0 – 26.9), and 22.2% (95% CI = 15.2 – 46.5), respectively.

**Conclusion:** The high hypertension prevalence in this rural community is associated with contrastingly low awareness, treatment, and control rates. Age, family history of hypertension, and obesity are the major drivers of hypertension in this community. Veracious policies are needed to improve awareness, prevention, diagnosis, treatment, and control of hypertension in this rural community.

## 1. Background

Hypertension is a major modifiable risk factor for cardiovascular diseases globally [1, 2] which is associated with increased cost on health systems, high morbidity and premature mortality. Globally, it is estimated that one billion adults live with hypertension; a figure which is projected to hit 1.5 billion by the year in 2025 [2, 3]. Furthermore, hypertension-related complications are responsible for over 50% of the 17.4 million deaths due to cardiovascular diseases every year worldwide [2]. At least 45% of deaths due to heart disease and 51% of deaths due to stroke are related to hypertension [2].

Cardiovascular diseases are the second commonest cause of premature disability and death in sub-Saharan Africa (SSA) [4–6]. A greater bulk of the burden of heart disease, kidney failure, stroke and premature death in this region is accounted for by hypertension [7–10]. The prevalence of hypertension in SSA is estimated at about 30% with disproportionately low awareness, treatment and control rates [7].

About 29.7% of the general Cameroonian population are affected by hypertension [8] while a significant number of cases of hypertension still remain undiagnosed and untreated, and even those on treatment hardly achieve a controlled blood pressure [9–12].

With the increasing prevalence of hypertension globally, including the rural areas, a continuous evaluation of the burden of hypertension in these areas is needed to plan prevention and control strategies. A limited number of community-based studies have assessed epidemiology of hypertension in rural areas in Cameroon with significant disparities across regions. Cognizant to this, we sought to assess the prevalence and factors associated with hypertension among adults in selected health areas in a rural health district in the West Region of Cameroon. In addition, we evaluated the rates of awareness, treatment, and control of hypertension in the same population.

## 2. Methods

### 2.1 Study design, setting and duration

This was a community-based, cross-sectional study conducted between August and October 2018 in the Baham Health District (BHD), as part of the University of Bamenda Medical Students Association (UBaMSA) annual community health campaign. The study was conducted in 5 of the 9 health areas of the BHD including the Hiala Cheffou, Bapa, Baham and Ngougoua health areas. Baham is a rural community located in the West Region of Cameroon. The Baham Health District had an estimated population of 51500 in 2001 [13] whose major activity is farming. It is made up of 9 health areas with a district hospital.

### 2.2 Study population and sampling

The five health areas in which our study was conducted were selected based on ease of accessibility. Consenting participants aged 18 years and older were consecutively recruited into the study. Participants with documented or reported diagnosis of chronic kidney disease, those who had taken cardiostimulants such as alcohol, “kola nut” (a caffeine containing fruit of the Kola tree; a genus of trees that are native to the tropical rainforests of Africa) and caffeine at least 30 minutes prior to the study, and pregnant women were excluded from the study.

The sample size was estimated using the following formula:

$$n = \frac{Z^2 P(1-P)}{d^2}$$

where n is the sample size (number of adult participants), P is the expected prevalence of HTN in an adult population (P= 0.378) [12], and d is the precision (if 5%, d= 0.05). Z statistics (Z): For the level of confidence of 95%, which is conventional, Z value is 1.96 for a 95% confidence interval (CI). A minimum of 361 adult participants was required for this study.

## 2.3 Study procedure and data collection

One month prior to the UBaMSA health campaign, members of the community were informed by mass communication (through the local radio stations), and interpersonal communication on the dates retained for activities of the campaign. The data collection process was guided by the World Health Organization (WHO) STEPwise approach to Surveillance (STEPS). Data was collected by trained medical students and medical doctors. Information on the participants' socioeconomic status (like age, sex, and education), lifestyle (fruits and vegetable consumption, smoking status, and physical activity), and medical history (family history of hypertension). In cases where participants did not understand English or French, a translator was used.

Blood pressure was measured using a reference protocol in which participants were seated, and measurements were taken after at least 10 minutes of rest. This was done using the auscultatory method with a calibrated sphygmomanometer placed at least 0.5cm above the elbow joint, covering at least 80% of the arm, and a stethoscope was used to detect the sounds. The analysis was done for the average of two measures performed at least five minutes apart.

Height was measured using a calibrated stadiometer to the nearest 0.1 cm. Weights were measured to the nearest 0.5kg with the use of a scale, and the participants mounted the scale only with light clothes on. Abdominal circumference was measured to the nearest 0.5cm with a measuring tape placed all around the bare abdomen at the level of the umbilicus.

### Definitions

1. Respondents were considered as hypertensive if they had an average SBP of 140 mm Hg or higher, or DBP of 90 mmHg or greater, or reported current use of anti-hypertensive medication [14].
2. Hypertension awareness rate was defined as the proportion of individuals who responded by a "yes" to have either been diagnosed with hypertension by a healthcare professional and/or "yes" to taking medication for hypertension.
3. The rate of hypertension treatment included the proportion of participants who were diagnosed with hypertension and reported being on treatment for hypertension.
4. Hypertension control was defined as the proportion of individuals on either pharmacotherapy or lifestyle modification or both for hypertension and who had an average SBP < 140 mmHg and DBP < 90 mmHg.
5. Occupational level was classified into "low" (no technical know-how or expert training required, e.g. manual workers), "medium" (requiring a degree of technical know-how but no expert training, like salesmen, and bike and taxi drivers) and "high" (major professionals requiring advanced training like teachers, health personnel, and accountants).
6. We defined an ex-smoker as someone who has smoked at least 100 cigarettes in their lifetime but had stopped smoking at least 28 days before the interview. A smoker was defined as someone who has smoked at least 100 cigarettes in their lifetime and are still regular smokers at the time of the

interview. Those who had never smoked or smoked less 100 cigarettes in their lifetime were classified as non-smokers.

7. Alcohol units per week = (number of bottles of beer consumed per week) x 5% x 650ml/1000 [15]. The routine beer bottle in Cameroon occupies 650ml of beer with a concentration of alcohol of 5%.
8. The intensity of physical activity was classified as “moderate” (e.g. brisk walking, moderate farm work like weeding and harvesting, haunting, lifting masses < 20kg, housework and domestic chores, and general building tasks such as roofing and painting) and “vigorous” (running, briskly ascending and descending hills, intense farm work such as manual tilling of the soil, digging ditches and carrying masses  $\geq$  20kg) [16]. Sedentary lifestyles at work and home were classified as “No physical activity”.
9. The body mass index (BMI) was calculated as the ratio of the weight in kilograms and the square of the height in metres. BMI based body habitus (in Kg/m<sup>2</sup>) was classified as underweight (BMI < 18.5), normal weight (BMI = 18.5-24.9), overweight (BMI = 25.0-29.9), and obese (BMI  $\geq$ 30) [17].
10. Abdominal obesity was defined as an abdominal circumference  $\geq$  102 cm in men or  $\geq$  88 cm in women [18].

## 2.4 Data analysis

Data was analysed with Stata v.16 (StataCorp 2019, College Station, TX: StataCorp LLC). Qualitative variables were reported using counts and percentages. Quantitative variables were summarised as means and medians with their corresponding standard deviation (SD) and interquartile range (IQR), respectively. We computed direct age-standardised prevalence of hypertension using the 2011 population structure of Cameroon [19]. For univariate analyses, the Pearson  $\chi^2$  test was used to compare categorical variables while the Wilcoxon rank sum test was used to compare medians across independent groups. Independent factors associated with hypertension were determined using unconditional maximum likelihood multivariable logistic regression models. Variables with a p-value  $\leq$  0.1 on univariate analysis qualified for inclusion in the multivariable model. We sequentially adjusted for socioeconomic factors (like age, gender, occupation, and education), lifestyle factors (smoking status, alcohol consumption, fruit consumption, and physical activity) and clinical characteristics (family history of hypertension and BMI). The maximum likelihood ratio test was used to evaluate model fit and select variables for the final multivariable model. Gender, alcohol consumption, and smoking status were retained in the final model as they have been reported as factors associated with hypertension in literature. Body mass index was retained in the final model over abdominal obesity to facilitate comparison of our findings with previously published studies and to prevent multicollinearity. Ordinal variables were assessed for linear trend using the  $\chi^2$  test for linear trend. The  $\chi^2$  test for heterogeneity was used to evaluate departures from linearity. Measures of association are reported as odds ratio (OR) with corresponding 95% confidence interval (CI). Missing data was handled using simple mean, median and mode imputation where appropriate. Two-tailed p-values below 0.05 were considered statistically significant.

## 3. Results

In total, 526 participants with a median age of 53.0 (IQR = 35 – 65) years were included in this study. The ages of the participants ranged from 18 – 99 years. About 67% of the participants were females and 76.6% were married, Table 1. A little over half of the participants were Catholic Christians and about three-quarters of them had at least a primary education. The average BMI was 27.2 (SD=5.2), and about 44% of the participants had android obesity.

### 3.1 Prevalence of hypertension

Of the 526 participants, 215 were classified as hypertensive giving an overall crude prevalence of 40.9% (95% CI = 36.7 – 45.1). Figure 1 shows the gender-specific prevalence of hypertension (with their 95% CI) across different age groups. There was a linear increase in the prevalence of hypertension among older participants, with no gender disparity. The overall age-standardised prevalence of hypertension was 23.9% (95% CI = 20.3 – 27.5).

### 3.2 Factors associated with hypertension

On univariate analysis, participants with hypertension were significantly older (median age in years = 64.0 versus 42.0 years) and consumed fruits less regularly (median daily fruit consumption per week = 2.0 versus 4.0) compared to those without hypertension, Table 2. There was strong evidence against the null hypothesis of no difference in marital status, occupation, level of education, family history of hypertension, and intensity of physical activity between participants with and without hypertension. There was weak evidence against the null hypothesis of no difference in exposure to wood smoke between participants with and without hypertension. There was a moderate positive correlation between BMI and abdominal circumference ( $r = 0.60$ ,  $p < 0.001$ ).

Figure 2 displays the final multivariable logistic regression model (without abdominal obesity). There was strong evidence of a 34% increase in the odds of hypertension for every five-year increase in age (adjusted odd's ratio [AOR] = 1.34; 95% CI = 1.23 – 1.44;  $p < 0.001$ ). Family history of hypertension associated with 2.22 times higher odds of hypertension (AOR = 2.22; 95% CI = 1.37 – 3.60;  $p < 0.001$ ). Obesity was associated with 2.57 times higher odds of hypertension (AOR = 2.57; 95% CI = 1.40 – 4.69;  $P_{\text{trend}} < 0.001$ ).

### 3.3 Awareness, treatment, and control of hypertension

Table 3 depicts the percentage of hypertension awareness, treatment, and control among our study participants. Of the 215 participants diagnosed with hypertension, 37.2% (95% CI = 31.0 – 43.9) were aware of their hypertensive status, while 20.9% (95% CI = 16.0 – 26.9) reported being on any form of treatment for hypertension. Of the 45 participants who were on treatment for hypertension, 22.2% (95% CI = 12.2 – 37.0) had a controlled BP.

## 4. Discussion

We report a prevalence of hypertension of 40.9% (age-standardised prevalence = 23.9%) with associated with low awareness, treatment, and control rates in the BHD. Increasing age, family history of hypertension and obesity were drivers of hypertension in this population.

#### 4.1 Prevalence of hypertension and associated factors

The crude prevalence of hypertension in our study is higher than the crude prevalence of 33.9% and 31.1% in rural areas of the Far North Region [12] and South West Region of Cameroon [20], respectively. This higher prevalence of hypertension in our study can be attributed to the older age of our study population compared with those of previous studies. Indeed, the median age of our study participants was 53 years compared to a mean age of 39 years reported by Lemogoum *et al* [12]. In addition, over 65% of the participants in the study by Arrey *et al* were between 20-29 years old [20]. Older age is a strong determinant of hypertension. We noted a strong positive linear trend between older age and hypertension as has been observed in other studies in Cameroon [8, 12, 20, 21] and elsewhere [26]. Age-standardisation with Cameroon's population of 2011 permitted comparison of our results with those of Lemogoum *et al* [12]. The age-standardised prevalence of hypertension in our study was half of the crude prevalence, indicating the contribution of age in the overall crude prevalence in this study. The age-standardised prevalence in our study was lower than that reported by Lemogoum *et al* [12].

Differences in ethnicity, socioeconomic, and lifestyle factors could account for the variation in the prevalence of hypertension in our study compared to previous studies [8, 12, 20, 21]. Our study recruited participants from the Bamileke ethnic group. In a recent publication by Defo *et al*, participants recruited from this ethnic group had the highest prevalence of hypertension in Cameroon [22]. Our study suggests that genetic predisposition to hypertension is a significant determinant of hypertension in our study population. History of hypertension was associated with over two-times increase in the odds of hypertension. In addition, difference in BMI could explain, in part, the variation in the prevalence of hypertension in this study compared to other studies reporting on the prevalence of hypertension in rural Cameroon [12, 20, 21]. Over 60% of our study participants had a BMI over the normal range, and there was a strong positive linear relationship between hypertension and BMI. Adiposity is a strong risk factor for hypertension and an important driver of the prevalence of hypertension in our study [23].

In Cameroon, there has been a roll up in the prevalence of hypertension in the general population from 16.4% in 1998 [24] to 29.7% in 2015 [8], with recent projections estimating an increase by 40% in 2025 and 95% in 2035 [25]. The prevalence of hypertension in this study approximates the overall prevalence of 47.5% reported in four urban areas in Cameroon a few years back [11] and a rural community in South Africa [26]. Such high prevalence of hypertension, especially among the elderly, in this rural communities warrants further investigation to ascertain the burden of the disease and plan effective prevention and management strategies.

We found no independent association between hypertension sex, education, marital status, and physical activity reported in previous studies [22] nor with the male sex as reported by Lemogoum *et al* in Far North Cameroon [16].

## 4.2 Awareness, treatment, and control of hypertension

We report a low awareness rate and even lower treatment and control rates. This is in line with findings reported in rural areas of the Far North and South West Regions of Cameroon [12, 20], and a systematic review by Ataklte *et al* [23]. In the Mafia Island of Tanzania, a low control rate of 20.5% was recorded despite the very high treatment rate [27]. This is in contrast to the relatively higher control rates (44.7%) reported in a rural community in Ghana [28].

The low awareness, treatment and control rates reported in this study could be explained by inadequate patient information on the disease, its risk factors, and consequences in the long run. Low awareness is a major barrier to effective management which will result in the development of hypertension-related complications. A paucity of health care professionals at the primary health care level and the absence of a hypertension clinic at the district hospital in this setting for limits awareness, treatment, and control of hypertension in this population. Implementing policies to improve population education on hypertension, and the importance of regular follow up by a trained nurse or physician to prevent long-term complications would go a long way to improve awareness, treatment and control of hypertension in our setting [29]. Other cost-effectiveness measurements like the use home BP monitoring could go a long way to improve adherence and control of hypertension in Cameroon [30].

## 5 Conclusion

About two in five participants in our study population had hypertension. The high prevalence of hypertension in this study was contrasted by low awareness, treatment, and control rates. In a bit to curb the burden of hypertension in Cameroon, national policies need to adopt measures to address obesity and its risk factors. Measures to improve awareness of hypertension like regulation community education, diagnosis, treatments, and control could go a long way to reduce the burden of hypertension in this rural community.

## Abbreviations

AOR: Adjusted odds ratio; BHD: Baham Health District; BMI: Body mass index; BP: Blood pressure; SD: Standard deviation; OR: odds ratio; UBaMSA: University of Bamenda Medical Students Association.

## Declarations

### Acknowledgements

The authors are grateful for the invaluable contributions of all those who participated in the University of Bamenda Medical Student (UBaMSA) health campaign during the September 2018 edition.

### Availability of data and materials

All data presented in the manuscript results are in possession of the corresponding author and will be made available on reasonable demand.

### **Authors' contributions**

Study design and conception: LPS, VNA and JJN; data collection: LPS, OPN, PNS, AAF, JNM, NFB, HTG; data analysis and interpretation: VNA and LPS; manuscript drafting: LPS and VNA; Revision of the manuscript: LPS, OPN, PNS, AAF, JNN, JNM, NFB, HTG. Critical revision: VNA, JJN and DM. All authors read and approved the final manuscript.

### **Ethics approval and consent to participate**

Ethical approval was obtained from the Faculty of Health Sciences Institutional Review Board, the University of Bamenda and all participants provided written informed consent prior to recruitment to the study.

**Funding:** Dr Agbor and Dr Noubiap are supported by postgraduate scholarship from the University of Oxford and the University of Adelaide, respectively.

### **Competing interests**

The authors declare that they have no competing interests.

## **References**

1. Kearney PM, Whelton M, Reynolds K, Muntner P, Whelton PK, He J. Global burden of hypertension: analysis of worldwide data. *The Lancet*. 2005;365:217–23. doi:10.1016/S0140-6736(05)17741-1.
2. Who A. Global brief on hypertension. World Health Organization. 2013.
3. Lim SS, Vos T, Flaxman AD, Danaei G, Shibuya K, Adair-Rohani H, et al. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *The Lancet*. 2012;380:2224–60. doi:10.1016/S0140-6736(12)61766-8.
4. Naghavi M, Abajobir AA, Abbafati C, Abbas KM, Abd-Allah F, Abera SF, et al. Global, regional, and national age-sex specific mortality for 264 causes of death, 1980–2016: a systematic analysis for the Global Burden of Disease Study 2016. *The Lancet*. 2017;390:1151–210.
5. Noubiap JJ, Agbor VN, Ndoadougou AL, Nkeck JR, Kamguia A, Nyaga UF, et al. Epidemiology of pericardial diseases in Africa: a systematic scoping review. *Heart*. 2019;105:180–8.
6. Danwang C, Temgoua MN, Agbor VN, Tankeu AT, Noubiap JJ. Epidemiology of venous thromboembolism in Africa: a systematic review. *Journal of Thrombosis and Haemostasis*. 2017;15:1770–81.

7. Adeloye D, Basquill C. Estimating the Prevalence and Awareness Rates of Hypertension in Africa: A Systematic Analysis. *PLOS ONE*. 2014;9:e104300. doi:10.1371/journal.pone.0104300.
8. Kingue S, Ngoe CN, Menanga AP, Jingi AM, Noubiap JJN, Fesuh B, et al. Prevalence and Risk Factors of Hypertension in Urban Areas of Cameroon: A Nationwide Population-Based Cross-Sectional Study. *The Journal of Clinical Hypertension*. 2015;17:819–24. doi:10.1111/jch.12604.
9. Tianyi F, Agbor V, Njamnshi A. Prevalence, awareness, treatment, and control of hypertension in Cameroonians aged 50 years and older: A community-based study. *Health Science Reports*. 2018.
10. Adidja NM, Agbor VN, Aminde JA, Ngwasiri CA, Ngu KB, Aminde LN. Non-adherence to antihypertensive pharmacotherapy in Buea, Cameroon: a cross-sectional community-based study. *BMC Cardiovascular Disorders*. 2018;18:150. doi:10.1186/s12872-018-0888-z.
11. Dzudie A, Kengne AP, Muna WFT, Ba H, Menanga A, Kouam CK, et al. Prevalence, awareness, treatment and control of hypertension in a self-selected sub-Saharan African urban population: a cross-sectional study. *BMJ Open*. 2012;2:e001217. doi:10.1136/bmjopen-2012-001217.
12. Lemogoum D, Van de Borne P, Lele CEB, Damasceno A, Ngatchou W, Amta P, et al. Prevalence, awareness, treatment, and control of hypertension among rural and urban dwellers of the Far North Region of Cameroon. 2018. doi:info:doi/10.1097/HJH.0000000000001513.
13. Baham, Cameroon. Wikipedia. 2018. [https://en.wikipedia.org/w/index.php?title=Baham,\\_Cameroon&oldid=866664358](https://en.wikipedia.org/w/index.php?title=Baham,_Cameroon&oldid=866664358). Accessed 14 Dec 2018.
14. Hernandez-Vila E. A Review of the JNC 8 Blood Pressure Guideline. *Texas Heart Institute Journal*. 2015;42:226–8. doi:10.14503/THIJ-15-5067.
15. Alcohol units - NHS. <https://www.nhs.uk/live-well/alcohol-support/calculating-alcohol-units/>. Accessed 14 Dec 2018.
16. WHO | What is Moderate-intensity and Vigorous-intensity Physical Activity? WHO. [https://www.who.int/dietphysicalactivity/physical\\_activity\\_intensity/en/](https://www.who.int/dietphysicalactivity/physical_activity_intensity/en/). Accessed 19 Dec 2018.
17. Body mass index - BMI. 2018. <http://www.euro.who.int/en/health-topics/disease-prevention/nutrition/a-healthy-lifestyle/body-mass-index-bmi>. Accessed 19 Dec 2018.
18. Okosun I, Choi S, Dent M, Jobin T, Dever G. Abdominal obesity defined as a larger than expected waist girth is associated with racial/ethnic differences in risk of hypertension. *Journal of Human Hypertension*. 2001;15:307–12.
19. Cameroon's National Institute of Statistics. *Statistics year book 2010*. Chapter 4: Characteristics of the population. 2011. <http://www.statistics-cameroon.org/downloads/annuaire2010/chap4.pdf>.
20. Arrey WT, Dimala CA, Atashili J, Mbuagbaw J, Monekosso GL. Hypertension, an Emerging Problem in Rural Cameroon: Prevalence, Risk Factors, and Control. *International Journal of Hypertension*. 2016. doi:10.1155/2016/5639146.
21. Princewel F, Cumber SN, Kimbi JA, Nkfusai CN, Keka EI, Viyoff VZ, et al. Prevalence and risk factors associated with hypertension among adults in a rural setting: the case of Ombe, Cameroon. *Pan Afr Med J*. 2019;34. doi:10.11604/pamj.2019.34.147.17518.

22. Kuate Defo B, Mbanya JC, Kingue S, Tardif J-C, Choukem SP, Perreault S, et al. Blood pressure and burden of hypertension in Cameroon, a microcosm of Africa. *Journal of Hypertension*. 2019;37:2190–9.
23. Aronow WS. Association of obesity with hypertension. *Ann Transl Med*. 2017;5. doi:10.21037/atm.2017.06.69.
24. Mbanya JCN, Minkoulou EM, Salah JN, Balkau B. The prevalence of hypertension in rural and urban Cameroon. *Int J Epidemiol*. 1998;27:181–5. doi:10.1093/ije/27.2.181.
25. Jingi AM, Dzudie A, Noubiap JJ, Menanga AP, Aminde LN, Fesuh B, et al. PT075 Trend in the Prevalence, Awareness, and Control of Hypertension in Cameroon: A Systematic Review and Projections for 2025 and 2035. *Global Heart*. 2016;2 Supplement:e138–9. doi:10.1016/j.gheart.2016.03.482.
26. Ntuli ST, Maimela E, Alberts M, Choma S, Dikotope S. Prevalence and associated risk factors of hypertension amongst adults in a rural community of Limpopo Province, South Africa. *African Journal of Primary Health Care & Family Medicine*. 2015;7. doi:10.4102/phcfm.v7i1.847.
27. Muhamedhussein MS, Nagri ZI, Manji KP. Prevalence, Risk Factors, Awareness, and Treatment and Control of Hypertension in Mafia Island, Tanzania. *International Journal of Hypertension*. 2016. doi:10.1155/2016/1281384.
28. Kweku M, Takramah W, Parby P, Axame W, Takase M, Adjuik M, et al. PREVALENCE AND CONTROL OF HYPERTENSION AMONG RURAL ADULTS IN THE LIKPE SUB-MUNICIPALITY OF THE HOHOE MUNICIPALITY OF GHANA. 2017;4:16.
29. Dzudie A, Kingue S, Dzudie A, Sliwa K, Mayosi B, Dzudie A, et al. Roadmap to achieve 25% hypertension control in Africa by 2025. *Cardiovasc J Afr*. 2017;28:261–72.
30. Ndip Agbor V, Temgoua MN, Noubiap JJN. Scaling up the use of home blood pressure monitoring in the management of hypertension in low-income countries: A step towards curbing the burden of hypertension. *The Journal of Clinical Hypertension*. 2017;19:786–9.

## Tables

**Table 1. Sociodemographic characteristics of the study population**

Participants' characteristics	Total (N=526)
Age (in years) <sup>§</sup>	53 (35 - 65)
Age group (in years)	
18-39	157 (29.8%)
40-59	170 (32.3%)
60 and over	199 (37.8%)
Gender (Female)	353 (67.1%)
Marital Status	
Married	403 (76.6%)
Single	123 (23.4%)
Occupation	
High	22 (4.2%)
Medium	113 (21.5%)
Low	391 (74.3%)
Religion	
Baptist	13 (2.5%)
Catholic	275 (52.3%)
Muslim	14 (2.7%)
Others	119 (22.6%)
None	19 (3.6%)
Presbyterian	86 (16.3%)
Level of education	
No formal education	131 (24.9%)
Primary	155 (29.5%)
Secondary	183 (34.8%)
Tertiary	57 (10.8%)
Family history of hypertension (Yes)	141 (26.8%)
Smoking status	
Non-smoker	435 (82.7%)
Ex-smoker	49 (9.3%)
Current smoker	42 (8.0%)
Alcohol units per week	
Non-drinker	163 (31.0%)
(0.01, 6.49]	205 (39.0%)
(6.49, 117]	158 (30.0%)
Body mass index (in kg/m <sup>2</sup> )*	27.2 (5.2)
Abdominal obesity (Yes)	233 (44.3%)
Systolic blood pressure (in mmHg)*	134.1 (24.5)
Diastolic blood pressure (in mmHg)*	83.0 (14.5)

<sup>§</sup> Summarised as median and interquartile range; \* Data summarised as mean (standard deviation); N = frequency

Table 2: Factors associated with hypertension in the Baham Health District on univariate analysis

Participants' characteristics	No hypertension	Hypertension	Total	p-value
	N=311	N=215	N=526	
Age group (in years)	42.0 (28.0-58.0)	64.0 (53.0-73.0)	53.0 (35.0-65.0)	<0.001 <sup>a,*</sup>
Gender				0.370 <sup>b</sup>
Female	204 (65.6%)	149 (69.3%)	353 (67.1%)	
Male	107 (34.4%)	66 (30.7%)	173 (32.9%)	
Marital Status				<0.001 <sup>b,*</sup>
Married	212 (68.2%)	191 (88.8%)	403 (76.6%)	
Single	99 (31.8%)	24 (11.2%)	123 (23.4%)	
Occupation				0.017 <sup>b,*</sup>
Low/unemployed	218 (70.1%)	173 (80.5%)	391 (74.3%)	
Medium	80 (25.7%)	33 (15.3%)	113 (21.5%)	
High	13 (4.2%)	9 (4.2%)	22 (4.2%)	
Religion				0.180 <sup>b</sup>
Baptist	11 (3.5%)	2 (0.9%)	13 (2.5%)	
Catholic	154 (49.5%)	121 (56.3%)	275 (52.3%)	
Muslim	11 (3.5%)	3 (1.4%)	14 (2.7%)	
Others	74 (23.8%)	45 (20.9%)	119 (22.6%)	
None	12 (3.9%)	7 (3.3%)	19 (3.6%)	
Presbyterian	49 (15.8%)	37 (17.2%)	86 (16.3%)	
Level of education				<0.001 <sup>b,*</sup>
None	55 (17.7%)	76 (35.3%)	131 (24.9%)	
Primary	81 (26.0%)	74 (34.4%)	155 (29.5%)	
Secondary	130 (41.8%)	53 (24.7%)	183 (34.8%)	
Tertiary	45 (14.5%)	12 (5.6%)	57 (10.8%)	
Family history of hypertension				0.002 <sup>b,*</sup>
No	243 (78.1%)	142 (66.0%)	385 (73.2%)	
Yes	68 (21.9%)	73 (34.0%)	141 (26.8%)	
Smoking status				0.760 <sup>b</sup>
Non-smoker	256 (82.3%)	179 (83.3%)	435 (82.7%)	
Ex-smoker	28 (9.0%)	21 (9.8%)	49 (9.3%)	
Current smoker	27 (8.7%)	15 (7.0%)	42 (8.0%)	
Exposure to wood smoke				0.048 <sup>b,*</sup>
≥ 4 days/week	200 (64.3%)	160 (74.4%)	360 (68.4%)	
< 4 days/week	75 (24.1%)	36 (16.7%)	111 (21.1%)	
Never	36 (11.6%)	19 (8.8%)	55 (10.5%)	
Alcohol units per week				0.510 <sup>b</sup>
Non-drinker	92 (29.6%)	71 (33.0%)	163 (31.0%)	
(0.01, 6.49]	120 (38.6%)	85 (39.5%)	205 (39.0%)	
(6.49, 117]	99 (31.8%)	59 (27.4%)	158 (30.0%)	

Daily consumption of vegetable per week	1.0 (1.0-2.0)	1.0 (1.0-2.0)	1.0 (1.0-2.0)	0.230 <sup>a</sup>
Daily consumption of fruit per week	4.0 (2.0-6.0)	2.0 (2.0-4.0)	2.0 (2.0-6.0)	<0.001 <sup>a,*</sup>
Intensity of daily physical activity				<0.001 <sup>b,*</sup>
Low	134 (43.1%)	134 (62.3%)	268 (51.0%)	
Moderate	123 (39.5%)	72 (33.5%)	195 (37.1%)	
Vigorous	54 (17.4%)	9 (4.2%)	63 (12.0%)	
Body mass index categories				0.053 <sup>b</sup>
Normal	119 (38.3%)	70 (32.6%)	189 (35.9%)	
Overweight	134 (43.1%)	86 (40.0%)	220 (41.8%)	
Obese	58 (18.6%)	59 (27.4%)	117 (22.2%)	
Abdominal obesity				<0.001 <sup>b,*</sup>
Yes	115 (37.0%)	118 (54.9%)	233 (44.3%)	
No	196 (63.0%)	97 (45.1%)	293 (55.7%)	


<sup>a</sup>P-value from Wilcoxon rank sum test, <sup>b</sup> P-value from Chi-square test.

Table 3. Awareness, treatment and control of hypertension, Baham Health District, 2018

	Frequency	Percentage (95% CI)
Hypertension awareness (N=215)	80	37.2 (31.0 - 43.9)
Hypertension treatment (N=215)	45	20.9 (16.0 - 26.9)
Treated and controlled (N=45)	10	22.2 (12.2 - 37.0)


N = Frequency, CI = confidence interval

## Figures



**Figure 1**

Prevalence (and 95% confidence interval [CI]) of hypertension stratified by age and gender. The red circle and black square represent the point estimate of the prevalence for females and males, respectively. The spikes represent the limits of the 95% CI.



**Figure 2**

Factors associated with Hypertension in the Baham Health District multivariable logistic regression analysis. Measures of associations are displayed as odds ratio (OR), black squares, with the 95% confidence interval (CI), horizontal spikes. Significant p-values are shown in bold. The red dashed line refers to the null value of 1.0. a= p-value for trend.