

Effect of Small and Large Energy Surpluses on Strength, Muscle, and Skinfold Thickness

Eric R Helms (✉ eric.helms@aut.ac.nz)

Auckland University of Technology <https://orcid.org/0000-0002-4849-9336>

Alyssa-Joy Spence

Auckland University of Technology

Colby Sousa

Auckland University of Technology

James Krieger

Weightology, LLC

Steve Taylor

unaffiliated

Dustin J Oranchuk

University of Colorado

Brad P Dieter

Macros Inc

Casey M Watkins

Seattle University

Research Article

Keywords: resistance training, body composition, hypertrophy, energy surplus

Posted Date: August 2nd, 2023

DOI: <https://doi.org/10.21203/rs.3.rs-3184470/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Sports Medicine-Open on November 2nd, 2023. See the published version at <https://doi.org/10.1186/s40798-023-00651-y>.

Abstract

Background

A goal of many who perform resistance training (RT) is to increase muscle mass and strength. An energy surplus is commonly advised to support such gains; however, if too large, it could cause unwanted fat gain without augmenting adaptation. Thus, we randomized 21 trained lifters into maintenance energy (MAIN), 5% (MOD), and 15% (HIGH) energy surplus groups while performing RT 3d/wk for eight weeks to determine if skinfold thicknesses (ST), squat and bench one-repetition maximum (1-RM), or biceps brachii or quadriceps muscle thicknesses (MT) were influenced by group assignment. COVID-19 complications reduced our ability to recruit and retain participants, leaving 17 completers. Thus, in addition to Bayesian ANCOVA group comparisons, we regressed outcomes against body mass (BM) changes to analyze the relationship between BM gain, ST increases, and increase in 1-RM and MT. For all outcomes we reported Bayes Factors (BF_{10}) indicating the odds ratio of the likelihood of one hypothesis relative to another (e.g., a $BF_{10} = 2$ indicates the hypothesis is twice as likely as another) and coefficients of determination (R^2) for regressions.

Results

ANCOVAs provided no evidence in support of the group model for MT or squat 1-RM. However, there was moderate ($BF_{10} = 9.9$) and strong evidence ($BF_{10} = 14.5$) that MOD increased bench 1-RM more than HIGH and MAIN, respectively. Further, there was moderate evidence ($BF_{10} = 4.2$) MOD increased ST more than MAIN and weak evidence ($BF_{10} = 2.4$) HIGH increased ST more than MAIN. The regression provided strong evidence that BM changes predicted changes in ST ($BF_{10} = 14.3$, $R^2 = 0.49$) and weak evidence that BM changes predicted changes in biceps MT ($BF_{10} = 1.4$, $R^2 = 0.24$), but no evidence any other variable was predicted by BM changes.

Conclusions

While some group-based differences were found, our larger N regression provides the most generalizable evidence. Therefore, we conclude faster rates of BM gain (and by proxy larger surpluses) primarily increase rates of ST gain rather than augmenting 1-RM or MT. However, the biceps, the muscle group which received the greatest stimulus in this study, may have been positively impacted by greater BM gain, albeit slightly. Our findings are limited to the confines of this study, where a group of lifters with mixed training experience performed moderate volumes 3d/wk for eight weeks. Thus, future work is needed to evaluate the relationship between BM gains, increases in ST and RT adaptations in other contexts.

Key Points

1. When assigning intended energy surplus sizes of 5-15%, faster rates of body mass gain primarily serve to increase the rate that fat mass accumulates, rather than increasing rates of hypertrophy or strength gain.
2. It is possible, however, that faster rates of body mass gain could enhance hypertrophy to some degree if a sufficient training stimulus is provided.
3. While further work is needed, if a sufficient training stimulus is provided, the rate of body mass gain that will best support hypertrophy is likely individual. Rates may be influenced by prior training experience, hereditary factors which influence one's potential maximum rate of muscle gain, and other variables (sleep, stress, etc.).

1 Background

Resistance Training (RT) is widely used in strength and conditioning, fitness, and clinical settings to elicit muscular adaptations, including skeletal muscle hypertrophy. In strength sports, such as powerlifting, lean mass is strongly related to competitive success [1]. In elderly populations, those with higher amounts of lean mass live longer [2] and score higher on functional tests [3] and in competitive bodybuilding, a more muscular physique contributes largely to one's score [4]. Therefore, athletes, the general population, personal trainers, strength and conditioning coaches, and health professionals can all benefit from a better understanding of how to increase muscular hypertrophy.

While RT provides the initial stimulus that can result in changes in muscle mass, other factors can influence the magnitude of these changes. Specifically, research over the last several decades has demonstrated that nutritional factors can have substantial effects on muscle mass accrual [5–8]. Multiple aspects of nutrition can influence muscular hypertrophy. For example, consuming a sufficiently high daily protein intake can augment gains in lean mass in response to RT [9–11]. Further, trainees may be able to perform greater volumes of RT, which are associated with greater hypertrophy [12] when consuming sufficient carbohydrate before training [13]. More broadly, an energy surplus could theoretically aid hypertrophy considering it is an energetically expensive process, that RT itself increases energy expenditure, and since energy expenditure can increase in response to overfeeding [8]. Conversely, sufficiently large energy deficits are associated with blunted hypertrophy [14]. Further, insufficient intakes of carbohydrate [15] or fat [16] - which are more likely when consuming insufficient energy - may result in hormonal environments potentially less conducive for hypertrophy.

While these findings may prompt a recommendation to practice overfeeding to optimize hypertrophy, doing so may not be required as increases in muscle mass have been observed while fat loss occurs [17]. Therefore, while not a prerequisite, an energy surplus (or at least the absence of a deficit) may support hypertrophy. What is not yet clear, is whether a larger surplus is more effective than a smaller surplus for supporting hypertrophy, or if simply performing RT while at energy balance is sufficient. Indeed, when different magnitudes of overfeeding are experimentally induced, hypertrophy is inconsistently enhanced, with surpluses that are too large sometimes resulting in increases in fat mass without additional muscle

gain [18, 19]. Thus, while prescribing a large surplus may ensure RT adaptation is supported, too large of a surplus may result in unwanted fat mass gain, potentially prompting a subsequent fat loss phase depending on individual goals. An important next step in applied research is to determine the optimal surplus size which maximizes muscle gain while minimizing fat gain.

Therefore, the purpose of this study was to compare RT-induced hypertrophy and strength gains in three groups of resistance-trained participants after eight weeks of supervised training while consuming three different energy intakes: 1) an estimated 15% surplus (HIGH), 2) an estimated 5% surplus (MOD), and 3) estimated maintenance (MAIN). We hypothesized that both surplus groups would increase muscle size and strength to a greater degree than the MAIN but would make similar gains to one another. Further, we hypothesized that increases in body fat would scale with the size of the energy surplus.

2 Methods

2.1 Participants

Twenty-one well-trained individuals (2 female, 19 male; Table 1) participated in this study. Four male participants were excluded from analysis due to inability to complete the study protocol. Therefore, the final sample size was 17. Reasons for discontinuation included COVID-19 regulations and sickness. One participant experienced pain during squat post testing which prevented completion of the post-test squat 1RM, all other data for this participant was used. Data were included only if participants completed pre- and post-testing, all participants completed > 90% of training sessions. For inclusion, participants were required to have at least one year of RT experience, defined as performing RT sessions at least twice weekly including the bench press and squat exercises each at least once per week with no complete breaks from training longer than two weeks in succession. Further, participants needed a minimum strength level of a 1x or 0.75x bodyweight bench press, and a 1.5x or 1.25x bodyweight squat for men and women, respectively. These criteria, as well as the ability to perform the back squat and bench press required technique, were confirmed during the familiarization session. All participants were injury-free at the time of inclusion. Moreover, all participants were informed of the risks and benefits associated with this study, after which they signed an institutionally approved written consent prior to data collection. The methods and procedures used in this study were approved by the first author's University ethics board (approval number 18/53).

2.2 Procedures

Participants visited the laboratory 25 times over eight weeks, these visits included pre- and post-testing, and each supervised training session. Nutrition intake was monitored throughout. All groups performed a thrice-weekly, full-body, hypertrophy-oriented supervised RT program. Group allocation was blinded for researchers supervising training sessions to mitigate any bias surrounding testing and training. Pre- and post-testing included a calliper skinfold assessment, an ultrasound assessment of upper and lower body muscle thickness (MT), and a one-repetition maximum (1-RM) test for bench press and back squat

exercises. Testing sessions were completed 48–72 hours before the first training session and 48–72 hours following the last training session.

2.3 Pre-intervention Maintenance and Familiarization Phase

Before the intervention, all participants underwent a maintenance and familiarization phase. The goals of this phase were for participants to establish the habit of weighing themselves every morning, tracking their food intake in a food tracking app in real-time (i.e., as they ate throughout the day), and to achieve weight stability.

Before beginning the maintenance and familiarization phase, participants received nutrition guidance from one of two research team members with concentrated experience regularly working with remote clients seeking to alter their body composition. Of the two nutrition research team members, one is a registered dietitian and the other an experienced clinical nutrition researcher with a PhD in exercise physiology. These research team members monitored nutrition intake via an online food tracking software (MyFitnessPal, California, USA) and conducted weekly check-ins. Initially, all participants met via video call with one of the two nutrition research team members. During this call, participants were instructed how to weigh themselves (e.g., first thing in the morning, after going to the bathroom, in a fasted state, and nude), given a detailed video tutorial demonstrating the proper way to track macronutrients, as well as instructions for completing their weekly check-ins. Participants remained in the maintenance and familiarization phase until they achieved weight stability. Weight-stable was defined as having the same weekly average body weight ($\pm 1\%$) for two successive weeks.

2.4 Nutrition Interventions

Once a participant was determined weight-stable, they were randomly assigned to a group. Only the nutrition supervising researcher was aware of group assignment, while the researchers supervising training were blinded to group assignment. Further, participants were asked to keep their group assignment to themselves when interacting with the researchers supervising training. Groups differed by their energy intake, with MAIN assigned an energy intake target predicted to keep their initial weight stable, within $\pm 1\%$, as defined during the maintenance phase. The MOD was assigned an energy intake target predicted to increase body weight by 0.4–0.6% every two weeks. Finally, HIGH was assigned an energy intake target predicted to increase body weight by 1.4–1.6% every two weeks.

All groups were given the same instructions regarding their macronutrient intake. Instructions required participants to consume a minimum of 1.8 grams of protein per kilogram of body weight, a minimum of 20% of energy from fat, and a minimum of 40% of energy from carbohydrate sources. Outside of these constraints, participants could modify their macronutrient distributions to individual preferences. Further, all participants were instructed to consume their food during three to five meals spread relatively evenly throughout the day, while ingesting at least 20 grams of protein within two hours of finishing their training sessions. Participants were instructed to maintain their supplement usage for the duration of the study and not to change supplement strategy, or doses consumed during the maintenance and

familiarization phases. Participants using anabolic steroids or other anabolic performance-enhancing drugs were excluded from the study.

2.5 Weekly Nutrition Check-ins and Adjustments

Once per week, each participant would check in with their nutrition supervising researcher. For each check-in, participants were instructed to verify accuracy for their spreadsheet where they recorded their weight, energy, protein, fat, and carbohydrate intake each day. Additionally, participants were directed to send a video report (or occasionally an email when unable to record a video) in which they would summarize their week and ask any pertinent questions regarding the nutrition intervention. Common questions were typically related to how to track various foods as accurately as possible and requests for advice on how to achieve energy and macronutrient targets.

Upon review, if a participant's body weight aligned with the desired rate of weight gain (or lack thereof) for their assigned group, no nutritional adjustments were made. However, if a participant's body weight did not align with the desired rate of weight gain for their group (i.e., fell below or above the target rate of weight gain, or weight stability) adjustments to the assigned energy intake (and thus macronutrients) were made based on estimated relationships between weight gain, loss, and energy intake target [20]. The nutrition supervising researcher responded to participants via video within 24 hours of receiving weekly information, answering questions, providing guidance, and detailing instructions for any changes to their energy or macronutrient goals.

2.6 Ultrasound collection and Analysis

For pre- and post-testing, each participant's height and body mass were taken upon arrival. Thereafter, participants lay supine on a massage table with their knees and hips fully extended for 10 minutes, to allow for inter- and intra-cellular fluid re-distribution [21]. The length of the lateral aspect of the thigh was measured as the distance from the superior border of the greater trochanter to the inferior border of the lateral condyle of the femur [22, 23]. The anterior aspect of the thigh was measured as the distance between the superior border of the patella and the inferior border of the anterior, superior iliac spine [22, 23]. Thigh lengths were recorded and marked with an indelible pen, 50% distance between the lateral and anterior borders [22, 23]. The vastus lateralis (VL) and lateral vastus intermedius (VI) were collected in one image, and the rectus femoris (RF) and anterior vastus intermedius were collected in one image [22, 23]. The vastus medialis was excluded as it can be further broken down into the obliquus and longus portions, with deep and superficial fiber bundles making consistent collection challenging [24]

In-vivo muscle architecture was determined via 2-dimensional B-mode ultrasonography using an ultrasound transducer and built-in software (45 mm linear array, 10 MHz; GE Healthcare, Vivid S5, Chicago, IL, USA). On each occasion, two images were captured and averaged to provide mean MT. A water-soluble gel was applied to the scanning head of the ultrasound probe to achieve acoustic coupling, with care taken to avoid the deformation of muscle architecture [25]. The transducer was positioned in the longitudinal plane to increase ease and reduce the time required to collect ultrasound images [26] and transducer tilting was carefully avoided [27]. The lateral thigh images were collected before the anterior

thigh. Ultrasound settings (Frequency: 12MHz, Brightness: maximum, gain: 60 dB, dynamic range: 70) were kept consistent across all participants. Due to large differences in MT, scanning depth was individualized for each participant and muscle, whereby settings were recorded and maintained through all collections [22, 23, 25]. Immediately following the lower body ultrasound collection, the participant stood up and the anthropometrist marked 50% of the length of the humerus as determined by the halfway point between the olecranon process and the lateral border of the acromion process. The center of the biceps and triceps brachii were marked with an indelible pen. Biceps and triceps images were collected in the transverse plane as no reliability or variability data exists for longitudinally collected images. All ultrasound settings and practices from the lower body assessment were applied to the biceps and triceps brachii.

Images were analyzed via digitizing software (ImageJ; National Institutes of Health, USA). MT (cm) was defined as the perpendicular distance between the deep and superficial aponeurosis, and the deep aponeurosis. All images were inspected and analyzed by the same experienced sonographer [22, 23].

2.7 Skinfold Assessment

Once all ultrasound imaging protocols were completed, body composition assessments were completed via an ISAK certified level 1 anthropometrist and included the standardized eight-site skinfold profile and associated girth assessments. Skinfold assessments were measured with Harpenden calipers (Baty International, England, UK) to the nearest 0.1 mm wherein measurement calibration was confirmed pre-trial. All anatomical reference points were measured according to standardized ISAK protocols, followed by skinfold assessments for triceps, subscapular, biceps, iliac crest, abdominal, front thigh, and medial calf regions. The sum of all eight sites was used in analysis. A complete profile was performed and repeated to verify measurement accuracy and mitigate any fluctuations in body fat pliability. A third measurement was taken for any variable which resulted in a difference in measurement scores greater than 5% or 1 mm, whichever value was less.

2.8 One Repetition Maximum Testing

Participants performed a standardized dynamic warm-up, followed by a specific warm-up based on individual 1-RM estimations. The specific warm-up included one set of 10 repetitions with the barbell, followed by one set each of 5 repetitions with 50%, 4 repetitions with 60%, 3 repetitions with 70%, 2 repetitions with 80%, and 1 repetition with 90% of estimated 1-RM load. Rest periods were three to five minutes long and given after each warm-up set and 1-RM attempt, starting after 70%. Participant repetition in reserve-based rating of perceived exertion (RPE) score [28] and their average concentric velocity (PowerTool, GymAware, Kinetic Performance Technology, Canberra, Australia) were used to direct 1-RM attempt selection. A 1-RM was recorded if the participant successfully completed a lift at a 10 RPE (maximal effort) with technical proficiency, or successfully completed a lift at a lower RPE but failed the subsequent attempt. The squat was performed to International Powerlifting Federation standards which required participants to reach a depth where the hip crease was below the top of the knee joint [29]. The bench press was performed in a touch-and-go style, where participants were required to maintain five

points of contact, including their head, shoulders, and buttocks in contact with the bench, and both their feet on the ground for the duration of the lift.

2.9 Training Protocol

Training sessions were supervised, in person when possible, and via video chat when necessary due to COVID-19 restrictions. Training sessions were completed thrice weekly on non-consecutive days at the same time of day whenever possible (occasionally participants rescheduled session times) for eight weeks (23 sessions total). Session layout (Table 2), main lift (Table 3) and accessory lift (Table 4) progressions can be seen in Tables 2–4. Warm-up and working sets of squat and bench press were calculated from pre-test 1-RM scores. General main lift progressions maintained three working sets where repetitions per set gradually decreased, while percentage of 1-RM increased throughout most of the program. The first and last week included only two sets as an introductory and tapering stimulus. Additionally, RPE was used to modify RT loads to ensure the percentage of 1-RMs assigned maintained the intended proximity to failure. Accessory exercise loads were determined by the previous week's training, with the goal of reaching volitional failure within a specified repetition range during each set. Rest periods were set at three minutes between working sets of squats and bench press, and two minutes between accessory circuits.

Table 1
Participant Characteristics

	All (n = 17)		5 (n = 5)		10 (n = 6) *n = 5		0 (n = 6)	
	Pre	Post	Pre	Post	Pre	Post	Pre	Post
Age (y)	27.2 ± 3.8		28.6 ± 3.4		28.8 ± 2.6		24.3 ± 3.9	
Height (cm)	173.2 ± 7.5		174.6 ± 6.2		174.0 ± 7.2		171.3 ± 9.5	
Body mass (kg)	77.5 ± 11.7	79.8 ± 12.1	82.6 ± 7.5	85.9 ± 6.9	79.2 ± 7.2	82.5 ± 7.3	71.7 ± 16.3	72.1 ± 16.1
Bench 1RM (kg)	98.4 ± 22.8	106.9 ± 23.9	103.6 ± 11.6	117.0 ± 11.1	100.6 ± 29.3	107.9 ± 31.6	91.8 ± 24.8	97.5 ± 22.9
Relative Bench (1RM/bm)	1.27 ± 0.21	1.34 ± 0.22	1.26 ± 0.11	1.36 ± 0.09	1.26 ± 0.33	1.3 ± 0.35	1.28 ± 0.15	1.36 ± 0.16
Squat 1RM (kg)	140.9 ± 33.1	150.4 ± 33.4	147.3 ± 25.6	158.0 ± 27.1	140.8 ± 40.9	148.6 ± 44.5*	135.7 ± 35.4	145.7 ± 33.0
Relative Squat 1RM (1RM/bm)	1.80 ± 0.33	1.89 ± 0.33	1.78 ± 0.19	1.83 ± 0.22	1.77 ± 0.48	1.8 ± 0.50*	1.89 ± 0.29	2.02 ± 0.22
*See subsection 2.1, one participant was unable to complete squat 1RM post-test.								
1RM = one-repetition maximum; bm = body mass								

Table 2
Training Program

Day 1	Day 2	Day 3
Back Squat	Back Squat	Back Squat
Bench Press	Bench Press	Bench Press
Lat Pulldown	Dumbbell Row	Lat Pulldown
Dumbbell Shoulder Press	Dumbbell Lateral Raise	Dumbbell Shoulder Press
Barbell Curl	Dumbbell Hammer Curl	Barbell Curl

Table 3
Squat and Bench Progression

Week	Day 1	Day 2	Day 3
0 – pre-test	x	x	Pre-testing
1 – intro	2x10x60% (5–7 RPE)	2x8x65% (5–7 RPE)	2x6x70% (5–7 RPE)
2	3x10x65% (6–8 RPE)	3x8x70% (6–8 RPE)	3x6x75% (6–8 RPE)
3*	3x10x67.5% (6–8 RPE)	3x8x72.5% (6–8 RPE)	3x6x77.5% (6–8 RPE)
4	3x9x70% (7–9 RPE)	3x7x75% (7–9 RPE)	3x5x80% (7–9 RPE)
5*	3x9x72.5% (7–9 RPE)	3x7x77.5% (7–9 RPE)	3x5x82.5% (7–9 RPE)
6	3x8x75% (8–10 RPE)	3x6x80% (8–10 RPE)	3x4x85% (8–10 RPE)
7*	3x8x77.5% (8–10 RPE)	3x6x82.5% (8–10 RPE)	3x4x87.5% (8–10 RPE)
8 – taper	2x6x80% (7–9 RPE)	2x4x85% (7–9 RPE)	post testing
RPE = rating of perceived exertion based on repetitions in reserve.			
First set's load dictated by percentage of pre-test one-repetition maximum. Then adjusted on subsequent sets by RPE; by participant if within range, increased or decreased by 2% for every 0.5 RPE below or above RPE range, respectively.			
*First set's load is determined by the percentage listed only if initial set was completed on the same day of prior week within or below RPE range. If repetitions were missed or RPE was above range, last week's load is repeated.			

Table 4
Accessory Lift Progression (all except back squat and bench press)

Week	Day 1	Day 2	Day 3
0 – pre-test	x	x	Pre-testing
1 – intro*	2x10-15	2x8-12	2x8-12
2	3x10-15RM	3x8-12RM	3x8-12RM
3	3x10-15RM	3x8-12RM	3x8-12RM
4	3x10-15RM	3x8-12RM	3x8-12RM
5	3x8-12RM	3x6-10RM	3x6-10RM
6	3x8-12RM	3x6-10RM	3x6-10RM
7	3x8-12RM	3x6-10RM	3x6-10RM
8 – taper#	2x8-12	2x6-10	Post-testing
RM = repetition maximum.			
Weeks 2–7: previous week's training consulted to determine load. Goal to fall within repetition range while reaching failure each set. Load increased if more repetitions than target repetition range are completed.			
*Researcher guided load selection with gradual increase from set 1 to set 2 to assess what load can be used in subsequent weeks.			
#Previous weeks loads reduced one increment for machine and dumbbells or 5kg for barbell lifts.			

2.10 Test-retest reliability and variability

Intrasession variability of mid-region VL, RF, and lateral VI and anterior VI MT (ICC = 0.93–0.98, CV = 2.4–5.7%, TEM = 0.15–0.25) of resistance-trained men was previously determined in our laboratory [22, 23]. Additionally, Jenkins et al. [30] and Radaelli et al. [31] reported similar test-retest statistics for biceps brachii MT (ICC = 0.91–0.99, CV = 4.2%) in untrained men and women, respectively.

2.11 Statistics

The statistician was initially blinded to group assignment. Before unblinding, changes in outcomes were analyzed using a Bayesian ANCOVA, with change as the dependent variable, group as the independent variable, and baseline value as a covariate. All models (baseline value, group, group + baseline value) were compared to the null model to produce Bayes Factors (BF_{10}) which indicate the odds ratio of the likelihood of one hypothesis relative to another where $BF_{10} = 1$, with values higher than 1 favoring an alternative model and values less than 1 favoring the null model. Strength of evidence in favor of either the null model or alternative models were interpreted according to Andraszewicz et al. [32]. After the ANCOVA analysis, the statistician was unblinded and regressed each outcome for the entire final sample of 17 participants against the change in body mass using Bayesian linear regression. Data were analyzed

using JASP 0.16.4 (University of Amsterdam). Data are presented as means \pm 95% credible interval (CI) – the 95% probability that the true (unknown) effect estimate lies within the interval, given the evidence provided by the observed data [33]- unless otherwise specified.

3 Results

3.1 VL_{MT}

There was moderate evidence in favor of pre-training VL_{MT}, but not group, as a predictor of the change in VL_{MT} from pre- to post-assessments ($BF_{10} = 3.6$, Table 5). There was weak evidence in favor of the null model over group ($BF_{10} = 0.35$, Table 5). Changes in VL_{MT} were similar among groups (Fig. 1A) and change in body mass was not favored over the null model as a predictor of change in VL_{MT} ($BF_{10} = 0.45$, $R^2 = 0.01$).

3.2 Lateral VI_{MT}

Similar to VL_{MT}, there was moderate evidence in favor of pre-training MT, but not group, as a predictor of the change in lateral VI_{MT} ($BF_{10} = 3.3$, Table 5). There was weak evidence in favor of the null model over the group model ($BF_{10} = 0.32$, Table 5). Changes in lateral VI_{MT} were similar among groups (Fig. 1B) and change in body mass was not favored over the null model as a predictor of change in lateral VI_{MT} ($BF_{10} = 0.43$, $R^2 = 0.0$).

3.3 Anterior VI_{MT}

There was weak evidence in favor of pre-training MT, but not group, as a predictor of the change in anterior VI_{MT} ($BF_{10} = 1.6$, Table 5). There was weak evidence in favor of the null model over the group model ($BF_{10} = 0.33$, Table 5). Changes in anterior VI_{MT} were similar among groups (Fig. 1C) and change in body mass was not favored over the null model as a predictor of change in anterior VI_{MT} ($BF_{10} = 0.51$, $R^2 = 0.04$).

3.4 RF_{MT}

There was weak evidence in favor of pre-training MT, but not group, as a predictor of the change in anterior VI_{MT} ($BF_{10} = 1.5$, Table 5). There was weak evidence in favor of the null model over the group model ($BF_{10} = 0.77$, Table 5). Changes in RF_{MT} were similar among groups (Fig. 1D) and change in body mass was not favored over the null model as a predictor of change in RF_{MT} ($BF_{10} = 0.63$, $R^2 = 0.08$).

3.5 Triceps MT

Evidence weakly favored the null model over all other models (Table 5). Specifically for group, there was weak evidence in favor of the null model over the group model ($BF_{10} = 0.52$, Table 5). Changes in triceps

MT were similar among groups (Fig. 1E) and change in body mass was not favored over the null model as a predictor of change in triceps MT ($BF_{10} = 0.43$, $R^2 = 0.0$).

3.6 Biceps MT

Evidence weakly favored the null model over all other models (Table 5). Specifically for group, there was weak evidence in favor of the null model over the group model ($BF_{10} = 0.35$, Table 5). Changes in biceps MT were similar among groups (Fig. 1F) and change in body mass was only weakly favored over the null model as a predictor of change in biceps MT ($BF_{10} = 1.4$, $R^2 = 0.24$).

3.7 Squat 1-RM

Evidence weakly to moderately favored the null model over all other models (Table 5). Specifically for group, there was moderate evidence in favor of the null model over the group model ($BF_{10} = 0.31$, Table 5). Changes in squat performance were similar among groups (Fig. 2a) and change in body mass was not favored over the null model as a predictor of change in squat performance ($BF_{10} = 0.62$, $R^2 = 0.08$).

3.8 Bench 1-RM

There was strong evidence in favor of group over all other models ($BF_{10} = 13.1$, Table 5). Post-hoc comparisons indicated moderate evidence ($BF_{10} = 9.9$) for group MOD as having a greater increase compared to group HIGH (Fig. 2b), and strong evidence ($BF_{10} = 14.5$) for group MOD as having a greater increase compared to group MAIN (Fig. 2b). Change in body mass was not favored over the null model as a predictor of change in bench performance ($BF_{10} = 0.97$, $R^2 = 0.16$).

3.9 Sum of Skinfolts

Evidence moderately favored group over all other models ($BF_{10} = 3.0$, Table 5). Post-hoc comparisons indicated moderate evidence ($BF_{10} = 4.2$) for group MOD as having a greater increase as compared to group MAIN (Fig. 3), and weak evidence ($BF_{10} = 2.4$) for group HIGH as having a greater increase as compared to group MAIN (Fig. 3). There was strong evidence that change in body mass was a predictor of change in sum of skinfold thickness over the null model ($BF_{10} = 14.3$, $R^2 = 0.49$).

Table 5

Bayesian statistics with Bayes Factors (BF) with baseline, group, and baseline and group models for outcome variables.

Outcome	Model	BF ₁₀ (compared to null model where BF ₁₀ = 1, ordered from highest BF ₁₀ to lowest for each outcome)
VL _{MT}	Pre-Training	3.6
	Group + Pre-Training	2.2
	Group	0.35
Lateral VI _{MT}	Pre-Training	3.3
	Group + Pre-Training	1.0
	Group	0.32
Anterior VI _{MT}	Pre-Training Thickness	1.6
	Group + Pre-Training	0.54
	Group	0.33
RF MT	Pre-Training	1.5
	Group + Pre-Training	1.4
	Group	0.77
Triceps MT	Group	0.52
	Pre-Training	0.49
	Group + Pre-Training	0.24
Biceps MT	Pre-Training	0.81
	Group + Pre-Training	0.37
	Group	0.35
Squat	Pre-Training 1RM	0.49
	Group	0.31

Outcome	Model	BF ₁₀ (compared to null model where BF ₁₀ = 1, ordered from highest BF ₁₀ to lowest for each outcome)
	Group + Pre-Training 1RM	0.15
Bench	Group	13.1
	Group + Pre-Training 1RM	5.9
	Pre-Training 1RM	0.48
Sum of Skinfolds	Group	3.0
	Group + Pre-Training SS	1.9
	Pre-Training SS	0.65
BF = Bayes Factor; SS = sum of skinfolds; MT = muscle thickness; VL = vastus lateralis; VI = vastus intermedius; RF = rectus femoris.		

4 Discussion

The present investigation is one of the few existing studies to assess the relationship between the magnitude of energy surplus and change in maximal strength, skinfold thicknesses, and upper and lower body MT in a resistance-trained population. Notably, due to the COVID-19 epidemic and subsequent lockdowns, we were unable to recruit the target *N* before study completion. Therefore, in addition to our *a priori* planned group-based comparisons, after unblinding we opted to conduct a *post hoc* regression analysis based on changes in body mass to enhance our statistical power and provide more meaningful conclusions. Our hypotheses linked to our initial group-based statistical comparisons were largely not supported by our findings. Specifically, group-based comparisons for changes in MT weakly favored the null, with similar changes in MT occurring at all sites, in all groups. Similarly, squat 1-RM changes moderately favored the null, with similar increases between groups. However, in partial support of our hypotheses, group-based changes in bench press 1-RM favored the group model with moderate and strong evidence that MOD outperformed HIGH and MAIN, respectively. Finally, the group-based comparisons for changes in the sum of skinfold thicknesses moderately favored the group model, also only partially in line with our hypotheses. Specifically, there was moderate evidence that MOD increased their sum of skinfold thicknesses more than HIGH, no evidence of a difference between MOD and HIGH, and only weak evidence than HIGH increased their sum of skinfold thicknesses more than MAIN. However, our regression analysis based on body mass changes suggests more of our hypotheses were supported than indicated by our group-based comparisons. Notably, despite no evidence that biceps MT was influenced by group assignment, there was weak support for body mass changes as a predictor of changes in biceps MT. In contrast, despite group-based bench press 1-RM comparisons favoring MOD,

changes in body mass were not favored over the null model as a predictor of bench press 1-RM changes. Lastly, there was strong evidence that changes in body mass predicted changes in the sum of skinfold thicknesses. To summarise, in the context of the studied population following the assigned training protocol, despite 1 RM strength and quadriceps and triceps MT being seemingly unaffected by gains in body mass (and thus, surplus size), participants who gained the most body mass clearly gained the most body fat and (although far less clearly) also gained more biceps MT.

Regarding changes in MT, the only site in which all three groups experienced an arguably meaningful mean increase, was the biceps (~ 0.2-0.3cm). At all other sites, for all groups, mean changes mostly clustered around zero. Given the resistance-trained status of the study population, it is possible that the training protocol provided an insufficient stimulus to produce meaningful quadriceps and triceps hypertrophy for the observed eight-week period. Notably, the only exercise which trained the quadriceps was the squat, for a total of nine sets per week, and while the triceps were trained with 15 sets per week, they were only trained via the multi-joint bench press and shoulder press exercises. However, the biceps were trained for a total of 18 sets per week, with half of these sets coming from multi-joint exercises (row, lat pulldown) and half from isolated elbow flexion (barbell curl and hammer curl). Viewed in light of the most recent meta-analyses on the relationship between weekly set volume and hypertrophy (which also counted combined isolated and multi-joint exercises), it is perhaps unsurprising that biceps MT increased most consistently. Schoenfeld et al. [12] reported a significant ($p = 0.002$) relationship with hypertrophy and weekly muscle-specific set volume in a continuous regression, significantly greater hypertrophy favoring 9+ sets when using a two-category comparison with < 9 sets ($p = 0.03$), and a non-significant ($p = 0.074$) graded dose response using a three-category comparison of 1–4, 5–9, and 10+ weekly sets. Further, in a similar analysis of higher volume studies, Baz-Valle et al. [34] reported no significant differences between 12–20 and 20+ sets for both biceps and quadriceps, but significantly greater hypertrophy in the triceps when performing 20+ weekly sets compared to 12–20. Given these findings, one would expect the biceps to experience the most hypertrophy of the measured muscles in the present study based on the volumes performed.

Potentially also relevant to our MT findings, was the programmed proximity to failure. The bench press and squat exercises were prescribed via %1-RM with the intent to be challenging, yet submaximal. When sets were performed too far from failure (or too close) per the participants perceived repetitions in reserve (RIR) [35], load was adjusted to ensure the successful completion of all repetitions in each set as close to the intended proximity to failure as possible (Supplementary file x). However, participants were verbally encouraged to train to a 0 RIR on all sets for all other exercises. Indeed, in the most recent meta-analysis [36] on the relationship between proximity to failure and hypertrophy, while the effect size (ES) was trivial to small (0.19), hypertrophy was significantly ($p = 0.045$) greater in groups that trained to failure. Thus, not only were the biceps trained with the highest volume of all measured muscle groups, but they were also trained more intensely. Intriguingly, given that our regression showed weak evidence of a relationship between gains in body mass and biceps MT, it may be that hypertrophy can only be augmented by an energy surplus when the stimulus for a given muscle is sufficiently potent, which may have only been the case for the biceps in the present study. With that said, this supposition should be couched until future

confirmatory research is published as the evidence for the relationship between body mass gains and increases in biceps MT was weak ($BF_{10} = 1.4$, $R^2 = 0.24$).

All three groups increased squat and bench press 1-RM, and despite the group-based finding that MOD gained more bench press strength than HIGH and MAIN, our regression did not reveal any evidence of a relationship between squat or bench press 1 RM strength gains and increases in body mass. Thus, the training protocol - which was identical between groups - was sufficient to produce maximal strength gains, but an energy surplus (regardless of size) and any subsequent gains in body mass did not augment these gains. Given that, on average, meta-analytic data [14] indicate that even an energy deficit does not significantly impair strength gains ($ES = -0.31$, $p = 0.28$) – despite significantly impairing lean mass gains ($ES = -0.57$, $p = 0.02$) - our findings that an energy surplus doesn't augment strength gains are perhaps to be expected. Further, considering there were similar (negligible) increases in triceps and quadriceps MT among groups - the only measured muscles which contribute to squat and bench press performance – one would also not expect to observe any strength differences due to greater contractile tissue gains between groups.

The strongest evidence we observed of an effect related to an energy surplus, was the effect of body mass gains on the increase in the sum of skinfold thicknesses. While our group-based analyses roughly comported with our regression - as both surplus groups increased their sum of skinfolds more than MAIN - they did not align with our hypothesized relationship whereby increases in the sum of skinfolds would follow a pattern of HIGH > MOD > MAIN. Rather, our evidence suggests that both MOD and HIGH, on average, were in a similar energy surplus. This finding specifically highlights a challenge of conducting ecologically valid translational research, as the intended difference between groups did not occur despite regular contact between participants and a skilled researcher with clinical nutrition experience. Nonetheless, our regression sidesteps this challenge, highlighting the clearer, strong relationship between body mass gains (and thus, the individual energy surplus magnitude) and sum of skinfold thickness changes. To summarise, it seems the clearest and strongest impact of a larger energy surplus is an increase in body fat, at least in the context of the present RT protocol and study population. Given the aforementioned weak relationship between body mass increases and biceps MT increases, it's possible that had a more potent resistance training stimuli for all muscles been imposed, such a program could have mitigated these gains in body fat to some degree (as more of the energy surplus might have been partitioned towards lean tissue accrual). However, without further study it is difficult to confirm this possibility, or whether the trade-off between greater increases in body fat would be worth the likely proportionally smaller increases in muscle mass (based on the stronger relationship between body mass gains and increases in skinfolds rather than biceps thickness). In practice, the value of this trade off might depend on the context of the individual. For example, someone with body aesthetic goals might choose slower weight gain to mitigate gains in body fat, while an American football lineman who benefits not only from increases in muscle mass, but also increases in body mass (to some degree regardless of composition), may choose faster weight gain.

Ours is one of the few studies on the effect of variable energy surpluses among resistance-trained participants. Previously, Garthe and colleagues [19] conducted an individualized 8-12-week nutritional intervention in 39 resistance-training elite athletes designed to enhance muscle gain. Specifically, participants were divided between two groups, one group was guided by a dietitian to reach a specific, modest daily energy surplus while participants in the comparative group followed a self-guided nutritional approach. The dietitian intervention led to the athletes consuming 3585 ± 601 kcal/d; ~ 600 kcal greater than the comparative group. In line with our findings, despite a five-fold greater increase in fat mass in the dietitian guided group (15 ± 4 vs. $3 \pm 3\%$) there were no significant differences between groups in strength or lean body mass increases. In somewhat of a contrast, using a Bayesian modelling approach, Smith and colleagues [18] reported that gains in body mass were a significant predictor of fat-free mass increases in a group of 21 resistance-trained (minimum 6 months experience) adults during a concurrent overfeeding and resistance-training protocol. For six weeks participants consumed a high energy protein and carbohydrate supplement with the goal of gaining 0.45kg/wk – although actual changes in body mass varied between participants - while performing three supervised weekly RT sessions. Despite a great deal of interindividual variability, the authors' model predicted that a body mass gain of $\sim 0.55\%$ /week resulted was indicative of all body mass gains being fat-free mass ($R^2 = 0.36$). Ultimately, larger surpluses are likely to cause excess gains in body fat, but the degree to which relatively larger or smaller surpluses impact gains in muscle mass is variable between individuals, and possibly impacted by the quality of the training program, its appropriateness for a given study group, which may be at least in part based on their prior training experience and history.

The limitations of the present study are notable. Drops outs and delays due to COVID-19 resulted in an N roughly 60% of what we intended, and thus, our group-based comparisons might be inaccurate due to an insufficient sample size. Further, group assignment was based on the target energy surplus; however, despite competent and consistent monitoring, the intended energy surpluses were not consistently followed by all participants. However, we sought to mitigate these limitations by performing a *post hoc* regression analysis on body mass as a continuous variable which both strengthens our sample size and corrects these energy surplus discrepancies. In addition to these limitations, despite our best intentions, we also had a primarily male sample (two out of 17 participants were female). Thus, future research is required to elucidate any potential sex differences. Additionally, given the relatively large variances in changes in skinfold and muscle thickness, the individual responses to the nutritional and resistance training interventions suggest that a larger sample size may provide a clearer answer on population level responses. Finally, as mentioned, a different, more potent training protocol could have produced different results. While “resistance-trained” participants were recruited, our participants spanned a range of what many would consider late-stage novices all the way to advanced athletes. In many cases our participants habitually trained with more frequency, and in some cases more volume than they performed during this investigation. Thus, logistics permitting, in future study researchers should endeavor to conduct a “lead in” training period where all participants follow the same low to moderate volume protocol in attempt to homogenise individual differences in training status before starting the actual study protocol. Further, in well-trained overfeeding populations, higher frequency and volume protocols may prove produce more

favorable body composition changes than we observed. If more directly supervised training sessions are not feasible, we recommend perhaps including 1–2 self-guided sessions (or sessions supervised by video) in addition to the typically conducted 2–3 supervised, lab-based training sessions.

5 Conclusions

In the present sample individuals who consumed larger energy surpluses - thereby gaining more body mass - experienced similar increases in strength and triceps and quadriceps muscle size but increased their skinfold thicknesses more compared to those who consumed smaller energy surpluses or maintenance calories. There was, however, weak evidence that larger surpluses resulted in greater gains in biceps thickness, the muscle group trained with the most volume and intensity. Thus, if an overfeeding strategy is followed, it may be more successful from a body composition standpoint if accompanied by a more stimulative training protocol for all muscle groups. Ultimately, however, given clearer evidence and a much stronger relationship between body mass gains and increases in the sum of skinfold thicknesses, we recommend conservative energy surpluses scaled to RT experience of 5–20% over maintenance energy or rates of weight gain of 0.25–0.5% of body mass per week, scaled to RT experience such that more advanced trainees consume smaller surpluses and gain weight more slowly [5].

Abbreviations

RT: Resistance training

HIGH: 15% energy surplus with training group

MOD: 5% energy surplus with training group

MAIN: Maintenance energy with training group

MT: Muscle thickness

1-RM: One repetition maximum

VL: Vastus lateralis

VI: Vastus intermedius

RF: Rectus femoris

RPE: Rate of perceived exertion

CI: Credible interval

RIR: Repetitions in reserve

ES: Effect size

Declarations

Ethics Approval and Consent to Participate

All participants were informed of the risks and benefits in plain language and provided informed consent prior to participating in the research. The methods and procedures used in this study were approved by the Auckland University of Technology ethics committee, AUTEK Reference number 18/53.

Consent for Publication

Not applicable

Availability of data and material

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

Legion Athletics provided funding that was used exclusively to pay the research assistants for their time spent collecting data. Renaissance Periodization provided funding that was exclusively used to pay the registered dietitian.

Authors' contributions

Conceptualization: EH, JK, and BD; Methodology: EH; Funding acquisition: EH; Data analysis: JK; Data collection: AJS, CS, DO, and CW; Dietary guidance: ST; All authors assisted in writing and editing the manuscript. All authors read and approved the final manuscript.

Acknowledgements

We wish to thank both the research participants and assistants who made this project possible.

References

1. Brechue WF, Abe T. The role of FFM accumulation and skeletal muscle architecture in powerlifting performance. *Eur J Appl Physiol* [Internet]. 2002;86(4):327–36. Available from: <https://doi.org/10.1007/s00421-001-0543-7>
2. Wang H, Hai S, Liu Y, Liu Y, Dong B. Skeletal Muscle Mass as a Mortality Predictor among Nonagenarians and Centenarians: A Prospective Cohort Study. *Sci Rep*. 2019;9(1):2420.
3. Cawthon PM, Orwoll ES, Peters KE, Ensrud KE, Cauley JA, Kado DM, et al. Strong relation between muscle mass determined by D3-creatine dilution, physical performance, and incidence of falls and mobility limitations in a prospective cohort of older men. *J Gerontol A Biol Sci Med Sci*. 2019 May 16;74(6):844–52.
4. Helms ER, Prnjak K, Linardon J. Towards a Sustainable Nutrition Paradigm in Physique Sport: A Narrative Review. *Sports* (Basel). 2019 Jul 16;7(7).
5. Iraki J, Fitschen P, Espinar S, Helms E. Nutrition Recommendations for Bodybuilders in the Off-Season: A Narrative Review. *Sports* (Basel). 2019 Jun 26;7(7).
6. Lambert CP, Frank LL, Evans WJ. Macronutrient considerations for the sport of bodybuilding. *Sports Med*. 2004;34(5):317–27.
7. Slater G, Phillips SM. Nutrition guidelines for strength sports: Sprinting, weightlifting, throwing events, and bodybuilding. *J Sports Sci* [Internet]. 2011 Jan 1;29(sup1):S67–77. Available from: <https://doi.org/10.1080/02640414.2011.574722>
8. Slater GJ, Dieter BP, Marsh DJ, Helms ER, Shaw G, Iraki J. Is an Energy Surplus Required to Maximize Skeletal Muscle Hypertrophy Associated With Resistance Training. *Front Nutr* [Internet]. 2019;6. Available from: <https://www.frontiersin.org/articles/10.3389/fnut.2019.00131>
9. Morton RW, Murphy KT, McKellar SR, Schoenfeld BJ, Henselmans M, Helms E, et al. A systematic review, meta-analysis and meta-regression of the effect of protein supplementation on resistance training-induced gains in muscle mass and strength in healthy adults. *Br J Sports Med* [Internet]. 2018 Mar 1;52(6):376. Available from: <http://bjsm.bmj.com/content/52/6/376.abstract>
10. Tagawa R, Watanabe D, Ito K, Ueda K, Nakayama K, Sanbongi C, et al. Dose-response relationship between protein intake and muscle mass increase: a systematic review and meta-analysis of randomized controlled trials. *Nutr Rev*. 2020 Nov 4;79(1):66–75.
11. Nunes EA, Colenso-Semple L, McKellar SR, Yau T, Ali MU, Fitzpatrick-Lewis D, et al. Systematic review and meta-analysis of protein intake to support muscle mass and function in healthy adults. *J Cachexia Sarcopenia Muscle*. 2022 Apr;13(2):795–810.
12. Schoenfeld BJ, Ogborn D, Krieger JW. Dose-response relationship between weekly resistance training volume and increases in muscle mass: A systematic review and meta-analysis. *J Sports Sci* [Internet]. 2017 Jun 3;35(11):1073–82. Available from: <https://doi.org/10.1080/02640414.2016.1210197>

13. King A, Helms E, Zinn C, Jukic I. The Ergogenic Effects of Acute Carbohydrate Feeding on Resistance Exercise Performance: A Systematic Review and Meta-analysis. *Sports Medicine* [Internet]. 2022;52(11):2691–712. Available from: <https://doi.org/10.1007/s40279-022-01716-w>
14. Murphy C, Koehler K. Energy deficiency impairs resistance training gains in lean mass but not strength: A meta-analysis and meta-regression. *Scand J Med Sci Sports*. 2022 Jan;32(1):125–37.
15. Whittaker J, Harris M. Low-carbohydrate diets and men's cortisol and testosterone: Systematic review and meta-analysis. *Nutr Health*. 2022 Dec;28(4):543–54.
16. Whittaker J, Wu K. Low-fat diets and testosterone in men: Systematic review and meta-analysis of intervention studies. *J Steroid Biochem Mol Biol* [Internet]. 2021;210:105878. Available from: <https://www.sciencedirect.com/science/article/pii/S0960076021000716>
17. Barakat C, Pearson J, Escalante G, Campbell B, De Souza EO. Body recomposition: can trained individuals build muscle and lose fat at the same time? *Strength Cond J* [Internet]. 2020;42(5). Available from: https://journals.lww.com/nsca-scj/Fulltext/2020/10000/Body_Recomposition__Can_Trained_Individuals_Build.3.aspx
18. Smith RW, Harty PS, Stratton MT, Rafi Z, Rodriguez C, Dellinger JR, et al. Predicting Adaptations to Resistance Training Plus Overfeeding Using Bayesian Regression: A Preliminary Investigation. *J Funct Morphol Kinesiol*. 2021 Apr 21;6(2).
19. Garthe I, Raastad T, Refsnes PE, Sundgot-Borgen J. Effect of nutritional intervention on body composition and performance in elite athletes. *Eur J Sport Sci* [Internet]. 2013 May 1;13(3):295–303. Available from: <https://doi.org/10.1080/17461391.2011.643923>
20. Hall KD. What is the required energy deficit per unit weight loss? *Int J Obes*. 2008;32(3).
21. Tomko PM, Muddle TW, Magrini MA, Colquhoun RJ, Luera MJ, Jenkins ND. Reliability and differences in quadriceps femoris muscle morphology using ultrasonography: The effects of body position and rest time. *Ultrasound*. 2018 Nov;26(4):214–21.
22. Oranchuk DJ, Nelson AR, Storey AG, Cronin JB. Variability of Regional Quadriceps Architecture in Trained Men Assessed by B-Mode and Extended-Field-of-View Ultrasonography. *Int J Sports Physiol Perform* [Internet]. 2020;15(3):430–6. Available from: <https://journals.humankinetics.com/view/journals/ijsp/15/3/article-p430.xml>
23. Oranchuk DJ, Stock MS, Nelson AR, Storey AG, Cronin JB. Variability of regional quadriceps echo intensity in active young men with and without subcutaneous fat correction. *Applied Physiology, Nutrition, and Metabolism* [Internet]. 2020 Jan 9;45(7):745–52. Available from: <https://doi.org/10.1139/apnm-2019-0601>
24. Castanov V, Hassan SA, Shakeri S, Vienneau M, Zabjek K, Richardson D, et al. Muscle architecture of vastus medialis obliquus and longus and its functional implications: A three-dimensional investigation. *Clin Anat*. 2019 May;32(4):515–23.
25. Caresio C, Molinari F, Emanuel G, Minetto MA. Muscle echo intensity: reliability and conditioning factors. *Clin Physiol Funct Imaging*. 2015 Sep;35(5):393–403.

26. Varanoske AN, Fukuda DH, Boone CH, Beyer KS, Stout JR, Hoffman JR. Scanning plane comparison of ultrasound-derived morphological characteristics of the vastus lateralis. *Clin Anat*. 2017 May;30(4):533–42.
27. Dankel SJ, Abe T, Bell ZW, Jessee MB, Buckner SL, Mattocks KT, et al. The impact of ultrasound probe tilt on muscle thickness and echo-intensity: a cross-sectional study. *J Clin Densitom*. 2020;23(4):630–8.
28. Zourdos MC, Klemp A, Dolan C, Quiles JM, Schau KA, Jo E, et al. Novel Resistance Training-Specific Rating of Perceived Exertion Scale Measuring Repetitions in Reserve. *J Strength Cond Res*. 2016;30(1).
29. Technical rules book of the international powerlifting federation. 2022. p. 17–8.
30. Jenkins NDM, Miller JM, Buckner SL, Cochrane KC, Bergstrom HC, Hill EC, et al. Test–Retest Reliability of Single Transverse versus Panoramic Ultrasound Imaging for Muscle Size and Echo Intensity of the Biceps Brachii. *Ultrasound Med Biol* [Internet]. 2015;41(6):1584–91. Available from: <https://www.sciencedirect.com/science/article/pii/S0301562915000563>
31. Radaelli R, Bottaro M, Wilhelm EN, Wagner DR, Pinto RS. Time course of strength and echo intensity recovery after resistance exercise in women. *J Strength Cond Res*. 2012 Sep;26(9):2577–84.
32. Andraszewicz S, Scheibehenne B, Rieskamp J, Grasman R, Verhagen J, Wagenmakers EJ. An Introduction to Bayesian Hypothesis Testing for Management Research. *J Manage*. 2015 Feb 10;41(2):521–43.
33. Hespanhol L, Vallio CS, Costa LM, Saragiotto BT. Understanding and interpreting confidence and credible intervals around effect estimates. Vol. 23, *Brazilian Journal of Physical Therapy*. 2019.
34. Baz-Valle E, Balsalobre-Fernández C, Alix-Fages C, Santos-Concejero J. A systematic review of the effects of different resistance training volumes on muscle hypertrophy. *J Hum Kinet*. 2022 Jan;81:199–210.
35. Helms ER, Byrnes RK, Cooke DM, Haischer MH, Carzoli JP, Johnson TK, et al. RPE vs. Percentage 1RM Loading in Periodized Programs Matched for Sets and Repetitions. *Front Physiol*. 2018;9:247.
36. Refalo MC, Helms ER, Trexler EricT, Hamilton DL, Fyfe JJ. Influence of Resistance Training Proximity-to-Failure on Skeletal Muscle Hypertrophy: A Systematic Review with Meta-analysis. *Sports Medicine* [Internet]. 2023;53(3):649–65. Available from: <https://doi.org/10.1007/s40279-022-01784-y>

Supplementary File

Supplementary File is not available with this version.

Figures




Figure 1

Changes in muscle thickness from pre- to post-intervention. MOD=moderate surplus, MAIN=maintenance, HIGH=high surplus. MT=muscle thickness, Δ =change, VL=vastus lateralis, VI=vastus intermedius, RF=rectus femoris. Box plots illustrate the median, interquartile range, and minimum and maximum values. Individual data points beyond the whiskers are considered outliers (quartile $\pm 1.5 \times$ interquartile range).




Figure 2

Changes in one-repetition maximum (1-RM) from pre- to post-intervention. MOD=moderate surplus, MAIN=maintenance, HIGH=high surplus. Δ =change. Box plots illustrate the median, interquartile range, and minimum and maximum values. Individual data points beyond the whiskers are considered outliers (quartile \pm 1.5 \times interquartile range).




Figure 3

Changes in the sum of skinfolds from pre- to post-intervention. MOD=moderate surplus, MAIN=maintenance, HIGH=high surplus. Δ =change. Box plots illustrate the median, interquartile range, and minimum and maximum values.