

# Morphological Variation of the Deciduous Second Molars in the Baka Pygmies

**Petra G. Šimková**

University of Vienna

**Gerhard W. Weber**

University of Vienna

**Fernando V. Ramirez Rozzi**

UMR7206 Ecoanthropologie, MNHN, CNRS

**Lotfi Slimani**

Université de Paris

**Jérémy Sadoine**

Université de Paris

**Cinzia Fornai** (✉ [cinzia.fornai@univie.ac.at](mailto:cinzia.fornai@univie.ac.at))

University of Vienna

---

## Research Article

**Keywords:** Morphological variation, molars, Baka Pygmies, infancy, peculiar, African population, biology, South American

**Posted Date:** July 22nd, 2021

**DOI:** <https://doi.org/10.21203/rs.3.rs-322362/v2>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

**Version of Record:** A version of this preprint was published at Scientific Reports on August 13th, 2021. See the published version at <https://doi.org/10.1038/s41598-021-95524-3>.

# Abstract

The Baka Pygmies are known for their short stature resulting from a reduced growth rate during infancy. They are peculiar also for their teeth erupt earlier than in any other African population, and their posterior dentition is larger than in non-Pygmy populations. However, the Baka's dental morphology, like several other aspects of their biology, is still understudied. Here, we explore the variation of the Baka's deciduous upper and lower second molars (dm2s) in comparison to a geographically heterogeneous human sample by means of 3D geometric morphometrics and analysis of dental traits. Our results show that the different populations largely overlap based on the shape of their dm2s, especially the lower ones. Their distal region and the height of the dentinal crown differ the most, with the Baka showing the most extreme range of variation. Upper and lower dm2s covary to a great extent ( $RV=0.82$ ). The Baka's and South Americans' dm2s were confirmed among the largest in our sample. Despite the Baka's unique growth pattern, long-lasting isolation, and extreme dental variation, it is not possible to distinguish them from other populations based on their dm2s' morphology only.

## Introduction

The Baka people are a Central African population with an average male height under 155 cm<sup>1</sup>. Like many other Central African human groups, the Baka have chosen to refer to themselves as Pygmies<sup>2</sup>, accordingly, we will use the term Pygmy in this manuscript for the reasons detailed in Methods.

Until recently, most African Pygmies maintained semi-nomadic lifestyles based mainly on hunting and gathering<sup>1,3</sup>. From the mid-20<sup>th</sup> century, the missionary and sedentarization programs influenced the lifestyle of the Baka populations<sup>4-9</sup> which moved closer to village sites inhabited by Bantu-speaking agriculturists. Some of the Baka started practicing horticulture in particular seasons of the year and trade products with their neighbors<sup>5,7,8</sup>. Nowadays, the Baka's hunting camps in the rainforest last only for a few months per year during the major dry season<sup>5,8</sup>, but they remain an essential source of wild resources the Baka are highly depending on<sup>7</sup>.

Short average stature in Pygmy populations results from their particular growth patterns, but the actual underlying evolutionary mechanisms have been long debated<sup>10,11</sup>. Pygmies' small body size was mainly interpreted as an adaptation to environmental conditions, food shortage<sup>11-13</sup>, and thermoregulation<sup>14</sup>. It is now established that the genetic basis of this specific phenotype is polygenic, including genes related to skeletal growth, allometric patterning, immunity, and metabolism, with effects on the growth hormone axis, in particular the expression of the growth hormone-insulin-like growth factor I<sup>15-17</sup>.

The life history variables in the Baka Pygmies seem to correspond with those of other non-Pygmy populations, including puberty growth spurts and maturity reached at a very similar age<sup>10</sup>. Furthermore, their birth size and birth weight are within the standard limits of modern humans, differently from other Pygmy populations that are already born with smaller body size<sup>6,10</sup>. However, during the first two years of

life, the Baka experience a significant deceleration of growth which causes a long-lasting growth delay throughout their whole development and results in a short adult stature (average women's height = 146.7 cm; average men's height = 153.5 cm)<sup>10</sup>. Although the timing of tooth eruption is commonly strongly correlated with other life history variables, in the Baka Pygmies the permanent dentition erupts earlier than in any other African groups, including other Pygmy populations (in females upper and lower first molars erupt respectively 0.75 and 0.71 years earlier, while in males 0.63 and 0.39 years earlier)<sup>3</sup>. The Baka's permanent dentition has been studied with respect to other non-Pygmy populations as well as in relation to their body size<sup>3,7,18-20</sup>. Their post-canine dentition was found to be larger than in their Bantu neighbors<sup>18</sup> as well as in other non-Pygmy populations. Sexual dimorphism of the Baka's permanent molars has been identified, being male molars significantly larger than in females<sup>18</sup>. However, no consistent pattern of correlation was found between the Baka's dental size and body height or weight<sup>19</sup>.

Several important aspects of the Baka's biology are still unknown, and the morphological variation of the Baka's teeth is yet unexplored. Given the peculiar set of phenotypical expressions shown by the Baka, we want to contribute to the understanding of the Baka's morphological variation by investigating their deciduous upper and lower second molars (dm2) compared to a geographically diverse dm2 sample (Table 1). For this purpose, we use two approaches: 3D landmark- and semilandmark-based methods and analysis of qualitative dental traits. Working with  $\mu$ CT derived 3D surface models, the geometry of both the enamel-dentine junction (EDJ) and the outer enamel surface<sup>21</sup> can be safely captured, while the roots cannot be investigated since they are completely resorbed in our Baka sample. Similarly, the moderate degree of wear of the occlusal aspect of the crowns permits the investigation of a limited number of discrete dental traits. Based on their results, Sardi and Ramirez Rozzi<sup>22</sup> inferred that the genetic pathways regulating cranial growth and dental morphogenesis are most likely distinct. If this is the case, we can expect that a reduced growth rate during infancy, resulting in short adult stature in the Baka Pygmies, will not necessarily correspond to dental changes. Thus, we predict that the Baka's dental shape does not differ substantially from other human groups. Additionally, considering the outcomes of previous work<sup>18</sup>, we expect to find sex-related size differences within the Baka's dm2s.

## Methods

### On the usage of the term 'pygmy'

In spite of the fact that term pygmy has been used by some as pejorative, its etymology is not dubious nor derogatory, deriving from the Greek *pygmē* 'a cubit', the measure of length from the elbow to the extreme of the middle finger (for an historical account of the usage of the term pygmy see Ballabriga, 1981<sup>54</sup>; Bahuchet, 1993<sup>55</sup>; Ramirez Rozzi, 2015<sup>56</sup>; 2021<sup>57</sup>). From a scientific standpoint, population genetics has shown that all human groups living in the African Equatorial forest usually referred to as Pygmies, shared a common ancestor which split from non-Pygmy groups around 65-50,000 years ago<sup>51,53</sup>: The term pygmy identifies groups of humans based on their geographic distribution, morphology, adaptation to the environment, growth pattern, and cultural and social traits: the Pygmies

have a distinct population history which deserves to be acknowledged. In fact, adopting the term 'Pygmy', the Baka distinguish themselves from the Nzime, a group of Bantu ethnicity living in the surroundings of the Bosquet area. Some among the detractors of the term Pygmy have suggested the use of the term 'twa' (meaning 'dwarf') as an alternative. However, the Baka strongly oppose to defining themselves as such since non-Pygmy neighbor populations use it with disparaging intent. Others have proposed 'hunters-gatherers', but this term accounts only for the subsistence strategy thereby clouding all of the other distinctive traits of the Pygmies. The Pygmies have faced centuries of oppression which have led them to hide their identity to escape persecution<sup>58</sup>. Thus, referring to the Baka as Pygmies, is not only proper in scientific and anthropological terms, but is also ethically correct.

## Sample

Our sample consists of 60 upper and 59 lower modern human deciduous second molars (Table 1). Of these, 55 teeth, including 23 udm2s and 32 ldm2s, are from 39 Baka individuals. The Baka were collected by F.V.R.R. after the Baka children naturally shed and donated them with the consent of their families. The rest of our sample was composed of individuals from different geographical regions, including Europe (udm2 = 22; ldm2 = 11), Africa (udm2 = 6; ldm2 = 7), Asia (udm2 = 5; ldm2 = 3), the Near East (udm2 = 2; ldm2 = 4), and South America (udm2 = 2; ldm2 = 2), which is quite a large sample considering the paucity of infant and juvenile specimens in osteological collections. More so, the meta-data for the comparative sample is often incomplete since the individuals come from archaeological collections (Table 1). Other limitations of the sample size are the wear stage and state of preservation of the teeth. Dental specimens showing a moderate abrasion of the horn tips (stages 3 and 4) were virtually reconstructed (see *μCT acquisition and data segmentation* below) before data collection. To achieve a successful virtual reconstruction of the EDJ surface, wear cannot exceed stage 4<sup>23</sup>, thus specimens showing higher degree of wear or with extensive decay in the occlusal area were either excluded from the sample or used only for the analyses of the cervical and crown outlines (see below). The heavy wear, the poor state of preservation, and occasional dental treatment made a large number of the Baka dental collection unusable for our study. In fact, the Baka's traditional dental treatments are quite invasive and entail the drilling of large cavities later filled with natural substances<sup>9</sup>. The teeth were naturally shed and thus they were used approximately until the age of 11 which explains the advanced degree of wear. In some cases, the molars were kept by the Baka children or their families until the next visit of F.V.R.R., which likely led to the poor state of preservation of the teeth at the moment of image data acquisition (i.e., for the presence of numerous and deep cracks).

## μCT acquisition and data segmentation

The dental datasets were imaged at four different facilities. The Baka teeth were scanned mainly at the Plateforme Imageries du vivant, Université de Paris, using a Micro CT-scanner PerkinElmer, Quantum FX (voxel size 20 - 40/30 - 59/10 - 20 μm, 90 kV, 16 mA) and at the Hard Tissue Research Unit, College of Dentistry (NYU), with a SCANCO Scantron 40 Micro-CT scanner (voxel size 12 μm, 70 kV, 275 mA, 200 ms). The comparative sample was scanned at the Centre de Microtomographie of the Université de

Poitiers, with a VISCOM X8050-16 (voxel size 24 – 60  $\mu\text{m}$ , 95 – 110 kV, 0.5 mA), and at the Vienna Micro-CT Lab, Austria, with a VISCOM X8060 NDT scanner (voxel size 21 – 60  $\mu\text{m}$ , 110 – 140 kV, 280 – 410 mA, 1400 – 2000 ms, 0.75 mm copper filter). X-ray images were taken from 1440 different angles. Using filtered back-projection in VISCOM XVR-CT 1.07 software, these data were reconstructed as 3D volumes with a color depth of 16,384 grey values.

The  $\mu\text{CT}$  data were then imported into Amira software ([www.fei.com](http://www.fei.com)) and virtually segmented to separate the enamel from the dentine, pulp, and the surrounding material (i.e., air, alveolar bone). In case of a slight abrasion of the dentinal horn tips, the specimens were virtually reconstructed by using the “brush” tool and extending the contours of the existing dentine into the empty area. In case both left and right dm2s were available from one individual, we preferred the left ones among our whole sample. However, if better preserved, we used the right tooth after virtual mirroring to the left side. Since there is no scientific evidence indicating the existence of directional asymmetry in human dentition, we assume that the choice of left teeth should not affect the results of this study<sup>59–61</sup>.

### **Reorientation and outline collection**

After segmentation, the surface models of the crowns were consistently reoriented in Geomagic Design X 64 ([www.3dsystems.com](http://www.3dsystems.com)) following an established protocol<sup>62–64</sup>. The crown and cervical outlines were collected from the reoriented surface models and projected onto the cervical plane. Afterwards, the outlines were split into 24 segments by as many equiangular radial vectors originating from the centroid of the outline area, using Rhinoceros 6 ([www.rhino3d.com](http://www.rhino3d.com)). Twenty-four pseudo-landmarks were placed at the point of intersection of the radii and the outline.

### **Landmark collection on the enamel-dentine junction**

To collect the landmarks on the EDJ, we followed established protocols<sup>62–64</sup>. For the Idm2s, we placed a total of eight landmarks (LM) on the five main horn tips and three at the deepest points between metaconid and entoconid, protoconid and hypoconid, and between hypoconid and hypoconulid. Afterwards, 23 curve semilandmarks (sLM) were placed to represent the EDJ marginal edge. To ensure homology, we traced the EDJ occlusal edge by creating a spline curve ignoring all the accessory cusps. For the udm2s, seven LMs were placed on the four horn tips, and on the deepest points of the central fovea and distal fossa, and the deepest point of the lingual marginal ridge of the hypocone. The EDJ marginal edge was resampled by 47 sLMs. The LM collection was carried out in the EVAN Toolbox 1.75 ([www.evan-society.org](http://www.evan-society.org)), which uses the bending energy technique for sliding the sLM<sup>65–67</sup>.

### **Geometric Morphometric analysis**

The geometric morphometric analyses were performed using the Evan Toolbox separately for each set of landmarks, resulting in four different analyses per tooth type, namely: 1) cervical outlines; 2) crown outlines; 3) EDJ and 4) dentinal crown, combining the landmark configuration of the EDJ with the cervical outline. First, the landmark configurations had to be normalized via General Procrustes Analysis

(GPA)<sup>68,65,69</sup>. The translation and rotation of the landmark sets were not necessary for the outline configurations since they were manually aligned. We run the principal component analysis (PCA) on the Procrustes shape coordinates and visualized the shape changes along the principal components by means of warpings, using the Thin-Plate Spline technique<sup>70,71</sup>. The analyses of the size were carried out using the natural logarithm of Centroid Size (lnCS). The lnCS of the combined EDJ and cervical outline is an accurate measure of the dentinal crown size since it embeds information about the crown's height, too. We explored shape covariation between udm2s and ldm2s as well as between different features of the same dental types by means of the 2-block partial least squares analysis (2B-PLS). Using R Studio ([www.r-project.org](http://www.r-project.org)), we performed the analysis of variance on Procrustes shape coordinates of larger subsamples, i.e., the Baka and the Europeans, to assess their degree of variation in the morphological expression of udm2s and ldm2s. In addition, the group mean differences between male and female Baka dentinal crowns were assessed with a permutation test of the Procrustes distances (10,000 random permutations) using a R script written for this purpose. A Mann-Whitney U test was used to test significance in size differences between the Baka and the Europeans, and between the Baka and the rest of the sample. Another one was performed to the sex-related size differences within the Baka. This analysis could not be carried out on the other populations owing to the lack of associated meta-data. These tests have been carried out using PAST 4.03 ([www.softpedia.com](http://www.softpedia.com)), as well as the Kruskal-Wallis test used to assess the size differences on populational level across the whole sample. Moreover, we performed a multivariate regression to analyze size-related shape variation of the dentinal crown.

## **Non-metric traits**

The non-metric traits were evaluated based on the Arizona State University Dental Anthropology System (ASUDAS)<sup>24,72</sup>. Because of the high degree of wear of the Baka sample and root resorption, we focused on four among the most informative dental traits visible on the outer enamel surface, possibly reflecting neutral genetic variation<sup>21,73</sup>.

### **1. Hypocone**

The hypocone is the fourth cusp of the upper molars that forms a separate region of the occlusal aspect, the trigon. Its manifestation varies from grade 0 to 6<sup>24</sup>. In this study, we used dichotomous classes to represent the hypocone degree of expression: none/light (0-3) and moderate/heavy (4-6).

### **2. Carabelli cusp**

The Carabelli cusp is an upper molar's accessory cusp occurring on the protocone (mesio-lingual cusp). This trait has been used as a diagnostic trait for European populations<sup>42</sup>, however, according to more recent studies, no differences in prevalence and expression between populations are expected<sup>24</sup>. According to ASUDAS, the Carabelli cusp's expression varies between grades 0 to 7, but for this study, we used two categories: none/light (0-4) and moderate/heavy (5-7).

### **3. Entoconulid**

The entoconulid, or cusp 6, can be found in the distal area of the occlusal aspect of lower molars between the hypoconulid and the entoconid. The manifestation of the entoconulid can be expressed between grades 0 to 5, which are grouped into none/light (0-2) and moderate/heavy (3-5) in our study.

#### 4. Metaconulid

The metaconulid, or cusp 7, can be found on the lingual aspect of the lower molars between the entoconid and metaconid. Its manifestation can be expressed with 6 grades: 0, 1, 1A, 2, 3, and 4. We dichotomized the scoring values as none/light (0 to 1A) and moderate/heavy (2 to 4).

The prevalence of the non-metric traits was analyzed by  $\chi^2$  test using SPSS ([www.ibm.com](http://www.ibm.com)).

## Results

The most salient finding of originating from our geometric morphometric investigation the analysis of the upper and lower dm2s can be summarized as follows. The populations show an overlapping distribution (Fig. 1; Fig. S1, S2, S3, S4). The dm2s vary from low-crowned and broad to tall and narrow. This pattern of variation is concordant between the upper and lower dm2s. Additionally, a reversed pattern of variation was detected for the upper and lower dm2s: the Baka's Idm2s showed a wider range of variation in comparison to the udm2s, while the opposite was found for the European upper and lower dm2s. Depending on the analysis, 58 to 71% of the total variance can be explained by the first three principal components (PCs; Table 2).

**Lower dm2 shape variation.** The results of the Idm2's dentinal crowns analysis (using landmarks resampling the EDJ marginal edge and the cervical outline; Figure 1a), revealed a broad overlap of all populations with the Baka sample. Shape variation along PC1 (explaining 32% of the total variance) concerned the relative bucco-lingual position of the hypoconulid corresponding with the relative bucco-lingual distal expansion of the dentinal crown, as well as its height. Variation along PC2 (16%) was driven by the mesio-distal elongation of the dentinal crown. Both along PC1 and PC2, the Baka exceeded the full range of shape variations within this geographically diverse sample for their particularly short and mesio-distally elongated dentinal crowns, and relatively bucco-lingually narrow distal aspect. We observed a 20% higher variance for the Baka sample (0.006) than for the European Idm2s (0.005) (Supplementary Table 1 online). Supplementary Figure S1 shows the sample distribution along PC3. The Baka male and female means did not differ significantly ( $p = 0.758$ ) in terms of Idm2 dentinal crown shape. Sexual dimorphism could not be tested in the other populations since their individuals' sex was unknown. As expected, the analysis of the EDJ alone did not provide any additional information with respect to the analysis of the dentinal crown, therefore we find reporting on this analysis redundant.

The Idm2 cervical outline (see Supplementary Fig. S2a online) varied mainly between round and hourglass-shaped, with the Baka encompassing almost the full range of variation of the considered sample, along PC1 (30%). Along PC2 (26%), the cervical outlines varied from elongated and narrow, to short and broad. The Baka and the South American Idm2s reached the most extreme configurations in

terms of bucco-lingual constriction and mesio-distal elongation. Similar to the cervical outline, the crown outline (see Supplementary Fig. S2b online) varied from elongated and bucco-lingually constricted to rounded and bucco-lingually expanded along PC1 (32%), and from mesio-buccally expanded to bucco-distally expanded along PC2 (21%).

**Upper dm2 shape variation.** The results for the udm2s dental crowns (Figure 1b) showed again an overlapping distribution of Europeans and Southeast Asians with the Baka. Along PC1 (29%), the udm2 dental crown varied between short with squared occlusal aspects, and taller with lingually displaced distal cusps (i.e., metacone and hypocone). Along PC2 (20%), variation was driven by the relative expansion of the trigon (consisting of the three main cusps and forming the central fovea) with respect to the talon (consisting of the disto-lingual cusp and fossa). In other words, the dental crowns possessed variably expanded central fovea with respect to their base. The Baka tend to have low, square-shaped dental crowns, while Europeans are more variable and are characterized also by tall teeth with lingually shifted distal cusps and reduced metacone. A clear separation between the Baka and the Egyptians, Bedouins, and South Americans was found, although these groups were represented by only two to four specimens. The analysis of variance (see Supplementary Table 1 online) revealed a 40% higher variance in the Europeans' udm2s (0.007) with respect to the Baka (0.005). Shape variation along PC3 is shown in Supplementary Figure S3 online. The Baka male and female means did not differ significantly ( $p = 0.456$ ) in terms of ldm2 dental crown shape. The other groups could not be tested because the information on the individuals' sex was missing. Similar to the ldm2s, the EDJ analysis for the udm2s was not a source of additional information with respect to the analysis of the dental crown.

For the cervical outline (see Supplementary Fig. S4a online), PC1 (34%) reflects the relative expansion of the disto-lingual and disto-buccal aspects with respect to each other, while PC2 (23%) represents the relative mesio-distal elongation of the cervical outline. The crown outlines changed from oval to rounded along PC1 (37%; see Supplementary Fig. S4b online). This shape change was associated with the relative mesio-buccal position of the paracone. Variation along PC2 (18%) consisted in the relative expansion of the talon with respect to the trigon.

**Covariation.** The covariation between upper and lower dm2s could be assessed only for individuals represented by both tooth types ( $n = 20$  of which 13 Baka). For the dental crown, the percentage of total squared covariance of udm2s and ldm2s for the Singular Warp (SW) scores 1 was 66% and the pairwise shape correlation between the antagonists was  $r_1 = 0.82$  (Table 3). Upper and lower dm2s clearly showed common trends of variation (Figure 2), being either short-crowned with reduced central fossa (in udm2s) and distal fossa (in ldm2s), or taller with expanded opposing fossae. Moreover, lower dental crowns were associated with relatively higher horn tips than taller crowns. It was not possible to clearly separate the populations from each other based on the upper and lower dm2 dyads, although the Baka tend to show extreme expression towards low and elongated dental crowns. Within the same tooth type, the highest covariation coefficient was found between the ldm2s cervical and crown outlines ( $r_1 = 0.81$ ), while cervical and crown outlines in udm2s covaried the least ( $r_1 = 0.74$ ).

**Size.** The comparison of the Baka's natural logarithm of Centroid Size (lnCS) for the dentinal crown with respect to the rest of the sample, showed that the Baka's upper and lower dm2s, tend to be among the largest specimens within our sample following the South Americans (Figure 3). With respect to the various populations, the Baka's upper and lower dm2s differed significantly from Europeans, which similarly to the Egyptians were among the smallest specimens. Similar results were obtained by observing the crown outline. The cervical and crown outlines of the Baka's udm2s were significantly larger than the rest of the sample (Table 4). Interestingly, while the Baka possess the largest udm2s cervical outline, the Egyptians show the highest values for the ldm2s. However, the findings for the groups with the smallest sample size (i.e., Egyptians, South American, and Southeast Asian individuals) require caution.

upper second molars. South Americans and the Baka possess the largest deciduous second molars. The smallest specimens are within the Europeans and Egyptians.

The multivariate regression showed that only a low percentage of morphological variation of the dentinal crown could be explained by lnCS (ldm2: ~3%; udm2: ~9%). In ldm2s, we observed size-related shape variation mainly connected to the expansion and reduction of the distal and occlusal aspects, and the height of the dentinal crown. Larger ldm2s showed low dentinal crowns with expanded distal aspects and rather mesio-distally elongated occlusal aspects. This underlines the fact that the largest teeth (among which the Baka) are short, narrow, and elongated, while the smallest (including Europeans) tend to be taller, mesio-distally shorter, and distally broader. Similarly, size-related shape changes in udm2s affected the height of the dentinal crown, the relative buccal expansion of the metacone and the occlusal aspect, and the bucco-lingual position of the hypocone. Larger udm2s (including the Baka) tended to show lower dentinal crowns with expanded metacone and buccally shifted hypocone, resulting in a squared crown as seen in occlusal view. Conversely, smaller udm2 specimens, including the Europeans, tend to show high crown, reduced metacone and lingually shifted hypocone.

The dm2 size differences between the Baka males and females were not statistically supported (see Supplementary Fig. S5 and Supplementary Table S2 online) based on the lnCS of the dentinal crowns or crown outlines. These analyses could not be carried out on the whole sample due to the lack of necessary meta-data.

**Non-metric traits.** The prevalence of the hypocone and Carabelli cusp for the udm2s, and entoconulid and metaconulid for the ldm2s are presented in Table 5. The high degree of occlusal wear along with the resorption of the roots in the Baka specimens hampered the observation of further discrete features. The hypocone was present in each of the udm2 specimens in our sample, and was the most variable trait although it was highly expressed in most of the individuals (grade 4 to 6)<sup>24</sup>. The Baka showed a significantly higher frequency of a massive hypocone manifestation (96%) in comparison to the rest of the sample ( $\chi^2 = 11.7$ ;  $p = 0.001$ ). Most specimens, however, showed a medium grade of expression (3-4). In our sample, we found no or slight expression of the Carabelli cusp, of the entoconulid, and the metaconulid, with no significant differences between populations.

## Discussion

The Baka Pygmies from south-eastern Cameroon, western Central African Republic, and northern Congo<sup>5-8,10,18</sup> are characterized by their peculiar life history and growth pattern with respect to other Pygmy and non-Pygmy populations. They are also known for their rather large permanent molars and early eruption times of the permanent teeth. However, in terms of dental morphology, we found that their dm2s are indistinguishable from other non-Pygmy populations. In fact, we observed a great overlap between populations in our Idm2 sample, which hampered the possibility to distinguish the diverse geographical groups based only on their tooth shape. Similar findings were obtained for permanent upper and lower premolars<sup>25,26</sup>, and first and second molars<sup>27,28</sup> from non-Pygmy populations.

Based on dental crown diameters, most of the variation occurs within rather than between populations<sup>29,30</sup>. The high degree of shape variation observed in the Baka's Idm2s is compatible with previous observations on Sub-Saharan populations, which have been proven highly variable in terms of both dental<sup>29</sup> and cranial dimensions<sup>31,32</sup>. Surprisingly, the extended shape distribution of the Baka's lower dm2s is not paralleled by the upper dm2s. Interestingly, the pattern of variation in European dm2s follows the opposite trend, with less variable lower dm2s and highly variable upper dm2s. This finding is consistent with our knowledge of the European permanent upper molars showing a high degree of hypocone and metacone reduction<sup>28</sup>.

Given the morphological resemblances of the dm2s to first molars (M1s)<sup>33-37</sup>, we draw comparisons of our results with those previously published on M1s<sup>28,38,39</sup>. Distal and lingual aspects appear to be the most variable regions in both upper and lower dm2s. This finding is in agreement with Halász's<sup>28</sup> research on M1s from various populations, and with Polychronis et al.<sup>39</sup>, who studied M1s from a Greek populations. There seems to be a clear trend that points to an increased shape variation in the distal, bucco-distal, and disto-lingual aspects of the crown. The pattern of shape covariation observed between upper and lower dm2s, from tall and narrow-crowned with expanded central and distal fossae to short and wide-crowned with reduced opposing fossae and relatively higher horn tips, was also observed in upper and lower M1s<sup>28</sup>. High pairwise correlation between opposing teeth developing in different jaws can be interpreted as a strong hint to tight genetic control over the dental development and is explained by the fact that antagonists have to occlude optimally to facilitate effective mastication. Likely, these functional constraints concern all posterior teeth, in fact similar results were achieved for dyads of other dental types<sup>26,28</sup>.

In terms of non-metric traits, our results for the hypocone contrasted the findings of Edgar and Lease<sup>33</sup> who observed a high percentage of reduced hypocone expression in deciduous molars in general. Instead, we found expression of larger hypocones in the Baka than in Europeans, as Lease<sup>40</sup> found between African Americans and Europeans. Today it is known that the manifestation of the Carabelli cusp is equally common in all world populations<sup>41</sup>, which was confirmed by our findings. However differently from Halász<sup>28</sup> who observed the manifestation prevalence of 100% in African and Near

Eastern M1s, and in accordance to Scott<sup>42</sup> we found that most of the specimens lacked a Carabelli cusp. According to Harris<sup>43</sup> the grade of expression of the Carabelli cusp in M1s and the tooth size are connected, with larger teeth showing this trait more often. We cannot support this result, since we did not find any significant differences in Carabelli cusp manifestation between the populations in spite of the significant size differences. Likewise, the various populations used in our sample did not differ in terms of entoconulid and metaconulid frequency or degree of expression.

Variation in dental size of the Pygmies' permanent dentition in comparison to non-Pygmy populations was assessed previously for various samples testing different hypotheses<sup>18-20,44</sup>. The general findings include the tendency of sub-Saharan African populations to show larger permanent teeth than Asians or Europeans<sup>30</sup>, which was confirmed by our results. According to Hanihara and Ishida<sup>30</sup>, the modern Australian aboriginals show the largest dental diameters, followed by a group of average-sized dentitions of sub-Saharan Africans, Native Americans, and Southeast Asians. Europeans and populations of the Near East show the smallest dentition. We observed the smallest dm2s sizes in Europeans and Bedouins, followed by Egyptians, Southeast Asians, and sub-Saharan Baka Pygmies. South Americans showed the largest dm2 crowns. It is worth noting that our results use InCS rather than linear distances, a different measure of size, depending also on the landmark configuration<sup>26</sup>. The dental size of the Baka and other Pygmies was studied also in relation to their body size and developmental patterns<sup>19,20</sup>. The results for the Baka's dm2s match previous findings on permanent molars which are larger than in any other non-Pygmy population<sup>20</sup>, including the Bantu<sup>18</sup>. However, the Baka's dm2s do not show sex-related size variation that was previously reported for the permanent molars<sup>18</sup>.

Early dental eruption timing has not only been found in the Baka's dm2s<sup>3</sup> but also in other sub-Saharan populations' permanent<sup>45-48</sup> and deciduous teeth<sup>49</sup>. According to Lam et al. (2015)<sup>50</sup>, the earlier onset of tooth eruption can be associated with postnatal factors such as an increased rate of weight gain during the first three months of life. It is worth noting that, contrary to the Sua Pygmies, the Baka's weight gain during the first years of infancy does not differ from that of the Bantu<sup>3</sup>. However, considering that the time of dm2 formation partially overlaps with the time of growth decrease in the Baka, and the dm2s' size is not negatively affected by the decrease in body growth, we can assume no or only a very low correlation between odontogenesis and body growth in the Baka. Romero et al.<sup>18</sup> and Ramirez Rozzi<sup>19</sup> underline this finding stating that short stature is likely an adaptation not related to dental shape and size variation, and is genetically inherited by the Baka's ancestors<sup>51</sup>. Furthermore, no clear genetic pathway affecting both the teeth and the somatic growth that could explain the phenomena we observed in the Baka has yet been found<sup>19</sup>. Earlier tooth eruption has been associated with a lower degree of root growth<sup>52</sup>, although root development and morphology in the Baka have not been studied yet, therefore the mechanisms prompting an early eruption of their teeth remain unknown.

In conclusion, despite the Baka's long-lasting geographical isolation<sup>3,51</sup>, very low levels of genetic admixture with non-Pygmy neighboring populations<sup>51,53</sup>, specific growth pattern, and dental eruption

timing, we did not find a morphological difference distinguishing the Baka from other world populations based only on their dental morphology. As expected, we also did not observe any significant shape differences between sexes. Yet, the Baka's Idm2s can be particularly narrow and elongated with respect to the rest of the sample analyzed, and udm2s also clearly exceed the range of low and squared-crown variation in the sample, despite the fact that they are less variable than the Idm2s. Outstandingly, differential patterns of variation of udm2s and Idm2s in Central Europeans and the Baka might reflect their different population history reaching back to the migration of the European ancestors out of Africa<sup>17</sup>, a topic that deserves further attention. Still, assigning an individual tooth to a certain population based on its shape alone remains impossible.

## Declarations

### Availability of materials and data:

All data generated or analyzed during this study are either included in this published article (and its Supplementary Information file) or available from the corresponding author on reasonable request.

### Acknowledgements:

This study would have not been possible without the kind collaboration of the Baka Pygmies, and especially of the Baka children and their families who facilitated access to the milk teeth. We are grateful to Sabine Eggers and Karin Wilschke-Schrotta (Natural History Museum, Vienna, Austria), Israel Hershkovitz (Department of Anatomy and Anthropology, Sackler Faculty of Medicine, Tel Aviv University, Israel), Priscilla Bayle (Université de Poitiers, Centre de Microtomographie, France), Stefano Benazzi and Giorgio Gruppioni (University of Bologna, Department for the Cultural Heritage, Italy) who granted access to the comparative materials. Special thanks to Martin Dockner (Vienna  $\mu$ CT Lab) for his friendly and ready technical support.

### Author contributions:

C.F. and F.V.R.R. initiated the project. C.F. and G.W.W. designed and supervised the project. P.G.Š. and C.F. performed the image data processing. P.G.Š. collected landmark data. P.G.Š. carried out the analyses supported by C.F. and G.W.W. P.G.Š. wrote the paper with the contribution of C.F., G.W.W. and F.V.R.R. P.G.Š. compiled figures and tables. L.S., J.S. acquired the  $\mu$ CT scans.

### Additional information:

C.F. was financially supported by the Swiss National Science Foundation (grant No 31003A\_176319). The authors declare no potential conflict of interest.

### Ethical statement:

We did not use any physically invasive or destructive methods in our study. Our sample included only teeth obtained from archaeological collections or donated by the Baka children after they naturally shed them, and thus were not extracted. The Baka teeth were collected during field works carried out as part of an international agreement between the French National Research Institute for Development (IRD) and the Ministry of Scientific Research and Technology of Cameroon (Accord-cadre de Coopération en matière de Recherche Scientifique et Technique, 2004). Accordingly, informed consent was obtained from all participants and from both parents of any participants aged under 18.

## References

1. Cavalli-Sforza, L. L. *African pygmies*. (Academic Pr, 1986).
2. Ramirez Rozzi, F. V. Thoughts about responsibility in anthropological research, from a study with the Baka Pygmies. *Ethics Med. Public Health***18**, 100668 (2021).
3. Ramirez Rozzi, F. V. Diversity in tooth eruption and life history in humans: illustration from a Pygmy population. *Sci. Rep.***6**, 27405 (2016).
4. Althabe, G. Changements sociaux chez les Pygmées Baka de l'Est-Cameroun. *Cah. D'Études Afr.***5**, 561–592 (1965).
5. Tsuru, D. Diversity of ritual spirit performances among the Baka Pygmies in southeastern Cameroon. *Afr. Study Monogr. Suppl. Issue* 47–84 (1998).
6. Ramirez Rozzi, F. V. Reproduction in the Baka pygmies and drop in their fertility with the arrival of alcohol. *Proc. Natl. Acad. Sci.***115**, E6126 (2018).
7. Romero, A., Ramirez Rozzi, F. V., Cuesta-Torralvo, E. & Pérez-Pérez, A. Age-related tooth wear in African rainforest hunter-gatherers. *Am. J. Phys. Anthropol.***170**, 622–628 (2019).
8. Yamauchi, T., Sato, H. & Kwamura, K. Nutritional Status, Activity Pattern, and Dietary Intake among the Baka Hunter-Gatherers in the Village Camps in Cameroon. *Afr. Study Monogr.***21**, 67–82 (2000).
9. Hattori, S. Utilization of Marantaceae plants by the Baka hunter-gatherers in southeastern Cameroon. (2006).
10. Ramirez Rozzi, F. V., Koudou, Y., Froment, A., Le Bouc, Y. & Botton, J. Growth pattern from birth to adulthood in African pygmies of known age. *Nat. Commun.***6**, 1–12 (2015).
11. Meazza, C. *et al.* The Enigma of Pygmy short stature. *Endocrinology***1**, (2006).
12. Diamond, J. M. Why are pygmies small? *Nature***354**, 111–112 (1991).

13. Migliano, A. B., Vinicius, L. & Lahr, M. M. Life history trade-offs explain the evolution of human pygmies. *Proc. Natl. Acad. Sci.***104**, 20216–20219 (2007).
14. Perry, G. H. & Dominy, N. J. Evolution of the human pygmy phenotype. *Trends Ecol. Evol.***24**, 218–225 (2009).
15. Becker, N. S. A. *et al.* Indirect evidence for the genetic determination of short stature in African Pygmies. *Am. J. Phys. Anthropol.***145**, 390–401 (2011).
16. Becker, N. S. *et al.* The role of GHR and IGF1 genes in the genetic determination of African pygmies' short stature. *Eur. J. Hum. Genet.***21**, 653–658 (2013).
17. Pemberton, T. J. *et al.* A genome scan for genes underlying adult body size differences between Central African hunter-gatherers and farmers. *Hum. Genet.***137**, 487–509 (2018).
18. Romero, A., Ramirez Rozzi, F. V. & Pérez-Pérez, A. Dental size variability in Central African Pygmy hunter-gatherers and Bantu-speaking farmers. *Am. J. Phys. Anthropol.***166**, 671–681 (2018).
19. Ramirez Rozzi, F. V. & Romero, A. Tooth dimensions and body size in a Pygmy population. *Ann. Hum. Biol.***46**, 467–474 (2019).
20. Shea, B. T. & Gomez, A. M. Tooth scaling and evolutionary dwarfism: an investigation of allometry in human pygmies. *Am. J. Phys. Anthropol.***77**, 117–132 (1988).
21. Weber, G. W. & Bookstein, F. L. Virtual anthropology: a guide to a new interdisciplinary field. (2011).
22. Sardi, M. L. & Ramirez Rozzi, F. V. Developmental connections between cranial components and the emergence of the first permanent molar in humans. *J. Anat.***210**, 406–417 (2007).
23. Molnar, S. Human tooth wear, tooth function and cultural variability. *Am. J. Phys. Anthropol.***34**, 175–189 (1971).
24. Scott, G. R. & Irish, J. D. *Human tooth crown and root morphology*. (Cambridge University Press, 2017).
25. Buchegger, L. Variation of outer and inner crown morphology in upper premolars. (University of Vienna, 2015).
26. Krenn, V. A. *et al.* Variation of 3D outer and inner crown morphology in modern human mandibular premolars. *Am. J. Phys. Anthropol.***169**, 646–663 (2019).
27. Teplanova, D. Geometric morphometric analyses of modern human upper second molars. (University of Vienna, 2015).

28. Halász, V. Morphological assessment of modern human upper and lower first molars. (University of Vienna, 2019).
29. Hanihara, T. 19 Geographic structure of dental variation in the major human populations of the world. *Anthropol. Perspect. Tooth Morphol. Genet. Evol. Var.***66**, 479 (2013).
30. Hanihara, T. & Ishida, H. Metric dental variation of major human populations. *Am. J. Phys. Anthropol. Off. Publ. Am. Assoc. Phys. Anthropol.***128**, 287–298 (2005).
31. Relethford, D. J. H. & Harpending, H. C. Craniometric variation, genetic theory, and modern human origins. *Am. J. Phys. Anthropol.***95**, 249–270 (1994).
32. Relethford, J. H. Apportionment of global human genetic diversity based on craniometrics and skin color. *Am. J. Phys. Anthropol. Off. Publ. Am. Assoc. Phys. Anthropol.***118**, 393–398 (2002).
33. Edgar, H. & Lease, L. Correlations between deciduous and permanent tooth morphology in a European American sample. *Am. J. Phys. Anthropol.***133**, 726–734 (2007).
34. Dahlberg, A. A. Analysis of the American Indian dentition. in *Dental anthropology* 149–177 (Elsevier, 1963).
35. Hanihara, K. Standard models for classification of crown characters in the human deciduous dentition. *Mimeogr. Assoc. Plaster Casts* (1960).
36. Lease, L. R. Ancestral determination of African American and European American deciduous dentition using metric and non-metric analysis. (2003).
37. Saunders, S. R. & Mayhall, J. T. Fluctuating asymmetry of dental morphological traits: new interpretations. *Hum. Biol.* 789–799 (1982).
38. Gómez-Robles, A. *et al.* A geometric morphometric analysis of hominin upper first molar shape. *J. Hum. Evol.***53**, 272–285 (2007).
39. Polychronis, G., Christou, P., Mavragani, M. & Halazonetis, D. J. Geometric morphometric 3D shape analysis and covariation of human mandibular and maxillary first molars. *Am. J. Phys. Anthropol.***152**, 186–196 (2013).
40. Lease, L. R. A descriptive study of African American deciduous dentition. *Dent. Anthropol. J.***26**, 45–55 (2013).
41. Hanihara, T. Morphological variation of major human populations based on nonmetric dental traits. *Am. J. Phys. Anthropol. Off. Publ. Am. Assoc. Phys. Anthropol.***136**, 169–182 (2008).
42. Scott, G. R. Population variation of Carabelli's trait. *Hum. Biol.* 63–78 (1980).

43. Harris, E. F. Carabelli's trait and tooth size of human maxillary first molars. *Am. J. Phys. Anthropol.***132**, 238–246 (2007).
44. Irish, J. D. Characteristic high-and low-frequency dental traits in sub-Saharan African populations. *Am. J. Phys. Anthropol. Off. Publ. Am. Assoc. Phys. Anthropol.***102**, 455–467 (1997).
45. Garn, S. M., Sandusky, S. T., Nagy, J. M. & Trowbridge, F. Negro-Caucasoid differences in permanent tooth emergence at a constant income level. *Arch. Oral Biol.***18**, 609–615 (1973).
46. Hassanali, J. & Odhiambo, J. W. Ages of eruption of the permanent teeth in Kenyan African and Asian children. *Ann. Hum. Biol.***8**, 425–434 (1981).
47. Blankenstein, R., Cleaton-Jones, P., Maistry, P., Luk, K. & Fatti, L. The onset of eruption of permanent teeth amongst South African Indian children. *Ann. Hum. Biol.***17**, 515–521 (1990).
48. Reid, D. J. & Dean, M. C. Variation in modern human enamel formation times. *J. Hum. Evol.***50**, 329–346 (2006).
49. Liversidge, H. Variation in modern human dental development. *Camb. Stud. Biol. Evol. Anthropol.* 73–113 (2003).
50. Lam, C. U. *et al.* Influence of metabolic-linked early life factors on the eruption timing of the first primary tooth. *Clin. Oral Investig.***20**, 1871–1879 (2016).
51. Patin, E. *et al.* Inferring the demographic history of African farmers and Pygmy hunter–gatherers using a multilocus resequencing data set. *PLoS Genet.***5**, (2009).
52. Dean, M. C. & Cole, T. J. Human Life History Evolution Explains Dissociation between the Timing of Tooth Eruption and Peak Rates of Root Growth. *PLOS ONE***8**, e54534 (2013).
53. Verdu, P. *et al.* Sociocultural behavior, sex-biased admixture, and effective population sizes in Central African Pygmies and non-Pygmies. *Mol. Biol. Evol.***30**, 918–937 (2013).
54. Ballabriga, A. Le malheur des Nains. Quelques aspects du combat des grues contre les Pygmées dans la littérature grecque. *Rev. Études Anciennes* 57–74 (1981).
55. Bahuchet, S. L'invention des Pygmées (Inventing Pygmies). *Cah. D'Études Afr.***33**, 153–181 (1993).
56. Ramirez Rozzi, F. V. Les pygmées: histoire d'une rencontre, origines d'une dénomination, évolution des représentations. *Images Mém.***45**, 13–20 (2015).
57. Ramirez Rozzi, F. V. Chez les Pygmées Baka de Cameroun. Voyage dans l'altérité. (L'Harmattan, 2021).
58. Bacirongo, I. & Nest, M. *Still a pygmy*. (Finch Publishing, 2014).

59. Frederick, M. & Gallup, G. Fluctuating dental asymmetry in great apes, fossil hominins, and modern humans: Implications for changing stressors during human evolution. (2007).
60. Kegley, A. & Hemingway, J. Assessing fluctuating odontometric asymmetry among fossil hominin taxa through alternative measures of central tendency: effect of outliers and directional components on reported results. *Homo***58**, 33–52 (2007).
61. Moskona, D., Vainder, M., Hershkovitz, I. & Kobylansky, E. Bilateral asymmetry in dental discrete traits in human isolates: South Sinai Bedouin tribes. *Anthropol. Anz.* 289–306 (1996).
62. Fornai, C. *et al.* The Qesem Cave hominin material (part 2): A morphometric analysis of dm2-QC2 deciduous lower second molar. *State Art Multidiscip. Res. Middle Pleistocene Qesem Cave Isr. 2015***398**, 175–189 (2016).
63. Fornai, C., Bookstein, F. L. & Weber, G. W. Variability of Australopithecus second maxillary molars from Sterkfontein Member 4. *J. Hum. Evol.***85**, 181–192 (2015).
64. Hershkovitz, I. *et al.* The earliest modern humans outside Africa. *Science***359**, 456–459 (2018).
65. Bookstein, F. L. *Morphometric tools for landmark data: geometry and biology*. (Cambridge University Press, 1997).
66. Gunz, P., Mitteroecker, P. & Bookstein, F. L. Semilandmarks in three dimensions. in *Modern morphometrics in physical anthropology* 73–98 (Springer, 2005).
67. Gunz, P. & Mitteroecker, P. Semilandmarks: a method for quantifying curves and surfaces. *Hystrix Ital. J. Mammal.***24**, 103–109 (2013).
68. Gower, J. C. Generalized procrustes analysis. *Psychometrika***40**, 33–51 (1975).
69. Mitteroecker, P., Gunz, P., Windhager, S. & Schaefer, K. A brief review of shape, form, and allometry in geometric morphometrics, with applications to human facial morphology. *Hystrix Ital. J. Mammal.***24**, 59–66 (2013).
70. Bookstein, F. L. Principal warps: Thin-plate splines and the decomposition of deformations. *IEEE Trans. Pattern Anal. Mach. Intell.***11**, 567–585 (1989).
71. Bookstein, F. L. Thin-plate splines and the atlas problem for biomedical images. in 326–342 (Springer, 1991).
72. Scott, G. R. & Turner, C. G. Dental Anthropology. *Annu. Rev. Anthropol.***17**, 99–126 (1988).
73. Rathmann, H. & Reyes-Centeno, H. Testing the utility of dental morphological trait combinations for inferring human neutral genetic variation. *Proc. Natl. Acad. Sci.***117**, 10769–10777 (2020).

# Tables

Due to technical limitations, table 1-5 is only available as a download in the Supplemental Files section.

# Figures






Figure 1

PCA plots for deciduous lower and upper second molar dental crowns. a) PC1 - PC2 plot for ldm2s in shape space for the combined enamel-dentine junction (EDJ) and cervical outline (warpings at values  $\pm 0.15/\pm 0.07$ ). The Baka show a broader range of morphological variation than the Europeans; b) PC1 - PC2 plot for udm2s in shape space for the combined EDJ and cervical outline (warpings at values  $\pm 0.15$ ). The Baka show a less variable morphology, while the Europeans exceed the range of variation.



**Figure 2**

2B-PLS plot capturing the covariation between dental crowns (combined enamel-dentine junction and cervical outlines) of deciduous upper and lower second molars (dm2s; warpings at values  $\pm 0.15$ ). The upper and lower dm2 dyads vary from low-crowned with expanded opposing fossae to high-crowned with reduced opposing fossae. The various populations can be hardly distinguished, although the subsample sizes are too small to allow definite statements. Within the Baka sample the most extreme expression towards the low-crowned variation is observed.



**Figure 3**

Boxplots of the natural logarithm of Centroid Sizes from the dentinal crown (combining enamel-dentine junction and cervical outline) in deciduous a) lower and b) upper second molars. South Americans and the Baka possess the largest deciduous second molars. The smallest specimens are within the Europeans and Egyptians.

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SimkovaetalSupplements.docx](#)
- [Tables.pdf](#)