

RISK FACTORS FOR MORTALITY AND TREATMENT OUTCOMES OF PATIENTS WITH CEFTRIAXONE RESISTANT E. COLI BACTEREMIA RECEIVING CARBAPENEM VERSUS BETA LACTAM/BETA LACTAMASE INHIBITOR THERAPY

Nosheen Nasir (✉ nosheen.nasir@aku.edu)

Aga Khan University <https://orcid.org/0000-0003-1610-8748>

Sara Ahmed

Aga Khan University

Samrah Razi

Aga Khan University

Safia Awan

Aga Khan University

Faisal Mahmood

Aga Khan University

Research note

Keywords: Escherichia coli; Carbapenem; Beta lactam/Beta lactam inhibitor; mortality

Posted Date: August 6th, 2019

DOI: <https://doi.org/10.21203/rs.2.12475/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on September 23rd, 2019. See the published version at <https://doi.org/10.1186/s13104-019-4648-7>.

Abstract

Background: Extended spectrum β -lactamases (ESBL) producing Enterobacteriaceae predominantly *E. coli* and *K. pneumonia* bacteremia have limited treatment options and high mortality. **Objective:** To determine the risk factors for in-hospital mortality particularly treatment with Carbapenem versus Beta lactam/beta lactamase combination (BL/BLI) in patients with Ceftriaxone resistant *E. coli* bacteremia. **Methods:** A retrospective descriptive study was conducted at the Aga Khan University, Karachi, Pakistan. Adult patients with monomicrobial Ceftriaxone resistant *E. coli* bacteremia were enrolled. Factors associated with mortality in patients were determined using logistic regression analysis. **Results:** Mortality rate was 37% in those empirically treated with carbapenem compared to 20% treated with BL/BLI combination therapy (p-value: 0.012) and was 21% in those treated with a carbapenem compared to 13% in patients definitively treated with BL/BLI combination therapy (p-value: 0.152). In multivariable logistic regression analysis, only Pitt bacteremia score of \geq four was significantly associated with mortality (OR: 7.7 CI: 2.6-22.8) while a urinary source of bacteremia was protective (OR: 0.26 CI: 0.11-0.58). **Conclusions:** In-hospital mortality in patients with Ceftriaxone resistant *E. coli* bacteremia did not differ in patients treated with either a carbapenem or BL/BLI combination. However, Pitt bacteremia score of \geq 4 was strongly associated with mortality.

Background

Escherichia coli is a leading cause of intra-abdominal, urinary tract, and bloodstream infections encountered in routine clinical practice (1). It is a recognized pathogen for causing both community acquired and nosocomially acquired infections (1, 2). Over the past two decades, antibiotic-resistant strains that produce extended spectrum β -lactamases (ESBL) have emerged among the Enterobacteriaceae, predominantly in *E. coli* and *Klebsiella pneumoniae* (2-4). ESBLs are a group of hydrolyzing enzymes that provide resistance to third generation cephalosporins and aztreonam, but can in turn be hydrolyzed by clavulanic acid (5). Several epidemiological studies have investigated the prevalence of ESBLs and their clinical impact (2, 3, 6). There is considerable geographic variability in the prevalence of ESBLs with the prevalence being highest in from Latin America and Asia Pacific region followed by Europe and North America (4). In a recently conducted meta-analysis from Pakistan, the pooled prevalence of ESBL producing Enterobacteriaceae was reported to be 40% (7). This is similar to proportions reported from neighbouring countries of China (8) and India (9) and considerably higher than US and developed countries from Europe (5). Hence, Pakistan is a country with high endemicity of community acquired ESBL producing Enterobacteriaceae with major public health implications (10, 11).

Since ESBL-producing organisms are frequently resistant to multiple antimicrobial agents therapeutic options for these infections are severely limited (12). Moreover infections caused by ESBL producing organisms have been associated with high morbidity and mortality (13). Studies have shown high Pitt Bacteremia score and Charlson co-morbidity index to be predictive of death in patients with sepsis secondary to ESBL producing organisms (14). Bacteremia with ESBL producing organisms has been associated with a delay in the initiation of appropriate treatment regimens and leads to an increase in the

hospital length of stay and cost of overall treatment (14, 15). Carbapenems have been considered as drugs of choice for treating severe infections caused by ESBL producing Enterobacteriaceae as they are not affected by ESBLs in vitro and clinical efficacy has been demonstrated in several observational studies as well as clinical trials (5, 16, 17). A large multi-country open label non-inferiority randomized controlled (MERINO) trial which compared piperacillin tazobactam with meropenem as definitive treatment for Ceftriaxone resistant E.coli or K.pneumoniae bloodstream infection found significantly higher long term mortality in patients randomized to piperacillin tazobactam arm (16). However, prolonged use of carbapenems places the community at risk of development of carbapenem resistance due to antibiotic selection pressure (18). Thus it is imperative to find alternatives for treatment. Beta lactam/beta lactam inhibitor combinations (BL/BLI) have been widely studied both as empiric and definitive choice of antibiotics and while some studies have reported no difference in clinical outcomes with use of BL/BLI combination compared to carbapenem (19); pooled data from meta-analysis have recommended their use for only clinically stable patients with urinary tract infections as the source (17).

Existing studies from Pakistan have shown that ESBL producing strains of Enterobacteriaceae show greater than 90 percent susceptibility for BL/BLI and hence may be an effective alternative (10, 11). Carbapenem use has severe cost constraints in a low middle country like Pakistan and is reserved for very severe infections. Despite the high endemicity of ESBL producing organisms, data on risk factors for mortality and the optimal choice of antibiotic therapy is lacking. Our study aims to determine the risk factors for in-hospital mortality in patients with ESBL E.coli bacteremia using Ceftriaxone resistance as a marker of ESBL presence; with emphasis on difference in outcome when patients were treated with a Carbapenem compared to beta lactam/beta lactam inhibitor combination at a tertiary care hospital in Pakistan.

Methods

A retrospective observational study was conducted at the Aga Khan University (AKU) Hospital in Karachi, Pakistan from January 2015 to December 2017. All adult patients with age greater than or equal to 18 years with monomicrobial ceftriaxone resistant E.coli bacteremia and sepsis (based on established criteria) (20) in whom treatment with either a carbapenem or BL/BLI combination had been initiated within 48 hours after the blood culture had been drawn were included in the study. Patients were identified from the hospital information management system.

Data collected from medical records for all patients included age, sex, nosocomial or community-onset acquisition, type and severity of underlying conditions using the Charlson comorbidity index, source of bloodstream infection (BSI) according to clinical and microbiological data, Pitt Bacteremia score, antimicrobial therapy, in-hospital mortality, and length of stay after BSI. Antimicrobial therapy administered before susceptibility results were available was considered empirical and therapy administered afterwards was considered definitive.

Frequency and percentages were used to describe categorical variables and comparison between variables was determined using chi square test or Fischer exact test where appropriate. P-value of <0.05 was considered significant. Mortality rates of patients treated with BL/BLI combination or carbapenems were compared using logistic regression analysis. Data was analyzed using SPSS version 16.0.

This study was conducted after receiving exemption from ethical approval from Aga Khan University Ethical Review Committee (2734-MED-ERC-13).

Results

Out of a total of the 295 patients that were included, 47 % were males while 53% were females. The mean age of the patients was 57.71 years (SD: 17.16): Of the patients who died, the mean age was 55.3 years (SD: 17.9) years compared to 60.1 (SD: 16.5) years for those who survived (p-value: 0.046). The overall in-hospital mortality in patients with Ceftriaxone resistant E.coli bacteremia was 21%. The mean length of stay was 11.3 days (SD: 12.5) in those who died compared to 6.8 days (SD: 5.68) in those who survived (p-value <0.001) (Table 1).

In the univariate analysis to determine association between choice of empirical treatment and mortality; out of 60 patients who died, n= 20 had received a carbapenem and n=31 had received a BL/BLI combination therapy. Mortality rate was 37% in those treated with carbapenem compared to 20% in patients empirically treated with BL/BLI combination therapy (p-value: 0.012). For definitive treatment; n= 36 received a carbapenem whereas n=21 received BL/BLI combination therapy among those who died. Mortality rate was 21% in those treated with a carbapenem compared to 13% in patients definitively treated with BL/BLI combination therapy (p-value: 0.152). Other variables found to be significantly associated with mortality included increasing mean Charlson comorbidity index and increasing Pitt Bacteremia score (Table 1).

However, in multivariable logistic regression analysis; a Pitt bacteremia score of greater than or equal to 4 was significantly associated with mortality (OR:16.8 CI: 7.47-38.0) while presence of urinary tract infection being the source of bacteremia was protective (OR: 0.33(CI: 0.15-0.73)) after adjusting for confounding from Charlson comorbidity index of greater than or equal to 3 (Table 2). Neither the empirical nor definitive antibiotic therapy with carbapenem versus BL/BLI combination therapy was significantly associated with in-hospital mortality.

Discussion

Empiric and definitive choice of antimicrobial therapy for ESBL producing Enterobacteriaceae has been subject of debate since a long time because of the implications of using carbapenem for ESBL infections which while effective has the potential for promoting carbapenem resistance (18). Our study did not show an association of mortality with use of either a carbapenem or a BL/BLI combination treatment in Ceftriaxone resistant E.coli bacteremia whether it was used empirically or definitively. These

findings are similar to a post hoc analysis of prospective cohort conducted by Rodriguez-Bano, J. et al where neither carbapenem nor BL/BLI combination influenced mortality or hospital length of stay (19). However, the recently conducted multi-country open label non-inferiority randomized controlled (MERINO) trial which compared piperacillin tazobactam with meropenem as definitive treatment for Ceftriaxone resistant E.coli or K.pneumoniae bloodstream infection found significantly higher long term mortality in patients randomized to piperacillin tazobactam arm and concluded that piperacillin tazobactam was not non-inferior (risk difference: 8.6%; P = .90). The authors therefore recommended avoiding use of this combination for Ceftriaxone resistant bloodstream infections (16). However, the trial had several limitations owing to its pragmatic design with crossover in both arms and that the cause of 30-day mortality was unrelated to infection. Despite the criticism, the results were quite robust for the conclusion (21). Given the conclusion of this trial, the concern for increased carbapenem resistance with increasing use has left many experts to consider piperacillin tazobactam for less severe infections with ESBL producing enterobacteriaceae (21). As for treatment comparison between both antibiotics as empirical therapy, there has been a systematic review and meta-analysis of non-randomized studies which concluded that carbapenems should be given preference as drug of choice in patients with ESBL-producing Enterobacteriaceae bloodstream infections (17). However, there was substantial clinical and statistical heterogeneity in the studies included in this review and a more recent review conducted by Sfeir M.M. et al in 2018 have not found any difference in 30-day mortality with use of either carbapenem or BL/BLI combination (22). Comparison studies for appropriate choice of antibiotic for ESBL producing Enterobacteriaceae bacteremia are lacking from South Asian region which differs from the west in terms of geographical and economic diversity. A resource poor country like ours where carbapenem use has severe cost implications and is considered to be one of the last resort antibiotics, it is imperative to determine which antibiotic is better and we recommend future randomized controlled trials in this region based on our results.

Several predictors of mortality; apart from choice of antibiotics; have been identified in various studies for ESBL Enterobacteriaceae bloodstream infections. These include high Charlson comorbidity index, high Pitt bacteremia scores and source of bloodstream infection other than urinary tract (13, 14, 23). In our study also we found significant association of Pitt bacteremia score of greater than 4 to be significantly associated with in-hospital mortality in patients with Ceftriaxone resistant E.coli Bacteremia. Moreover, similar to findings of other studies, presence of urinary tract infection was favourable.

Limitations:

There are several limitations of our study. Firstly, we have used Ceftriaxone resistance as a marker of ESBL presence and we do not have genotypic and MIC data for our isolates. However, the study is intended to guide in real life scenarios encountered in clinical practice in our setting where molecular studies are not widely available. Moreover, large pragmatic trial has also utilized ceftriaxone resistance as surrogate marker for better feasibility (16). Secondly, our study is a single center study and hence the generalizability may be limited to our region.

Conclusion

In conclusion, our study shows that there was no difference in in-hospital mortality in patients with Ceftriaxone resistant E. coli bacteremia in patients treated with either a carbapenem or BL/BLI combination whether empirically or definitively. However, Pitt bacteremia score of greater than or equal to 4 was strongly associated with mortality while urinary tract as site of infection was protective. We recommend randomized controlled trials for further delineating choice of appropriate antibiotic in our resource poor region.

List Of Abbreviations

ESBL: Extended spectrum β -lactamases

BL/BLI: Beta lactam/beta lactamase combination

BSI: bloodstream infection

ERC: Ethical review Committee

Declarations

Ethics approval and consent to participate: Ethical exemption was obtained from the hospital's ethics review committee (2734-MED-ERC-13).

Consent for publication: Not applicable

Availability of data and material: All data generated or analysed during this study are included in this published article.

Competing interests: No conflict of interest

Funding: None

Authors' contributions:

Nosheen Nasir: Conceived idea, supervised data collection, major contributor of manuscript

Sara Ahmed: Data collection and contribution in writing manuscript

Samrah Razi: Data collection and contributed to manuscript

Safia Awan: Data Analysis and Reviewed manuscript

Faisal Mahmood: Conceived idea, reviewed manuscript, provided expert opinion and supervised the study

All authors discussed the results and contributed to the final manuscript

References

1. Allocati N, Masulli M, Alexeyev MF, Di Ilio C. Escherichia coli in Europe: an overview. *Int J Environ Res Public Health*. 2013;10(12):6235-54.
2. Oteo J, Perez-Vazquez M, Campos J. Extended-spectrum [beta]-lactamase producing Escherichia coli: changing epidemiology and clinical impact. *Current opinion in infectious diseases*. 2010;23(4):320-6.
3. Rawat D, Nair D. Extended-spectrum beta-lactamases in Gram Negative Bacteria. *J Glob Infect Dis*. 2010;2(3):263-74.
4. Falagas ME, Karageorgopoulos DE. Extended-spectrum beta-lactamase-producing organisms. *The Journal of hospital infection*. 2009;73(4):345-54.
5. Paterson DL, Bonomo RA. Extended-spectrum beta-lactamases: a clinical update. *Clin Microbiol Rev*. 2005;18(4):657-86.
6. Pitout JD, Laupland KB. Extended-spectrum beta-lactamase-producing Enterobacteriaceae: an emerging public-health concern. *The Lancet Infectious diseases*. 2008;8(3):159-66.
7. Abrar S, Hussain S, Khan RA, Ul Ain N, Haider H, Riaz S. Prevalence of extended-spectrum-beta-lactamase-producing Enterobacteriaceae: first systematic meta-analysis report from Pakistan. *Antimicrobial resistance and infection control*. 2018;7:26.
8. Zhang J, Zheng B, Zhao L, Wei Z, Ji J, Li L, et al. Nationwide high prevalence of CTX-M and an increase of CTX-M-55 in Escherichia coli isolated from patients with community-onset infections in Chinese county hospitals. *BMC Infect Dis*. 2014;14:659.
9. Ravikant, Kumar P, Ranotkar S, Zutshi S, Lahkar M, Phukan C, et al. Prevalence and identification of extended spectrum beta-lactamases (ESBL) in Escherichia coli isolated from a tertiary care hospital in North-East India. *Indian J Exp Biol*. 2016;54(2):108-14.
10. Fatima S, Muhammad IN, Khan MN, Jamil S. Phenotypic expression and prevalence of multi drug resistant extended spectrum beta-lactamase producing Escherichia coli and Klebsiella pneumoniae in Karachi, Pakistan. *Pakistan journal of pharmaceutical sciences*. 2018;31(4):1379-84.
11. Jabeen K, Zafar A, Hasan R. Frequency and sensitivity pattern of Extended Spectrum beta Lactamase producing isolates in a tertiary care hospital laboratory of Pakistan. *J Pak Med Assoc*. 2005;55(10):436-9.
12. Chopra T, Marchaim D, Johnson PC, Chalana IK, Tamam Z, Mohammed M, et al. Risk factors for bloodstream infection caused by extended-spectrum beta-lactamase-producing Escherichia coli and Klebsiella pneumoniae: A focus on antimicrobials including cefepime. *American journal of infection control*. 2015;43(7):719-23.
13. Kang CI, Kim SH, Park WB, Lee KD, Kim HB, Kim EC, et al. Bloodstream infections due to extended-spectrum beta-lactamase-producing Escherichia coli and Klebsiella pneumoniae: risk factors for

- mortality and treatment outcome, with special emphasis on antimicrobial therapy. *Antimicrobial agents and chemotherapy*. 2004;48(12):4574-81.
14. Russo A, Falcone M, Gutierrez-Gutierrez B, Calbo E, Almirante B, Viale PL, et al. Predictors of outcome in patients with severe sepsis or septic shock due to extended-spectrum beta-lactamase-producing Enterobacteriaceae. *International journal of antimicrobial agents*. 2018;52(5):577-85.
 15. Battle SE, Bookstaver PB, Justo JA, Kohn J, Albrecht H, Al-Hasan MN. Association between inappropriate empirical antimicrobial therapy and hospital length of stay in Gram-negative bloodstream infections: stratification by prognosis. *The Journal of antimicrobial chemotherapy*. 2017;72(1):299-304.
 16. Harris PNA, Tambyah PA, Lye DC, Mo Y, Lee TH, Yilmaz M, et al. Effect of Piperacillin-Tazobactam vs Meropenem on 30-Day Mortality for Patients With E coli or Klebsiella pneumoniae Bloodstream Infection and Ceftriaxone Resistance: A Randomized Clinical Trial. *JAMA*. 2018;320(10):984-94.
 17. Vardakas KZ, Tansarli GS, Rafailidis PI, Falagas ME. Carbapenems versus alternative antibiotics for the treatment of bacteraemia due to Enterobacteriaceae producing extended-spectrum beta-lactamases: a systematic review and meta-analysis. *The Journal of antimicrobial chemotherapy*. 2012;67(12):2793-803.
 18. McLaughlin M, Advincula MR, Malczynski M, Qi C, Bolon M, Scheetz MH. Correlations of antibiotic use and carbapenem resistance in enterobacteriaceae. *Antimicrobial agents and chemotherapy*. 2013;57(10):5131-3.
 19. Rodriguez-Bano J, Navarro MD, Retamar P, Picon E, Pascual A, Extended-Spectrum Beta-Lactamases-Red Espanola de Investigacion en Patologia Infecciosa/Grupo de Estudio de Infeccion Hospitalaria G. beta-Lactam/beta-lactam inhibitor combinations for the treatment of bacteremia due to extended-spectrum beta-lactamase-producing Escherichia coli: a post hoc analysis of prospective cohorts. *Clin Infect Dis*. 2012;54(2):167-74.
 20. Levy MM, Evans LE, Rhodes A. The Surviving Sepsis Campaign Bundle: 2018 Update. *Crit Care Med*. 2018;46(6):997-1000.
 21. Sfeir MM. Post-MERINO trial: Any role for piperacillin-tazobactam in treating bloodstream infections caused by extended-spectrum beta-lactamase producing Enterobacteriaceae? *International journal of antimicrobial agents*. 2019;53(5):557-8.
 22. Sfeir MM, Askin G, Christos P. Beta-lactam/beta-lactamase inhibitors versus carbapenem for bloodstream infections due to extended-spectrum beta-lactamase-producing Enterobacteriaceae: systematic review and meta-analysis. *International journal of antimicrobial agents*. 2018;52(5):554-70.
 23. Al-Hasan MN, Lahr BD, Eckel-Passow JE, Baddour LM. Predictive scoring model of mortality in Gram-negative bloodstream infection. *Clinical microbiology and infection : the official publication of the European Society of Clinical Microbiology and Infectious Diseases*. 2013;19(10):948-54.

Tables

Table 1: Comparison of demographics and clinical features of patients with Ceftriaxone resistant E.coli Bacteremia who died compared to those who survived

Patient Characteristics	Died (n=62)	Recovered (n=235)	p-value
Age (Mean \pm SD)	55.28 \pm 17.87	60.14 \pm 16.46	0.046
Gender			
Male	34	106	0.172
Female	28	129	
Charlson Comorbidity Index (CCI) Mean (SD)	2.98 (2.12)	1.93(1.73)	<0.001
Pitt Bacteremia Score Median (IQR)	3(4)	1(2)	<0.001
Site of Infection			
Intra-abdominal	24	44	<0.001
UTI	24	171	
CLABSI	4	4	
Gut translocation	4	5	
Source unclear	6	11	
Empirical Antibiotics*			0.012
Carbapenem	20	33	
BL/BLI	31	121	
Definitive Antibiotics¶			
Carbapenem	36	138	0.152
BL/BLI	12	77	
Length of stay (Mean \pmSD)	11.27 \pm 12.46	6.82 \pm 5.68	<0.001

*Empirical group includes only those patients who had been given either a carbapenem or BL/BLI combination treatment empirically. ¶ Definitive group includes only those patients who had been given either a carbapenem or BL/BLI combination treatment as definitive treatment

Table 2: Multivariable analysis showing factors associated with in-hospital mortality in patients with Ceftriaxone resistant E.coli Bacteremia

Variables	aOdd ratio(95% CI)	<i>p</i> -value
Pitt Bacteremia Score		
<4 (Ref)	1.0	
≥4	16.85(7.47-38.0)	<0.001
Source of Infection		
Intra-abdominal (Ref)	1.0	0.002
Urinary tract Infection	0.33(0.15-0.73)	0.006
CLABSI	3.03(0.62-14.66)	0.16
Gut translocation	2.97(0.50-17.43)	0.22
Source unclear	1.07(0.28-4.03)	0.91
Charlson Comorbidity Index score		
<3 (Ref)	1.0	
≥3	1.98(0.98-3.97)	0.055