

A multicentre parallel-group randomised trial assessing multiparametric MRI characterisation and image-guided biopsy of prostate in men suspected of having prostate cancer: MULTIPROS study protocol

Magdalena Szewczyk-Bieda

Ninewells Hospital

Cheng Wei

University of Dundee <https://orcid.org/0000-0001-5714-4863>

Katherine Coll

University of Dundee

Stephen Gandy

Ninewells Hospital

Peter Donnan

University of Dundee

Senthil Kumar Arcot Ragupathy

Aberdeen Royal Infirmary

Paras Singh

Royal Free Hospital

Jennifer Wilson

Ninewells Hospital

Ghulam Nabi (✉ g.nabi@dundee.ac.uk)

Study protocol

Keywords: prostate cancer, multiparametric magnetic resonance imaging, prostate biopsy

Posted Date: August 7th, 2019

DOI: <https://doi.org/10.21203/rs.2.12482/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on November 21st, 2019. See the published version at <https://doi.org/10.1186/s13063-019-3746-0>.

Abstract

Background There is growing evidence suggesting that multiparametric magnetic resonance imaging (mpMRI) is a marker for prostate cancer (PCa) aggressiveness and could be used to plan treatment. Improving early detection of clinically significant PCa with pre-biopsy mpMRI would very likely have advantages including optimising the diagnosis and treatment of diseases and diminishing patient anxiety. **Methods and materials** This is a prospective multicentre study of pre-biopsy mpMRI diagnostic test accuracy with subgroup randomisation at a 1:1 ratio with respect to transrectal ultrasound (TRUS) and MRI/US fusion-guided biopsy or TRUS-only biopsy. It is designed as a single-gate study with a single set of inclusion criteria. The total duration of the recruitment phase was 48 months; however, this has now been extended to 66 months. A sample size of 600 participants is required. **Discussion** The primary objective is to determine whether mpMRI can improve PCa detection and characterisation. The key secondary objective is to determine whether MRI/US fusion-guided biopsy can reduce the number of false-negative biopsies. Ethical approval was obtained from the East of Scotland Research Ethics Committee 1 (14/ES/1070) on 20 November 2014. The results of this study will be used for publication and presentation in national and international journals and scientific conferences.

Introduction

1.1 Introduction and rationale

Men suspected of having prostate cancer (PCa) because of high levels of serum prostate-specific antigen (PSA) and/or abnormal digital rectal examination (DRE) are offered prostate biopsies guided by transrectal ultrasound (TRUS). Several challenges are seen in this crucial step. Firstly, the incidence of false-negative biopsies, i.e. clinically significant tumours not being detected, may be as high as 35% in first-time biopsies¹; secondly, up to 44% of men younger than 70 years of age may need a second set of biopsies following initial negative results²; thirdly, up to 50% of PCa cases currently detected by these methods may not be clinically relevant when considering other patient-related factors; fourthly, data from UK Prostate Testing for Cancer and Treatment (ProtecT) has shown significant morbidity of the biopsy procedure, at times even leading to patient death³; finally, several studies have shown no improvement in PCa mortality, despite increased detection rates through screening programmes⁴, and the number of deaths from PCa in the UK sadly remains largely unchanged.

Post-biopsy standard of care magnetic resonance imaging (MRI) is offered to patients with positive biopsy results or when there is a continuous increase in PSA levels despite negative biopsies. The performance of post-biopsy MRI in the staging and detection of PCa is significantly degraded by post-biopsy artefact^{5, 6}. Therefore, it is recommended that MRI should be performed at least 10 weeks after biopsy, and if possible after 20 weeks⁷. Such a wait is unacceptable to patients and causes breaching of time with respect to treatment guidelines. Improving early detection of clinically significant PCa would very likely save many lives by refining the stratification of patients for optimising the treatment of

significant disease, whilst simultaneously diminishing patient anxiety and morbidity through overdiagnosis and overtreatment.

1.2 objectives

The primary objective is to determine whether using multiparametric MRI (mpMRI) can improve cancer detection and the characterisation of PCa, and primary outcome measurements include:

- Number of PCa detected by mpMRI when compared to gold-standard prostatectomy specimen.
- Number of clinically significant cancers detected by mpMRI when compared to gold-standard prostatectomy specimen.

The secondary objective is to assess whether MRI/US fusion-guided biopsy can reduce the number of false-negative biopsies, and key outcome measurements include:

- Number of cancers detected in each randomised group, namely the intervention group (TRUS and MRI/US fusion-guided biopsy) vs. the standard of care group (TRUS only biopsy).
- Number of clinically significant cancers detected in each randomised group.

1.3 Trial design

This is a prospective multicentre study of pre-biopsy mpMRI diagnostic test accuracy with subgroup randomisation at a 1:1 ratio with respect to TRUS and MRI/US fusion-guided biopsy or TRUS-only biopsy. This is the first large trial to randomise men into either targeted or non-targeted biopsy with pre-biopsy MRI and choose radical prostatectomy histopathology using 3D fabricated molds as gold standard (Figure 1).

Methods And Design

2.1 Study population and setting

Prospective recruitment of men referred with suspected clinically localised PCa will be carried out in urology clinics. The number of patients screened for eligibility over the recruitment period is estimated to be 1000. The number of patients not eligible for the study is estimated to be 350 and the number of patients who are likely to decline is estimated to be 50. The number of participants who are likely to progress to pre-biopsy MRI is estimated to be 600. Local audit information suggests that after a positive diagnosis of PCa, around 25% of men will opt for radical surgery. Based on the recruitment of 600 eligible men, the lowest amount of complete data for radical prostatectomy histology for analysis will be n~150. Our power calculation based on 80% sensitivity and a precision of +/- 9% and an AUROC = 0.9 suggests

that we need a minimum of 80 men with complete datasets from imaging and histopathology of radical prostatectomy to answer the primary objective. The dropout rate and incomplete datasets have been taken into consideration in designing this study.

Rais-Bahrami et al.⁸ suggest that 80% of all pre-biopsy MRI (n~600) will have lesions suspicious of PCa, regardless of the level of suspicion assigned by mpMRI (n~480). Based on such information, we calculated that each of the two randomisation arms will have approximately (n~240) patients.

Initially, the total duration of the recruitment phase was 48 months; however, this will now be extended to 66 months with approval from the funding body. It is estimated that approximately 600 participants will be recruited and will receive a pre-biopsy MRI scan.

Participants will then be divided into two groups based on the mpMRI results. The group with positive mpMRI (n~480) will be randomised to the intervention (TRUS and MRI/US fusion-guided biopsy) group (n~240) or the standard care (TRUS-guided biopsy) group (n~240). The group with negative mpMRI (n~120) will undergo standard care (TRUS-guided biopsy) with histological comparative analysis. Participants will be followed through to a definite treatment stage. At sites where local facilities allow and for those who will undergo radical prostatectomy, histological findings will be compared to the pre-biopsy MRI (n~80). The active trial participation for each participant will be varied depending on sites' local pathway.

Initially, the total duration of the recruitment phase was 48 months; however, this will now be extended to 66 months with approval from the funding body. It is estimated that approximately 600 participants will be recruited and will receive a pre-biopsy MRI scan.

2.2 Participant selection and enrolment

As standard care pathways differ among sites, the route of identification and screening of eligibility will differ in order to accommodate the local pathway. Potential participants will be identified by either the clinical team or the study team, with Caldicott Guardian Approval, depending on local practice.

2.2.1 Identifying potential participants

Potential participants may be identified through the following routes:

- At the urology clinic
- From GP referrals to the urology team
- By the urology team after standard care mpMRI (positive results only)

2.2.2 Contacting potential participants

Contact will be made by either the clinical team or the study team through the following routes:

- In person at clinics

The clinical team will assess whether the patient meets the eligibility criteria for the study. If so, the study patient information sheet (PIS) will be given and a contact telephone number will be requested for a follow-up phone call. A model PIS is shown in supplementary appendix 1.

- Via letter

The clinical urological team will review GP referral letters or positive mpMRI results and identify patients who meet trial eligibility criteria. Those who met the eligibility criteria will be sent a clinic appointment with the trial information leaflet.

- Via telephone call

The clinical urological team or delegate may contact the patient and check whether he has received his NHS clinic appointment before sending out the information pack. After confirmation of receipt, the urological team or delegate (with appropriate Caldicott Guardian Approval) will introduce the study to the patient and if they indicate interest, then an invitation pack will be sent.

If the patient declines the information pack or declines to provide a telephone number, no further attempt to recruit the patient will be made. The clinical or trial team will telephone patients who have agreed to be contacted. Any questions will be answered, and the patient will be asked whether he is interested in participating. The research team will arrange to see the patient for consent discussion. If required, the patient will be given additional time to consider participation before a final decision is made to arrange an appointment.

Patients will be advised to attend all their planned scheduled appointments in the invitation letter and study introduction call. Prospective participants will be encouraged to telephone if they have any questions that they would like answered prior to attendance or to inform the research team if they change their minds with regard to participating. Patients who do not wish to participate will follow their local standard of care pathway.

2.2.3 Inclusion criteria

- Males aged 40–75 years at referral
- Have an MRI scan in the past: for sites with pre-biopsy MRI included in standard of care
- With at least 10 years' life expectancy at clinical referral
- With clinically localised PCa: PSA \leq 20ng/ml
- **And/or** abnormal DRE but < T3 disease
- Ability to give informed consent

2.2.4 Exclusion criteria

- Unable to give informed consent
- Prior prostatic biopsy within 12 months
- Contraindications to biopsy
- Poor general health and life expectancy < 10 years
- Previous diagnosis of acute prostatitis within 12 months
- History of PCa
- Prior transurethral prostatectomy
- Contraindications to MRI including: cardiac pacemakers, allergic reaction to gadolinium-based contrast, renal failure, intracranial clips, claustrophobia (applicable to participants who do not receive MRI as part of standard care pathway)
- Previous hip replacement (applicable to participants who do not receive MRI as part of standard care pathway)

2.2.5 Ineligible and non-recruited participants

Ineligible patients or those who decide not to enter the study will receive standard of care. The reason(s) for ineligibility will be explained to all individuals who express interest in the study. Any questions that they have will be answered. A record of the number of patients screened will be kept that indicates the number of patients eligible/ineligible to participate.

2.2.6 Blinding

In this study, the blinding of the biopsy performer on men randomised to MR/US fusion-guided biopsy is not feasible, but radiologists and pathologists will be double blinded regarding to the patients' outcomes until the end of the statistic analyses.

2.3 Allocations and Interventions

All enrolled participants will be offered 1.5 Tesla or 3.0 Tesla mpMRI prior to prostate biopsy where this is not already within the standard care pathway. A subgroup of participants with positive pre-biopsy MRI will be randomised into two further subgroups at a 1:1 ratio. Randomisation will be via TRUST, a web-based, good clinic practice (GCP) - compliant randomisation system run by the HIC. Randomisation will be implemented with random block sizes stratified by site and minimised by: The Prostate Imaging Reporting and Data System (PI-RADS) suspicion score (3, 4 or 5); index lesion size (in two categories: above and below 6mm in maximal diameter on MRI); age (40–59, 60–75); PSA <10.1 or ≥10.1 to ≤20ng/ml). The first will undergo an TRUS and MRI/US fusion-guided biopsy and the second will

undergo standard of care TRUS-guide biopsies only. The MRI/US fusion-guided biopsy will be performed during the same attendance as the TRUS-guided biopsy. TRUS and MRI/US fusion-guided biopsies will be performed by NHS-accredited radiologists and/or urologists from the research team. The persons undertaking the biopsy will have prior experience and will have received appropriate training from clinical experts and/or the equipment supplier's application specialist(s). Those participants with negative pre-biopsy MRI will follow the local standard of care pathway; where this includes biopsy, data from the biopsy will be collected for analysis.

In a subgroup of patients who undergo radical prostatectomy the prostatectomy specimen will be sectioned using customised molds^{9, 10} allowing for direct comparison with mpMRI. This will be carried out in NHS Tayside, NHS Grampian and other sites where local facilities allow. Comparison of mpMRI with a reference standard of radical prostatectomy pathology will be performed as the primary outcome.

2.4 Withdrawal procedures

It will be made clear to patients that they can withdraw from the study at any time and return to standard of care. however, any data collected up to the time of withdrawal will be included if participants agree to it.

2.5 Proposed outcome and statistical analyses

2.5.1 Prediction Performance (Objective 1)

The diagnostic accuracy of each of the three mpMRI predictive variables (T2WI [five-point ordinal scale], DWI [five-point ordinal scale] and DCE-MRI [five-point ordinal scale]) will be evaluated through logistic regression with PCa as the binary outcome (Yes/No). The T2WI, DWI and DCE-MRI will be dichotomised by assigning 0 to scores 1–2, as well as 1 to scores 3–5¹¹, and presented as percentages and denominators in tables. The predictive performance of the dichotomised variables for the diagnosis of PCa will be analysed both individually and by considering all possible combinations among them. Additional variables as potential predictors in the logistic regression model will also include age, stage and comorbidity. Predictive ability will be estimated from the area under receiver operating characteristic (AUROC) curves drawn from the estimated probabilities given by the corresponding logistic regression models. Accuracy will be assessed by implementing the Hosmer–Lemeshow test for calibration. A score algorithm will be derived from the final logistic regression model.

2.5.2 Randomised Controlled Trial (Objective 2)

Analysis of the randomised comparison between MRI/US fusion-guided biopsy and standard TRUS-guided biopsy will be implemented according to the ICH E9 'Statistical Principles in Clinical Trials'.

Analysis will be based on the intention-to-treat principle. The key outcome for the randomised study is to measure whether the MRI/US fusion-guided biopsy can improve the detection of cancer foci and to establish the reliability of the pre-biopsy MRI for the localisation of cancer foci within the prostate gland.

Secondary outcomes will include:

- PCa detected on fusion biopsy which are missed on TRUS biopsy.
- Safety outcomes of death, and side effects such as pain and bleeding with the severity and duration of symptoms recorded in each randomised group.
- Comparison of MRI negative standard care TRUS-guided biopsies with MRI positive TRUS histopathology to facilitate analysis of the diagnostic accuracy of MRI in men suspected of a target condition.

Binary outcomes such as cancer (Yes/No) and pain (Yes/No) will be analysed using logistic regression with the intervention arm as a binary variable in the model. All analyses will be stratified by site and minimisation variables and adjusted for prostatic volume and maximum lesion size. Outcomes such as Gleason Score (GS) and the number of lesions (depending on the range of values obtained) will be analysed as continuous linear regression or ordinal logistic regression. Subgroup analyses will be carried out as secondary analyses by adding treatment by subgroup variable interaction terms in the regression model. Analyses will be conducted by the study statisticians utilising SAS (9.4).

2.6 Data collection and management

The case report form (CRF) will be developed together with the trial management team, statistician and data manager to ensure that the data management system supports the research aims of the study. The data management system will be fully validated, including the provision of test data and supporting documentation. It is the CI's responsibility to ensure the accuracy of all data entered and recorded in the CRF/Electronic Data Capture System (EDCS). Data will be stored on servers controlled by the HIC and housed at the University of Dundee. Back-up and disaster recovery will be provided by the HIC according to its standard operating procedures. The data in the CRF will be stored by the MULTIPROS study group for 5 years after the end of study. Future information on the EDCS is available upon request to the HIC.

2.7 Trial management, monitoring and auditing

2.7.1 Study management

Tayside Clinical Trial Unit, a UKCRC-registered trials unit will be responsible for overseeing management of the study. A dedicated trial specific manager will run the day-to-day work of the study and report to the CI. Specifically, the trial manager in consultation and close working with the CI, will ensure completeness

and consistency of CRFs. The study management team will discuss any query and resolve it with the input from the CI. The CI may choose to delegate some of the tasks to a member. A details trial-specific Delegation Log will be made available at each recruiting site, with clear mention of the tasks each member of staff will be taking part in. The study management group consists of the Chief and co-investigators and representative of each participating sites. The TMG meets monthly to keep a track of progress with recruitment and resolve challenges as and when they arise.

2.7.2 Joint data monitoring committee and trial steering group charter

A Trial Steering Committee (TSC) with an integrated Data Monitoring Committee (DMC) will be put in place. The combined group will perform a patient safety review and provide overall strategic direction and supervision during the progress of the study. The DMC is integrated because the foreseen risks profile of the study is minimal. Membership will consist of a lay-member of patients group as Chair and a minimum of two other experts in area of trial management or treatment of prostate cancer. The CI will participate as invited by the TSC/DMC. The independent committee members will ensure research governance, review data and suggest alteration to study based on a risk-benefit analyses of the data. The later may include early termination of recruitment and other modifications as desired. The terms of reference will be agreed upon to clarify the membership, roles and responsibilities of the joint committee. All the processes and procedures will be documented in the MULTIPROS Joint Committee Charter.

2.7.3 Inspection of records

The CI, PIs and all the participating sites to the recruitment in the study will undertake a commitment to allow free study-specific monitoring, periodic audits and a Research Ethics Committee (REC) review. It is expected that the CI will sign an agreement to allow direct access to all the documentations and study-related records and material to study-sponsor or representative of the sponsor whenever needed.

2.7.4 End of study

The end of study for a participant is a date at which biopsy procedure was performed plus 30 days. The definition of the end of the study date for this trial is database lock. The Sponsor, CI and/or the management group reserve the right to close the study to recruitment for any reasons including safety of the participants, less than satisfactory infra-structure or lack of progress. If the decision is taken to terminate the study prematurely, the decision of end of the study will be communicated to the Sponsor and REC within 90 days, or 15 days of the decision. The CI will be responsible for any further follow-up of the participants and their safety. The Sponsor and REC will be informed and a summary report of the study will be submitted to both within 1 year of the decision to end the study.

2.8 Harms

As the study does not employ an Investigational Medicinal Product, Adverse Events (AEs) and Serious Adverse Events (SAEs) will be recorded and reported as per Health Research Authority (HRA).

An SAE is defined as an untoward occurrence that:

- results in death;
- is life-threatening;
- requires hospitalisation or prolongation of existing hospitalisation;
- results in persistent or significant disability or incapacity;
- consists of a congenital anomaly or birth defect; or
- is otherwise considered medically significant by the investigator.

An SAE occurring to a research participant will be reported to the main REC and Sponsor where in the opinion of the CI the event is: Related – it results from administration of any of the research procedures and Unexpected – the type of event is not listed in the protocol as an expected occurrence.

Due to the clinical status of the study population we will record in the CRF but not report to REC or Sponsor, Serious Adverse Events (SAEs) in the following categories:

- Any death or hospitalisation due to new diagnosis or treatment of a cancer
- Any death or hospitalisation due to exacerbation of an existing medical condition
- SAEs due to expected side effects of biopsy as defined in the BAUS Transrectal Ultrasound-Guided Biopsies of the Prostate Gland Leaflet

AEs will only be recorded in relation to study procedures namely: MRI, biopsy procedure and post biopsy events. Participants will either be called or approached at the time of their routine clinic visit approximately 1-2 weeks following biopsy procedure, to record biopsy related AEs. It is anticipated that pain and bleeding will occur as a result of biopsy, but they will only be recorded as an AE if they continue for more than 4 days post biopsy.

2.9 Patient and public involvement

The research question was discussed in Tayside Urological Cancers Network (TUCAN) meeting with input from patients and public representation. The main attraction of proposal for patients and public was reducing the number of biopsies, improving detection of clinically significant cancers and avoiding having biopsies in men not needed. Patients helped in the preparation of lay-man summary during the grant application. They contributed to the design of information and consent forms. We have advertised study locally in the meetings with patients and public such as Dundee Prostate Cancer Group and Perth

and Kinross prostate cancer and prostate diseases groups. The results will be summarised in a layman summary (briefing document) and widely publicised in local and national patients' groups and prostate cancer charities. The advantages of intervention were discussed in patients and public engagement meetings.

Discussion

3 Discussion

3.1 Study conduct responsibilities

3.1.1 Protocol amendments, deviations and breaches

It is expected that the study protocol and other documentations may need further amendments during the life-time of the proposed research. The CI will have a responsibility to inform and seek approval for any such proposed alterations to the protocol from the Sponsor, REC and NHS R&D Office(s). The changes as thought to be necessary in the study protocol will not be executed without prior approval in place. The current protocol version is V10.0 issued on 24/01/2019.

It will be responsibility of the CI to ensure documentation of any deviation from the protocol that may take place during the conduct of the trial. In case there is any deviation necessary, a clear reason and nature of this should be documented and communicated to the Sponsor. In case deviation merits an amendment in the study protocol, documentations need further submission to the Sponsor for approval and further appropriate action by REC AND NHS R&D office to allow necessary reviews and approval.

Any serious breach of GCP during the study, if suspected, will be immediately reported to the Sponsor using the standard- form "Notification to Sponsor of Serious Breach or Serious Deviation".

3.1.2 Study record retention

Most of the source data within the study will form part of the patient's clinical records. Data generated specifically for the study but of potential clinical relevance will be stored in the patient records and is thus subject to NHS Scotland minimum retention periods. In accordance with this, pathology and genetic data and samples will be kept for a minimum period of 30 years.

The research database which will bring together all relevant data, and which will not contain personal identifiers, will be kept for a period of 5 years after the end of the study. Records will then be deleted unless a review at the time suggests that a longer retention period is indicated. Access and security arrangements for the database will be the same as those during the study.

3.2 Consenting participants

Timing of consent will differ depending on the local standard care pathway; however, at all sites, consent will be obtained prior to any study-related procedures commencing. Patient consent will be discussed with a suitably qualified member of the research team. Consent will only be taken if patients express a wish to participate. Consent will only be sought once a full discussion has taken place and the patient has expressed satisfaction with the information received. The person discussing the study with the patient will normally be the research nurse from the study team. A model consent form is shown in supplementary appendix 2.

3.3 Data access and confidentiality

The study team and the CI of the proposed trial will ensure compliance to the requirement of the Data Protection Act 1998. The core principles of the act such as the collection, processing of information, disclosure and long-term storage will be adhered to without exceptions. Given the study involves a close collaboration with the NHS, the CI and study team will ensure adherence to the latest version of the NHS Scotland code of practice and procedures on the Patient Confidentiality. The standard operating procedures such as limited access measures to computers via confidential login and password details will be followed to ensure restricted access to participant data by the CI. No personal data will be published through outcomes of the study and participant's confidentiality will be ensured by not revealing any identification details of the individuals.

Participants' identifiable details through imaging data, lab specimens, data entry forms, pathology reports and other records will be protected through institutional designed mechanisms. There will be an access on the principles of "limitation of purpose" basis to the study staff to all the records. Without prior written permission of participants, no clinical details except when absolutely necessary for the purpose of study monitoring or auditing by the Sponsor will be disclosed or released by the CI or study team. The Sponsor or its representative will be informed in writing to obtain permission of any disclosure to third party and without approval of the Sponsor no study confidential data, records of the participants or any other unpublished details will not be disclosed.

3.4 Ancillary and post-trial care

After the biopsy procedure the clinical/research team will assess and record any experience of pain, haematuria, and blood from the back passage. All criteria will be assessed during a telephone call or clinic visit approximately 7 to 14 days post-biopsy. Patients will be given a diary sheet as an aide-mémoire to record any symptoms which will be used to complete CRF via telephone call/clinic visit. In the event that a diary sheet is not completed/returned the clinical/research team will review the clinical records to assess if the participant has experienced any defined AEs and record in the CRF. A model of patient biopsy diary form is shown in supplementary appendix 3.

3.5 Dissemination

The results of this study will be used for publication and presentation in national and international journals and scientific conferences.

3.6 Roles and responsibilities

For roles and responsibilities of trial sponsor, CI, PI and committees in this study, please see supplementary appendix 4.

Trial Status

The favourable approval for this trial was obtained from the East of Scotland REC 1 (Ref: 14/ES/1070) on 20th November 2014. The first patient enrolled on 17th February 2015. 416 participants are recruited so far. The total duration of the recruitment phase was 48 months; however, this has now been extended to 66 months. The recruitment is scheduled to finish on 30th June 2020.

Abbreviations

AE	Adverse Event
CRF	Case Report Form
DRE	Digital Rectal Examination
HIC	Health Informatics Centre
mpMRI	Multiparametric Magnetic Resonance Imaging
PCa	Prostate cancer
PIS	Patient Information Sheet
PSA	Prostate Specific Antigen
R&D	Research & Development
REC	Research Ethics Committee
SAE	Serious Adverse Events
TCTU	Tayside Clinical Trial Unit
TRUS	Transrectal Ultrasound
US	Ultrasound

Declarations

6.1 Ethics approval and consent to participate

The favourable approval for this trial was obtained from the East of Scotland REC 1 (Ref: 14/ES/1070) on 20th November 2014. Every participating site will have their own local REC approval before recruiting participants. Informed consent will be obtained from all study participants.

6.2 Consent for publication

Not applicable

6.3 Availability of data and material

All datasets recorded and analysed in MULTIPROS study are available on reasonable requests from corresponding author.

6.4 Competing interests

The MULTIPROS study and all authors declare no competing interest.

6.5 Funding

This work is funded by Prostate Cancer UK (CSO-PG13-005). The funder was not involved in the design of the study and collection, analysis, and interpretation of data and in writing the manuscript.

6.6 Authors' contributions

All authors listed above (MSB, CW, KC, SG, PD, SKAR, PS, JW, GN) have made substantial contributions in this work: study conception and design (GN, MSB); drafting and interpretation of content (CW, MSB); data acquisition and critical revision (KC, SG, PD, SKAR, PS, JW, MSB); agreement of final version to be published (GN, CW, MSB, KC, SG, PD, SKAR, PS, JW).

6.7 Acknowledgements

Not applicable

References

1. Taira, A. V., Merrick, G. S., Galbreath, R. W. et al.: Performance of transperineal template-guided mapping biopsy in detecting prostate cancer in the initial and repeat biopsy setting. *Prostate Cancer and Prostatic Diseases*, **13**: 71, 2010
2. Welch, H. G., Fisher, E. S., Gottlieb, D. J. et al.: Detection of Prostate Cancer via Biopsy in the Medicare–SEER Population During the PSA Era. *JNCI: Journal of the National Cancer Institute*, **99**: 1395, 2007
3. Rosario, D. J., Lane, J. A., Metcalfe, C. et al.: Short term outcomes of prostate biopsy in men tested for cancer by prostate specific antigen: prospective evaluation within ProtecT study. *BMJ*, **344**: d7894, 2012
4. Schröder , F. H., Hugosson , J., Roobol , M. J. et al.: Screening and Prostate-Cancer Mortality in a Randomized European Study. *New England Journal of Medicine*, **360**: 1320, 2009
5. Tamada, T., Sone, T., Jo, Y. et al.: Prostate Cancer: Relationships between Postbiopsy Hemorrhage and Tumor Detectability at MR Diagnosis. *Radiology*, **248**: 531, 2008
6. Qayyum, A., Coakley, F. V., Lu, Y. et al.: Organ-Confined Prostate Cancer: Effect of Prior Transrectal Biopsy on Endorectal MRI and MR Spectroscopic Imaging. *American Journal of Roentgenology*, **183**: 1079, 2004
7. Kirkham, A. P. S., Haslam, P., Keanie, J. Y. et al.: Prostate MRI: Who, when, and how? Report from a UK consensus meeting. *Clinical Radiology*, **68**: 1016, 2013
8. Rais-Bahrami, S., Siddiqui, M. M., Turkbey, B. et al.: Utility of multiparametric magnetic resonance imaging suspicion levels for detecting prostate cancer. *J Urol*, **190**: 1721, 2013
9. Sheikh, N., Wei, C., Szewczyk-Bieda, M. et al.: Combined T2 and diffusion-weighted MR imaging with template prostate biopsies in men suspected with prostate cancer but negative transrectal ultrasound-guided biopsies. *World J Urol*, **35**: 213, 2017
10. Wei, C., Jin, B., Szewczyk-Bieda, M. et al.: Quantitative parameters in dynamic contrast-enhanced magnetic resonance imaging for the detection and characterization of prostate cancer. *Oncotarget*, **9**: 15997, 2018

11. Shah, V., Pohida, T., Turkbey, B. et al.: A method for correlating in vivo prostate magnetic resonance imaging and histopathology using individualized magnetic resonance-based molds. *Rev Sci Instrum*, **80**: 104301, 2009

Figures




Figure 1

This is the first large trial to randomise men into either targeted or non-targeted biopsy with pre-biopsy MRI and choose radical prostatectomy histopathology using 3D fabricated molds as gold standard (Figure 1).

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [supplement1.docx](#)
- [supplement2.pdf](#)
- [supplement3.pdf](#)
- [supplement4.docx](#)
- [supplement5.pdf](#)
- [supplement6.pdf](#)
- [supplement7.pdf](#)