

Effects of Moving Cupping Therapy for Plaque Psoriasis: A Protocol for a Randomised Multicentre Clinical Trial

Meng Xing

Yueyang Hospital of Integrated Traditional Chinese and Western Medicine, Shanghai University of Traditional Chinese Medicine

Xiaoning Yan

Shaanxi Traditional Chinese Medicine Hospital

Suqing Yang

First Affiliated Hospital of Heilongjiang University of Traditional Chinese Medicine

Linge Li

Shijiazhuang Hospital of Traditional Chinese Medicine

Liping Gong

Affiliated Hospital of Jiangxi University of Traditional Chinese Medicine

Hongxia Liu

Hospital of Traditional Chinese Medicine, Xinjiang Medicine University

Rong Xu

Yueyang Hospital of Integrated Traditional Chinese and Western Medicine, Shanghai University of Traditional Chinese Medicine

Jie Chen

Shanghai University of Traditional Chinese Medicine Yueyang Hospital of Integrated Traditional Chinese Medicine and Western Medicine, Shanghai University of Traditional Chinese Medicine

Luo Ying

Yueyang Hospital of Integrated Traditional Chinese and Western Medicine, Shanghai University of Traditional Chinese Medicine

Yiding Zhao

Shaanxi Traditional Chinese Medicine Hospital

Yuepeng An

First Affiliated Hospital of Heilongjiang University of Traditional Chinese Medicine

Yang Liu

Shijiazhuang Hospital of Traditional Chinese Medicine

Gang Huang

The Affiliated Hospital of Jiangxi University of Traditional Chinese Medicine

Fei Guo

Hospital of Traditional Chinese Medicine, Xinjiang Medicine University

Qingfeng Yin

Jiansu Famous Medical Technology Co. Ltd, Nanjing University of Traditional Chinese Medicine

Ruiping Wang

Yueyang Hospital of Integrated Traditional Chinese and Western Medicine, Shanghai University of Traditional Chinese Medicine

Bin Li

Yueyang Hospital of Integrated Traditional Chinese and Western Medicine, Shanghai University of Traditional Chinese Medicine

Xin Li (✉ 13661956326@163.com)

Shanghai University of Traditional Chinese Medicine Yueyang Hospital of Integrated Traditional Chinese Medicine and Western Medicine <https://orcid.org/0000-0003-2525-9679>

Study protocol

Keywords: Plaque Psoriasis; Moving Cupping; Protocol; Randomized controlled trial

Posted Date: February 5th, 2020

DOI: <https://doi.org/10.21203/rs.2.12490/v3>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on February 26th, 2020. See the published version at <https://doi.org/10.1186/s13063-020-4155-0>.

Abstract

Background: The clinical treatment of plaque psoriasis is based on comprehensive therapy, which is expensive and unsatisfactory, and some of the drugs currently used have serious side effects. Moving cupping therapy has been used clinically for thousands of years in China and has the advantage of being inexpensive and easy to perform. It is now widely used in public hospitals in China for the treatment of psoriasis. However, at present, a comprehensive evaluation of the current clinical evidence regarding its use is lacking. This study aims to evaluate the efficacy and safety of moving cupping in the treatment of plaque psoriasis. **Methods and analysis:** A multicentre, two-arm parallel group, single-blind randomised controlled trial will be conducted in six academic and non-academic hospitals in China. A total of 110 adult patients (aged 18-65 years) who meet the inclusion criteria are now being recruited. Participants will receive (1) basic treatment combined with moving cupping therapy or (2) basic treatment combined with moving cupping placebo. The treatment cycle will be 4 weeks, and the treatment efficacy will be assessed weekly using the Psoriasis Area and Severity Index during the treatment period and the follow-up visits at weeks 6 and 8. The body surface area, physician's global assessment, Dermatology Life Quality Index, patient-reported quality of life, visual analogy scale, traditional Chinese medication syndrome scoring scale, combined medication, and adverse events will also be recorded and compared to the baseline values. **Discussion:** The results of this trial may help make better decisions in the treatment of plaque psoriasis. If the results are considered to be favorable, this ancient Chinese medicine therapy may be worthier of promotion because of its convenience and cheap advantages to benefit patients. **Trial registration:** This study has been registered at ClinicalTrials.gov under the identifier number NCT03952676. Registered on 15 May 2019.

Background

Psoriasis is a chronic inflammatory skin disease that affects approximately 2-3% of the global population [1,2]. It is characterized by epidermal hyperproliferation, abnormal keratinocyte differentiation, angiogenesis with blood vessel dilatation, and excess T-helper cell type 1 (Th1) and Th17 inflammation [3]. The plaque type is the most common form of psoriasis, accounting for 85-90% of psoriasis cases [1]; it leads to detrimental physical effects and reduced psychological well-being [4,5]. It is also closely related to metabolic syndrome, cardiovascular disease, and chronic obstructive pulmonary disease [6]. For most patients, psoriasis results in the restriction of various aspects of everyday life, enormous personal costs, and mental stress [7].

The current treatment for psoriasis primarily comprises local and systemic treatments. Local treatment mainly comprises hormonal drugs and calcineurin inhibitors, and systemic treatment comprises etretin, immunosuppressants, and biological agents [8]. However, the various side effects and high economic costs limit the clinical application of these treatments [9]. Therefore, developing or identifying safe and effective treatments for psoriasis has major social and economic benefits.

Recently, complementary and alternative medicine (CAM) therapies have become an increasingly important area of dermatology, and cupping is becoming an important CAM therapy. Cupping is an ancient method that has been used worldwide. From ancient Egypt to the Han Dynasty in China [10,11], and from Hippocrates in Greece to the early Islamic period [12,13], there have been numerous descriptions of cupping treatments for various diseases. Cupping is also currently used to treat a wide range of medical conditions [14,15], and randomized controlled trials (RCTs) have confirmed the efficacy of cupping for certain pain-related diseases such as osteoarthritis of the knee and chronic low back pain [16,17]. There are two types of cupping methods; dry and wet. Moving cupping therapy is a unique dry method that has been used as a traditional treatment for thousands of years. This method involves the application of lubricant to the body part or to the mouth of the cup and using the flashing method or the cotton sticking method to adsorb the cup to the treatment area. The cotton sticking method involves pasting cotton soaked in alcohol on the inner wall of the cup; it is then ignited and adsorbed on the treatment area, but the skin may be scalded due to the excessive dripping of alcohol. The flashing method quickly withdraws the cotton soaked in alcohol. This is a safe and common method of cupping. The physician pushes the cup by hand to move it up and down and left and right, thus causing flushing, congestion, and even ecchymosis of the skin in the treatment area [18]. Therefore, moving cupping therapy integrates the functions of warm moxibustion (which involves lighting the moxa stick and hanging it at a certain distance from the skin), cupping, scraping, massage, and drug therapy, and it has a wide range of clinical applications. This therapy has the ability to regulate immune function [19,20], thereby improving skin tolerance [21,22]. Psoriasis damages the epidermis, which destroys the human skin barrier composed of keratinocytes, intercellular substance, natural moisturizing factors and sebum membranes. The Koebner phenomenon illustrates the important relationship between an impaired skin barrier and psoriasis [23,24]. Moving cupping treatment can cause the mechanical stimulation of pulling and is beneficial for the secretion of sebaceous glands and sweat glands, and the use of lubricants can significantly improve the skin barrier function. Psoriasis is also known to be closely related to lipid metabolism [25]. Cupping therapy can upregulate anti-inflammatory lipids, downregulate the function of pro-inflammatory lipids, improve the balance of lipid metabolites, and alleviate the inflammatory response [26].

Moving cupping therapy has been widely used for the treatment of plaque psoriasis, which is safe and associated with few adverse reactions, mainly local blisters. Additionally, it has been recognized by a large number of peers and patients. One study compared the efficacy of moving cupping and narrowband ultraviolet B based on oral Chinese herbs and found that moving cupping has clear advantages, but lacked high-quality medical evidence [27]. Therefore, we designed a multicenter, prospective, single-blind, placebo-controlled, randomized clinical trial to evaluate the safety and efficacy of moving cupping therapy for plaque psoriasis.

Objectives and hypotheses

Moving cupping is widely used as a CAM therapy. The aim of this RCT is to evaluate the efficacy of moving cupping as a plaque psoriasis treatment. The hypothesis of this study is that the Psoriasis Area and Severity Index (PASI), used as the primary endpoint to demonstrate the clinical efficacy of plaque psoriasis treatment, will be decreased significantly with moving cupping treatment compared to placebo treatment [28]. Secondary endpoints, including body surface area (BSA), physician's global assessment (PGA), traditional Chinese medicine syndrome scoring scale (TCMSSS), and patient-reported outcomes (Dermatology Life Quality Index [DLQI] and visual analogy scale [VAS]), for the evaluation of post-treatment clinical efficacy, psychology, quality of life, and degree of pruritus will also be considered. This study complies with the relevant Standard Protocol Items: Recommendations for Interventional Trials (SPIRIT) Checklist (Supplementary File 1) [29].

Methods

Study design

The proposed study is designed as a prospective, multicenter, two-arm parallel group, single-blind RCT. Patients with plaque psoriasis will be randomly assigned to the treatment group (moving cupping) or the placebo group (sham moving cupping). Eligible participants will receive 4 weeks of treatment and a total of 8 weeks of follow-up (Table 1).

Patient and public involvement

Patients were involved in the design of the RCT.

Recruitment

We plan to recruit 122 patients with plaque psoriasis for this study. They will be evenly distributed among six hospitals using fixed enrollment. Recruitment will occur at the Yueyang Hospital of Integrated Traditional Chinese and Western Medicine (Shanghai, China), Shaanxi Traditional Chinese Medicine Hospital (Shaanxi, China), First Affiliated Hospital of Heilongjiang University of Traditional Chinese Medicine (Heilongjiang, China), the Affiliated Hospital of Jiangxi University of Traditional Chinese Medicine (Jiangxi, China), Hospital of Traditional Chinese Medicine, Xinjiang Medicine University (Xinjiang, China), and Shijiazhuang Hospital of Traditional Chinese Medicine (Shijiazhuang, China). Eligible recipients will be recruited via social media, word of mouth, and dermatology outpatient clinics at local hospitals. Each hospital will designate three dermatologists for patient recruitment and one physician (dermatologists trained and qualified to perform moving cupping) for moving cupping treatment. Patients will be informed and asked to provide written informed consent prior to study commencement. Figure 1 presents a flowchart of participant enrollment and analysis throughout the course of the trial. Researchers will obtain recipient consent prior to trial participation. All potential and

enrolled recipients' personal information will be recorded, kept in a secure folder, and made accessible only to the researchers in order to protect their privacy.

Participant screening

All patients who are diagnosed with plaque psoriasis will undergo laboratory blood testing before inclusion, including a complete blood cell count, liver function test, renal function test, and pregnancy test. Additionally, routine urine tests, vital sign monitoring, and physical examinations will be performed (Table 1).

Criteria

The inclusion criteria are as follows: meet the 2018 China College of Dermatology diagnostic criteria for plaque psoriasis; skin lesions involve $\leq 10\%$ BSA (the lesions are mainly located on the torso/limbs, palm/sole, or face/scalp, with the vulva area being unaffected); aged 18-65 years; and provide consent to participate in the study and sign the informed consent form.

The exclusion criteria are as follows: any clinically active skin diseases other than moderate to severe psoriasis vulgaris that might confound or influence the study aim; patients who received any systemic treatments within 4 weeks before the baseline visit; patients who received topical treatment within 2 weeks before the baseline visit (e.g., corticosteroids, ultraviolet light therapy including sunbathing); patients with an active infectious disease that is difficult to control; patients with a history of severe systemic disease, an alanine aminotransferase or aspartate transaminase level greater than 1.5-fold that of the average; any of the main routine blood indices (white blood cell count, red blood cell count, hemoglobin level, platelet count) lower than the normal limit, or other laboratory abnormalities; immunocompromised patients who may experience skin allergies or infection with moving cupping therapy; pregnant or lactating women; patients with a history of alcohol or drug abuse; and patients with a history of serious mental illness.

The exit/termination criteria were as follows: patient lost to follow-up; violation of the research plan (including lack of compliance [medication, contraindication, interview plan]); adverse events (including worsening of psoriasis above baseline [for example, PASI $>125\%$]) [28]; and pregnancy.

Randomization, allocation, and blinding

Once eligibility is confirmed and consent is obtained, participants will be block-randomized using a non-stratified, permuted block with varying lengths ($n = 6$ blocks; patient distribution = 1:1) for each arm/group. Independent statisticians will provide computer-generated random sequences. Distribution

will be conducted by a central web-based interactive randomization service system. When a participant is eligible, the researcher who conducts the random grouping will need to log onto the website to view the grouping situation of the participant. Due to the nature of moving cupping, it is not possible to blind the operating physician involved in treatment application. To ensure blinding, we will use a placebo cup to simulate the treatment of moving cupping and shield the patient's eyes with a black opaque eye mask. The treatment device will be of the same size and material as well, and the treatment method will maintain consistent frequency and intensity, but with no adsorption force. Each hospital will designate one operating physician who is a dermatologist trained to perform moving cupping therapy and has obtained the relevant operational qualifications. The operating physician will not participate in the statistical analysis, and the treatment plan and grouping will be known only by the statistical analyst.

Intervention

Moving cupping intervention

The cup has the texture of transparent glass. For convenience, different cup sizes will be selected according to the site of the patient's skin lesions (Figure 2).

Moving cupping involves standardized manipulations. First, Vaseline is applied to the skin lesion area. Using the flashing method, a 95% ethanol-soaked cotton ball is held with tweezers and the cup is held upside-down. After the cotton ball is ignited, it is immediately moved down inside the cup and removed; the cup is then quickly placed on the skin lesion area. After using the cup to absorb the skin lesion area, the cup body is held in one hand and pushed and pulled according to the skin lesion area while applying light force, such that the skin of the treatment area turns purple. Even force is applied when pushing the cup to prevent the cup from falling off due to air leakage. This is repeated on the skin lesion area 30 times [30]. The cup is changed five times per push and pull, with an interval of no more than 10 s. This will be performed once every other day for 4 weeks.

Moving cupping placebo

A special perforated cup made of the same material as that used for the intervention group will be selected and used during the same manipulation method as described for the intervention. Due to the perforated design, the fire cannot burn the air in the cup; therefore, it cannot form a negative pressure adsorption force. This simulates the form of moving cupping therapy without the therapeutic effects (Figure 2).

Basic treatment

1. Increase moisturizing of the skin: As a basic treatment, the use of a moisturizer at all times is mandatory. The moisturizer should be applied to the skin areas that are free from erosion and

exudation, as well as the dry non-lesion areas. A soft, fragrance-free moisturizer should be used. Yuze moisturizer will be recommended to patients with sensitive dry skin.

2. Standard bath: Generally, patients should rinse quickly with warm water (35-39°C) once per day for approximately 5 min. Moisturizer should be applied within 2 min of bathing to avoid dehydration of the epidermis. The use of alkaline detergents to clean the skin should be avoided.
3. Avoidance of induced and aggravated factors: For some patients, certain foods can aggravate or induce disease. These foods should be avoided.
4. Some patients may have food allergies. When food allergies are identified, such foods should be avoided to prevent inducing or aggravating the condition.
5. Maintain a reasonably healthy lifestyle: Patients should avoid staying up late and becoming overly stressed. Spicy, irritating foods should be avoided, and appropriate exercise should be performed. Patients should attempt to maintain normal bowel movements.

Measurement

Baseline measurements

Vital signs and disease severity will be monitored by physical examinations, biochemical examinations, and various evaluation indices before treatment. Foods that cause allergic reactions will be identified by food allergen testing. Data from these measures will be used as baseline moving cupping therapy data.

Examination during the interview

Interview plan

All patients will be interviewed at weeks 1, 2, 3, and 4 during treatment. They will be reminded of the interview by phone 1 day before it is scheduled to occur. During weeks 1, 2, 3, and 4 of treatment, the psoriatic lesions will be measured by PASI, PGA, BSA, and TCMSSS, and the DLQI and VAS will be used to assess the quality of life and psychological status, and degree of pruritis, respectively. Their moisturizing, diet, and bathing habits will be monitored via patient diaries. In addition, during weeks 6 and 8 of the follow-up, the PASI, PGA, BSA, and VAS will be recorded again to evaluate the efficacy of moving cupping. Possible adverse events and combined medications should be accurately recorded at all of these time points.

Examination during follow-up

Follow-up plan

All patients will be followed-up by phone at weeks 6 and 8 after treatment. During each follow-up session, the treatment effects will be assessed using the PASI, PGA, BSA, and VAS.

Primary parameters

PASI: Incorporates the extent of psoriasis at four anatomical sites by evaluating signs of erythema, scale, and elevation. PASI scores range from 0 to 72. The primary outcome of the RCT is the proportion of patients with a reduction in PASI scores of $\geq 75\%$ compared to baseline at the 4 week follow-up visit [28].

Secondary parameters

The secondary parameters of this study include the following:

1. PGA: The PGA is scored using a 5-point scale reflecting the overall degree of erythema, infiltration, and desquamation across all psoriatic lesions [28].
2. BSA: The BSA involved in psoriasis is estimated by fingerprinting, whereby the entire palm of the patient represents approximately 1% of the total BSA. The number of handprints on the psoriatic skin on a body part is used to determine the extent to which the body part is affected by psoriasis (%) [28].
3. DLQI: The DLQI is a participant-reported questionnaire used to measure the health-related quality of life of adults with skin diseases. Scores range from 0 to 30, with a higher score indicating a greater impact on the participant's quality of life [31].
4. VAS: The VAS is used to measure lesion pruritus from 0 to 100 mm at each visit (with 0 indicating no pruritus and 100 indicating maximum pruritus) [32].
5. TCMSSS: According to the different TCM syndromes, comprehensive evaluations will be performed to determine the conditions of the tongue, pulse, and skin. The score ranges from 0 to 6 [33].

Safety monitoring

All participants will be advised to remain under supervision in the clinical research unit for 15 min after treatment. In addition, data of the participants will be monitored by the research team to detect any adverse events throughout the study, such as skin damage or potential allergies. All potential adverse events will be recorded. If any participant experiences an adverse effect due to trial participation, they will receive free treatment and compensation accordingly. If any concerns of the participants are identified during screening or clinical assessment, further clinical evaluations and/or investigations will be performed immediately. If concerns are identified during the study, the participants will be withdrawn.

Sample size

The sample size of the current trial was calculated based on the following formula [34]: (see Formula 1 in the Supplementary Files)

According to recently published clinical trial results [28], the PASI-50 reached 62.1% (P_1) in the study group and 35.3% in the control group (P_2). Therefore, we assumed that the inspection level (α) was 0.05 and the power was 0.8 ($\beta=1-\text{power}=0.2$). For the two-sided tests, 51 participants will be required for each group. Given a 20% loss to follow-up, we expect to require 61 participants for each group. Therefore, this trial will require at least 122 participants.

Timeline

Recruitment began in August 2019, and the intervention period will end in December 2020. Table 1 provides the schedule of enrollment, intervention, and assessment.

Data collection and management

Each center will designate a full-time researcher, and data collection will be performed in a dermatology clinic using a unified data collection system (provided by Nanjing Ningqi Medical Technology Co., Ltd. Data Management Center). With the exception of the DLQI, R-QoL, and VAS, the evaluation indicators will be completed by data collectors. All data collectors will be trained to implement the PASI and other scales.

Data monitoring and auditing

This study will establish the Data and Safety Monitoring Board (DSMB), which is composed of three members, including an acupuncturist, a dermatologist, and a statistician. All members must declare any conflict of interest during the trial. The DSMB will monitor the progress of the trial and review the safety and quality of the data. They will meet quarterly to review any adverse events and safety issues. All adverse events and severe adverse events will be reported to the main researchers, IRB, and DSMB within 24 h. The primary investigator will receive the interim results and make the final decision regarding study termination.

Statistical analysis

In this study, quantitative variables (PASI, DLQI, R-QoL, VAS) with normal distributions will be described as means and standard deviation; non-normal distribution will be summarized as median and interquartile range. Quantitative variables will be compared using a Student's t test for normally-distributed variables and the Wilcoxon rank-sum test for non-normally-distributed variables. Qualitative variables (sex, age) will be summarized as frequency and proportions, and the chi-square or Fisher's exact test will be used to test the differences between qualitative variables. Quantitative variables with repeated measures will be applied with general linear models. $P < 0.05$ will be considered statistically significant. All statistical analyses will be performed with SAS version 9.4 statistical package. The measurer will be blinded to the results.

Missing data

The possibility of loss to follow-up has been considered and will be calculated as a part of the sample size estimation. The dropout rates and reason will be recorded. In addition, we will account for other types of randomly missing data by treating dropouts as non-success or non-survival using the intention-to-treat principle.

Ethics and dissemination

The study has been approved by the ethics committee of Yueyang Hospital of Integrated Traditional Chinese and Western Medicine (ref. approval no. 2019-003). Patients will be informed at the beginning of the study that they have the right to withdraw from the study at any time without providing a reason. Even in the event of a withdrawal, the required treatment will be provided to the patient. The results of the study will be published in an international peer-reviewed journal.

Discussion

Psoriasis often requires multiple methods of combination therapy in the clinic; however, this does not always result in satisfactory outcomes. Therefore, there is increasing concern regarding the current combinations of CAM in modern medical practice. Cupping is a form of CAM that has existed for thousands of years in various civilizations. It plays a unique role in various diseases, including dermatology. Studies have found that cupping can lower the level of superoxide dismutase in the blood, which has a role in reducing oxidative stress [35,36]. Cupping can also significantly reduce the hemoglobin level in the cupping area and increase the level of oxyhemoglobin. In addition, it can increase HSP-70 and β -endorphins to relieve pain [37]. Although the mechanism of moving cupping treatment for psoriasis is not clear, there are indications that moving cupping can alleviate plaque psoriatic skin inflammation and excessive thickening of skin lesions. Conventional treatment combined with moving

cupping has a better curative effect. However, it is true that the therapeutic effect still requires further rigorous scientific verification.

Currently, only small-scale clinical research has been performed for moving cupping treatment of psoriasis. Relevant RCT studies did not adopt blinding methods, lacked placebo control, did not follow the consort statement, and were published in languages other than English [27,30]. Therefore, large RCTs are needed to assess the role and possible adverse outcomes of moving cupping for the treatment of plaque psoriasis.

This study will be the first placebo-controlled study of moving cupping for the treatment of plaque psoriasis with multiple centers, double-arm parallel groups, and a single-blind RCT. Furthermore, the study has several advantages. The first is that it will use a placebo control to exclude the placebo effect, reduce the risk of bias, and define the clinical efficacy of moving cupping for the treatment of plaque psoriasis. The second advantage is that, because of the large land area of China, the study will include a total of six hospitals in different regions, thus including various ethnic and socio-economic populations. As such, its experimental design will increase the generalizability of the trial outcomes. However, this study also has some limitations. Due to the particularity of moving cupping therapy, it is difficult for different operators to maintain the same strength when moving the cup. Furthermore, because different cup sizes will be selected for the different treatment sites, adsorption forces will be inconsistent. All of these limitations may lead to the risk of bias. The placebo cupping therapy method has been used previously in only two clinical trials [38,39]. Although the moving cupping placebo lacks absorption during treatment, we found that most patients do not understand the specific operation of the treatment method; therefore, they cannot distinguish between the moving cupping placebo and moving cupping. Patients were blinded to whether they are accepting real or placebo cupping. In conclusion, this study aims to verify whether moving cupping therapy is effective for treating plaque psoriasis. We also hope to provide a safe and effective treatment for plaque psoriasis through this study.

Trial Status

The protocol version number is 2.0 and the version date is May 15, 2019. The study plans to start recruiting participants in August 2019 and complete the recruitment in December 2020.

Abbreviations

PASI: Psoriasis Area and Severity Index

PGA: Physician's global assessment

BSA: Body surface area

DLQI: Dermatology Life Quality Index

VAS: Visual analogue scale

TCMSSS: Traditional Chinese medicine syndrome scoring scale

RCT: Randomized controlled trial

Declarations

Ethics approval and consent to participate

Central ethical approval has been confirmed by the Yueyang Hospital of Integrated Traditional Chinese and Western Medicine (ref approval no. 2019-003). We will not begin recruiting at other centers for the trial until local ethical approval has been obtained. Informed consent will be obtained from all participants before study initiation.

Consent for publication

Not applicable.

Availability of data and material

We declare that the materials described in the manuscript, including all relevant raw data, will be freely available to any scientist wishing to use them for non-commercial purposes without breaching participant confidentiality.

Competing interests

None declared.

Funding

This study was supported by the National Key Research and Development Program of China (No. 2018YFC1705303), the NSFC of China (No. 81874470, 81973860), the Shanghai Development Office of TCM [Nos. ZY(2018-2020)-FWTX-1008, ZY(2018-2020)-CCCX-2004-08, and ZY(2018-2020)-FWTX-4010], the Shanghai Science and Technology Committee (No. 18401932300).

Author contributions

XL, MX, and BL conceived the study. XY, YL, XY, LL, and HL designed the study. The study protocol was drafted by MX and XL and was revised by RW, BL, and QY. SY, LL, LG, HL, RX, JC, LY, YZ, YY, FG. and LH will implement the study. All authors approved the final manuscript of this study protocol.

Acknowledgements

None.

References

- [1] Parisi R, Symmons DP, Griffiths CE, et al. Global epidemiology of psoriasis: a systematic review of incidence and prevalence. *J Invest Dermatol.* 2013; 133:377–85.
- [2] Chandran V, Raychaudhuri SP. Geoepidemiology and environmental factors of psoriasis and psoriatic arthritis. *J Autoimmun.* 2010;34: J314–21.
- [3] Kurd SK, Richardson S, Gelfand JM. Update on the epidemiology and systemic treatment of psoriasis. *Expert Rev Clin Immunol.* 2007; 3:171–85.
- [4] Boehncke WH, Schön MP. Psoriasis. *Lancet* 2015; 386: 983–94.
- [5] Zhao Y, Fischelevich R, Petrali JP, et al. Activation of keratinocyte protein kinase C zeta in psoriasis plaques. *J Invest Dermatol* 2008; 128: 2190–97.
- [6] Li Xin, Kong Lingjun, Li Fulun et al. Association between Psoriasis and Chronic Obstructive Pulmonary Disease: A Systematic Review and Meta-analysis. [J]. *PLoS ONE*, 2015, 10: e0145221.
- [7] Schöffski O, Augustin M, Prinz J, et al. Costs and quality of life in patients with moderate to severe plaque-type psoriasis in Germany: a multi-center study. [J]. *Jddg Journal Der Deutschen Dermatologischen Gesellschaft*, 2007, 5(3):209–218.
- [8] Nast A, Jacobs A, Rosumeck S, et al. Methods Report: European S3-Guidelines on the systemic treatment of psoriasis vulgaris – update 2015 – EDF in cooperation with EADV and IPC[J]. *Journal of the European Academy of Dermatology & Venereology Jeadv*, 2015, 29(12):2277.
- [9] Schöffski O, Augustin M, Prinz J, et al. Costs and quality of life in patients with moderate to severe plaque-type psoriasis in Germany: a multi-center study. [J]. *Jddg Journal Der Deutschen Dermatologischen Gesellschaft*, 2010, 5(3):209-218.
- [10] Haller JS, Jr. The glass leech. Wet and dry cupping practices in the nineteenth century. *N Y State J Med.* 1973; 4:583–92.
- [11] Chirali IZ. Traditional Chinese medicine cupping therapy. London: Elsevier Churchill Livingstone; 2007.
- [12] Kouskoukis CE, Leider M. Cupping: the art and the value. *Am J Dermatopathol* 1983; 5:235–40.
- [13] AlRawi SN, Khidir A, Elnashar MS, Abdelrahim HA, Killawi AK, Hammoud MM, et al. Traditional Arabic & Islamic medicine: validation and empirical assessment of a conceptual model in Qatar. *BMC Complement Altern Med.* 2017; 17:157.
- [14] Kordafshari G, Ardakani M R S, Keshavarz M, et al. The Role of Phlebotomy (Fasd) and Wet Cupping (Hijamat) to Manage Dizziness and Vertigo from the Viewpoint of Persian Medicine[J]. *Journal of*

Evidence-Based Complementary and Alternative Medicine, 2016, 22(3):369-373.

[15] Lu S, Du S, Fish A, et al. Wet cupping for hypertension: a systematic review and meta-analysis[J]. *Clinical and Experimental Hypertension*, 2019, 41(5):474-480.

[16] Teut M, Ullmann A, Ortiz M, et al. Pulsatile dry cupping in chronic low back pain – a randomized three-armed controlled clinical trial[J]. *BMC Complementary and Alternative Medicine*, 2018, 18(1):115.

[17] Teut M, Kaiser S, Ortiz M, et al. Pulsatile dry cupping in patients with osteoarthritis of the knee – a randomized controlled exploratory trial[J]. *BMC Complementary and Alternative Medicine*, 2012, 12(1).

[18] Acupuncture Technical Practice Part 5: Cupping. Project group. National Standard of the People's Republic of China (GB/T21709.5-2008) Acupuncture Technical Practice Part 5: Cupping [J]. *Chin Acup Moxib*, 2010, 30(2):160-162.[19] Meng F, Gao ZL, Li L, et al. Reconstruction of lymphatic vessels in the mouse tail after cupping therapy[J]. *Folia Morphol*, 2019, 4:1–19.

[20] Dons"Koi B V, Chernyshov V P, Osypchuk D V, et al. Repeated cupping manipulation temporary decreases natural killer lymphocyte frequency, activity and cytotoxicity[J]. *Journal of Integrative Medicine*, 2016, 14(3):197-202.

[21] Lowe DT. Cupping therapy: An analysis of the effects of suction on skin and the possible influence on human health. *Complementary Therapies in Clinical Practice*. 2017; 29:162–168.

[22] Lee H Y, Lee H J, Kim G C, et al. Plasma cupping induces VEGF expression in skin cells through nitric oxide-mediated activation of hypoxia inducible factor 1[J]. *Scientific Reports*, 2019, 9(1).

[23] Gruber R, Schmuth M. The barrier function of normal skin. Morphologic and functional aspects of the skin barrier [J]. *Hautarzt*, 2014. 65 (3) : 234-240

[24] Bergboer JG, Zeeuwen PL, Schalkwijk J. Genetics of psoriasis: evidence for epistatic interaction between skin barrier abnormalities and immune deviation [J], *J Invest Dermatol*, 2012 132 (10) : 2320-2321

[25] Kim HN, Han K, Park YG, et al. Metabolic syndrome is associated with an increased risk of psoriasis: A nationwide population-based study[J]. *Metabolism Clinical and Experimental*, 2019,10(11):19 -24.

[26] Zhang Q, Wang X, Yan G, et al. Anti- Versus Pro-Inflammatory Metabololipidome Upon Cupping Treatment [J]. *Cellular Physiology and Biochemistry*, 2018:1377-1389.

[27] Liu HX, Zhang CH, Liu ZX, et al. Observation on the clinical efficacy of moving cupping therapy for plaque psoriasis [C]. *Compilation of Papers of 2010 National Conference on Dermatology and Venereology of Integrated Traditional Chinese and Western Medicine*. 2010.

- [28] European Medicines Agency. Committee for medicinal products for human use (CHMP). Guideline on clinical investigation of medicinal products indicated for the treatment of psoriasis. London, UK: EMEA; 2004. http://www.ema.europa.eu/docs/en_GB/document_library/Scientific_guideline/2009/09/WC500003329.pdf. Accessed August 31, 2017.
- [29] Chan AW, Tetzlaff JM, Altman DG, et al. SPIRIT 2013 statement: defining standard protocol items for clinical trials. *Ann Intern Med* 2013; 158:200–7.
- [30] Feng L, Guo F, Ji Y. Clinical study on the treatment of plaque psoriasis of blood stasis syndrome based on the treatment of moving cupping [J]. *Chinese Journal of Traditional Chinese Medicine*, 2014 (10): 3343-3345.
- [31] Finlay A Y, Khan G K. Dermatology Life Quality Index (DLQI)—a simple practical measure for routine clinical use. [J]. *Clinical & Experimental Dermatology*, 2010, 19(3):210-216.
- [32] Price D D, Mcgrath P A, Rafii A, et al. The validation of visual analogue scales as ratio scale measures for chronic and experimental pain. [J]. *Pain*, 1983, 17(1):45-56.
- [33] Department of Dermatology, Chinese Medicine Association of China. Psoriasis vulgaris (white peony) Chinese medicine evidence-based clinical practice guide (2013 edition) [J]. *Journal of Traditional Chinese Medicine*, 2014, 55(1): 76-82.
- [34] Deng W, He J. Design and statistically analysis of clinical trials [M]. Beijing: People’s Medical Publishing House, 2012
- [35] Tagi SM, Celik HT, Ciftci S, Kazanci FH, Arslan M, Erdamar N, et al. Wet-cupping removes oxidants and decreases oxidative stress. *Complement Ther Med* 2014; 22:1032–6.
- [36] Li T, Li Y, Lin Y, Li K. Significant and sustaining elevation of blood oxygen induced by Chinese cupping therapy as assessed by near-infrared spectroscopy. *Biomed Opt Express*. 2017; 8:223–9.
- [37] Subadi I, Nugraha B, Laswati H, Josomuljono H. Pain relief with wet cupping therapy in rats is mediated by heat shock protein 70 and β -endorphin. *Iran J Med Sci*. 2017; 42:384.
- [38] Volpato MP, Albano Breda IC, Carolina de Carvalho R, de Castro Moura C, Ferreira LL, Silva ML, Torres Silva JR. Single Cupping Therapy Session Improves Pain, Sleep And Disability In Patients With Nonspecific Chronic Low Back Pain[J]. *Journal of acupuncture and meridian studies*, 2019,
- [39] Lauche R, Spitzer J, Schwahn B, Ostermann T, Bernardy K, Cramer H, Dobos G, Langhorst J. Efficacy of cupping therapy in patients with the fibromyalgia syndrome—a randomised placebo controlled trial[J]. *Scientific reports*, 2016, 6 37316.

Table

Table 1 Study schedules of enrollment, intervention, and assessment

	Study Period							
	Enrollment	Allocation	Treatment period				Follow-up period	
Time point	-week ₁	0	week ₁	week ₂	week ₃	week ₄	week ₆	week ₈
Enrollment								
Eligibility screen	×							
Informed consent	×							
List other procedures	×							
Allocation		×						
Interventions:								
Basic treatment plus moving cupping therapy								
Basic treatment plus moving cupping placebo								
Assessments:								
PASI	×	×	×	×	×	×	×	×
BSA	×	×	×	×	×	×	×	×
PGA	×	×	×	×	×	×	×	×
DLQI		×	×	×	×	×		
VAS		×	×	×	×	×	×	×
TCMSSS		×	×	×	×	×		
Vital signs	×	×	×	×	×	×		
Routine blood test	×					×		
Blood biochemical test	×					×		
Routine urine test	×					×		
Food allergen test	×							
Combined medications	×	×	×	×	×	×		
Pregnancy test	×							
Physical examination	×					×		
Adverse event		×	×	×	×	×		
Serious adverse events		×	×	×	×	×		

PASI, Psoriasis Area and Severity Index; PGA, physician’s global assessment; BSA, body surface area; DLQI, Dermatology Life Quality Index; PR-QoL, patient-reported quality of life; VAS, visual analogue scale; TCMSSS, traditional Chinese medicine syndrome scoring scale.

Figures





Figure 1

Consolidated Standards of Reporting Trials diagram of participant enrollment and analysis



A: The cups used for the intervention group can produce a negative pressure adsorption force when the air in the cup is burned. B: The cup used for the placebo group has a special perforation design, and no negative pressure adsorption was formed after combustion.

Figure 2

Different cups used for the intervention group and the control group when implementing moving cupping therapy

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Ethicalmaterial.pdf](#)
- [Funding.pdf](#)
- [SPIRITchecklist.pdf](#)
- [Formula1.jpg](#)
- [Appendix.pdf](#)

- [renamed1c3ba.doc](#)