

Partnering with General Practitioners to Optimize Survivorship for Patients with Lymphoma: A phase II Randomized Controlled Trial (The GOSPEL 1 Trial)

Raymond Chan (✉ Raymond.Chan@qut.edu.au)

Queensland University of Technology (QUT) <https://orcid.org/0000-0003-0248-7046>

Stephanie Buhagiar

Division of Cancer Services, princess Alexandra Hospital

Laisa Teleni

School of Nursing, Queensland University of Technology (QUT)

Camilla Simonsen

Division of Cancer Services, Princess Alexandra Hospital and School of Nursing, Queensland University of Technology (QUT)

Jane Turner

Faculty of Medicine, University of Queensland and Royal Brisbane and Women's Hospital

Lee Jones

Queensland University of Technology (QUT)

Louisa Gordon

QIMR Berghofer Medical Research Institute

Ria Joseph

School of Nursing, Queensland University of Technology (QUT)

Oluwaseyifunmi Andi Agbejule

Queensland University of Technology

Fiona Henderson

Division of Cancer Services, Princess Alexandra Hospital

Joel Rhee

University of Wollongong

Sally Mapp

Princess Alexandra Hospital

Study protocol

Keywords: Lymphoma, hematology, shared-care, quality of life, protocol, randomized controlled trial

Posted Date: August 25th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-32868/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on January 6th, 2021. See the published version at <https://doi.org/10.1186/s13063-020-04945-4>.

Abstract

Background: Survival rates for lymphoma are highest amongst hematological malignancies. In 2019, it was estimated that over 6,400 Australians were diagnosed with lymphoma, a group of hematological malignancies with a high 5-year survival rate of ~76%. There is an increased focus on the promotion of wellness in survivorship and active approaches to reducing morbidity related to treatment; however, current models of follow-up care heavily rely on hospital-based specialist-led care.

Maximizing the potential of general practitioners (GPs) in the ongoing management of cancer is consistent with the national health reform principles, and the Cancer Council Australia's Optimal Care Pathways. GPs are well positioned to provide guideline-based follow-up care and are more likely to address comorbidities, psychosocial issues and promote healthy lifestyle behaviors. This study aims to test the feasibility of the GOSPEL 1 intervention for implementing an integrated, shared-care model in which cancer center specialists and community-based GPs collaborate to provide survivorship care for patients with lymphoma.

Methods: We describe a protocol for a phase II, randomized controlled trial with two parallel arms and a 1:1 allocation. Sixty patients with Hodgkin's and Non-Hodgkin's lymphoma will be randomized to usual specialist-led follow-up care (as determined by the treating hematologists) or a shared follow-up care intervention (i.e., GOSPEL 1). GOSPEL 1 is a nurse-enabled, pre-specified shared-care pathway with follow-up responsibilities shared between cancer center specialists (i.e., hematologists and specialist cancer nurses) and GPs. Outcome measures assess feasibility as well as a range of patient reported outcomes including health-related quality of life as measured by the Functional Assessment of Cancer Therapy – Lymphoma, patient experience of care, symptom distress, comorbidity burden, dietary intake, physical activity behaviors, financial distress/interference and satisfaction of care. Safety indicators including hospital admission and unscheduled lymphoma clinic visits as well as process outcomes such as intervention fidelity and economic indicators will be analyzed.

Discussion: This trial is designed to explore the feasibility and acceptability of a new model of shared-care for lymphoma survivors. Patient reported outcomes as well as potential barriers to implementation will be also analyzed to inform a larger definitive clinical trial testing the effects of a shared-care model on health-related quality of life of lymphoma survivors.

Trial registration: Australia and New Zealand Clinical Trials Registry, Registration number: ACTRN12620000594921, Date registered: 22/05/2020, registration link: <https://www.anzctr.org.au/ACTRN12620000594921.aspx>

Introduction

Background and rationale {6a}

In 2019, it was estimated that over 6,400 Australians were diagnosed with Hodgkin's Lymphoma (HL) and Non-Hodgkin's Lymphoma (NHL) (1). Lymphomas are hematological malignancies with high overall five year survival rates (~76%) resulting in a large cohort of people post therapy or undergoing regular surveillance for incurable but indolent disease not yet warranting treatment. Lymphoma survivors require ongoing survivorship care including monitoring for cancer progression or recurrence, surveillance for secondary/new primary cancer, and management of a range of long-term bio-psycho-social effects from their cancer diagnosis and treatment. Moreover, many cancer survivors are also in need for management of comorbidities (2). Compared to people without cancer, people with cancer are more likely to develop mental and behavioral problems (2.5 times), circulatory conditions (1.3 times), musculoskeletal conditions (1.4 times), and endocrine system disorders (1.2 times) (2). These health concerns highlight the importance of a well-integrated, patient-centered model of care that addresses cancer-related survivorship needs as well as comorbidities for people who have been diagnosed with lymphoma.

The current models of cancer care in Australia are mostly specialist-led and focus on surveillance for disease recurrence, rather than holistic care needs. These models of care limit integration between specialist institutions and the patient's community-based general practitioner (GP), also called family physician or primary care physician in some countries. With the ever-growing and ageing population of cancer survivors, specialist-led follow-up is not optimal as it does not encourage an integrated, patient-centered approach in which comorbidities and long-term effects experienced by cancer survivors can be effectively managed. Specialist-led follow-up does not capitalize on the expertise of GPs, and it requires patients living in the rural/regional areas to travel long distances to the cancer centers.

Ongoing care of patients with chronic illnesses such as cancer is core business for general practice. Maximizing the potential of GPs in the ongoing management of cancer is consistent with the national health reform principles (3-5), and the Optimal Care Pathways (6, 7). In the breast cancer setting, it is clear that GPs can provide guideline-based follow-up care and are more likely to address comorbidities, psychosocial issues and promote healthy lifestyle behaviors compared to cancer specialists (8). Shared-care arrangements might also facilitate a smoother transition to 'mainstream' clinical care for patients who have successfully undergone treatment and had several years of follow-up without recurrence. A shared-care approach is likely to be more cost-effective than a specialist-led model due to the added benefits from harnessing the expertise of both the cancer specialist as well as the GP. Such an approach, if effective, may further relieve pressures in health services to meet the ever-increasing demand. The recent COVID-19 pandemic has further highlighted the need to explore, implement, and evaluate alternative models of care, enabling a reduction in acute hospital setting presentations and enhanced community GP involvement (9).

Although implementing an integrated, shared-care follow-up model is the next logical step for cancer patients with high survival rates, the evidence base supporting this model is less established for lymphoma compared to breast cancer. There are two key reasons that a shared follow-up care model between specialists and GPs is not routinely implemented for lymphoma survivors across Australia. Firstly, randomized controlled trials (RCT) evidence for the effectiveness and cost-effectiveness of a

shared-care model between specialists and GPs for lymphoma are *not* yet established (10). Secondly, a number of barriers to shared-care between the acute cancer care setting and GPs have been reported (11, 12). These include, but are not limited to, the lack of a coordinator who drives a shared model involving multiple providers, lack of patient and provider knowledge about the benefits of shared-care and how to implement it, insufficient or delayed communication between cancer specialists and GPs, and lack of awareness of available support such as funding models, tools and resources (11-13). These barriers might be overcome if a specialist cancer nurse (SCN) advises stakeholders of the benefits of shared-care (patient and GPs) and facilitates effective and timely care coordination and communication by acting as the conduit between the specialist cancer multidisciplinary team and the GPs at key transition time points (such as completion of treatment) (14).

Objectives {7}

The objective of the study is to test the feasibility of a prospective, pragmatic, randomized controlled trial (RCT) of the *GOSPEL* /intervention - an integrated, shared-care model involving hematologists and GPs for lymphoma follow-up.

Trial design {8}

This phase II pilot RCT aims to assess the feasibility of a larger definitive clinical trial. Outcome data will be collected at three points: (t_1) baseline (at enrolment); (t_2) 6 months; and (t_3) 12 months.

Methods: Participants, Interventions And Outcomes

Study setting {9}

This study is to be conducted in a large, Australian metropolitan tertiary teaching hospital and general practices.

Eligibility criteria {10}

Due to the varying follow-up requirements for patients with lymphoma the study population consists of two groups of patients who require two distinct follow-up pathways. Group one (post-treatment follow up) consists of patients with a histopathologically-confirmed diagnosis of aggressive or indolent lymphoma in the acute post-treatment phase (i.e., within three months of completion of chemotherapy). Group two (Observation follow-up) consists of patients with a histopathologically-confirmed diagnosis of indolent lymphoma followed up in the surveillance clinic who are at least 2 years post-treatment or treatment naïve. In addition to meeting one of the group descriptions above, participants must also meet all of the following criteria to be eligible for inclusion: ≥ 18 years of age, have an Eastern Cooperative Oncology Group (ECOG) performance status < 2 , be an ambulatory outpatient at the time of recruitment, be able to

nominate a GP or GP clinic to be involved in their follow-up, have access to a telephone, be able to speak and read English. Patients meeting any of the following criteria are excluded: the presence of severe mental, cognitive or physical conditions that would limit the patient's ability to participate as per treating clinician, lymphoma not in remission (applicable for group 1 only).

Who will take informed consent? {26a}

A Research Nurse will obtain informed consent.

Additional consent provisions for collection and use of participant data and biological specimens {26b}

Not applicable. This trial does not involve collecting additional participant data or biological samples for storage.

Interventions

Explanation for the choice of comparators {6b}

With the ever-growing population of cancer survivors, the specialist model of follow-up is insufficient and unsustainable. The shared-care model between specialists and GPs focusses on the complex care needs of lymphoma survivors, encompassing the strengths and expertise of multiple providers (10). It has the potential to contribute to better patient health outcomes as well as reducing the strain on overcrowded hospital-based outpatient services. Effective shared care in the hematology setting relies on establishing and maintaining ongoing communications channels (10). It is also proposed that transition to primary care occurs via a specialist nurse to ensure key components of survivorship care are addressed and communicated to health care providers (15). These key considerations have been incorporated in the intervention described below.

Intervention description {11a}

Arm 1 - Participants who are randomized to the usual care arm will receive standard follow-up care plus a survivorship booklet on "Living Well After Treatment – A guide for patients and families" published by Leukaemia Foundation (16). This booklet was designed by our research team in collaboration with expert clinicians and consumers via the Leukaemia Foundation. The current follow-up arrangement is a specialist-led model as determined by the treating Hematologist.

Arm 2 – Gospel 1. Participants who are randomized to the GOSPEL I arm will receive a multi-faceted intervention that includes a pre-specified shared-care pathway for follow-up. The design of the GOSPEL I is informed by a number of clinical guidelines [i.e., National Comprehensive Cancer Network (17) and European Society of Medical Oncology (18-20)], the Optimal Care Pathway for HL and Diffuse Large B-Cell Lymphoma (DLBCL) (21) the self-efficacy model (22, 23), and the Capabilities for Supporting Prevention and Chronic Condition Self-Management framework.(24) Table 1 outlines the active ingredients of the intervention.

After enrolment, participants in the intervention arm will receive a consultation with a Specialist Cancer Nurse to provide a treatment summary, the shared follow-up care appointment schedule, survivorship patient education (including the survivorship booklet on “Living well After Treatment” published by the Leukaemia Foundation)(16) and co-develop a draft Survivorship Care Plan (SCP) with the patient. The SCP will include up to three SMART (Specific, Measurable, Achievable, Realistic and Timely) goals using motivational interviewing and self-efficacy techniques. The treatment summary and draft SCP will be provided to the GP.

Within four weeks of the Specialist Cancer Nurse consultation, a case-conference at a mutually agreed time (maximum of 40 minutes) between the Specialist Cancer Nurse and the patient’s nominated GP will be completed to communicate the treatment summary, negotiate responsibilities of the shared-follow-up care schedule and negotiate the GP’s role in facilitating the SCP goals. The GP may propose changes or express if they are unable to take part in specific care activities outlined in the SCP. The finalized SCP will be filed in the patient’s medical records and provided to the patient and the GP. As a safety measure the SCN will ensure that surveillance activities are conducted as per clinical guidelines. Where the GP is unable to complete any of these activities, the SCN will ensure these surveillance activities are transferred to the acute cancer center/specialist. This will be documented as a process measure for future studies in this area.

The GP will also be provided with an escalation pathway which utilizes the SCN as the immediate access point for rapid re-entry to the acute setting and point of contact for the GP for support and resources. The SCN thus plays a key role in providing resources to support the GP +/- practice nurse in carrying out all activities outlined in the SCP.

Table 1 Active Ingredients of GOSPEL I model of care intervention for patients allocated to receive the GOSPEL 1 intervention.

Active Ingredient	Personnel Involved	Activities
<p>SCN-led clinic</p> <p>Duration: 60 min</p> <p>Mode: Teleconference or face to face</p>	<p>Specialist Cancer Nurse and the patient</p>	<ul style="list-style-type: none"> · Treatment summary including follow-up appointment schedule · Co-developing the SCP (including planning for health goals) · Post-treatment education <p>Note: The completed draft SCP will be sent to the GP prior to the case-conference. The research team or SCN will organize the case conference with the GP.</p>
<p>GP Case conference</p> <p>Duration: Maximum 40 mins</p> <p>Mode: Teleconference</p>	<p>SCN, GP (± one more healthcare professional).</p>	<ul style="list-style-type: none"> · The SCN will present the Treatment Summary, and the draft Survivorship Care Plan. · The SCN will negotiate follow-up responsibilities with the GP and will · Answer any questions the GP may have. · Additional education and support to the GP +/- Practice Nurse if applicable regarding physical examination and blood analysis. <p>Note: A copy of the completed and agreed Survivorship Care Plan provided to the GP, the patient and scanned for leMR. Where a full case conferencing is not possible, all efforts will be made to facilitate a teleconference of a shorter duration (~5 minutes) to delivery all key information on the Survivorship Care Plan</p>
<p>Standardized shared follow-up care</p> <p>Mode: Face to face</p>	<p>GP, SCN, Patient, Hematologist</p>	<p>Note: The GP will be given the direct telephone number of the SCN responsible for the patient. At any time if the GP becomes concerned about the patient, he/she can ring the SCN for advice or request escalation to acute care for review.</p>

Abbreviations: SCN, specialist cancer nurse; SCP, Survivorship Care Plan; GP, General Practitioner; leMR, integrated electronic medical records; FBC, full blood count; U&E, urea and electrolytes; LFT, liver function tests; LDH, lactate dehydrogenase

Criteria for discontinuing or modifying allocated interventions {11b}

The presence of any of the following criteria constitutes cause for the withdrawal of the participant: altered mental capacity resulting in inability to provide continuing informed consent; notification from treating oncologist and or GP that participant is not deemed to have capacity to consent; recurrence or progressive disease or death.

Strategies to improve adherence to interventions {11c}

Fidelity of the intervention will be assessed using the framework for behavioral interventions recommended by NIH (25, 26) as outlined in Table 2.

Table 2 Framework for behavioral interventions recommended by National Institute of Health

Goal	Strategies
Provider requirements	<p>Intervention nurses must be SCNs. SCNs work in dedicated cancer services and are primarily responsible for care of people at a specific phase or across all phases of the cancer journey, or work in a broader context but provide a specialist resource in cancer control to a range of generalist providers (for example, a cancer nurse coordinator). SCNs meet the minimum standard required for specialist practice in cancer nursing as set out in the competency standards from Cancer Australia⁽²⁷⁾. Therefore, the nurse delivering the GOSPEL I intervention requires the critical thinking, coordination and collaboration competencies defined in the SCN competency standards.</p>
Training Providers	<p>Training will be provided to the SCNs to ensure standardization of intervention delivery. Provision of a study manual to all SCNs which includes:</p> <ul style="list-style-type: none"> · Generic study related information: standard operating procedures, study overview, reporting/documentation guidelines, communication flowchart, rationale for the study treatment, completion of survivorship care plan, self-management goal setting and health coaching (including motivational interviewing) resources · Interventionist specific information: Job description, intervention protocol, quality assurance and monitoring processes <p>Completion of the eviQ Cancer Survivorship Introductory Course (~4.5 hours over 6 modules). This course was developed by the Australian Cancer Survivorship Centre in collaboration with Cancer Australia, Queensland University of Technology and the University of Sydney and is available online free of charge.</p> <p>Completion of face-to-face training from the research team which includes self-management support in cancer care, motivational interviewing techniques, setting SMART goals. Intervention-specific procedures required for this trial. Education and resources regarding MBS item numbers that facilitate the proposed Model of Care. Resources to support GPs in performing physical examination and blood analysis.</p>
Delivery of Intervention	<p>Intervention procedures are monitored through completion of intervention component checklists to ensure that the intervention is delivered as intended. Intervention checklists are completing during the SCN-led clinic and GP Case Conferences to track protocol deviations. All Nurse-Led clinics will be recorded and checked by another member of the research team against the clinic checklist to allow protocol deviation tracking across interventionists and conditions.</p> <p>Minimizing contamination between conditions by training SCNs to address participant questions about randomization and their assigned condition using non-biased explanations.</p> <p>SCNs will be supported during a weekly 15-30-minute meeting for the first 3 months of the trial between the SCN and CIA or Project Manager. In addition to ongoing troubleshooting and support, intervention fidelity will be closely monitored and will be discussed during this meeting.</p>
Receipt of Intervention	<p>The SCP serves as a resource for a participant to understand and refer to whenever they are unsure of follow-up schedule and collaborative goal setting.</p>
Enactment of Treatment Skills	<p>Enactment of treatment skills includes processes to monitor and improve participant ability to perform treatment-related behavioral skills and cognitive strategies in relevant real-life settings as intended. This goal will be achieved by:</p> <ul style="list-style-type: none"> - ensuring participants are aware of the follow up schedules and responsibilities of all health professionals

- ensuring participants will have a copy of the completed self-management care plan including all care responsibilities and goals set for the individual

Relevant concomitant care permitted or prohibited during the trial {11d}

No concomitant care or intervention is prohibited during the trial.

Provisions for post-trial care {30}

There is no specified ancillary or post-trial care for participants in this trial. However, it is expected that the SCP generated will have the value of informing longer term updates of the SCP and future survivorship care.

Outcomes {12}

The feasibility outcomes are recruitment and acceptability of the intervention based on completion rates and semi-structured interviews. Participants or healthcare providers who opt into the 12-month semi-structured interview will be interviewed either face-to-face, by telephone or through videoconferencing as per interviewee preference.

A range of patient-reported, and process outcome measures will also be collected. Health-related quality of life as measured by the Functional Assessment of Cancer Therapy - Lymphoma (FACT-Lym) will be collected at baseline, 6 and 12 months post enrolment (28). This valid and reliable tool captures key domains of health-related quality of life relevant to lymphoma and key symptoms that are relevant to the study population and sensitive to the GOSPEL 1 intervention. Additional outcomes include a range of patient-reported and process outcomes related to implementation as shown in Table 3

Participant timeline {13}

Table 3 Schedule for data collection during the GOSPEL 1 trial

	Study Period					
	Enrolment	Allocation	Post-allocation	Close-out		
Timepoint	$-t_1$	0	t_1	t_2	t_3	t_x
Enrolment						
Eligibility screen	X					
Informed consent	X					
Allocation		X				
Interventions						
GOSPEL 1						
Usual Care						
Outcomes						
Recruitment (feasibility)			X			
Barriers and facilitators of care: optional interviews (feasibility)						
				X		
Satisfaction of care						X
HRQoL (FACT-Lym)			X	X		X
Patient experience of care			X	X		X
Symptom Distress			X	X		X
Comorbidity burden			X			X
Dietary intake			X	X		X
Physical activity			X	X		X
Financial distress			X	X		X
Employment interference			X	X		X
Participant Characteristics						
Demographics			X			
Clinical characteristics			X			
Process Outcomes						
Intervention fidelity: completion of						X

checklists	
Clinical encounters at cancer center	X
Cost analysis: resources to conduct the intervention	X
Safety indicators: clinical encounters, unscheduled clinic visits and rapid referrals back to acute care	X

t₁: baseline, t₂: 6 months, t₃: 12 months

Sample size {14}

In this pilot study, we will recruit 30 patients per arm in order to provide initial insights into the intervention feasibility and protocol as well as preliminary effect size estimates. The aim of this study is not hypothesis testing, the power level is therefore not a valid consideration for sample size.(29, 30) The sample size for this study (n=60) falls within the range of sample size recommendations for pilot studies of this nature.(29, 30)

Recruitment {15}

Potentially eligible patients will be identified by clinicians and/or the research nurse through attendance at multidisciplinary team meetings and utilizing an existing clinical database. Potentially eligible patients will be identified, the treating clinician will ask the potential participant whether they agree to being contacted by the research nurse and will advise the research team accordingly. A patient brochure has been produced for clinicians to use when first discussing the study with potential participants. Only patients who have agreed to be contacted will be approached by the research nurse. Participants are given as much time as possible to consider their participation and are encouraged to take the information away and discuss joining the trial with family, friends and their GP if they so wish to. Participants are also encouraged to ask the research nurses, their treating doctors or nursing staff any questions in relation to their participation

Assignment of interventions: allocation

Sequence generation {16a}

Permuted block randomization (using blocks of 4 and 8) will be conducted to assign participants to the control or intervention arms. To ensure equal distribution of patients with different follow-up schedules, patients will be stratified by diagnosis (NHL vs HL) and follow-up pathway (group one: post-treatment vs group two: surveillance clinic).

Concealment mechanism {16b}

Allocation sequence is implemented using sequentially numbered opaque, sealed envelopes. Envelopes are only accessed by the research nurse to randomize the patient once recruitment and baseline data has been collected.

Implementation {16c}

Allocation sequence is generated by a researcher not involved in recruitment or data collection. Patients are enrolled by a research nurse who collects baseline data prior to randomization. Enrolling nurses assign participants to the intervention after baseline data collection.

Assignment of interventions: Blinding

Who will be blinded {17a}

After assignment to the intervention, only outcome assessors and data analysts are blinded to group allocation. Where participants opt to complete their data collection by phone, they are advised not to reveal their group allocation to the outcome assessor. Due to the nature of the intervention, no participants or treating clinicians are blinded.

Procedure for unblinding if needed {17b}

No unblinding procedures required as only outcome assessors and data analysts are blinded.

Data collection and management

Plans for assessment and collection of outcomes {18a}

Patient-reported outcomes are self-administered using online surveys or administered in person or via telephone with an outcome assessor trained in the administration of the study instruments.

A range of feasibility, patient-reported, and process outcomes related to implementation will be collected as shown in Table 3.

- Barriers and facilitators of care as measured by voluntary participation in semi structured interviews 12 months post enrolment. All intervention arm patients, lymphoma cancer nurses, GPs, and other nurses, and hospital- and community-based rehabilitation providers will be invited to participate in a one off semi-structured interview to discuss factors that facilitated or hindered the implementation of the GOSPEL I intervention
- Health-related Quality of Life as measured by The Functional Assessment of Cancer Therapy - Lymphoma (FACT-Lym) will be collected at baseline, 6- and 12-months post enrolment (28). This valid and reliable tool captures key domains of Health-related quality of life relevant to lymphoma and key symptoms that are relevant to the study population and sensitive to the GOSPEL 1 intervention
- Patient experience of care as measured by the Patient Assessment of Care for Chronic Conditions (PACIC). This is a 20-item instrument derived from the '5As' model, a patient-centered model of behavioral counselling congruent with the Chronic Care Model and has been frequently used to enhance self-management support and linkages to community resources.
- Symptom distress as measured by the Memorial Symptom Assessment Scale (MSAS) (31). This is valid and reliable in measuring symptom prevalence, characteristics and distress in patients with cancer (31).
- Comorbidity burden as measured by The Charleston Comorbidity Index (CCI) supplemented with items from Self-Administered Comorbidity Questionnaire (SCQ) (32).
- Physical activity as measured by the Active Australia Survey (33) which is designed to measure participation in leisure-time physical activity, and a single item from the International Physical Activity Questionnaire (34) will be used to measure sedentary behaviors.
- Dietary behaviors, specifically usual vegetable intake and usual fruit intake, as measured by two short dietary questions from the National Nutrition Survey (35), which have been validated in the Australian population. Both questions discriminate between groups with significantly different fruit and vegetable intakes. In administering these questions, information about which foods are included as vegetables and fruits are provided and serve sizes are described.
- Financial distress as measured by a 0-10 numerical analogue scale (where 10 is "a great deal" and 0 is "none") describing financial distress as a result of cancer and treatment (36)
- Employment interference as measured by a 0-10 numerical analogue scale (where 10 is "a great deal" and 0 is "none") describing employment interference as a result of cancer and treatment (37)
- Satisfaction of care as measured by the 0-10 numerical analogue scale (with 10 being the most satisfied) supplemented with short, structured qualitative questions.
- Process outcomes, including completion of intervention components, as measured by completion of Intervention materials such as clinic checklists, audio recordings of SCN-led clinics and Research

Nurse (RN) records in database will be checked for completion (Y/N) and duration (min) of SCN-led clinics.

- Cost analysis will be conducted using hospital casemix data describing occasions of service including duration and indication, RN records in database. All resources required to conduct the intervention (e.g., staff, training, materials, communications, office space, utilities) will be monitored
- Safety indicators will be measured via hospital records to query the number of hospital clinical encounters, unscheduled lymphoma clinic visits and rapid referrals back to acute care

Plans to promote participant retention and complete follow-up {18b}

Participants who deviate from the protocol will not be withdrawn from the trial. Participants who withdraw from the trial nominate the degree to which they withdraw (i.e., whether they withdraw from active data collection ± passive data collection such as hospital records).

Data management {19}

All participant characteristic and outcome data are entered directly into REDCap (Research Electronic Data CAPture – Vanderbilt University, hosted at Queensland University of Technology) by the research nurse and the participants through self-administered online survey. To ensure data quality, the database is designed with branching logic, data validation and range checks for data values, where possible.

All source data, clinical records and laboratory data relating to the study will be archived at the clinical site as appropriate for 15 years after the completion of the study. All data will be available for retrospective review or audit. No study document will be destroyed without prior written agreement between the responsible organization and the investigator. If the investigator wishes to assign the study records to another party or move them to another location, he/she must notify the responsible organization in writing of the new responsible person and/or the new location.

Confidentiality {27}

Data on potential participants is recorded, including reasons for ineligibility or refusal to participate. Participants are only identified by a unique participant study number on the case report forms and other study documents. Other study-related documents (e.g., signed consent form, participant log) are kept in strict confidence by the Investigator.

Participants are informed that data is held on file by the responsible organizations and that these data may be viewed by staff including the study project manager and by external auditors on behalf of the responsible organizations and appropriate regulatory authorities (to include reviewing Human Research

Ethics Committee (HREC) and the Research Governance Officers). Participants data in publications and conference presentation reports will only be presented in aggregated form. All participant data will be held in strict confidence.

Plans for collection, laboratory evaluation and storage of biological specimens for genetic or molecular analysis in this trial/future use {33}

Not applicable. There is no collection of biological specimens in the current trial.

Statistical methods

Statistical methods for primary and secondary outcomes {20a}

Descriptive statistics will be used to report on feasibility and process-related elements (e.g., recruitment, intervention, retention rates) as well as clinical and resource outcomes. Preliminary effect size estimates for patient and resource use outcomes will be calculated following intention-to-treat principles using linear mixed models. Models will include group, time and their interaction and be adjusted by diagnosis and age. Balance of demographic variables between usual care and intervention group will be investigated using chi-square and t-test and will be included in the model if found to be both significantly associated with the outcome and confounding the intervention. Assumptions of all models (normality, linearity, homoscedasticity) will be examined using the residuals of the model and will be described using mean, median, skewness, kurtosis and plots such as histograms and QQ-plots. If assumptions are violated, models will be either bootstrapped or log transformation as appropriate.

Interim analyses {21b}

Not applicable. No interim analysis is planned.

Methods for additional analyses (e.g. subgroup analyses) {20b}

Patients allocated to the GOSPEL I arm will be invited to participate in an interview at the 12-month time point. Guiding questions and analysis of the interviews will be guided by the Consolidated Framework for Implementation Research (*CFIR*). Based on our previous qualitative work, we expect that the number of

interviews will be approximately 24 (GPs, patients, Hematologists). All interviews will be recorded and transcribed verbatim for analysis.

Methods in analysis to handle protocol non-adherence and any statistical methods to handle missing data {20c}

Any discrepancies and missing data will be alerted and resolved with the relevant research team member(s) as soon as practical. All electronic CRFs will be maintained on the system with details of any changes logged accordingly. Preliminary effect size estimates for patient and resource use outcomes will be calculated following intention-to-treat principles using linear mixed models. Patterns of missing data will be examined using chi-square and t-tests. Missing data for the outcomes will be accounted for by using mixed models allowing the use each available case by computing maximum likelihood estimates.

Plans to give access to the full protocol, participant level-data and statistical code {31c}

Not applicable. There are no plans for granting public access of the full protocol, participant level dataset or statistical code

Oversight and monitoring

Composition of the coordinating centre and trial steering committee {5d}

The Chief investigators are the trial steering committee that will provide all governance to the conduct of the study.

Composition of the data monitoring committee, its role and reporting structure {21a}

Not applicable. There is no data monitoring committee established for this pilot trial.

Adverse event reporting and harms {22}

An adverse event (AE) is any event, side effect, or other untoward medical occurrence that occurs in conjunction with the use of the study intervention in humans, whether or not considered to have a causal relationship to the interventions. An AE can, therefore, be any unfavorable and unintended sign (that could

include a clinically significant abnormal laboratory finding), symptom, or disease temporally associated with the use of the study intervention, whether or not considered related to the intervention. Conditions recognized as being excluded from AE reporting are as follows: any event, side effect, or other medical occurrence that is anticipated because of the normal course of treatment (standard care). There are no known side effects/adverse events associated with the proposed model of care intervention (38). Due to the nature of this intervention, there will be no reporting of AE.

Frequency and plans for auditing trial conduct {23}

There are no plans for auditing trial conduct beyond the independent research governance requirements and annual reporting to the HREC.

Plans for communicating important protocol amendments to relevant parties (e.g. trial participants, ethical committees) {25}

All agreed protocol amendments are clearly recorded on a protocol amendment form and are signed and dated by the original protocol approving signatories. All protocol amendments will be submitted to the institutional HREC for approval before implementation. The only exception will be when the amendment is necessary to eliminate an immediate hazard to the trial participants. In this case, the necessary action will be taken first, with the relevant protocol amendment following shortly thereafter. Once HREC approval has been granted, investigators and the ANZCTR will be updated

Dissemination plans {31 a}

It is intended that the findings from this trial will be disseminated at academic and professional conferences and via a manuscript submission to a peer-reviewed journal. Participants will be identified in such reports only in aggregate or by study identification number, gender and age. There are no publication restrictions.

Discussion

Despite the strong case for shared, follow-up care model for lymphoma survivors involving cancer specialists and GPs, barriers to shared-care mean that it is not routinely implemented across Australia. These include the need for coordination across multiple providers, the need for improved patient and provider knowledge about the benefits of shared-care and how to implement it, insufficient or delayed communication between cancer specialists and GPs, and lack of awareness of available support such as funding models, tools and resources (11, 12). Facilitated by the Specialist Cancer Nurse the current study aims to help stakeholders (patient, hematologist and GP's) realize the benefits of shared-care (patient and

GPs), facilitating effective and timely care coordination, and acting as the conduit between the specialist cancer multidisciplinary team and the GPs.

Practical issues for this trial include estimating the time required to coordinate the trial across multiple providers including engaging GPs and fidelity with the intervention components. The proposed study will provide important information on the feasibility of a definitive phase 3 trial to advance the evidence to support an integrated model of care that will ultimately optimize outcomes for patients with lymphoma. The information collected through the trial, qualitative interviews and economic evaluations are crucial in guiding the development of such a trial.

Trial status

The protocol published here is version 1.3 dated 31 March 2020. The trial has not yet commenced recruitment. Estimated start date of recruitment is 13th July 2020. Estimated end date of recruitment is 13th July 2021.

Trial registration: Australia and New Zealand Clinical Trials Registry, Registration number: ACTRN12620000594921, Date registered: 22/05/2020, registration link: <https://www.anzctr.org.au/ACTRN12620000594921.aspx>

Abbreviations

AE	Adverse event
CCI	Charleston Comorbidity Index
CRF	Case report form
ECOG	Eastern Cooperative Oncology Group
FACT-Lym	Functional Assessment of Cancer Therapy - Lymphoma
HREC	Human Research Ethics Committee
HRQoL	Health-related quality of life
HL	Hodgkin's Lymphoma
ieMR	Integrated electronic medical record
IPAQ	International Physical Activity Questionnaire
MSAS	Memorial Symptom Assessment Scale
NHL	Non-Hodgkin's Lymphoma
PACIC	Patient Assessment of Care for Chronic Conditions
PICF	Patient Information and Consent Form
QUT	Queensland University of Technology
RCT	Randomised Controlled Trial
REDCap	Research Electronic Data Capture
RN	Research Nurse
SCP	Survivorship Care Plan
SCQ	Self-Administered Comorbidity Questionnaire
T1	Timepoint 1 (baseline)
T2	Timepoint 2 (6 months)
T3	Timepoint 3 (12 months)
T4	Timepoint 4 (24 months)
Y/N	Yes/No

Declarations

Acknowledgements

The authors acknowledge support from service executives including Professor Euan Walpole, Adj Associate Professor Leanne Stone, and Dr Doreen Tapsall.

Authors' contributions {31b}

RC is the chief investigator; he conceived the study, secured funding, and led the proposal and protocol development. SM, SB, CS, LT, LG, FH, JR contributed to study design and to the development of the protocol. JT is responsible for training of intervention nurses. CS, SB are responsible for delivering the intervention. CS, RJ and OAA are responsible for data collection and administrative support to the operationalization of the trial. All authors have provided input and have read and approved the final manuscript.

Funding {4}

This study is funded by a Health Innovation, Investment and Research Office (HIIRO) Queensland Advancing Clinical Research Fellowship. The funding body had no role in the design of the study and will not have a role in collection, analysis, and interpretation of data or in writing the manuscript.

Availability of data and materials {29}

There are no limitations on investigator access to the trial dataset. The datasets generated and/or analyzed during the current study are not going to be made publicly available but will be made available from the corresponding author on reasonable request.

Ethics approval and consent to participate {24}

This study is approved by the Metro South Hospital and Health Services Human Research Ethics Committee (HREC/2020/QMS/61872). Written informed consent will be obtained from all participants.

Consent for publication {32}

Not applicable. No details, images or videos relating to an individual person will be published, as all data will be presented in aggregate.

Competing interests {28}

The authors declare that they have no competing interests.

References

1. Australian Institute of Health and Welfare. Cancer in Australia 2017. Canberra: AIHW; 2017.
2. Ng HS, Roder D, Koczwara B, Vitry A. Comorbidity, physical and mental health among cancer patients and survivors: An Australian population-based study. *Asia Pac J Clin Oncol*. 2017.
3. Commonwealth of Australia. National Primary Health Care Strategic Framework. Canberra, ACT.; 2013.
4. Emery J. Cancer survivorship—the role of the GP. *Aust Fam Physician*. 2014;43(8):521-5.
5. Emery JD, Shaw K, Williams B, Mazza D, Fallon-Ferguson J, Varlow M, et al. The role of primary care in early detection and follow-up of cancer. *Nat Rev Clin Oncol*. 2014;11(1):38-48.
6. Cancer Australia. Influencing best practice in breast cancer. 2016.
7. DoH Victoria. Optimal care pathway for women with breast cancer.
8. Der Vartanian C, Anderson K, Evans K, Anderiesz C, Giles C, Zorbas H. A new model supporting best practice follow-up care for early breast cancer in Australia: shared follow-up care for early breast cancer. *Cancer Survivorship 2017; South Australia*.2017.
9. Chan A, Koczwara B, Fitch M, Ashbury F, Chan R. Cancer survivors care during COVID-19 – perspectives and recommendations from the MASCC survivorship study group. *Supportive Care in Cancer*. 2020;Inpress.
10. Taylor K, Chan RJ, Monterosso L. Models of survivorship care provision in adult patients with haematological cancer: an integrative literature review. *Support Care Cancer*. 2015;23(5):1447-58.
11. Hudson SV, Miller SM, Hemler J, Ferrante JM, Lyle J, Oeffinger KC, et al. Adult cancer survivors discuss follow-up in primary care: 'not what i want, but maybe what i need'. *Ann Fam Med*. 2012;10(5):418-27.
12. Neuman HB, Jacobs EA, Steffens NM, Jacobson N, Tevaarwerk A, Wilke LG, et al. Oncologists' perceived barriers to an expanded role for primary care in breast cancer survivorship care. *Cancer Med*. 2016;5(9):2198-204.
13. Jefford M, Koczwara B, Emery J, Thornton-Benko E, Vardy JL. The important role of general practice in the care of cancer survivors. *Aust J Gen Pract*. 2020;49(5):288-92.
14. Chan RJ, Downer TR. Cancer Nurses Can Bridge the Gap Between the Specialist Cancer Care and Primary Care Settings to Facilitate Shared-Care Models. *Cancer Nurs*. 2018;41(2):89-90.
15. Cooper JM, Loeb SJ, Smith CA. The primary care nurse practitioner and cancer survivorship care. *J Am Acad Nurse Pract*. 2010;22(8):394-402.
16. Leukaemia Foundation. Living Well After Treatment - A guide for patients and families: Leukaemia Foundation; 2018.
17. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology [Available from: https://www.nccn.org/professionals/physician_gls/default.aspx#site.

18. Dreyling M, Ghielmini M, Rule S, Salles G, Vitolo U, Ladetto M. Newly diagnosed and relapsed follicular lymphoma: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol.* 2017;28(12):3109.
19. Eichenauer DA, Aleman BMP, Andre M, Federico M, Hutchings M, Illidge T, et al. Hodgkin lymphoma: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol.* 2018;29(Suppl 4):iv19-iv29.
20. Vitolo U, Seymour JF, Martelli M, Illerhaus G, Illidge T, Zucca E, et al. Extranodal diffuse large B-cell lymphoma (DLBCL) and primary mediastinal B-cell lymphoma: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol.* 2016;27(suppl 5):v91-v102.
21. Cancer Council. Optimal cancer care pathway for people with Hodgkin and diffuse large B-cell lymphomas 2019 [updated September 17, 2019. Available from: <https://www.cancer.org.au/health-professionals/optimal-cancer-care-pathways.html>.
22. Bandura A. *Social Foundation of Thought and Action*: Prentice-Hall; 1986.
23. Bandura A, Adams NE, Beyer J. Cognitive processes mediating behavioral change. *Journal of personality and social psychology.* 1977;35(3):125-39.
24. DoHA, Flinders University. *Capabilities for Supporting Prevention and Chronic Condition Self-Management*. Canberra: Department of Health and Ageing and Flinders University; 2009.
25. Bellg AJ, Borrelli B, Resnick B, Hecht J, Minicucci DS, Ory M, et al. Enhancing treatment fidelity in health behavior change studies: best practices and recommendations from the NIH Behavior Change Consortium. *Health psychology : official journal of the Division of Health Psychology, American Psychological Association.* 2004;23(5):443-51.
26. Robb SL, Burns DS, Docherty SL, Haase JE. Ensuring treatment fidelity in a multi-site behavioral intervention study: implementing NIH Behavior Change Consortium recommendations in the SMART trial. *Psycho-oncology.* 2011;20(11):1193-201.
27. Aranda S, Yates P. *A National Professional Development Framework for Cancer Nursing (EdCaN)*. Canberra: Cancer Australia; 2009.
28. Hlubocky FJ, Webster K, Cashy J, Beaumont J, Cella D. The Development and Validation of a Measure of Health-Related Quality of Life for Non-Hodgkin's Lymphoma: The Functional Assessment of Cancer Therapy's Lymphoma (FACT-Lym). *Lymphoma.* 2013;2013:9.
29. Hertzog MA. Considerations in determining sample size for pilot studies. *Research in nursing & health.* 2008;31(2):180-91.
30. Julious S. Sample size of 12 per group rule of thumb for pilot study. *Pharmaceutical Statistics.* 2005;4:287-91.
31. Portenoy RK, Thaler HT, Kornblith AB, Lepore JM, Friedlander-Klar H, Kiyasu E, et al. The Memorial Symptom Assessment Scale: an instrument for the evaluation of symptom prevalence, characteristics and distress. *European journal of cancer (Oxford, England : 1990).* 1994;30a(9):1326-36.

32. Sangha O, Stucki G, Liang MH, Fossel AH, Katz JN. The Self-Administered Comorbidity Questionnaire: a new method to assess comorbidity for clinical and health services research. *Arthritis and rheumatism*. 2003;49(2):156-63.
33. Australian Institute of Health and Welfare. *The Active Australia Survey: a guide and manual for implementation, analysis and reporting*. . Canberra: AIHW; 2003.
34. Craig CL, Marshall AL, Sjostrom M, Bauman AE, Booth ML, Ainsworth BE, et al. International physical activity questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc*. 2003;35(8):1381-95.
35. Rutishauser I, Webb K, Abraham B, Allsopp R. *Evaluation of short dietary questions from the 1995 National Nutrition Survey*. Canberra: Australian Government Department of Health and Ageing; 2001.
36. Chan R, Cooper B, Koczwara B, Gordon L, Chan A, CJ T, et al. Phenotypic and symptom characteristics associated with inter-individual variability in financial distress in patients with breast cancer: a longitudinal study. *Cancer Nurs*. 2020;Inpress.
37. Chan RJ, Cooper B, Koczwara B, Chan A, Tan CJ, Paul SM, et al. A longitudinal analysis of phenotypic and symptom characteristics associated with inter-individual variability in employment interference in patients with breast cancer. *Support Care Cancer*. 2020.
38. Grunfeld E, Levine MN, Julian JA, Coyle D, Szechtman B, Mirsky D, et al. Randomized trial of long-term follow-up for early-stage breast cancer: a comparison of family physician versus specialist care. *J Clin Oncol*. 2006;24(6):848-55.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [61872EthicalClearance2.pdf](#)