

ICD-11 Complex Posttraumatic Stress Disorder and Subclasses of Borderline Personality Disorder in a Korean General Population Sample with Childhood Abuse Experiences: A Latent Class Analysis

Jisu Lee

Chungbuk National University

Hyunjung Choi (✉ hchoi@g.cbnu.ac.kr)

Chungbuk National University

Research Article

Keywords: complex posttraumatic stress disorder, borderline personality disorder, latent class analysis, pathological personality traits, childhood abuse

Posted Date: September 26th, 2023

DOI: <https://doi.org/10.21203/rs.3.rs-3342037/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Additional Declarations: No competing interests reported.

Version of Record: A version of this preprint was published at Borderline Personality Disorder and Emotion Dysregulation on December 15th, 2023. See the published version at <https://doi.org/10.1186/s40479-023-00242-z>.

Abstract

Background Complex posttraumatic stress disorder (CPTSD) and borderline personality disorder (BPD), which are distinctive diagnoses, share the common risk factor of childhood abuse experiences. However, additional evidence is needed to determine which factors contribute to the manifestation of different symptoms.

Method Participants were 499 Korean adults sampled from an online panel of a general population sample who reported experiences of childhood abuse. A latent class analysis (LCA) was conducted to identify the patterns of CPTSD and BPD symptoms. We adopted a three-step LCA to compare types of childhood abuse, invalidating environments, attachment styles, and pathological personality traits among different classes.

Results The LCA revealed four classes: a CPTSD and BPD “comorbid” class, an “externalizing BPD” class, an “avoidant BPD” class, and a low symptom class. The “comorbid” class showed the highest scores in all symptoms and risk factors. The “externalizing BPD” class was distinguished from the “avoidant BPD” class by the externalizing versus internalizing associated pathological personality traits. The “avoidant BPD” class experienced high emotional neglect in addition to other types of abuse and it also showed an additional avoidant attachment style.

Conclusion Childhood abuse may heighten the risk for high comorbidity of CPTSD and BPD as well as externalizing-internalizing subgroups of BPD. Beyond the identification of CPTSD and BPD, assessing attachment styles and pathological personality traits based on dimensional approaches would benefit the tailoring of effective treatment.

Background

Childhood abuse and neglect have been shown to serve as risk factors for a range of psychopathologies [1], including complex posttraumatic stress disorder (CPTSD) [2, 3] and borderline personality disorder (BPD) [4–6]. *The International Classification of Diseases* (ICD-11) CPTSD consists of posttraumatic stress disorder (PTSD) symptoms such as re-experiencing, avoidance, and heightened sense of threat, along with disturbances in self-organization (DSO) symptoms, which are characterized by affect dysregulation, negative self-concept, and disturbed relationships [7]. Meanwhile, BPD is characterized by pervasive dysregulation in relationships, emotions, cognition, and identity [8].

While the symptoms of BPD and DSO overlap with each other, recent discussions have agreed that CPTSD and BPD can be differentiated. Using an exploratory factor analysis, the ICD-11 PTSD, DSO, and BPD symptoms were shown to have distinct constructs [9, 10]. In a network analysis, CPTSD and BPD were clearly separated, as they only overlapped in terms of the affect dysregulation symptom [11]. In the results of a latent class analysis (LCA) conducted among women with childhood abuse histories, despite the presence of overlapping symptoms, BPD was distinguished from CPTSD [3]. CPTSD, PTSD, and BPD were all distinguished from each other in a latent profile analysis (LPA) among community sample

women with at least one potential lifetime trauma experience [12]. Studies have shown that, whereas DSO is related to emotional and interpersonal avoidance, BPD is more characterized by unstable interpersonal relationships and sense of self, impulsivity, and reactive emotional responses [3, 9]. Consistently, dissociative symptoms in CPTSD have referred to relatively prolonged emotional numbing, which are at higher levels than in BPD, while they are responses to temporary stress in BPD [12].

Some studies have shown that CPTSD and BPD can be highly comorbid. For example, in an LCA with a general population sample with sexual trauma, a distinctive CPTSD and a PTSD class was distinguished in addition to classes of those comorbid with BPD [13]. Other studies have found comorbid classes of CPTSD with varying symptom severity in BPD, where a distinguished class has not been found, both in a multiple traumatized treatment-seeking people [14] and a young people sample with lifetime trauma [15].

The results showed that CPTSD and BPD were distinctive, while the phenomena of the symptoms differed even for the similarities of the symptoms [3, 16]. However, no population characteristics of type of trauma emerged as distinctive risk factors for CPTSD and BPD. For instance, childhood interpersonal violence is a common risk factor for both CPTSD and BPD [3, 9], and most populations are multiply traumatized, because trauma experience itself may be a risk factor for further traumatization later in life [e.g., 13–15]. It is also necessary to understand the differential trajectories of how CPTSD and BPD develop. Therefore, in the present work, we focused on attachment styles and pathological personality traits based on the dimensional perspective of psychopathology [e.g., 17, 18] in people who have experienced childhood abuse.

Childhood abuse is a risk factor for insecure attachment, including both anxious and avoidant attachment. Attachment style has also been shown to mediate interpersonal problems and emotional maladjustment in later life [19–22]. Anxious attachment is characterized by both a strong desire to be close and concerns about being rejected in relationships. Meanwhile, avoidant attachment is associated with fear and avoidance of getting close to or depending on others [19]. The fear of abandonment that is commonly seen in BPD closely resembles the pattern of anxious attachment. A meta-analysis showed that BPD is more strongly associated with anxious than avoidant attachment [23]. The attachment style in CPTSD symptoms appear to be more complicated. In one study, childhood trauma-related CPTSD symptoms have been found to be associated with both anxious and avoidant attachment [24], while multiple studies have repeatedly confirmed the strong relationship between CPTSD and anxious attachment [10, 25–27].

Differences in the personality traits of CPTSD and BPD can be identified based on the Hierarchical Taxonomy of Psychopathology (HiTOP) model developed by Kotov and colleagues [17, 18]. The HiTOP directs a paradigm shift from a categorical model to the dimensional model of psychopathology and aim to identify a transdiagnostic variable to be targeted in treatment. This model explains the dimensional features related to PTSD and BPD according to the *Diagnostic and Statistical Manual of Mental Disorders* (DSM) [8]. PTSD was explained in terms of the internalizing spectrum, while BPD was classified along both the internalizing and antagonistic externalizing dimensions [18]. HiTOP contains spectra that

correspond to the five domains of the pathological personality traits presented in the Alternative Model for Personality Disorders (AMPD) in DSM-5 [8]. The AMPD concept has previously been used to understand not only personality disorders but also other mental disorders [28, 29]. In Møller et al, DSO symptoms showed strong correlations with the maladaptive personality trait domains of negative affectivity, detachment, and psychoticism [29]. Gamache and colleagues conducted an LPA using AMPD concepts as indicators and identified subprofiles of BPD along a dimension of severity and different categories of personality traits [30].

The current study aimed to investigate how the symptoms of CPTSD and BPD manifest in a generalized population sample with childhood abuse experiences. The attachment styles and pathological personality traits associated with each latent syndrome were compared. We also compared being raised in an invalidating environment which has been discussed as a risk factor for BPD involving neglect and mislabeling of a child's emotions, thoughts, and behaviors; reinforcement of maladaptive emotional expression; and a lack of opportunities to develop problem-solving strategies [31]. The research hypotheses are as follows: First, the CPTSD and BPD classes are expected to be distinguishable using LCA. Second, both CPTSD and BPD are expected to be highly related to child abuse and invalidating environments. Third, the CPTSD and BPD classes are each expected to show both high avoidant and anxious attachment, while the BPD class is expected to show less avoidant attachment. Finally, the two classes are expected to differ in terms of pathological personality traits. CPTSD is expected to show high scores in the negative affectivity domain corresponding to the internalizing spectra as well as the detachment domain in relation to chronic interpersonal avoidance. In particular, BPD is expected to be associated with the disinhibition domain.

Methods

Participants and procedures

The present study was approved by the Institutional Review Board of the authors' institution. Data were collected from an online panel of adults who reported having at least one childhood abuse experience. Stratified sampling was conducted while considering gender, age, and residence information [32] based on South Korean population data from Statistics Korea. The exclusion criteria were individuals who were under the age of 19, those who did not use Korean as their first language, and those with no experiences of childhood abuse. We excluded data based on a logical criterion that informs the validity of the sample, as suggested by Kramer and colleagues [33]. We considered data to be invalid when responding to extreme values in opposite directions to items with identical content regarding childhood abuse and an invalidating environment. Out of 14,220 people, 13,511 were excluded due to the absence of a history of childhood abuse, and invalid data from 172 individuals were excluded. An additional 171 data were collected, and a total of 499 participants were ultimately included.

There were 256 men (51.3%) and 243 women (48.7%) in the sample. In terms of age, participants were in their 20s through 50s, with the proportions being 22.2% in their 20s, 22% in their 30s, 27.5% in their 40s,

and 28.3% in their 50s. In terms of education, 107 people were high school graduates or lower (21.4%) whereas 392 had college degrees and above (78.5%). Regarding employment conditions, there were 330 regular workers (66.1%), 47 nonregular workers (9.4%), 34 self-employed workers (6.8%), 66 unemployed individuals (13.2%), and 22 students (4.4%).

Using the cutoff criteria [34], 31.7% of the sample reported physical abuse (PA), 28.3% reported emotional abuse (EA), 29.7% reported sexual abuse (SA), 28.5% reported physical neglect (PN), and 37.5% reported emotional neglect (EN).

Measures

International Trauma Questionnaire, ITQ

The International Trauma Questionnaire (ITQ) measures PTSD and DSO symptoms according to ICD-11 using a 5-point Likert scale. The reliability coefficient Cronbach's α was above .77 for all PTSD and DSO symptoms, except for the avoidance items [35]. Cronbach's α values for the Korean version were .92 for PTSD symptoms and .91 for DSO symptoms [36]; in the present study, they were .95 and .94, respectively.

Borderline Symptom List-short form, BSL-23

The Borderline Symptom List-short form (BSL-23) measures symptoms of BPD based on DSM-IV rated on a 5-point Likert scale [37]. The items capture symptoms such as self-perception, affect regulation, self-destruction, dysphoria, loneliness, intrusions, and hostility. We used the Korean version validated by Kang et al. [38]. Cronbach's α was .97 in the development study [39] and .98 in both Kang et al. [38] and the current study.

Child Trauma Questionnaire-Short Form, CTQ-SF

The Child Trauma Questionnaire-Short Form measures the frequency of five types of child abuse, including PA, EA, SA, PN, and EN, on a 4-point Likert scale [40]. The Cronbach's α values of the Korean version were .88 for PA, .82 for EA, .87 for SA, .68 for PN, and .86 for EN [41]. In the current study, the Cronbach's α values were .85 for PA, .84 for EA, .93 for SA, .67 for PN, and .92 for EN.

Invalidating Childhood Environment Scale, ICES

The Invalidating Childhood Environment Scale (ICES) assesses parental invalidation and invalidating family types on a 5-point Likert scale [42]. We used the parental invalidation scales from the Korean version [43]. The paternal and maternal Cronbach's α values were respectively .80 and .77 in the original version, .87 and .86 in Korean version, and .75 and .82 in the current study.

Experiences in Close Relationships-revised, ECR-R

The Experiences in Close Relationships-Revised (ECR-R) measures adult attachment styles [44]. The current study used the Korean version validated by Kim [45]. The ECR-R consists of 18 items for anxious attachment and 18 items for avoidant attachment, which are responded to on a 5-point Likert scale. In the

Korean version, the Cronbach's α values were .89 for anxious attachment and .85 for avoidant attachment; in the current study, they were .93 and .91, respectively.

Personality Inventory for DSM-5, PID-5

The Personality Inventory for DSM-5 (PID-5) measures the pathological personality traits presented in the DSM-5 AMPD [46]. We used the Korean version validated by Shin and Hwang [47], which consists of 25 facets organized into five domains (negative affectivity, detachment, antagonism, disinhibition, and psychoticism). In total, 220 items are rated on a 4-point Likert scale (0: not at all ~ 3: very much). The reliability coefficients for each domain were as follows: Negative Affectivity .93, Detachment .96, Antagonism .95, Disinhibition .84, Psychoticism .96 [46]. In the current study, the coefficients were .97, .93, .96, .94, and .96, respectively.

Analyses

An LCA, which is a structural equation modeling method that identifies unobserved structures in multivariate categorical data [48], was conducted using M plus 8.0. Each of the three symptoms of PTSD (reexperience, avoidance, and heightened sense of threat) and DSO (affect dysregulation, negative self-concept, and disturbed relationship) were scored as present and non-present according to the algorithm suggested by Cloitre and colleagues [35], where any of the two items from each symptom is rated 2 or higher, meaning that the symptom is present. The latent variable of BPD symptoms was composed of each item of the BSL-23, where symptoms were defined as present when the score was 2 or higher.

To identify the optimal model, we compared the Akaike's Information Criteria (AIC) [49], the Bayesian Information Criteria (BIC) [50], the Sample-Size Adjusted BIC (SSA-BIC) [51], and entropy values. When the AIC, BIC, and SSA-BIC values are smaller [52] and the entropy value is 0.8 or higher, the classification accuracy is considered to be good [53]. If the Lo-Mendell-Rubin Adjusted likelihood ratio test (LMR-A) and the Bootstrap Likelihood Ratio Test (BLRT) values are significant, then the k-level model is considered to be a better fit than the k-1 level model [54].

We used a three-step LCA to examine differences in the type of childhood abuse, invalidating environments, attachment style, and pathological personality traits across latent classes. A three-step method improves statistical power by using auxiliary observed variables to adjust classification errors in the parameter estimation process [55].

Results

Latent class analysis

The fit scores of each latent class model are presented in Table 1. We excluded the five-class model because the maximum likelihood was not repeatedly extracted in that model. The four-class model was selected, because it had the lowest values for BIC, SSA-BIC, and AIC, as well as significant relative fit indices.

Table 1
Latent class models and fit indices

Model	Log-likelihood	BIC	ssa-BIC	AIC	Entropy	LMR-A p-value	BLRT p-value
1 class	-9851.951	19884.068	19792.020	19761.902			
2 class	-6711.054	13788.652	13601.383	13540.109	0.979	0.000	0.000
3 class	-6184.318	12921.558	12639.067	12546.636	0.958	0.000	0.000
4 class	-5951.737	12642.773	12265.061	12141.473	0.976	0.003	0.000
5 class	-5768.696	12463.071	11990.137	11835.393	0.963	0.013	0.000
<p>Note. BIC = Bayesian Information Criterion; ssa-BIC = sample-size adjusted BIC; AIC = Akaike Information Criterion; LMR-A = Lo–Mendell–Rubin Adjusted likelihood ratio test; BLRT = Bootstrap Likelihood Ratio Test.</p> <p>*5 class was excluded because the best log-likelihood value was not replicated.</p>							

Table 1 Latent class models and fit indices

Each class of symptom patterns is presented in Fig. 1, and the conditional response probabilities are presented in Table 2. If the response probabilities are above 70% or below 30%, the consistency within that class was considered to be high [56].

Table 2
Conditional response probabilities in each class

Symptoms		Class 1	Class 2	Class 3	Class 4
		Comorbid (<i>n</i> = 189)	Externalizing BPD (<i>n</i> = 53)	Avoidant BPD (<i>n</i> = 137)	Low Symptoms (<i>n</i> = 120)
PTSD 1	Re-experiencing	0.91	0.13	0.30	0.06
PTSD 2	Avoidance	1.00	0.11	0.38	0.12
PTSD 3	Sense of threat	0.97	0.08	0.49	0.10
DSO 1	Affect dysregulation	1.00	0.59	0.60	0.16
DSO 2	Negative self-concept	0.99	0.54	0.46	0.10
DSO 3	Disturbed relationship	0.97	0.61	0.57	0.14
BSL-1	Difficulty in concentration	0.91	0.79	0.59	0.17
BSL-2	Helplessness	0.90	0.93	0.56	0.10
BSL-3	Losing mind and memory	0.94	0.74	0.37	0.02
BSL-4	Disgust	0.92	0.89	0.41	0.05
BSL-5	Thoughts of hurting oneself	0.78	0.83	0.20	0.00
BSL-6	Not trusting others	0.88	0.90	0.52	0.18
BSL-7	No right to live	0.89	0.97	0.33	0.02
BSL-8	Lonely	0.90	0.91	0.71	0.12
BSL-9	Pressured by inner tension	0.96	0.94	0.59	0.07
BSL-10	Scary images	0.81	0.73	0.53	0.00
BSL-11	Hated oneself	0.90	0.87	0.42	0.00
BSL-12	Wants to punish oneself	0.79	0.77	0.12	0.00
BSL-13	Distressful shame	0.86	0.88	0.21	0.00

Symptoms		Class 1	Class 2	Class 3	Class 4
		Comorbid (<i>n</i> = 189)	Externalizing BPD (<i>n</i> = 53)	Avoidant BPD (<i>n</i> = 137)	Low Symptoms (<i>n</i> = 120)
BSL-14	Mood fluctuation	0.89	0.87	0.53	0.03
BSL-15	Distressed by voices or noises	0.83	0.97	0.25	0.01
BSL-16	Horrible effects being judged	0.88	0.88	0.32	0.03
BSL-17	Felt vulnerable	0.89	0.93	0.51	0.01
BSL-18	Fascinated with death	0.86	0.84	0.20	0.00
BSL-19	Felt meaningless	0.90	0.91	0.39	0.04
BSL-20	Fear of losing control	0.93	1.00	0.18	0.03
BSL-21	Disgusted with oneself	0.90	0.73	0.23	0.01
BSL-22	Felt distancing from oneself	0.90	0.85	0.16	0.00
BSL-23	Felt worthless	0.88	0.98	0.46	0.02
Mean of probabilities		0.90	0.76	0.40	0.05
Note. BSL-23 = Borderline Symptom List-short form.					
*The p-values of all post-hoc tests were less than .001 (<i>df</i> = 3).					

Table 2 Conditional response probabilities in each class

To facilitate the identification of each class, the probable PTSD, CPTSD [35] and BPD [38] diagnoses among each class were calculated based on the diagnostic algorithm and cutoff points, and these are presented in Table 3. The scores of PTSD, DSO, and BPD symptoms were described together.

Table 3

Mean score of the ITQ and BSL-23 and probable diagnoses of CPTSD and BPD among classes

		Class 1	Class 2	Class 3	Class 4
		Comorbid	Externalizing BPD	Avoidant BPD	Low Symptoms
		(<i>n</i> = 189)	(<i>n</i> = 53)	(<i>n</i> = 137)	(<i>n</i> = 120)
<i>M</i> (<i>SD</i>)	ITQ (PTSD)	2.42 (0.61)	0.71 (0.45)	1.02 (0.74)	0.33 (0.38)
	ITQ (DSO)	2.57 (0.55)	1.34 (0.77)	1.46 (0.77)	0.43 (0.41)
	BSL-23	2.50 (0.42)	2.43 (0.45)	1.30 (0.39)	0.36 (0.25)
<i>n</i> (%)	PTSD	8 (4.2)	0 (0.0)	15 (10.9)	0 (0.0)
	CPTSD	153 (81.0)	0 (0.0)	2 (1.5)	0 (0.0)
	BPD	189 (100.0)	53 (100.0)	133 (97.1)	19 (15.8)

Note. ITQ = International Trauma Questionnaire; BSL-23 = Borderline Symptom List-short form.

Table 3 Mean score of the ITQ and BSL-23 and probable diagnoses of CPTSD and BPD among classes

Class 1 showed high probabilities for symptoms of PTSD, DSO, and BPD. The majority of the people ($n = 153, 81\%$) in Class 1 satisfied a probable diagnosis for CPTSD and 100% of those in Class 1 satisfied a probable diagnosis for BPD. Therefore, it was named the “comorbid” class. Class 2 showed very low probabilities for PTSD symptoms and a high probability of 70% or more for all BPD symptoms. All people in Class 2 showed a probable diagnosis of BPD, but no CPTSD or PTSD. Class 3 showed moderate probabilities for PTSD symptoms and a high probability for the BPD symptom of loneliness. People in Class 3 showed low probabilities for other BPD symptoms, such as thoughts of hurting oneself, wanting to punish oneself, fascination with death, distressful shame, self-disgust, distressful voices and noises, fear of losing control, and distancing oneself. In Class 3, the majority of people ($n = 133, 97.1\%$) had probable BPD and only two (1.5%) had CPTSD. Thus, we named Class 2 the “externalizing BPD” class and Class 3 the “avoidant BPD” class. Class 4 showed low probabilities of all symptoms, so we named it the “low symptoms” class.

Analyses of differences among classes

Childhood abuse and invalidation environments

Table 4 presents the differences in childhood abuse and invalidating environment among latent classes. The comorbid class showed significantly higher PA, SA, EA, and PN scores. The externalizing BPD class was significantly higher than the low symptoms class in both SA and PN scores. The avoidant BPD class

showed particularly high EN compared to the low symptom class, although there were no significant differences between it and the comorbid and externalizing BPD classes. The avoidant class also showed higher scores in PA and EA compared to the externalizing BPD class. The comorbid class showed the highest scores in the invalidating environment by both mother and father, followed by the two BPD classes and the low symptom class.

Table 4

Mean score and differences in childhood abuse, invalidating environment, and attachment styles

		Class 1	Class 2	Class 3	Class 4	Chi-Square	Post-hoc
		Comorbid	Externalizing	Avoidant	Low		
		(n = 189)	BPD	BPD	Symptoms		
			(n = 53)	(n = 137)	(n = 120)		
Childhood abuse	Physical abuse	8.00	3.32	4.34	2.74	257.17	1 > 3 > 2,4
	Emotional abuse	8.66	3.86	5.68	2.98	306.65	1 > 3 > 2,4
	Sexual abuse	6.87	1.67	1.60	0.57	331.83	1 > 2,3 > 4
	Physical neglect	7.83	4.78	5.19	3.30	244.58	1 > 2,3 > 4
	Emotional neglect	7.89	8.09	8.34	7.08	5.18	3 > 4
Invalidating environment	Maternal	26.23	18.31	19.69	15.42	198.61	1 > 2,3 > 4
	Paternal	27.36	18.28	19.38	15.37	206.94	1 > 2,3 > 4
Attachment style	Anxious	65.56	50.28	53.63	37.08	249.28	1 > 2,3 > 4
	Avoidant	58.62	54.13	61.30	53.03	15.89	1,3 > 4

*All post-hoc tests, except for emotion regulation, had p-values less than .001 (df = 3).

Attachment styles

The comorbid class showed the highest scores in anxious attachment, followed by the two BPD classes and the low symptom class. As for avoidant attachment, the avoidant BPD class showed the highest score, as it showed a similar level as the comorbid class.

Table 4 Mean score and differences in childhood abuse, invalidating environment, and attachment styles

Table 5 Mean score and differences in pathological personality traits

Table 5
Mean score and differences in pathological personality traits

	Class 1 Comorbid (n = 189)	Class 2 Externalizing BPD (n = 53)	Class 3 Avoidant BPD (n = 137)	Class 4 Low Symptoms (n = 120)	Chi- Square	post- hoc
Negative affectivity	1.74	1.27	1.34	0.80	423.32	1 > 2,3 > 4
Emotional lability	1.78	1.34	1.49	0.94	223.91	1 > 2,3 > 4
Anxiousness	1.76	1.36	1.55	0.89	211.37	1 > 3 > 2 > 4
Depressivity	1.75	1.34	1.32	0.65	402.42	1 > 2,3 > 4
Hostility	1.74	1.24	1.36	0.88	197.86	1 > 2,3 > 4
Perseveration	1.77	1.14	1.22	0.84	316.54	1 > 2,3 > 4
Separation insecurity	1.61	1.20	1.08	0.61	248.09	1 > 2,3 > 4
Submissiveness	1.76	1.20	1.34	0.96	155.46	1 > 2,3 > 4
Detachment	1.74	1.30	1.42	1.03	248.01	1 > 2,3 > 4
Withdrawal	1.82	1.31	1.60	1.22	109.55	1 > 3 > 2,4
Intimacy avoidance	1.68	1.26	1.23	0.92	186.60	1 > 2,3 > 4
Anhedonia	1.72	1.37	1.44	0.93	226.71	1 > 2,3 > 4
Restricted affectivity	1.67	1.29	1.23	1.09	129.54	1 > 2,3 > 4
Suspiciousness	1.80	1.26	1.48	0.91	223.98	1 > 3 > 2 > 4
Antagonism	1.58	1.23	1.06	0.81	234.81	1 > 2 > 3 > 4
Deceitfulness	1.62	1.30	1.16	0.85	178.35	1 > 2,3 > 4

	Class 1 Comorbid (n= 189)	Class 2 Externalizing BPD (n= 53)	Class 3 Avoidant BPD (n= 137)	Class 4 Low Symptoms (n= 120)	Chi- Square	post- hoc
Manipulativeness	1.61	1.18	1.17	0.93	102.50	1 > 2,3 > 4
Callousness	1.58	1.20	0.96	0.70	273.12	1 > 2 > 3 > 4
Grandiosity	1.57	1.34	1.12	1.07	74.63	1 > 2 > 3,4
Attention seeking	1.54	1.15	0.97	0.69	156.01	1 > 2,3 > 4
Disinhibition	1.64	1.28	1.18	0.87	350.43	1 > 2,3 > 4
Rigid perfectionism	1.76	1.31	1.43	1.13	112.26	1 > 2,3,4 3 > 4
Distractibility	1.73	1.30	1.20	0.69	302.45	1 > 2,3 > 4
Impulsivity	1.66	1.15	1.06	0.76	253.33	1 > 2,3 > 4
Irresponsibility	1.62	1.28	1.16	0.78	232.71	1 > 2,3 > 4
Risk taking	1.51	1.30	1.05	0.89	151.54	1 > 2 > 3 > 4
Psychoticism	1.65	1.21	1.08	0.57	434.56	1 > 2,3 > 4
Unusual beliefs and experiences	1.55	1.21	0.98	0.51	306.14	1 > 2 > 3 > 4
Eccentricity	1.71	1.25	1.12	0.64	340.80	1 > 2,3 > 4
Cognitive and perceptual dysregulation	1.67	1.18	1.09	0.54	405.23	1 > 2,3 > 4
*The p-values of all post-hoc tests were less than .001 (df = 3).						

Pathological personality traits

Table 5 lists the differences in pathological personality traits among classes. The comorbid class showed significantly higher scores in all domains and facets. Meanwhile, the externalizing BPD class showed significantly higher antagonism domain scores than the avoidant BPD and low symptoms classes. Additionally, the externalizing BPD class was higher in some facets than the avoidant BPD class, such as callousness and grandiosity in the antagonism domain, risk taking in the disinhibition domain, and unusual beliefs and experiences in the psychoticism domain. By contrast, the avoidant BPD class showed higher scores than the externalizing BPD class in other facets, such as anxiousness in the negative affectivity domain as well as withdrawal and suspiciousness in the detachment domain.

Discussion

This study identified CPTSD and BPD symptoms in an adult general population sample with childhood abuse experiences with the aim of comparing attachment styles and pathological personality traits that may differ in the development of each diagnosis in addition to variant aversive childhood experiences. Four classes were identified: a comorbid class, an externalizing BPD class, an avoidant BPD class, and a low symptoms class.

Our sample did not show a distinct CPTSD class but a comorbid class, which may be because childhood abuse is a common risk factor for PTSD, CPTSD, and BPD [9], and because high comorbidity of CPTSD and BPD has been reported for adults who experienced childhood abuse [3, 16]. These results were similar to those of previous studies, where the CPTSD and BPD comorbid class was shown to be dominant with varying symptom levels of BPD [14, 15].

Among the four groups, the comorbid class had the highest number of individuals, with a total of 189 (37.9%). This finding supports the high comorbidity of CPTSD and BPD [13–15] and the idea that childhood abuse is a common risk factor for both CPTSD and BPD [3, 9]. Overall, showing similar symptomatic proportions as Cloitre et al. [3], the majority of individuals were in the BPD classes when adding up the three symptomatic classes, thus leaving 120 individuals (24.0%) with low symptom probabilities. Some individuals who have experienced childhood abuse may not develop severe psychopathology due to various protective factors [57]; our study showed that these included low levels of adverse experiences, unstable attachment, and pathological personality traits.

The comorbid class had the highest probabilities for all symptoms along with higher scores for childhood abuse and invalidating environment, thus supporting the idea that the accumulation of these risk factors leads to the comorbidity of CPTSD and BPD [13, 16]. Both anxious and avoidant attachment styles were dominant in this class, confirming the findings of previous research [10, 23–27]. This class also had the highest scores on all domains and facets of pathological personality traits, thus indicating the severity of psychopathology [17]. We may therefore conclude that the most severe adulthood mental health

consequences for childhood abuse are comorbid CPTSD and BPD, which require intensive psychosocial treatment [e.g., 58, 59].

In line with previous discussions showing that the presentation of BPD is highly heterogeneous [30, 60–62], our study identified subclasses of BPD. The two BPD classes showed moderate probabilities for DSO as well as a low probability for PTSD symptoms. The proportion of probable diagnoses for BPD was dominant, while those for CPTSD and PTSD were also relatively low, supporting that this group refers to subgroups of BPD. While the externalizing BPD class had a high probability of all BPD symptoms, the avoidant BPD showed loneliness as a dominant symptom while showing low probability for symptoms representing dissociation, shame or self-disgust, and suicidal or self-harm urges.

Both BPD classes showed similar levels of childhood sexual abuse and physical neglect and invalidating childhood environments, while the avoidant BPD class showed a substantially higher levels of emotional neglect and physical and emotional abuse than the externalizing BPD class. This is consistent with the findings of Müller et al. [63], which suggested that emotional neglect leads to social anxiety and avoidance. Emotional abuse or neglect was detected as a critical risk factor for people with BPD [5], which is consistent with the avoidant subgroup identified in our study, where this subgroup was also associated with avoidant attachment. In anxious attachment, both BPD classes showed high scores, which was also shown in previous studies, indicating that the ambivalent interpersonal relationship patterns in BPD are similar to the characteristics of anxious attachment [23].

The externalizing BPD class was associated with the pathological personality traits of the antagonism and disinhibition domains that make up the externalizing dimension [18], with significantly higher scores than the avoidant BPD class in callousness and grandiosity from the antagonism domain and in risk-taking from the disinhibition domain. This indicates that the externalizing BPD class can be understood as a group with a relatively deficient ability to empathize, an exaggerated sense of self, and a tendency to act according to their urges without regard for danger [8].

The avoidant BPD class scored higher than the externalizing BPD class in the pathological personality traits of anxiousness from the negative affectivity domain as well as those of withdrawal and suspiciousness from the detachment domain. The avoidant BPD class can be understood as a group of individuals with BPD who experience loneliness and anxiety related to negative perceptions of other people and environments who are detached and withdrawn from socio-interpersonal relationships [8]. Additionally, unusual beliefs and experiences from the psychoticism domain were significantly lower in the avoidant BPD class than they were in the externalizing BPD class, thus supporting that the avoidant BPD class may have lower levels of dissociative symptoms and uncontrollability.

Millon and colleagues [64] classified the following subtypes of BPD: discouraged BPD, impulsive BPD, petulant BPD, and self-destructive BPD. The results of the current study partially support this classification. While the externalizing subtype may represent impulsive, petulant, or self-destructive BPD, the avoidant subclass resembles the discouraged BPD subtype, which is characterized by passivity, separation anxiety, and a tendency to avoid others or social activities, ultimately causing one to

experience loneliness and emptiness [64]. Prior empirical studies have also repeatedly identified subgroups of BPD: an “unstable” subtype and an “empty” subtype in a large undergraduate sample [65] as well as an “extravert/externalizing” and a “schizotypal/paranoid” subtype in a treatment study sample [62]. Gamache and colleagues [30] identified an externalizing (“impulsive”) and an internalizing (“depressivity”) subprofile. Our sample identified similar subclasses, namely, externalizing and avoidant, and these findings are consistent with the HiTOP model, as BPD is characterized by both externalizing and internalizing spectra [17, 18]. The heterogeneous expressions of BPD may vary depending on the spectrum of externalization-internalization and associated personality traits (antagonism vs detachment); therefore, tailoring treatment according to the particular traits would have the largest treatment effects. For example, while stage-oriented treatment is needed in comorbid CPTSD/BPD and externalizing BPD starting from decreasing life-threatening behaviors, avoidant BPD may benefit from targeting inhibited grief starting from childhood abuse-associated stage 2 DBT [31]. Treatment of those in the CPTSD comorbid group should target PTSD symptoms in stage 2, and both BPD subgroups may benefit from targeting invalidating experiences or emotional abuse/neglect in stage 2 DBT, although PTSD symptoms are not present [e.g., 66].

The limitations of and suggestions for future research in this study are as follows: First, we were unable to compare the risk factors and pathological personality traits of CPTSD and BPD due to the absence of a distinct CPTSD class. Therefore, it is necessary to expand the discussion through variant sample studies including a clinical sample in the future. We also did not assess trauma in adulthood, nor did we assess the effect of polyvictimization in one’s lifetime. Moreover, as a cross-sectional study, the developmental course of CPTSD and BPD could not be confirmed. Further, using an online panel has the disadvantage of a high possibility of invalid participation [33]. However, although there were limitations, we applied a logical criterion when making judgments and excluded invalid responses.

Conclusion

Childhood abuse and an invalidating environment were found to lead to devastating symptomatology, eventually yielding high comorbidity of CPTSD and BPD and externalizing-internalizing subgroups of BPD. In addition to assessing CPTSD and BPD symptoms, assessing attachment styles and pathological personality traits based on dimensional approaches may be the key to tailoring different treatment approaches for individuals and to anticipate treatment trajectories.

Declarations

Ethics approval and consent to participate

The present study was approved by the Institutional Review Board of Chungbuk National University. All participants agreed to an informed consent before participating.

Consent for publication

Not applicable.

Availability of data and materials

The datasets generated and/or analysed during the current study are not publicly available because the participants of this study did not agree for their data to be shared publicly but are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This research was supported by a grant of the Korea Health Technology R&D Project through the Korea Health Industry Development Institute (KHIDI), funded by the Ministry of Health & Welfare, Republic of Korea (grant number: HC20C0140).

Authors' contributions

JL conceived the study, collected and analyzed the data, and developed the manuscript. HC conceived the study and revised the manuscript. All authors contributed to the submission of this study and approved the final manuscript.

Acknowledgements

Not applicable.

References

1. Keyes KM, Eaton NR, Krueger RF, McLaughlin KA, Wall MM, Grant BF, et al. Childhood maltreatment and the structure of common psychiatric disorders. *Br J Psychiatry*. 2012;200(2):107–15. <https://doi.org/10.1192/bjp.bp.111.093062>.
2. Cloitre M, Stolbach BC, Herman JL, Kolk BVD, Pynoos R, Wang J, et al. A developmental approach to complex PTSD: childhood and adult cumulative trauma as predictors of symptom complexity. *J Trauma Stress*. 2009;22(5):399–408. <https://doi.org/10.1002/jts.20444>.
3. Cloitre M, Garvert DW, Weiss B, Carlson EB, Bryant RA, Distinguishing PTSD, complex PTSD, and borderline personality disorder: a latent class analysis. *Eur J Psychotraumatol*. 2014;5(1):25097. <https://doi.org/10.3402/ejpt.v5.25097>.
4. Ibrahim J, Cosgrave N, Woolgar M. Childhood maltreatment and its link to borderline personality disorder features in children: a systematic review approach. *Clin Child Psychol Psychiatry*. 2018;23(1):57–76. <https://doi.org/10.1177/1359104517712778>.

5. Porter C, Palmier-Claus J, Branitsky A, Mansell W, Warwick H, Varese F. Childhood adversity and borderline personality disorder: a meta-analysis. *Acta Psychiatr Scand*. 2020;141(1):6–20. <https://doi.org/10.1111/acps.13118>.
6. Widom CS, Czaja SJ, Paris J. A prospective investigation of borderline personality disorder in abused and neglected children followed up into adulthood. *J Pers Disord*. 2009;23(5):433–46. <https://doi.org/10.1521/pedi.2009.23.5.433>.
7. World Health Organization. ICD-11: International Classification of Diseases 11th Revision. 2018. Available at: <https://icd.who.int/en>.
8. American Psychiatric Association. Diagnostic and statistical manual of mental disorders (5th Text Revision). Washington: American Psychiatric Association. ; 2022. <https://doi.org/10.1176/appi.books.9780890425787>.
9. Hyland P, Karatzias T, Shevlin M, Cloitre M. Examining the discriminant validity of complex posttraumatic stress disorder and borderline personality disorder symptoms: results from a united kingdom population sample. *J Trauma Stress*. 2019;32(6):855–63. <https://doi.org/10.1002/jts.22444>.
10. Powers A, Petri JM, Sleep C, Mekawi Y, Lathan EC, Shebuski K, et al. Distinguishing PTSD, complex PTSD, and borderline personality disorder using exploratory structural equation modeling in a trauma-exposed urban sample. *J Anxiety Disord*. 2022;88:102558. <https://doi.org/10.1016/j.janxdis.2022.102558>.
11. Owczarek M, Karatzias T, McElroy E, Hyland P, Cloitre M, Kratzer L, et al. Borderline Personality Disorder (BPD) and Complex Posttraumatic Stress Disorder (CPTSD): a network analysis in a highly traumatized clinical sample. *J Pers Disord*. 2023;37(1):112–29. <https://doi.org/10.1521/pedi.2023.37.1.112>.
12. Cyr G, Godbout N, Cloitre M, Bélanger C. Distinguishing among symptoms of posttraumatic stress disorder, complex posttraumatic stress disorder, and borderline personality disorder in a community sample of women. *J Trauma Stress*. 2022;35(1):186–96. <https://doi.org/10.1002/jts.22719>.
13. Frost R, Hyland P, Shevlin M, Murphy J. Distinguishing complex PTSD from borderline personality disorder among individuals with a history of sexual trauma: a latent class analysis. *Eur J Trauma Dissociation*. 2020;4(1):100080. <https://doi.org/10.1016/j.ejtd.2018.08.004>.
14. Jowett S, Karatzias T, Shevlin M, Albert I. Differentiating symptom profiles of ICD-11 PTSD, complex PTSD, and borderline personality disorder: a latent class analysis in a multiply traumatized sample. *Pers Disord Theory Res Treat*. 2020;11(1):36. <https://doi.org/10.1037/per0000346>.
15. Saraiya TC, Fitzpatrick S, Zumberg-Smith K, López-Castro T, Back E, Hien SA. Social-emotional profiles of PTSD, complex PTSD, and borderline personality disorder among racially and ethnically diverse young adults: a latent class analysis. *J Trauma Stress*. 2021;34(1):56–68. <https://doi.org/10.1002/jts.22590>.
16. Ford JD, Courtois CA. Complex PTSD and borderline personality disorder. *Borderline Personal Disord Emot Dysregul*. 2021;8(1):16. <https://doi.org/10.1186/s40479-021-00155-9>.

17. Kotov R, Krueger RF, Watson D, Achenbach TM, Althoff RR, Bagby RM, et al. The hierarchical taxonomy of psychopathology (HiTOP): a dimensional alternative to traditional nosologies. *J Abnorm Psychol.* 2017;126(4):454. <https://doi.org/10.1037/abn0000258>.
18. Kotov R, Krueger RF, Watson D, Cicero DC, Conway CC, DeYoung CG, et al. The hierarchical taxonomy of psychopathology (HiTOP): a quantitative nosology based on consensus of evidence. *Annu Rev Clin Psychol.* 2021;17:83–108. <https://doi.org/10.1146/annurev-clinpsy-081219-093304>.
19. Bretherton I, Ridgeway D, Cassidy J. Assessing internal working models of the attachment relationship: an attachment story completion task for 3-year-olds. *Attachment in the preschool years: Theory, research, and intervention.* Chicago: University of Chicago Press; 1990. 273–308.
20. Mosquera D, Gonzalez A, Leeds AM. Early experience, structural dissociation, and emotional dysregulation in borderline personality disorder: the role of insecure and disorganized attachment. *Borderline Personal Disord Emot Dysregul.* 2014;1(1):1–8. <https://doi.org/10.1186/2051-6673-1-15>.
21. Peng W, Liu Z, Liu Q, Chu J, Zheng K, Wang J, et al. Insecure attachment and maladaptive emotion regulation mediating the relationship between childhood trauma and borderline personality features. *Depress Anxiety.* 2021;38(1):28–39. <https://doi.org/10.1002/da.23082>.
22. Schneider BH, Atkinson L, Tardif C. Child-parent attachment and children's peer relations: a quantitative review. *Dev Psychol.* 2001;37(1):86. <https://doi.org/10.1037/0012-1649.37.1.86>.
23. Smith M, South S. Romantic attachment style and borderline personality pathology: a meta-analysis. *Clin Psychol Rev.* 2020;75:101781. <https://doi.org/10.1016/j.cpr.2019.101781>.
24. van Dijke A, Hopman JA, Ford JD. Affect dysregulation, psychoform dissociation, and adult relational fears mediate the relationship between childhood trauma and complex posttraumatic stress disorder independent of the symptoms of borderline personality disorder. *Eur J Psychotraumatol.* 2018;9(1):1400878. <https://doi.org/10.1080/20008198.2017.1400878>.
25. Karatzias T, Shevlin M, Hyland P, Brewin CR, Cloitre M, Bradley A, et al. The role of negative cognitions, emotion regulation strategies, and attachment style in complex post-traumatic stress disorder: implications for new and existing therapies. *Br J Clin Psychol.* 2018;57(2):177–85. <https://doi.org/10.1111/bjc.12172>.
26. Maercker A, Bernays F, Rohner SL, Thoma MV. A cascade model of complex posttraumatic stress disorder centered on childhood trauma and maltreatment, attachment, and socio-interpersonal factors. *J Trauma Stress.* 2022;35(2):446–60. <https://doi.org/10.1002/jts.22756>.
27. Sandberg DA, Refrea V. Adult attachment as a mediator of the link between interpersonal trauma and International Classification of Diseases (ICD)-11 complex posttraumatic stress disorder symptoms among college men and women. *J Interpers Violence.* 2022;37:23–4. <https://doi.org/10.1177/08862605211072168>.
28. Fowler JC, Madan A, Allen JG, Oldham JM, Frueh BC. Differentiating bipolar disorder from borderline personality disorder: diagnostic accuracy of the difficulty in emotion regulation scale and personality inventory for DSM-5. *J Affect Disord.* 2019;245:856–60. <https://doi.org/10.1016/j.jad.2018.11.079>.

29. Møller L, Meisner MW, Søgaaard U, Elklit A, Simonsen E. Assessment of personality functioning in ICD-11 posttraumatic stress disorder and complex posttraumatic stress disorder. *Pers Disord Theory Res Treat.* 2021;12(5):466. <https://doi.org/10.1037/per0000491>.
30. Gamache D, Savard C, Leclerc P, Payant M, Côté A, Faucher J, et al. Latent profiles of patients with borderline pathology based on the alternative DSM-5 model for personality disorders. *Borderline Personal Disord Emot Dysregul.* 2021;8(1):1–13. <https://doi.org/10.1186/s40479-021-00146-w>.
31. Linehan MM. *Cognitive-behavioral treatment of borderline personality disorder.* New York: Guilford Press; 1993.
32. Neyman J. On the two different aspects of the representative method: the method of stratified sampling and the method of purposive selection. *Breakthroughs in statistics.* New York: Springer; 1992. 123–50.
33. Kramer J, Rubin A, Coster W, Helmuth E, Hermos J, Rosenbloom D, et al. Strategies to address participant misrepresentation for eligibility in web-based research. *Int J Methods Psychiatr Res.* 2014;23(1):120–9. <https://doi.org/10.1002/mpr.1415>.
34. Bernstein D, Fink L. *Manual for the childhood trauma questionnaire.* New York: The Psychological Corporation; 1998.
35. Cloitre M, Shevlin M, Brewin CR, Bisson JI, Roberts NP, Maercker A, et al. The international trauma questionnaire: development of a self-report measure of ICD-11 PTSD and complex PTSD. *Acta Psychiatr Scand.* 2018;138(6):536–46. <https://doi.org/10.1111/acps.12956>.
36. Choi H, Lee W, Hyland P. Factor structure and symptom classes of ICD-11 complex posttraumatic stress disorder in a South Korean general population sample with adverse childhood experiences. *Child Abuse Negl.* 2021;114:104982. <https://doi.org/10.1016/j.chiabu.2021.104982>.
37. Bohus M, Limberger MF, Frank U, Sender I, Gratwohl T, Stieglitz R-D. Development of the borderline symptom list. *Psychother Psychosom Med Psychol.* 2001;51(5):201–11. <https://doi.org/10.1055/s-2001-13281>.
38. Kang J, Kim J, Seok J, Koo B, Ryu JS, Shin H, et al. Validation of the Korean version of the Borderline Symptom List Short Version (K-BSL-23). *Korean J Clin Psychol.* 2023;42(2):23–34. <https://doi.org/10.15842/KJCP.PUB.42.2.23>.
39. Bohus M, Kleindienst N, Limberger MF, Stieglitz RD, Domsalla M, Chapman AL, et al. The short version of the borderline symptom list (BSL-23): development and initial data on psychometric properties. *Psychopathology.* 2009;42(1):32–9. <https://doi.org/10.1159/000173701>.
40. Bernstein DP, Stein JA, Newcomb MD, Walker E, Pogge D, Ahluvalia T, et al. Development and validation of a brief screening version of the childhood trauma questionnaire. *Child Abuse Negl.* 2003;27(2):169–90. [https://doi.org/10.1016/S0145-2134\(02\)00541-0](https://doi.org/10.1016/S0145-2134(02)00541-0).
41. Kim D, Park SC, Yang H, Oh DH. Reliability and validity of the Korean version of the childhood trauma questionnaire-short form for psychiatric outpatients. *Psychiatry Investig.* 2011;8(4):305. <https://doi.org/10.4306/pi.2011.8.4.305>.

42. Mountford V, Corstorphine E, Tomlinson S, Waller G. Development of a measure to assess invalidating childhood environments in the eating disorders. *Eat Behav.* 2007;8(1):48–58. <https://doi.org/10.1016/j.eatbeh.2006.01.003>.
43. Khu B, Song J. Validation of Korean version of Invalidating Childhood Environment Scale (ICES). *Korean J Youth Stud.* 2019;26(10):157–80. <https://doi.org/10.21509/KJYS.2019.10.26.10.157>.
44. Fraley RC, Waller NG, Brennan KA. An item response theory analysis of self-report measures of adult attachment. *J Pers Soc Psychol.* 2000;78(2):350. <https://doi.org/10.1037/0022-3514.78.2.350>.
45. Kim SH. Adaptation of the experiences in close relationships-revised scale into Korean: confirmatory factor analysis and item response theory approaches [master's thesis]. Seoul, Korea: Seoul National University; 2004.
46. Krueger RF, Derringer J, Markon KE, Watson D, Skodol AE. Initial construction of a maladaptive personality trait model and inventory for DSM-5. *Psychol Med.* 2012;42(9):1879–90. <https://doi.org/10.1017/S0033291711002674>.
47. Shin S, Hwang S. Reliability and validity of the Korean version Personality Inventory for DSM-5 (PID-5). *Korean J Soc Personal Psychol.* 2016;30(3):1–24.
48. Lanza ST, Cooper BR. Latent class analysis for developmental research. *Child Dev Perspect.* 2016;10(1):59–64. <https://doi.org/10.1111/cdep.12163>.
49. Akaike H. Factor analysis and AIC. *Psychometrika.* 1987;52:317–32.
50. Schwarz G. Estimating the dimension of a model. *Ann Stat.* 1978;6(2):461–4.
51. Sclove SL. Application of model-selection criteria to some problems in multivariate analysis. *Psychometrika.* 1987;52(3):333–43.
52. Nylund KL, Asparouhov T, Muthén BO. Deciding on the number of classes in latent class analysis and growth mixture modeling: a monte carlo simulation study. *Struct Equ Modeling.* 2007;14(4):535–69. <https://doi.org/10.1080/10705510701575396>.
53. Clark SL, Muthén B. Relating latent class analysis results to variables not included in the analysis. *Statmodel.* 2009. Available at: <https://www.statmodel.com/download/relatinglca.pdf>. Accessed 15 Oct 2013.
54. Lo Y, Mendell NR, Rubin DB. Testing the number of components in a normal mixture. *Biometrika.* 2001;88(3):767–78. <https://doi.org/10.1093/biomet/88.3.767>.
55. Asparouhov T, Muthén B. Auxiliary variables in mixture modeling: three-step approaches using M plus. *Struct Equation Modeling: Multidiscip J.* 2014;21(3):329–41. <https://doi.org/10.1080/10705511.2014.915181>.
56. Nylund-Gibson K, Choi AY. Ten frequently asked questions about latent class analysis. *Transl Issues Psychol Sci.* 2018;4(4):440. <https://doi.org/10.1037/tps0000176>.
57. Afifi TO, MacMillan HL. Resilience following child maltreatment: a review of protective factors. *Can J Psychiatry.* 2011;56(5):266–72. <https://doi.org/10.1177/070674371105600505>.

58. Bohus M, Kleindienst N, Hahn C, Müller-Engelmann M, Ludäscher P, Steil R, et al. Dialectical behavior therapy for posttraumatic stress disorder (DBT-PTSD) compared with cognitive processing therapy (CPT) in complex presentations of PTSD in women survivors of childhood abuse: a randomized clinical trial. *JAMA psychiatry*. 2020;77(12):1235–45. <https://doi.org/10.1001/jamapsychiatry.2020.2148>.
59. Harned MS, Wilks CR, Schmidt SC, Coyle TN. Improving functional outcomes in women with borderline personality disorder and PTSD by changing PTSD severity and post-traumatic cognitions. *Behav Res Ther*. 2018;103:53–61. <https://doi.org/10.1016/j.brat.2018.02.002>.
60. Antoine SM, Fredborg BK, Streiner D, Guimond T, Dixon-Gordon KL, Chapman AL, et al. Subgroups of borderline personality disorder: a latent class analysis. *Psychiatry Res*. 2023;323:115131. <https://doi.org/10.1016/j.psychres.2023.115131>.
61. Bohus M, Stoffers-Winterling J, Sharp C, Krause-Utz A, Schmahl C, Lieb K. Borderline personality disorder. *Lancet*. 2021;398(10310):1528–40. [https://doi.org/10.1016/S0140-6736\(21\)00476-1](https://doi.org/10.1016/S0140-6736(21)00476-1).
62. Smits ML, Feenstra DJ, Bales DL, de Vos J, Lucas Z, Verheul R, et al. Subtypes of borderline personality disorder patients: a cluster-analytic approach. *Borderline Personal Disord Emot Dysregul*. 2017;4:1–15. <https://doi.org/10.1186/s40479-017-0066-4>.
63. Müller LE, Bertsch K, Bülau K, Herpertz SC, Buchheim A. Emotional neglect in childhood shapes social dysfunctioning in adults by influencing the oxytocin and the attachment system: results from a population-based study. *Int J Psychophysiol*. 2019;136:73–80. <https://doi.org/10.1016/j.ijpsycho.2018.05.011>.
64. Millon T, Millon CM, Meagher SE, Grossman SD, Ramnath R. *Personality disorders in modern life*. New Jersey: John Wiley & Sons; 2012.
65. Johnson BN, Levy KN. Identifying unstable and empty phenotypes of borderline personality through factor mixture modeling in a large nonclinical sample. *Pers Disord Theory Res Treat*. 2020;11(2):141. <https://doi.org/10.1037/per0000360>.
66. Harned MS. *Treating trauma in dialectical behavior therapy: the DBT prolonged exposure protocol (DBT PE)*. New York: Guilford Publications; 2022.

Figures




Figure 1

Symptom endorsement of complex posttraumatic stress disorder and borderline personality disorder items