

Risk perception related to COVID-19 among the Iranian general population: An application of the extended parallel process model

Leila Jahangiry (✉ jahangiry@razi.tums.ac.ir)

Tabriz University of Medical Sciences Faculty of Health and Nutrition <https://orcid.org/0000-0002-0491-5764>

Fatemeh Bakhtari

Tabriz University of Medical Sciences

Zahara Sohrabi

Tabriz University of Medical Sciences

Parvin Reihani

Tabriz University of Medical Sciences

Sirous Samei

Tabriz University of Medical Sciences

Koen Ponnet

Universiteit Gent

Ali Montazeri

AECER

Research article

Keywords: extended parallel process model, coronavirus disease 2019, Fear Control, Danger control, Behavior response

Posted Date: August 10th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-33493/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published on October 19th, 2020. See the published version at <https://doi.org/10.1186/s12889-020-09681-7>.

Abstract

Background: The novel coronavirus disease 2019 (COVID-19) has emerged as a major global public health challenge. This study aimed to investigate how people perceived the COVID-19 outbreak using the components of the Extended Parallel Process Model (EPPM) and to find out how this might contribute to possible behavioral responses to the prevention and control of the disease.

Methods: This cross-sectional study was conducted online in Iran in March and April 2020. The Data were collected using an electronic questionnaire via Porsline. Participants were recruited using online applications and posts on a number of platforms such as Telegram, WhatsApp, and Instagram asking people to take part in the study. To collect data a self-designed questionnaire based on the EPPM was used in order to measure the risk perception (efficacy, defensive responses, perceived treat) related to the COVID-19. Descriptive statics, chi-square, t-test and analysis of variance (ANOVA), were used to explore the data.

Results: A total of 3,727 individuals with a mean age (SD) of 37.0 (11.1) years participated in the study. The results revealed significant progressive increases in efficacy, defensive responses and perceived treat particularly among those aged 60 and over. Women had significantly higher scores than men on some aspects such as self-efficacy, reactance, and avoidance but men had higher perceived susceptibility scores for COVID-19 than women. Overall 56.4% of participants were engaged in danger control (preventive behavior) while the remaining 43.6% were engaged in fear control (non-preventive behavior) process.

Conclusion: More than half of all participants motivated by danger control. This indicated that more than half of participants had high perceived efficacy (i.e., self-efficacy and response efficacy). Self-efficacy scores were significantly higher among participants who were older, female, single, lived in rural areas, and had good economic status. The results suggest that socioeconomic and demographic factors are the main determinants of risk perceptions for controlling the COVID-19 pandemic.

Introduction

The novel coronavirus disease 2019 (COVID-19) has emerged as a major global public health challenge [1]. This potentially fatal infectious disease, which has affected most countries worldwide, is characterized by a steady speed of spread, leading to the World Health Organization (WHO) classifying it as a pandemic [2]. The WHO early report on Feb 11, 2020 for the onset of the COVID-19 outbreak announced that the distribution of infected people by coronavirus was dramatically increased. In Iran, the first official report of a COVID-19-related death was on February 19, 2020. As of April 13, 73,303 Iranian people have been infected with the COVID-19. In total, 4,585 have died and 45,983 have recovered when we were carrying this study [3].

The COVID-19 is transmitted from human to human through respiratory droplets or direct contact. Available findings show that avoiding exposure is the key to preventing COVID-19 infection, which is why quarantining, physical distancing, and isolation have become the primary strategies for reducing COVID-19 incidence and mortality. Quarantine restricts the movement of people and reduces the infection rate for contagious diseases [4]; physical or social distancing involves staying at least two meters away from other persons; and isolation is the state of separating patients with COVID-19 from otherwise healthy people [5]. Maintaining hygiene practices, such as proper hand washing, wearing a mask in crowded places, and staying at home, controlling the spread of the disease and breaking the transmission chain [6].

During the early stages of the coronavirus pandemic in Iran, several social media campaigns were launched to inform people about the risks of coronavirus and to persuade them to follow health care recommendations. Therefore, providing necessary information about people's perception related to COVID-19 is important for health policy makers in implementing effective and appropriate strategies for prevention and control of the disease.

Psychosocial and cultural factors affect adherence to health advice [7]. There are several factors affecting COVID-19 response behaviors. Multiple health models suggest that risk perception of COVID-19 is a vital component for behavior change [8], one of which is the extended parallel process model (EPPM) [9]. According to EPPM, when people are exposed to a risky situation, they produce two cognitive appraisals: one of the efficacy of the recommended response and one of the threat [10]. EPPM therefore indicates that the perception of risk would act together with self-efficacy levels for prevention [11]. It is important to understand how people are being presented with risky situations during the COVID-19 pandemic, how people are assessing these risks, and how such assessments lead them to changing their behaviors in order to follow the COVID-19 health recommendations. This study aimed to investigate how people have perceived the COVID-19 outbreak using the components of the EPPM (efficacy, defensive responses, perceived threat) and how these might contribute to possible behavioral responses to the prevention and control of the disease.

Materials And Methods

Study design

This was a cross sectional online study. The Ethics Committee of Tabriz University of Medical Sciences approved the study (No: IR.TBZMED.REC.1398.1307). We obtained informed consent from all participants. This cross-sectional study was conducted online in Iran in March and April 2020.

Data collection and participants

Data were collected using an electronic questionnaire via Porsline. Participants were recruited using online applications and posts on several platforms such as Telegram, WhatsApp, and Instagram. We asked people for sharing the study announcements and to participate in the study. We also send several messages to significant others, virtual groups to share the study announcements. The posts asked people aged 15 years and over to take part in a study that investigates on the COVID-19. Those interested in participating were directed to complete the online questionnaire.

Sample size estimation

The sample size for the study was estimated using the following formula [12]:

$$n = Z^2P(1-P)/d^2$$

where Z for 95% confidence interval is 1.96, P = 0.5 (assuming that 50% of people would either be engaged in danger control or fear control processes), and d = 0.02 (precision = 2%). As such we estimated at least 2400 participants would be requiring for the study to have a power of 80% at 5% significant level. However, in practice 3727 individuals participated in the study.

The study questionnaire

The Risk-Percept COVID-19 was used to collect the data. This questionnaire was developed based on a literature review of the EPPM-based risk perception assessments of other infectious diseases. The questionnaire was subjected to psychometric evaluation before the data collection and found to be a valid and reliable measure. The Cronbach's alpha coefficients for the dimensions ranged from 0.69 to 0.79, which indicated acceptable internal consistency for the questionnaire. In addition, the stability of the questionnaire as assessed by Intraclass Correlation Coefficient (ICC) showed satisfactory results (ICC ranged from 0.71 to 0.80). However, the final questionnaire consisted of 29 items tapping into three pre-defined dimensions:

1. Efficacy (perceived self-efficacy and perceived response efficacy): This dimension included 13 items.

1.1 The perceived self-efficacy included six items measuring participants' beliefs about their ability to perform the recommended responses to the COVID-19. A sample item reads as follows: "I am able to use a mask in closed places with more than two people."

1.2 The response efficacy included seven items measuring participants' beliefs about the effectiveness of the recommended preventative responses to the COVID-19. A sample item reads as follows: "I believe that by disinfecting surfaces, I am less likely to get coronavirus."

2. Defensive response (denial, reactance and avoidance): This dimension included eight items measuring people's beliefs about their perception of the risk of COVID-19.

2.1 A sample item for denial: "I think sickness or death are in God's hands and following the precautionary measures is not so important".

2.2 A sample item for reactance: "I believe that health staff significantly exaggerate about the disease".

2.3 A sample item for avoidance: "When TV or radio talks about coronavirus, I flip the channel".

3. Perceived threats (susceptibility and severity). This dimension included eight items measuring people's beliefs about the magnitude of the COVID-19 and about their risk of experiencing the disease.

3.1 A sample item for susceptibility: "I am at risk for getting the coronavirus".

3.2. A sample item for severity: "The coronavirus is a lethal threat".

Each item was rated on a 5-point Likert scale (1=strongly disagree, 2=disagree, 3=neutral, 4=agree, 5=strongly agree) giving an overall row score ranging from 29 to 145. Additionally, the following demographic data were collected for all participants: age, gender, education, marital status, economic status, history of coronavirus, family history of coronavirus, having chronic diseases, and living condition (urban vs. rural).

Scoring

1. Risk perception: Using the following simple linear transformation [$\text{Row score} - \text{the lowest possible raw score} / \text{highest possible raw score} - \text{the lowest possible raw score}$] x 100, the row scores were converted into a score of 0 to 100 where lower scores indicated lower risk perception and the higher scores indicated higher risk perception.

2. Danger control and fear control: Consistent with previous studies [9], we subtracted the perceived threat score from the perceived efficacy score, resulting in a discriminating value. The discriminating value could be either positive or negative. A positive value meant that a person was engaging in danger control processes because their perceived efficacy was stronger than their threat perceptions. In other words, a person was likely to engage in some level of protective behaviors with regard to the specific health threat. A negative value meant that a person was engaging in fear control processes because their threat perceptions were stronger than their perceptions of efficacy. In these cases, a person was likely to engage in fear control processes and was probably not protecting himself or herself against the specific health threat.

Statistical analysis

Statistical analyses were performed using the Statistical Package for Social Science, Version 18, for Windows (SPSS Inc., Chicago, IL, USA). The normality of the data was analyzed using a Kolmogorov–Smirnov test. The characteristics of the participants were summarized as numbers, percentages, or means with standard deviations, where appropriate. Analysis of variance (ANOVA), t-tests, and chi-square were used to compare the study sub-groups. In all tests, a value of $p < 0.05$ was considered statistically significant.

Results

Demographic characteristics

Table 1 shows the sociodemographic characteristics of the respondents (1933 male, and 1794 female). The mean age of the respondents was 37.0 (SD = 11.1) years. More than half of all participants (52%) were married. In all 40 respondents (1.1%) reported that they had coronavirus and 69 (1.9%) stated that they had a family member with a confirmed case of the disease. The majority of participants (96.1%) lived in urban areas.

Perceived risk of the coronavirus disease

The perceived risk for all participants based on sociodemographic characteristics are shown in Table 2. The results by age revealed that as age increased, the significant progressive increase in perceived self-efficacy, avoidance response and perceived susceptibility scores were observed. This applied particularly for participants aged 60 and over. Women had significantly higher scores than men for self-efficacy, reactance, and avoidance defensive responses. Men had higher perceived susceptibility scores for the COVID-19 than women. The average scores across all dimensions showed significant increases as education levels increased, as well as for participants who were married or had good economic status. Perceived risk scores showed that participants with confirmed coronavirus cases had significantly higher scores except for the perceived susceptibility scores. Participants without any family history of coronavirus had higher perceived scores for response efficacy and self-efficacy. We found significantly higher scores for perceived risk among respondents with no chronic diseases (Table 3).

Danger control and fear control

Table 4 shows the discriminating values indicating danger control and fear control scores based on different sociodemographic characteristics. A total of 56.4% of participants were engaging in danger control processes and 43.6% in fear control processes. The respondents in former group were more likely to engage in preventive behaviors while those in the latter group were more likely to delay recommended responses for preventing

themselves from the COVID-19. There were significant differences in danger and fear control scores by age, gender, education, economic status, and having chronic diseases.

Discussion

This EPPM-based study was conducted to assess the risk perceptions, overall perceived danger and fear control processes among Iranian people during the early stages of the COVID-19 pandemic. The study provides a timely assessment and initial evidence related to the risk perceptions and psychological responses of more than 3,700 individuals across the country who took part in the study.

According to the EPPM, two cognitive appraisals might initiate after a person learns about a health risk: one related to the threat it poses and a second related to the efficacy to follow the recommended responses. When the threat of COVID-19 is perceived to be more significant and efficacy is low, people are usually act to protect themselves from the fear rather than the danger itself (fear control process). Conversely when perceived efficacy is significantly high, even if the perceived treat would be high, people usually are motivated to protect themselves from the danger and could manage the threat (danger control process) [13]. Our results showed that 56.4% of respondents were motivated by danger control responses and 43.6% by fear control responses. This indicates that more than half of all participants had high perceived efficacy (i.e., self-efficacy and response efficacy). Self-efficacy scores were significantly higher among participants who were older, female, single, lived in rural areas, or had good economic status.

A current study from China showed an association between self-efficacy and social support among patients who had been treated for coronavirus [14]. This is inconsistent with our results, where it was found that participants with a history of coronavirus had lower self-efficacy scores. Respondents who had a family member with coronavirus and those with three or more comorbidities had lower self-efficacy scores for controlling COVID-19. Self-efficacy is a positive mental state that is part of the cognitive appraisal process reducing stress and tension [15]. The results showed that respondents with high self-efficacy were better able to control their emotions. We found that efficacy was significantly higher among respondents who were well-educated and had good economic status. It seems that these individuals believed that they can carry out the recommended responses to protect themselves from the COVID-19.

The results also showed that respondents who were male, older, well-educated, and married had significantly higher perceptions of susceptibility. In fact, these individuals were simply thinking about the threat of the COVID-19 and believed that the threat was relevant to themselves. According to the WHO, older people are at higher risk of contracting COVID-19 [16]. The Iranian health care system and media provided significant coverage of the COVID-19 pandemic, recommending that all people, especially older people take good care of themselves. This likely resulted in older people learning that they were more susceptible to the disease.

Individuals usually use psychological defense strategies to control their fears. These strategies include denial, avoidance, and reactance. Our results showed that lower defensive response scores correlated with better responses from participants. Defensive avoidance occurs when individuals block out feelings and thoughts about a threat or ignore further information about it, for example, switching the television channel or skipping COVID-19-related news. People in younger age groups (15-29 years) had higher avoidance scores and lower self-efficacy scores, indicating that younger people tended to take more risks and ignore health recommendations [17]. The

results of our study suggest that socioeconomic and demographic factors are the main determinants of psychological responses to controlling the COVID-19 pandemic.

Strengths and limitations

This study benefited from a relatively good sample size and using a theory based questionnaire for data collection was an advantage. The greatest strength of this study was its format. The online method allowed for the timely collection of information from a wide range of community groups. Since the pandemic feature of the COVID-19 made other data collection methods unsafe and difficult for both the researchers and the study participants, the online sampling method was particularly convenient. Another strength could be related to the fact that we collected the data before blow up of news and information about the COVID-19. However, because of the online nature of the study, we were unable to reach people who did not have access to the Internet or online applications. In addition, it is necessary to mention that online survey during the early stage of a pandemic was relatively new and therefore although received the invitation to participate, did not attend to respond to the questionnaire. In fact, we could not identify none responders. Furthermore, most participants were relatively well educated. Thus the findings might not be generalized to all population.

Conclusion

More than half of all participants motivated by danger control. This indicated that more than half of participants had high perceived efficacy (i.e., self-efficacy and response efficacy). Self-efficacy scores were significantly higher among participants who were older, female, single, lived in rural areas, and had good economic status. The results suggest that socioeconomic and demographic factors are the main determinants of possible behavioral responses to controlling the COVID-19 pandemic.

Abbreviations

WHO: World Health Organization; **EPPM:** extended parallel process model; **ANOVA:** Analysis of variance; **COVID-19:** coronavirus disease 2019

Declarations

Ethics approval and consent to participate

The study received ethical approval from the Ethics Committee of Tabriz University of Medical Sciences (NO: IR.TBZMED.REC. 1398.1307). We obtained informed consent and parental consent from participants and participants aged 15-18 years, respectively.

Consent for publication

The authors have agreed on the content of the manuscript.

Availability of data and material

The questionnaires and datasets generated and/or analyzed during the current study are available from the corresponding authors on reasonable request.

Competing interests

The authors declare no conflicts of interest.

Funding

Tabriz University of Medical Sciences provided facilities.

Authors' contributions

LJ and AM were responsible for the study design. LJ did the analyses. LJ and AM were responsible for data interpretation. FB, PR, SS, and ZS helped in the study design and data collection. KP and AM helped in the drafting of the manuscript. AM critically reviewed the manuscript and provided the final manuscript. All authors have read and approved the final manuscript.

Acknowledgments

We acknowledge the contributions of Tabriz University of Medical Sciences, Tabriz, Iran for providing facilities to the study. Our thanks go to all the respondents of the study for their valuable co-operation.

References

1. Contreras, G.W., *Getting ready for the next pandemic COVID-19: Why we need to be more prepared and less scared*. J Emerg Manag, 2020. **18**(2): p. 87-89.
2. World Health Organization. Coronavirus disease (COVID-19) advice for the public.
3. Basic protective measures against the new coronavirus. Accessed 25 Mar 2020.3.
4. Iranian Ministry of Health and Medical Education COVID-19 daily epidemiology journal.
5. Brooks, S.K., et al., *The psychological impact of quarantine and how to reduce it: rapid review of the evidence*. The Lancet, 2020. **395**(10227): p. 912-920.
6. <https://www.cdc.gov/coronavirus/2019-ncov/prevent-getting-sick/social-distancing.html>.
7. public., W.H.O.C.d.C.-a.f.t. and B.p.m.a.t.n.c.A.M. 2020.
8. Pulcini, C., et al., *Factors associated with adherence to infectious diseases advice in two intensive care units*. Journal of Antimicrobial Chemotherapy, 2006. **57**(3): p. 546-550.
9. Napper, L.E., D.G. Fisher, and G.L. Reynolds, *Development of the perceived risk of HIV scale*. AIDS and behavior, 2012. **16**(4): p. 1075-1083.
10. Popova, L., *The extended parallel process model: Illuminating the gaps in research*. Health Education & Behavior, 2012. **39**(4): p. 455-473.
11. McMahan, S., K. Witte, and J.a. Meyer, *The perception of risk messages regarding electromagnetic fields: extending the extended parallel process model to an unknown risk*. Health communication, 1998. **10**(3): p. 247-259.
12. Chi, A.I., et al., *Learning AIDS in Singapore: Examining the effectiveness of HIV/AIDS efficacy messages for adolescents using ICTs*. Knowledge Management & E-Learning: An International Journal, 2010. **2**(2): p. 169-187.

13. Daniel WW. *Biostatistics: A Foundation for Analysis in the Health Sciences*. 7th edition. New York: John Wiley & Sons; 1999.
14. Witte, K., *Fear control and danger control: A test of the extended parallel process model (EPPM)*. Communications Monographs, 1994. **61**(2): p. 113-134.
15. Xiao, H., et al., *The Effects of Social Support on Sleep Quality of Medical Staff Treating Patients with Coronavirus Disease 2019 (COVID-19) in January and February 2020 in China*. Medical science monitor : international medical journal of experimental and clinical research, 2020. **26**: p. e923549-e923549.
16. Schönfeld, P., et al., *The effects of daily stress on positive and negative mental health: Mediation through self-efficacy*. International Journal of Clinical and Health Psychology, 2016. **16**(1): p. 1-10.
17. <http://www.euro.who.int/en/health-topics/health-emergencies/coronavirus-covid-19/statements/statement-older-people-are-at-highest-risk-from-covid-19>.
18. Dillon, L., et al., *Risk, protective factors and resilience to drug use: identifying resilient young people and learning from their experiences*. 2007: Home Office London.

Tables

Table 1: Sociodemographic characteristics of the study sample (N = 3727)

	Male (= 1933)	Female (n =1794)	Total
	No. (%)	No. (%)	No. (%)
Age groups			
15-29	376 (19.5)	489 (27.3)	865 (23.2)
30-44	1051 (54.4)	944 (52.6)	1995 (53.5)
45-59	419 (21.7)	315 (17.6)	734 (19.7)
60+	87 (4.5)	46 (2.6)	133 (3.6)
Educational status			
Primary	99 (5.1)	103 (5.7)	202 (5.4)
Secondary	458 (23.7)	465 (25.9)	923 (24.8)
Higher	1376 (71.2)	1226 (68.3)	2602 (69.8)
Marital status			
Single	439 (49.8)	442 (50.2)	881 (23.6)
Married	1483 (52.7)	1329 (47.3)	2812 (75.4)
Widowed/divorced	11 (0.6)	23 (1.3)	34 (0.9)
Economic status			
Good	378 (19.6)	547 (30.5)	925 (24.8)
Intermediate	1084 (56.1)	1049 (58.5)	2153 (57.2)
Poor	471 (24.4)	198 (11.0)	669 (17.9)
History of coronavirus			
Yes	16(0.9)	21 (1.1)	37 (1.0)
No	1778 (99.1)	1912 (98.9)	3690 (99.0)
History of coronavirus in family members			
Yes	33 (1.7)	36 (2.0)	69 (1.9)
No	1758 (98.0)	1900 (98.3)	3658 (98.1)
Having hypertension			
Yes	171 (8.8)	116 (6.5)	287 (7.7)
No	1762 (91.2)	1678 (93.5)	3440 (92.3)
Having diabetes			
Yes	98 (5.1)	51 (2.8)	149 (4.0)
No	1835 (97.2)	1743 (94.9)	3578 (96.0)

Having respiratory diseases			
Yes	65 (3.4)	45 (2.5)	110 (3.0)
No	1868 (96.6)	1749 (97.5)	3617 (97.0)
Having cardiovascular diseases			
Yes	76 (3.9)	42 (2.3)	118 (3.2)
No	1857 (96.1)	1752 (97.7)	3609 (96.8)
Having other diseases			
Yes	177 (9.1)	196 (10.8)	373(9.9)
No	1760 (91.1)	1600 (89.2)	3360 (90.2)
Living condition			
Urban	1862 (96.4)	1717 (95.7)	3579 (96.1)
Rural	71 (3.6)	77 (4.3)	148 (3.9)

Table 2: Risk perception based on the EPPM by demographic characteristic (N = 3727)

	Efficacy		Defensive response			Threats	
	Self-efficacy	Response efficacy	Denial	Reactance	Avoidance	Susceptibility	Severity
	Mean (SD)	Mean (SD)	Mean (SD)	Mean (SD)	Mean (SD)	Mean (SD)	Mean (SD)
Age (years)							
15-29	65.6 (17.2)	84.0 (11.3)	23.2 (18.9)	19.8 (20.2)	31.1 (27.2)	65.5 (19.6)	76.4 (14.5)
30-44	65.6 (17.2)	83.5 (11.1)	22.0 (16.0)	19.2 (17.9)	26.6 (22.3)	69.1 (17.7)	77.1 (13.4)
45-59	68.3 (16.5)	84.2 (10.9)	21.3 (16.2)	18.9 (17.2)	24.3 (20.5)	67.7 (16.8)	77.0 (13.3)
60+	69.4 (16.9)	83.7 (11.0)	23.0 (17.5)	22.0 (17.6)	24.1 (16.9)	68.8 (16.3)	76.0 (14.0)
<i>P-value</i>	<.0001	0.473	0.134	0.309	<.0001	<.0001	0.539
Gender							
Female	67.6 (17.4)	85.0 (10.7)	22.0 (16.5)	17.5 (17.0)	26.0 (23.0)	66.1 (19.1)	76.9 (13.8)
Male	66.4 (16.8)	82.7 (11.3)	22.3 (17.3)	21.1 (19.4)	28.0 (23.4)	69.3 (17.2)	76.9 (14.1)
<i>P-value</i>	0.055	0.087	0.366	0.001	0.015	<.0001	0.56
Educational status							
Primary	68.1 (18.3)	81.7 (13.3)	29.2 (21.7)	25.8 (22.6)	31.9 (27.4)	59.2 (21.0)	74.3 (15.5)
Secondary	68.7 (16.5)	83.5 (10.9)	24.8 (18.0)	20.5 (19.0)	28.0 (24.1)	64.5 (19.1)	76.0 (13.8)
Higher	66.3 (17.2)	84.0 (10.9)	20.7 (15.9)	18.4 (17.6)	27.2 (23.2)	69.6 (17.3)	77.4 (13.9)
<i>P-value</i>	0.001	0.011	<.0001	<.0001	0.002	<.0001	0.001
Marital status							
Single	68.3 (16.9)	83.8 (10.9)	23.3 (18.4)	20.0 (20.1)	30.7 (25.9)	65.2 (20.2)	76.4 (14.3)
Married	66.6 (17.2)	83.9 (11.0)	21.7 (16.4)	19.1 (17.7)	25.9 (22.2)	68.6 (17.4)	77.1 (13.8)
Widowed/Divorced	63.3 (13.5)	78.5 (13.8)	26.9 (22.6)	21.5 (20.4)	32.7 (24.4)	62.7 (20.0)	74.4 (14.8)
<i>P-value</i>	0.024	0.021	0.017	0.355	<.0001	<.0001	0.224
Economic status							
Poor	63.1	81.6	23.9	20.8	27.8	67.5 (18.6)	77.9

	(18.0)	(13.0)	(16.9)	(18.7)	(24.4)		(14.1)
Intermediate	66.4 (16.5)	83.6 (10.3)	22.0 (16.4)	19.6 (18.2)	27.1 (22.7)	67.9 (17.7)	76.5 (13.6)
Good	71.1 (17.0)	85.9 (10.7)	21.2 (18.1)	17.7 (17.7)	26.6 (23.2)	67.6 (19.0)	77.2 (14.7)
<i>P-value</i>	<.0001	<.0001	0.007	0.003	0.603	0.841	0.050
Living condition							
Urban	67.0 (17.1)	83.7 (11.1)	22.1 (16.8)	19.5 (18.3)	27.3 (23.2)	67.7 (18.2)	77.0 (14.0)
Rural	67.8 (18.5)	84.7 (10.1)	22.1 (19.5)	16.5 (18.3)	23.1 (22.3)	68.1 (18.2)	74.8 (13.2)
<i>P-value</i>	.051	.55	1.00	.051	0.033	0.817	0.066

Table 3: Risk perception based on the EPPM by disease status (N = 3727)

	Efficacy		Defensive response			Threats	
	Self-efficacy	Response efficacy	Denial	Reactance	Avoidance	Susceptibility	Severity
	Mean (SD)	Mean (SD)	Mean (SD)	Mean (SD)	Mean (SD)	Mean (SD)	Mean (SD)
History of coronavirus							
Yes	57.7 (21.9)	74.9 (23.5)	32.8 (23.7)	30.1 (26.8)	41.5 (28.7)	70.0 (22.5)	67.4 (21.3)
No	67.1 (17.7)	83.9 (10.8)	22.0 (16.8)	19.3 (18.2)	27.0 (23.1)	67.7 (18.2)	77.9 (13.9)
<i>P-value</i>	0.036	<0.0001	<0.0001	0.009	<0.0001	0.442	0.006
History of coronavirus in family members							
Yes	64.5 (20.9)	81.5 (18.9)	24.5 (19.7)	17.7 (20.5)	30.0 (27.8)	68.6 (22.1)	72.9 (17.4)
No	67.0 (17.0)	83.8 (10.8)	22.1 (16.9)	19.4 (18.3)	27.1 (23.1)	67.7 (18.1)	77.0 (13.9)
<i>P-value</i>	0.018	<0.0001	0.25	0.32	0.45	0.031	0.142
Frequency of chronic diseases							
0	67.5 (16.9)	84.1 (10.8)	21.8 (16.9)	19.3 (18.3)	27.1 (23.5)	69.3 (18.6)	76.5 (13.9)
1	65.2 (17.7)	83.1 (11.5)	23.0 (17.0)	19.3 (18.1)	27.3 (22.4)	67.6 (16.7)	78.4 (14.0)
2	65.7 (16.4)	82.8 (11.6)	26.1 (19.0)	22.7 (20.4)	27.9 (21.0)	69.0 (15.7)	77.2 (12.8)
3 and more	60.0 (21.3)	77.6 (18.3)	20.8 (14.6)	17.2 (16.2)	20.0 (22.1)	66.6 (22.2)	74.8 (19.7)
<i>P-value</i>	0.001	0.002	0.026	0.224	0.390	0.021	0.016

Table 4: Danger control and fear control by demographic characteristics of the study sample (N= 3727)

	Danger control	Fear control	P-value
	No. (%)	No. (%)	
Age			<0.0001
15-29	531 (25.3)	334 (20.5)	
30-44	1061 (50.5)	934 (57.4)	
45-59	434 (20.7)	300 (18.5)	
60+	75 (3.6)	58 (3.6)	
Gender			<0.0001
Female	1079 (51.4)	715 (44.0)	
Male	1022 (48.6)	911 (56.0)	
Educational status			<0.0001
Primary	142 (6.8)	60 (3.7)	
Secondary	595 (28.3)	328 (20.2)	
Higher	1238 (76.1)	1364 (64.9)	
Marital status			0.133
Single	519 (24.7)	362 (22.3)	
Married	1566 (74.5)	1246 (76.6)	
Widowed/Divorced	16 (0.8)	18 (1.1)	
Economic status			<.0001
Good	582 (27.7)	343 (21.1)	
Not good, not bad	1194 (56.8)	939 (57.7)	
Poor	325 (15.5)	344 (21.2)	
History of coronavirus			0.132
Yes	17 (0.8)	20 (1.2)	
No	2084 (99.2)	1606 (98.8)	
History of coronavirus in a family member			0.141
Yes	34 (49.6)	35 (2.2)	
No	2067 (98.4)	1591 (97.8)	
Disease status			<0.0001
0	1677 (79.8)	1202 (73.9)	

1	346 (16.5)	357 (22.0)
2	66 (3.1)	49 (3.0)
3 and more	12 (0.6)	18 (1.1)
Living condition		0.555
Urban	2014 (95.9)	1565 (96.2)
Rural	87 (4.1)	61 (3.8)
All	2101 (56.4)	1626 (43.6)